



# Bluetongue virus seroprevalence and risk factor analysis in cattle and water buffalo in southern Italy (Campania region)

Gianmarco Ferrara<sup>1</sup> · Elvira Improda<sup>1</sup> · Federica Piscopo<sup>1</sup> · Riccardo Esposito<sup>1</sup> · Giuseppe Iovane<sup>1</sup> · Ugo Pagnini<sup>1</sup> · Serena Montagnaro<sup>1</sup>

Received: 27 July 2023 / Accepted: 31 August 2023 / Published online: 8 September 2023  
© The Author(s) 2023

## Abstract

Bluetongue is an arthropod-borne viral infection that is notifiable in several countries and causes significant economic losses and major concerns for ruminant trade. In this study, we investigated bluetongue 1 seroprevalence in the Campania region, southern Italy, in cattle and buffalo populations, and assessed which factors were correlated with a high risk of exposure. The infection was widespread, as evidenced by the high individual (43.6%) and herd prevalence (85.4%). The highest prevalence was found in adult animals. Among the climatic factors analyzed, average temperature played a prominent role, being capable of affecting the probability of being positive for this infection. Surprisingly, exposure to Schmallenberg virus did not predispose animals to be positive for bluetongue virus, even though these infections share the same vector (*Culicoides*). Our data, consistent with those in the literature, suggest the transversal spread of bluetongue virus in the Mediterranean area, and indicate a limited co-exposure rate between Bluetongue and Schmallenberg viruses.

**Keywords** Bluetongue · Co-exposure · Water Buffalo · Ruminants · Italy · Orbivirus · Arbovirus · Bunyavirus · Schmallenberg virus

## Introduction

Arboviruses pose a threat to both human and animal health, causing infections that have spread rapidly in recent decades, reaching areas where they were previously considered tropical (Socha et al. 2022). The name of this virus group, which has no taxonomic meaning, derives from the common transmission of its members (arthropod-borne), and consequently, the spread of these viruses depends on the distribution of vectors (Koenraad et al. 2014). This group includes viruses responsible for concerning zoonoses (such as dengue, yellow fever, or zika), as well as several viruses of veterinary importance (such as Schmallenberg and bluetongue) (Socha et al. 2022). Bluetongue virus (BTV) is a non-contagious RNA virus

spread by the bite of *Culicoides*, that has significantly invaded Europe in recent decades (Mignotte et al. 2021).

BTV was responsible for well-known outbreaks in the late 1990s and early 2000s and is now considered an endemic and re-emerging virus throughout Europe. Because of the severe symptoms exhibited by infected animals, BTV infection causes economic losses and impacts the international cattle trade (estimated to be up to US\$ 3 billion) (Alkhamis et al. 2020). Although BTV mostly results in clinical symptoms in sheep, cattle act as reservoir hosts and play an important role in BT transmission and epidemiology because they exhibit prolonged viremia (rarely associated with reproductive failure and low milk production) (Alkhamis et al. 2020).

Surveillance for this virus is based on the identification of the pathogen in vectors and symptomatic hosts, as well as the detection of specific antibodies. Bluetongue is endemic among ruminant populations in Italy, and repeated outbreaks are described, especially in small ruminants, while information on circulation in other species is more lacking (Carvelli et al. 2019). The aim of this study was to investigate the BTV seroprevalence in the Campania region, southern Italy.

✉ Gianmarco Ferrara  
gianmarco.ferrara@unina.it

<sup>1</sup> Department of Veterinary Medicine and Animal Productions, University of Naples Federico II, Via Federico Delpino N.1, 80137 Naples, Italy

## Materials and methods

### Sampling and study area

The current study was conducted in Campania (410000000 N-143000000 E), a southern Italian region overlooking the Mediterranean Sea. Due to the promiscuity of ruminants and meteorological conditions, this area favors the circulation of arboviruses. Samples were selected from those collected in a previous study (approved by the Institutional Ethics Committee of the Department of Veterinary Medicine and Animal Production, Centro Servizi Veterinari, Federico II University of Naples, PG/2022/0093419, 20 July 2022) (Ferrara et al. 2023a, b). 218 buffaloes and 250 cattle were sampled. Animals came from 31 separate districts and four provinces (21 cattle herds and 20 water buffalo herds), none of which practiced BTV immunization. Samples were collected at the end of the 2022 vector season (October–November 2022). Data on potential risk variables (species, age, and housing) were collected. Environmental data (annual average temperature, altitude, and distance from the coast) were calculated using the farm's geographical coordinates and data obtained by the Italian National Meteorological Service from meteorological stations located near the sampling farms (<http://centrofunzionale.regione.campania.it/#/pages/dashboard>).

### Serological analysis

Sera were separated from blood samples by centrifugation at 1000 g for 10 min and kept at -20 °C until testing. A commercial assay (ID Screen® Bluetongue Competition, ID Vet) was used to detect BTV antibodies in each sample. This competitive assay detects antibodies against all BTV serotypes (using the VP7 core protein as antigen) and is selective (does not cross-react with antibodies against the closely related epizootic hemorrhagic disease virus). The assay was performed following the manufacturer's instructions and optical density values at 450 nm were determined using a spectrophotometer. Results were calculated for each serum as a ratio to the mean of negative controls (Sample/Negative control ratios, or S/N) and compared to cut-off values (S/N < 40% are defined as positives).

### Statistical analysis

Prevalence and information on sampled animals were employed for risk factor analysis. The chi-square test was used to carry out a bivariate analysis of the putative risk factors for BTV seropositivity at the animal level. The serological result (positive or negative) was regarded as dependent variable, whereas the information gathered through the

epidemiological questionnaires was considered as independent variables. Species, age, province, type of housing, altitude, mean annual temperature, total annual rainfall, and distance from the coast were the independent variables. The correlation between dependent and independent variables was investigated using Chi-square statistics. A p-value of less than 0.05 was considered significant, and a threshold of 0.2 was utilized to select the variables to be included in the multivariate logistic regression (using the forward elimination strategy). Odds ratios (OR) and 95% confidence intervals were calculated to analyze the degree of correlation between independent factors and BTV seropositivity. Fit models were assessed using the Akaike Information Criterion (AIC), and those that best fit the data were chosen. The Variance inflation factor (VIF) was used to test for collinearity. For statistical analysis, MedCalc Statistical Software version 16.4.3 (MedCalc Software, Ostend, Belgium; [www.medcalc.org](http://www.medcalc.org)) and JMP version 14.1.0 (SAS Institute Inc.) were used.

## Results

A competitive ELISA was used to screen 468 ruminants for the presence of BTV antibodies and 204 animals tested positive. A prevalence of 43.6% was therefore reported at the individual level, but it was significantly greater at the herd level (85.4%), given that only 6 out of 41 farms did not have any positive animals. No substantial differences were detected between species at the herd or individual level. In fact, seroprevalence varied between 45.2% and 90.5% in cattle and between 41.7% and 80% in buffalo at the individual and herd levels, respectively. The province of Benevento had fewer positive animals than any other province (with a prevalence of approximately 20% compared to a prevalence ranging from 40 to 52% in the other provinces). There were no relationships between high seropositivity and the type of housing evaluated (although grazing animals had greater percentages of positives). Furthermore, animals that were exposed to SBV were no more likely to be positive for BTV (only 21.8% of animals seropositive for SBV testing were also exposed to BTV). Age was significantly associated with a higher probability of infection among the individual risk variables (Table 1). Among the environmental parameters included (average annual temperature, altitude, yearly rainfall, and distance from the coast), the mean annual temperature plays the most important role and is the only factor associated with the greatest seroprevalences (Table 1). Multivariate logistic regression included all variables with a p-value of 0.2 or lower. All variables were included because they satisfied the multicollinearity requirement (VIF less than 2.5).

**Table 1** Univariate investigation (chi-square) assessed individual and environmental possible risk variables for bluetongue virus seropositivity (location, species, housing, outcome to SBV ELISA, annual mean temperature, altitude, yearly rainfall, distance from the coast)

Factor	<i>n</i>	Positive	%	95%CI	$\chi^2$	<i>p</i> -value
Total	468	204	43.6	39.1 – 48.08		
Species						
Cattle	250	113	45.2	39.03 – 51.37	0.56	0.45
Water buffalo	218	91	41.7	35.2 – 48.29		
Age						
Adult	389	179	46	41.06 – 50.97	5.51	0.019
Young	79	25	31.7	21–39 – 41.90		
Location						
Avellino	63	31	49.2	36.86 – 61.55	17.81	<0.001
Benevento	54	11	20.4	9.63 – 31.11		
Caserta	168	68	40.5	33.05 – 47.90		
Salerno	183	94	51.4	44.12 – 58.61		
Housing						
Fully stallfed	422	178	42.2	37.47 – 46.89	3.47	0.063
Partly grazed	46	26	56.5	42.2 – 70.85		
Coexposure with SBV						
Yes	236	102	43.2	36.9 – 49.54	0.03	0.871
No	232	102	44	37.58 – 50.35		
Mean annual t°						
≤ 16 °C	221	83	37.6	31.2 – 43.94	6.2	0.013
> 16 °C	247	121	49	42.75 – 55.22		
Altitude (a.s.l.)						
≤ 250 m	331	138	41.7	36.38 – 47.00	1.65	0.198
> 250 m	137	66	48.2	39.81 – 56.54		
Annual rainwater (cm)						
≤ 1000	124	63	50.8	42.01 – 59.61	3.57	0.059
> 1000	344	141	41	35.79 – 46.19		
Distance from the coast						
≤ 20 km	183	73	39.9	32.8 – 46.9	1.67	0.196
> 20 km	285	131	46	40.18 – 51.75		

Mean annual temperature was confirmed as the primary risk factor by multivariate analysis. Age was also identified as a risk factor for BTV positivity in multivariate analysis. Adult animals were 1.48 times more likely to

test positive for BTV antibodies than younger animals (Table 2). Furthermore, animals raised in regions with an average annual temperature above 16 °C were 1.64 times more likely to test positive (Table 2).

**Table 2** A logistic regression model was used to examine the relationship between putative risk variables (*p* < 0.2), and bluetongue virus seropositivity

Factor	Coefficient (β)	Standard error	OR	OR CI%	<i>p</i> -value
Mean annual temperature (> 16 °C)	0.492	0.19	1.64	1.13–2.4	0.01
Age (> 24 months)	0.644	0.265	1.48	1.12–1.69	0.015

## Discussion

The results of this study confirm the trend observed in tropical and temperate regions where BTV has become endemic. In fact, a high prevalence has been observed in countries where the host vector cycle is conserved (as in Italy), not only due to its wide distribution but also as a result of a long-lasting antibody response. When ruminants are infected with BTV, they produce antibodies against both structural (on which the assay used in this study is based) and nonstructural viral proteins (Zhugunissov et al. 2015). BTV-specific neutralizing antibodies are detectable 5–6 days after the infection and can persist for several months, providing long-term resistance to reinfection with the homologous BTV serotype (representing the basis for vaccination) (Ramakrishnan et al. 2006; Maclachlan et al. 2014). Since our sampling coincided with the end of the vector season, we ensured to identify the largest number of positive animals.

The prevalence we observed is in line with that found in other countries where BT is endemic. Great part of the studies were conducted on cattle, describing exposure rates ranging from 27 to 67% in African and Asian countries (Hwang et al. 2019; Bakhshesh et al. 2020). All these studies reported age as a significant risk factor since adult animals are not protected by colostral immunity and interact with new vector seasons during their lives.

However, there is limited information on the diffusion of BT in buffalo in Europe. In fact, most of the studies in this sense are conducted on other continents, finding prevalence very similar to that observed in other ruminants, such as in Asia (Sohail et al. 2019; Rupner et al. 2020). All these data support the central role of buffalo in the epidemiology of this infection, even if there is no data on the duration of the viraemia. The differences observed in these studies could be due not only to a different epidemiological situation but also to the approach used and the timing of sampling (environmental factors can influence the vector season and thus the spread of these infections).

Numerous descriptive studies in the literature provide data about co-infections and co-exposures with BTV and SBV or other arboviruses. In regions where both infections are endemic, the prevalence of SBV is usually higher than that of BTV and the co-exposure rate is rather low (this is described in both domestic and wild animals in several countries). SBV has been described as the predominant viral exposure in captive fallow deer (Wernike et al. 2022). This trend is also confirmed in the roe deer in Flanders (27.9% for SBV and 0% for BTV), in Poland in the wild European bison (76.1% for SBV and 24.7% for BTV), and in Ireland in the

red deer (1.5% for SBV and 0% for BTV) (Tavernier et al. 2015; Graham et al. 2017; Krzysiak et al. 2017).

Co-infection with BTV-SBV, as well as with other arboviruses, has been documented in domestic ruminants in Israel (Behar et al. 2022). In contrast, the only European study that has described the degree of co-exposure to SBV and BTV in domestic animals is a 6-year study conducted in Poland, in which, among other factors, exposure to BTV in cattle was a risk factor for SBV positivity (Kęsik-Maliszewska et al. 2021). The lack of association between BTV and SBV exposure and the low rate of co-exposure are highly interesting, especially considering that these viruses share the same vector. Even more surprising, there is no evidence in the literature at the molecular level, and co-infection has not been described in either *Culicoides* or animals. In the absence of scientific confirmation, plausible explanations for these results involving the host or the vector can be proposed. For example, we can consider the degree of competition between the two viruses to infect the vector.

Regarding environmental risk factors, as seen in previous studies carried out on other arboviruses, average temperature is the most impactful factor in determining the effective spread of this infection. It is not surprising that other environmental factors do not affect the prevalence of BTV, as *Culicoides* are now described in Italy at any altitude and in any ecosystem. The success of these infections must be sought in the stratification and in the dispersal of the vector, which can be transmitted over hundreds of kilometers (Mignotte et al. 2021; Karthikeyan et al. 2022).

Certainly, the fact that these arbovirus infections are so common, with even modest rates of co-exposure, should be a reason for significant worry, considering that many nations have extremely strict rules regarding the purchase and sale of genetic material from other countries. Embryos, sperm, and oocytes from seropositive animals cannot be distributed in several European countries. We should thus consider the true damage that these diseases bring to the overall cattle sector rather than focusing on the minor harm they cause to symptomatic animals. Surveillance of this disease is critical for understanding the level of spread in a territory, taking control measures, and therefore accomplishing prevention. In Italy, and in particular in the Campania region, evidence of arbovirus disease in different populations has been described (Montagnaro et al. 2019; Petruccelli et al. 2020) and probably more will be described in the future. Cattle and buffaloes play a fundamental role in the maintenance of some arboviruses, and at the same time, they are important sentinels to prevent any outbreaks. Future investigations are necessary in order to understand the main serotypes circulating in the area as well as the characteristics of the infection in buffalo.

## Conclusions

Bluetongue is widespread among buffalo and cattle herds in the Campania region. Age and mean annual temperature were the most significant risk variables in both univariate and multivariate analyses. BTV prevalence is not affected by SBV seropositivity. Cattle serve as key sentinels to anticipate outbreaks in sheep and goats, in addition to being the major reservoir of infection due to extensive viremia. Further studies are needed to evaluate the duration of viraemia in buffalo and to establish the precise epidemiological role of this species.

**Author contribution** All authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by Gianmarco Ferrara and Elvira Improda. Federica Piscopo and Riccardo Esposito have coordinated the sampling procedures. Serena Montagnaro performed the statistical analysis. Giuseppe Iovane and Ugo Pagnini supervised all the experiments. The first draft of the manuscript was written by Gianmarco Ferrara and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

**Funding** Open access funding provided by Università degli Studi di Napoli Federico II within the CRUI-CARE Agreement.

**Data availability** Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

## Declarations

**Ethics approval** This study was performed in line with the principles of the Declaration of Helsinki. Approval was granted by the Institutional Ethics Committee of Department of Veterinary Medicine and Animal production (Centro Servizi Veterinari), University of Naples, Federico II (PG/2022/0093419 20 July 2022). Animal handling was performed following the guidelines of the guiding principles for biomedical research involving animals. Owners were informed of the objectives of the study through their informed consent. Blood samples from cattle and water buffaloes were collected by veterinarians according to the good practices of veterinary medicine.

**Competing interests** The authors have no relevant financial or non-financial interests to disclose.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- Alkhamis MA, Aguilar-Vega C, Fountain-Jones NM, et al (2020) Global emergence and evolutionary dynamics of bluetongue virus. *Sci Rep* 10. <https://doi.org/10.1038/s41598-020-78673-9>
- Bakhshesh M, Otarod V, Fallah Mehrabadi MH (2020) Large-scale seroprevalence and risk factors associated with Bluetongue virus in Iran. *Prev Vet Med* 179. <https://doi.org/10.1016/j.prevetmed.2020.104994>
- Behar A, Friedgut O, Rotenberg D, et al (2022) Insights on Transmission, Spread, and Possible Endemization of Selected Arboviruses in Israel—Interim Results from Five-Year Surveillance. *Vet Sci* 9. <https://doi.org/10.3390/vetsci9020065>
- Carvelli A, Sala M, Autorino GL, et al (2019) A cross-sectional serosurvey in a sheep population in central Italy following a bluetongue epidemic. *PLoS One* 14. <https://doi.org/10.1371/journal.pone.0208074>
- Ferrara G, Iovane V, Improda E et al (2023a) Seroprevalence and risk factors for bovine coronavirus infection among dairy cattle and water buffalo in Campania Region, Southern Italy. *Animals* 13. <https://doi.org/10.3390/ani13050772>
- Ferrara G, Wernike K, Iovane G, Pagnini U, Montagnaro S (2023b) First evidence of schmallenberg virus infection in southern Italy. *BMC Vet Res* 19:91–95. <https://doi.org/10.1186/s12917-023-03666-5>
- Graham DA, Gallagher C, Carden RF, et al (2017) A survey of free-ranging deer in Ireland for serological evidence of exposure to bovine viral diarrhoea virus, bovine herpes virus-1, bluetongue virus and Schmallenberg virus. *Ir Vet J* 70. <https://doi.org/10.1186/s13620-017-0091-z>
- Hwang JM, Kim JG, Yeh JY (2019) Serological evidence of bluetongue virus infection and serotype distribution in dairy cattle in South Korea. *BMC Vet Res* 15. <https://doi.org/10.1186/s12917-019-2000-z>
- Karthikeyan R, Rupner RN, Koti SR et al (2022) Analysis of bluetongue disease epizootics in sheep of Andhra Pradesh, India using spatial and temporal autocorrelation. *Vet Res Commun* 46:967–978. <https://doi.org/10.1007/s11259-022-09902-x>
- Keşik-Maliszewska J, Collins ÁB, Rola J et al (2021) Schmallenberg virus in Poland endemic or re-emerging? A six-year serosurvey. *Transbound Emerg Dis* 68:2188–2198. <https://doi.org/10.1111/tbed.13870>
- Koenraadt CJM, Balenghien T, Carpenter S, et al (2014) Bluetongue, Schmallenberg - what is next? Culicoides-borne viral diseases in the 21st century. *BMC Vet Res* 10. <https://doi.org/10.1186/1746-6148-10-77>
- Krzyśiak MK, Iwaniak W, Keşik-Maliszewska J et al (2017) Serological Study of Exposure to Selected Arthropod-Borne Pathogens in European Bison (*Bison bonasus*) in Poland. *Transbound Emerg Dis* 64:1411–1423. <https://doi.org/10.1111/tbed.12524>
- Maclachlan NJ, Henderson C, Schwartz-Cornil I, Zientara S (2014) The immune response of ruminant livestock to bluetongue virus: From type I interferon to antibody. *Virus Res* 182:71–77. <https://doi.org/10.1016/j.virusres.2013.09.040>
- Mignotte A, Garros C, Dellicour S, et al (2021) High dispersal capacity of *Culicoides obsoletus* (Diptera: Ceratopogonidae), vector of bluetongue and Schmallenberg viruses, revealed by landscape genetic analyses. *Parasit Vectors* 14. <https://doi.org/10.1186/s13071-020-04522-3>
- Montagnaro S, Piantedosi D, Ciarcia R et al (2019) Serological Evidence of Mosquito-Borne Flaviviruses Circulation in Hunting Dogs in Campania Region, Italy. *Vector-Borne Zoonotic Dis* 19:142–147. <https://doi.org/10.1089/vbz.2018.2337>

- Petrucelli A, Zottola T, Ferrara G, et al (2020) West nile virus and related flavivirus in european wild boar (*Sus scrofa*), latium region, Italy: A retrospective study. *Animals* 10. <https://doi.org/10.3390/ani10030494>
- Ramakrishnan MA, Pandey AB, Singh KP et al (2006) Immune responses and protective efficacy of binary ethylenimine (BEI)-inactivated bluetongue virus vaccines in sheep. *Vet Res Commun* 30:873–880. <https://doi.org/10.1007/s11259-006-3313-5>
- Rupner RN, VinodhKumar OR, Karthikeyan R et al (2020) Bluetongue in India: a systematic review and meta-analysis with emphasis on diagnosis and seroprevalence. *Vet Quart* 40:229–242. <https://doi.org/10.1080/01652176.2020.1810356>
- Socha W, Kwasnik M, Larska M, et al (2022) Vector-Borne Viral Diseases as a Current Threat for Human and Animal Health—One Health Perspective. *J Clin Med* 11. <https://doi.org/10.3390/jcm11113026>
- Sohail T, Yaqub T, Abbas T et al (2019) Seroprevalence of Bluetongue virus in small and large ruminants in Punjab province, Pakistan. *Acta Trop* 189:22–29. <https://doi.org/10.1016/j.actatropica.2018.09.020>
- Tavernier P, Sys SU, De Clercq K, et al (2015) Serologic screening for 13 infectious agents in roe deer (*Capreolus capreolus*) in Flanders. *Infect Ecol Epidemiol* 5. <https://doi.org/10.3402/IEE.V5.29862>
- Wernike K, Fischer L, Holsteg M et al (2022) Serological screening in wild ruminants in Germany, 2021/2022: No evidence of SARS-CoV-2, bluetongue virus or pestivirus spread but high seroprevalences against Schmallenberg virus. *Transbound Emerg Dis* 69:e3289–e3296. <https://doi.org/10.1111/tbed.14600>
- Zhugunissov K, Yershebulov Z, Barakbayev K et al (2015) Duration of protective immunity after a single vaccination with a live attenuated bivalent bluetongue vaccine. *Vet Res Commun* 39:203–210. <https://doi.org/10.1007/s11259-015-9643-4>

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.