

Erratum to: Macronutrient (N, P, K) and Redoximorphic Metal (Fe, Mn) Allocation in *Leersia oryzoides* (Rice Cutgrass) Grown Under Different Flood Regimes

Samuel C. Pierce · Matt T. Moore · Dan Larsen ·
S. R. Pezeshki

Published online: 1 September 2012
© Springer Science+Business Media B.V. 2012

Erratum to: Water Air Soil Pollut DOI 10.1007/s11270-009-0120-y

The original publication of this article (doi:10.1007/s11270-009-0120-y) unfortunately contained errors.

Table 2. Values for K, Fe, and Mn are all an order of magnitude too high. The values have been decreased by a factor of ten, and rounded to under four significant figures.

Figure 2. Aboveground Fe concentrations are actually two orders of magnitude higher than shown in the original figure. For the sake of simplicity, all units have been changed to mg/L and the height of the bars reduced in the aboveground treatment to reflect this change. The lowercase letters elucidating treatment

differences in shoot tissue were not included in the updated graph, as there was no treatment effect.

Figure 3. Mn concentrations are actually two orders of magnitude higher than shown. The units have been changed to mg/L and the scale on the Y-axis has been decreased by an order of magnitude.

The following text should be removed: Discussion, page 81, column 2, paragraph 2, sentence 3 (running from lines 22–25 of right column). “Root Fe concentrations in Chen et al., (2005) were, however, more than an order of magnitude higher than those in the present study.”

The corrected Table 2, Fig. 2, and Fig. 3 are reproduced in the next page.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11270-009-0120-y>.

S. C. Pierce (✉) · S. R. Pezeshki
Department of Biology, The University of Memphis,
Memphis, TN 38152, USA
e-mail: spierce1@memphis.edu

S. C. Pierce
e-mail: spierce@cfr.msstate.edu



M. T. Moore
USDA-Agricultural Research Service,
National Sedimentation Laboratory,
P.O. Box 1157, Oxford, MS 38655, USA

D. Larsen
Department of Earth Sciences, The University of Memphis,
Memphis, TN 38152, USA

Present Address:
S. C. Pierce
Department of Wildlife, Fisheries, and Aquaculture,
Mississippi State University,
Mississippi State, MS 39762, USA

Table 2 Mean biomass and total nutrient uptake across treatments

	Control	Intermittently flooded	Partially flooded	Continuously flooded
Biomass (g)				
Belowground	23.3±6.6	22.2±6.4	23±9.2	19.8±5.6
Aboveground	38.3±12	38.8±9.1	37.3±8.9	45.7±12
N (mg)				
Belowground	124.0±68	120.4±66	109.1±54	98.6±46
Aboveground	277.7±88	225±99	264±100	303±134
P (mg)				
Belowground	35.3±15	49.4±29	57.0±29	53.0±16
Aboveground	82.1±20	67.7±24	70.2±24	94.8±38
K (mg)				
Belowground	143±39	138±61	103±30	99±45
Aboveground	356±129	347±108	263±84	349±91
Fe (mg)				
Belowground	47.3±20	68.9±43	113±74	130±45 ^a
Aboveground	5.95±2.8	8.29±4.1	8.11±8.1	8.21±3.7
Mn (mg)				
Belowground	2.67±0.8	4.53±2.0	5.88±3.8	5.71±2.4
Aboveground	3.16±1.4	8.07±2.7	10.7±4.2 ^a	13.5±5.8 ^a

**Fig. 2** Fe tissue concentrations**Fig. 3** Mn tissue concentrations