

Anthropogenic Platinum Enrichment in the Vicinity of Mines in the Bushveld Igneous Complex, South Africa

Sebastien Rauch · Olalekan S. Fatoki

Received: 9 April 2012 / Accepted: 12 November 2012 / Published online: 30 November 2012
© Springer Science+Business Media Dordrecht 2012

Abstract South Africa is the world's single largest platinum (Pt) producer owing to important resources in the Bushveld Igneous Complex (BIC). Little attention has been paid to the environmental impacts of mining in the region despite the extent of mining activities. Here, we present a study on the occurrence of Pt in the vicinity of mines in the BIC. Elevated concentrations were found at all sampling sites in the mining area. The highest Pt concentration in soil ($698 \pm 178 \text{ ng g}^{-1}$, $< 125 \mu\text{m}$) was found at a smelter, while processing plants, shafts and nearby towns had lower concentrations. Elevated Pt concentrations were also found in grass with a maximum of $256 \pm 122 \text{ ng g}^{-1}$ at the smelter and a significant correlation between soil and grass concentrations. In comparison, Pt concentrations in soil and grass at a background site were 2.7 ± 0.4 and $0.6 \pm 0.2 \text{ ng g}^{-1}$, respectively. Laboratory uptake experiments indicate that atmospheric deposition is the main source of Pt in grass. The occurrence of elevated Pt concentrations and its accumulation in vegetation raise concern over exposure of the local

population through inhalation, skin exposure or dietary intake.

Keywords Platinum · Mining · Smelter · South Africa · Soil · Grass · Deposition

1 Introduction

The availability of abundant natural resources is a major component of the South African economy (Vermaak 1997). Platinum (Pt) is one of South Africa's most valuable resources owing to the general scarcity of this element in the Earth's crust (Peucker-Ehrenbrink and Jahn 2001) and its increasing use (Johnson-Matthey 2003). In 2010, South Africa produced 143 tons Pt, representing 76 % of the global Pt supply (Johnson-Matthey 2010).

South Africa's Pt resources are located in the Bushveld Igneous Complex (BIC), North of Pretoria (Fig. 1). The BIC, which was formed by the repeated injection of lava (or magma) into a sub-volcanic, shallow-level chamber, is the world's largest mafic-ultramafic intrusion with an area of 65,000 km² and 75 % of the world's Pt resources (Cawthorn 1999). Pt is mainly mined from two ore layers, i.e. the Merensky Reef and the Upper Group 2 (UG2) chromitite layer. Until recently, mining was concentrated on the Merensky Reef in the western limb of the BIC because of higher grade, lower chromite content and easier access to mining sites. However, increasing demand and decreasing reserves

S. Rauch (✉)
Department of Civil and Environmental Engineering,
Chalmers University of Technology,
412 96 Göteborg, Sweden
e-mail: Sebastien.rauch@chalmers.se

O. S. Fatoki
Faculty of Applied Sciences, Cape Peninsula University
Technology,
Cape Town, South Africa

have resulted in increasing mining of the deeper UG2 layer and of the eastern limb of the BIC. In 2006, mining of UG2 was expected to represent as much as 60 % of the total ore processed in the BIC, and 18 % of the production was expected to take place in the eastern limb (Johnson-Matthey 2003).

Most of the South African Pt production is exported for use in jewellery, automobile exhaust catalysts and industrial applications (Johnson-Matthey 2010). Studies on the environmental relevance of Pt have focused on urban environments where elevated Pt concentrations have been measured as a result of automobile catalyst emissions (Ravindra et al. 2004; Rauch and Morrison 2008). Although metal production activities are recognised to be major sources of metals into the environment (Nriagu and Pacyna 1988; Dudka and Adriano 1997), potential Pt emissions from mining and ore processing have only drawn little interest until now. Studies in Russia have shown that smelters which process Pt-containing ores are an important source of Pt into the regional environment (Boyd et al. 1997; Gregurek et al. 1998; Gregurek et al. 1999; Niskavaara et al. 2004) and may contribute to the cycle of Pt in the Northern Hemisphere (Rauch et al. 2005; Moldovan et al. 2007). The most significant risk from exposure to platinum is sensitization of the airways (Lindell 1997). Exposure to platinum can also cause sensitization of the skin. Mutagenic effects have been demonstrated *in vitro*, but genotoxic risks in humans remain uncertain (Lindell 1997).

Mining and ore processing activities are major sources of sulphur dioxide and airborne particles in the BIC (Steyn 2005). Air quality monitoring has recently been put in place in an effort to control emissions (Steyn 2005), but potential Pt emission from mining and ore processing in the BIC has not been investigated. In this study, we investigate the hypothesis that mining activities are causing increased environmental Pt concentrations in the BIC, and that local populations are exposed to this metal as a result of mining emissions.

2 Experimental

2.1 Soil and Grass Sampling

Soil and grass samples were collected in the vicinity of Pt mines in the BIC. Details of the sampling and location of

selected mines are provided in Table 1 and Fig. 1, respectively. Soil collection was performed using a gardening-type trowel, and stainless steel scissors were used for grass collection. Samples were collected as close as possible to the mining complexes and at least 5 m from roads in order to limit the potential impact of automobile traffic emissions (Schäfer et al. 1999). All samples were kept in ziplock PE bags until return to the laboratory.

In a first step, a screening was performed by the collection of soil samples at six Pt mines (labelled M1–M6) in August 2006. To provide information relevant to the determination of the source of Pt, samples were obtained from the surface soil (0–2 cm) and from a deeper layer (4–6 cm). Further sampling was then performed at mine M4 (where the highest concentration was found during the screening) in August 2007 and December 2008. Surface soil (0–2 cm) was collected near an underground shaft, a processing plant and a smelter (two samples), as well as in two towns (three samples for town A and two samples for town B) located within 5 km of the smelter. In addition, grass (*Aristida bipartita*, *Bothriochloa insculpta*, *Digitaria eriantha*, *Ischaemum afrum* or *Panicum maximum*, depending on availability at the sampling site) was collected in August 2007.

2.2 Uptake Experiment

Common oat (*Avena sativa*) was grown on selected, unsieved soil samples to assess potential Pt bioavailability. We note that all samples used in this study are true grasses (*Poaceae*), and the uptake experiment supports the study of Pt bioavailability by plants, despite possible difference in uptake rate for different species. Soil samples used in the uptake experiment were collected in 2008 at mine M4 (near the smelter and the underground shaft) and in nearby towns. Approx. 10 ml Milli-Q (18 M Ω cm, Millipore, USA) water was added daily. Obtained plants were harvested after 4 weeks and prepared as described below, along with respective soil samples.

2.3 Samples Preparation and Analysis

The soil samples were dried overnight at 105 °C in acid washed beakers and were homogenised using a mortar and pestle. The samples were then sieved, and the <125- μ m fraction was retained for analysis. Selected soil samples were sieved into five fractions




Fig. 1 Schematic map of the Bushveld Igneous Complex with selected Pt mines

Table 1 Pt concentration in soil and grass at selected mines and in towns in the vicinity of mine M4

Sampling location	Main feature in vicinity of sampling	Sampling year	Number of sampling sites	Pt concentration (ng g^{-1} , number of replicates in brackets)		
				Topsoil (0–2 cm)	Deeper soil (4–6 cm)	Grass
Background		2007	1	2.7±0.4 (4)	–	0.6±0.2 (3)
Eastern Limb						
M1	Processing plant	2006	1	41±9 (3)	25±4 (3)	–
M2	Underground shaft	2006	1	12±1 (3)	6.6±1.8 (3)	–
M3	Processing plant	2006	1	147±36 (3)	62±7 (3)	–
Western Limb						
M4	Smelter	2006	1	653±40 (3)	107±26 (3)	–
	Smelter	2007	2	698±178 (7)	–	256±122 (7)
	Smelter	2008	1	625±45 (3)	–	–
	Underground shaft	2007	1	91±10 (3)	–	17±7 (3)
	Processing plant	2007	1	110±30 (3)	–	25±7 (4)
	Processing plant	2008	1	123±69 (3)	–	–
	Town A	2007	3	85±33 (8)	–	65±43 (5)
	Town A	2008	1	60±78 (3)	–	–
M5	Processing plant	2006	1	104±17 (3)	7.8±0.8 (3)	–
		2007	1	144±91 (7)	–	43±20 (4)
M6	Processing plant	2007	2	73±18 (3)	–	–
		2008	1	21±3 (3)	3.2±1.5 (3)	–

Concentrations are presented as average ± standard deviation

(<125, 125–250, 250–500, 500–1,000 and >1,000 μm), and all fractions <1,000 μm were analysed. Grass samples were thoroughly washed in Milli-Q water before drying to remove particles deposited onto the grass surface, dried overnight at 105 °C and cut into small pieces using stainless steel scissors.

The samples were prepared by closed-vessel microwave digestion (Mars5, CEM Corporation, Matthews, USA) using Aqua regia (3:1; HCl/HNO₃; v/v; HCl, Suprapure grade, Merck, Darmstadt, Germany) as extractant. Approximately 0.5 g of soil or 0.3 g of grass was weighed into HP500 Teflon vessels, and 8 ml of Aqua regia was added to the samples. The temperature programme consisted of an 8-min ramp to 165 °C followed by a 2-min holding period and a second 5-min ramp to 175 °C with a 5-min holding period with a maximum allowed pressure of 200 psi. The obtained solutions were transferred to Erlenmeyer flasks and slowly taken to dryness on a hotplate. The residues were dissolved in 10 ml 2 % HCl. Samples were kept at 8 °C and diluted prior to analysis.

Platinum, Cr and Cu concentrations were determined by inductively coupled plasma mass spectrometry (ICP-MS, Elan6000, PE Sciex, Canada) using standard operating conditions (Rauch et al. 2001). Concentrations were determined by external calibration using Pt solutions prepared from single-element standards (High Purity Standards, Promochem AB, Ulricehamn, Sweden) and multi-element standards for Cr and Cu (CertiPUR XI, Merck, Darmstadt, Germany). Interference from HfO on Pt was checked and corrected when necessary using a Hf standard (Parent et al. 1997; Rauch et al. 2001). Accuracy of the analytical procedure was checked for Pt using a road dust reference material (BCR-723, IRMM, Geel, Belgium) for which the measured Pt concentration was 75 ng g⁻¹ for a certified concentration of 81.3 ± 2.5 ng g⁻¹. All results are presented as average ± standard deviation.

3 Results and Discussion

Average Pt concentrations for all analysed samples, including soil and grass collected at the background site, mining sites and nearby towns, are presented in Table 1.

3.1 Pt Concentrations in Soil at Mining Sites

Platinum concentrations in topsoil collected in the vicinity of selected mines, in 2006, ranged from 9.9 ± 0.7 ng g⁻¹ near an underground shaft at M2 to 653 ± 40 ng g⁻¹ near a smelter at M4 (Fig. 2). Significant contribution of the smelter to local Pt concentrations was confirmed by further sampling at mine M4 in 2007 and 2008. Concentration at the smelter was 698 ± 178 ng g⁻¹ in 2007 and 625 ± 45 ng g⁻¹ in 2008, whereas lower concentrations were found at a shaft and a processing plant at mine M4 (Fig. 3). Platinum concentration at the smelter was approximately 250 times larger than at a background site where a concentration of 2.7 ± 0.4 ng g⁻¹ was found, 1,400 times larger than average upper continental crust concentrations (Peucker-Ehrenbrink and Jahn 2001) and approximately five times lower than in Merensky Reef or UG2 ores (Cawthorn 1999). In comparison, typical Pt concentrations in urban and roadside soil are in the 25–250 ng g⁻¹ range (Rauch and Morrison 2008).

Similarly, high concentrations have been reported in the context of geochemical exploration. Platinum concentrations were found to exceed 180 ng g⁻¹ in soil in part of the BIC including in the Rustenburg area, and a causal relationship between Pt concentrations and mining operations has been suggested (Wilhelm et al. 1997). In addition, Pt concentrations measured in the BIC are in the same order as concentrations near plants processing Pt-containing ores in Russia. The average Pt concentration in the topsoil from a catchment located 5 km South of the Monchegorsk plants




Fig. 2 Platinum concentrations in soil (<125 μm) collected at selected mines in 2006, including topsoil (0–2 cm) and the deeper soil layer (4–6 cm). Background concentration obtained from samples collected in 2007 is provided for comparison. Error bars represent 1 SD

was 49.6 ng g^{-1} with a maximum concentration of 466 ng g^{-1} (Boyd et al. 1997).

3.2 Platinum Concentrations in Soil in Nearby Towns

Soil samples collected in two towns located in the vicinity of mine M4 had elevated Pt concentrations relative to the background site. Platinum concentrations in town A (1 km N of the smelter) were $85 \pm 33 \text{ ng g}^{-1}$ in 2007 and $60 \pm 78 \text{ ng g}^{-1}$ in 2008. Concentrations in town B (3 km SW of the smelter) were $144 \pm 91 \text{ ng g}^{-1}$ in 2007 and $73 \pm 18 \text{ ng g}^{-1}$ in 2008. These concentrations are approximately 20- to 50-fold higher than the measured background concentration and in the same order as at the processing plant and the underground shaft at mine M4 (Fig. 3). There was no significant difference (*t* test at 90 % confidence level) between soil concentrations in the two towns, despite town B being further from the smelter, and no correlation between Pt concentrations and distance to the smelter ($R^2=0.3$).

3.3 Source of Pt in Soil and Atmospheric Dispersion

Although elevated concentrations are often indicative of an anthropogenic source, elevated Pt concentrations in the BIC might also be the result of natural occurrence. To assess the source of Pt at selected mines, soil samples were collected at two depths (0–2 and 4–6 cm) in 2006. Average platinum concentrations were higher in the top soil layer (0–2 cm) than those in the deeper layer (4–6 cm) at all mines (Fig. 2). Although the soil layers are not dated, the higher concentrations

in the top soil layers indicate a recent increase in Pt concentrations that may be linked to Pt mining and the reported increase in Pt production in the BIC (Johnson-Matthey 2003).

Platinum concentrations in soil were found to vary with mining site (Fig. 2), possibly due to differences in Pt production, mining activity and/or location of the sampling site relative to the mine. The smelter at M4, where the highest concentrations were found at two separate sampling sites, is the mine with the largest production volume. However, there is no correlation between soil concentrations at the different mines and Pt production, indicating that production volume only has a limited influence on soil concentrations. This is supported by variations at mine M4 (Fig. 3) where the highest concentrations were found near the smelter. Russian smelters processing Pt-containing ores have been reported to be an important regional source of this metal (Gregurek et al. 1998, 1999; Niskavaara et al. 2004), and the high concentrations observed at the smelter (Figs. 2 and 3) indicate that Pt smelters are an important source of Pt in the BIC. Other sources, including processing plants and shafts, also contribute to Pt emissions.

The particle size distribution of Pt at the smelter and processing plant reveals differences in the source of Pt. The highest concentration at the smelter was found in the finest fraction ($<125 \mu\text{m}$), whereas Pt was more evenly distributed between the different fractions at the processing plant (Fig. 4). Ore processing essentially consists of milling and separation by flotation and may, therefore, only result in the emission coarse particles. In




Fig. 3 Platinum concentrations in soil ($<125 \mu\text{m}$) and grass at mine M4 and in nearby towns for samples collected in 2007 and 2008. Error bars represent 1 SD




Fig. 4 Particle size distribution (125, 125–250, 250–500 and 500–1,000 μm) of Pt in selected soil samples. Error bars represent 1 SD

contrast, smelting involves high-temperature processes that may result in the formation of finer particles. The highest concentration in soil from nearby towns was measured in the smallest particle size fraction (Fig. 4), indicating that the smelter contributes to elevated Pt levels in the towns, possibly due to the emission of Pt in fine particles.

Elevated concentrations in the nearby towns indicate an efficient atmospheric transport. Elevated Pt

concentrations in road dust have been reported as far as Pretoria and attributed to atmospheric dispersion from Pt mines in the BIC (Rauch and Fatoki 2010). Pretoria is located >50 km from Pt mines in the BIC and >100 km East of mine M4. These results support a widespread contamination, and we suggest that the emission of fine Pt particles from the smelters in the BIC is resulting in the dispersion of Pt in the region.

3.4 Platinum Concentration in Grass

Elevated Pt concentrations were found in grass collected at mine M4 and in nearby towns in 2007 (Fig. 3). The highest Pt concentration was measured near the smelter ($256 \pm 122 \text{ ng g}^{-1}$). Elevated Pt concentrations were also found in towns A and B (64 ± 43 and $43 \pm 20 \text{ ng g}^{-1}$, respectively). In comparison, a Pt concentration of $0.6 \pm 0.2 \text{ ng g}^{-1}$ was measured at the background site. A significant correlation was found between Pt concentrations in soil and in grass, with grass concentrations approximately three times smaller than soil concentrations (Fig. 5a).

The occurrence of Pt in grass represents both atmospheric deposition and uptake from bioavailable Pt in soil. Studies on the occurrence of Pt in the roadside environment show that Pt emitted from automobile catalyst is bioavailable (Zimmermann and Sures 2004; Ek et al. 2004). Platinum bioavailability was assessed in laboratory uptake experiments. Grass grown on soil collected near Pt mines contained elevated Pt concentrations with a significant correlation between soil and grass




Fig. 5 Correlation between Pt (a), Cr (b) and Cu (c) concentrations in soil and grass mine at M4, nearby towns and the background site. Error bars represent 1 SD




Fig. 6 Platinum uptake by plants grown on selected unsieved soils. Presented soil concentrations correspond to the <1,000- μm particle size fraction

Table 2 Copper and chromium concentrations in topsoil and grass at mine M4 and nearby towns in 2007 samples

Sampling location	Number of sampling sites	Cu concentration ($\mu\text{g g}^{-1}$)		Cr concentration ($\mu\text{g g}^{-1}$)	
		Topsoil	Grass	Topsoil	Grass
Background	1	22±1	3±1	43±3	7±6
Smelter	2	200±62	71±24	756±202	29±13
Underground shaft	1	40±7	10±1	217±40	17±6
Processing plant	1	136±45	12±1	340±87	22±12
Town A	3	55±19	26±12	271±105	13±4
Town B	2	65±14	19±7	232±76	21±11

Concentrations are presented as average \pm standard deviation

concentrations (Fig. 6), but at concentrations much lower than in grass collected at the mining sites. These results indicate that atmospheric deposition is the main source of Pt in grass, while uptake of bioavailable Pt is likely to be a smaller contributor to Pt levels in grass at the mining sites. The occurrence of Pt in grass indicates that elevated Pt concentrations are also present in crops from local agricultural fields [approximately 400,000 km² of cultivated land in the Rustenburg area (Steyn 2005)] and gardens, raising concern over potential exposure through dietary intake.

3.5 Comparison with Copper and Chromium

Chromium and copper were analysed in 2007 samples (mine M4) because these elements might be associated with platinum emissions. Platinum is mined from chromium-rich minerals (Cawthorn 1999), and elevated Cu concentrations have been reported in soil from a tailing disposal facility at a South African Pt mine (Maboeta et al. 2006). Measured Cr and Cu concentrations are provided in Table 2. Copper and Cr concentrations were lowest at the background site and highest at the smelter. Positive correlations were found between Pt and Cu concentrations and between Pt and Cr concentrations in both soil and grass, indicating that mining is a common source for these elements. Chromium concentrations exceed the expected values for uncontaminated soils (9.9–121 $\mu\text{g g}^{-1}$, Ali et al. 2004) at all sites, except the background site. In contrast, Cu concentrations do not exceed the expected values for uncontaminated soils (2–250 $\mu\text{g g}^{-1}$, Ali et al. 2004) at any of the sites, indicating that Cu contamination by mining activities is more limited.

Positive correlations were also found between soil and grass concentrations for both Cr and Cu (Fig. 5). The outlier for Cu corresponds to the processing plant and may be due to differences in soil properties or Cu speciation at this site. Similar slopes for Cu and Pt indicate that the accumulation rates for these two elements are similar, although accumulation mechanisms might be different. Cu is an essential element, and therefore, similar accumulation rates indicate that Pt accumulation rates are relatively high.

4 Conclusion

Elevated Pt concentrations were measured in soil and grass in the vicinity of Pt mines in the BIC and attributed to mining activities. Elevated Pt concentrations were also found in two towns located within a few kilometres of a smelter where the highest Pt concentrations were measured. These results raise concern over exposure of the local population through inhalation, skin contact or dietary intake. Several towns and cities, including Rustenburg (400,000 inhabitants), and local food production activities are located in the Pt mining region. As Pt emissions are expected to increase as a result of the current expansion of Pt mining in the BIC, further work is needed to quantify Pt emissions from South African mines, understand the behaviour of these elements in the environment, characterise human exposure and determine its impact on the local population.

Acknowledgments This research was funded by the Swedish International Development Agency (SIDA) under the Swedish

Research Links programme and the South African National Research Foundation (NRF).

References

- Ali, N. A., Ater, M., Sunahara, G. I., & Robidoux, P. Y. (2004). Phytotoxicity and bioaccumulation of copper and chromium using barley (*Hordeum vulgare* L.) in spiked artificial and natural forest soils. *Ecotoxicology and Environmental Safety*, *57*, 363–374.
- Boyd, R., Niskavaara, H., Kontas, E., Chekushin, V., Pavlov, V., Often, M., & Reimann, C. (1997). Anthropogenic noble-metal enrichment of topsoil in the Monchegorsk area, Kola peninsula, northwest Russia. *Geochemical Exploration*, *58*, 283–289.
- Cawthorn, R. G. (1999). The platinum and palladium resources of the Bushveld Complex. *South African Journal of Science*, *95*, 481–489.
- Dudka, S., & Adriano, D. C. (1997). Environmental impacts of metal ore mining and processing: a review. *Journal of Environmental Quality*, *26*, 590–602.
- Ek, K. H., Morrison, G. M., & Rauch, S. (2004). Environmental routes for platinum group elements to biological materials—a review. *The Science of the Total Environment*, *334*, 21–38.
- Gregurek, D., Melcher, F., Niskavaara, H., Pavlov, V. A., Reimann, C., & Stumpfl, E. F. (1999). Platinum-group elements (Rh, Pt, Pd) and Au distribution in snow samples from the Kola Peninsula, NW Russia. *Atmospheric Environment*, *33*, 3281–3290.
- Gregurek, D., Reimann, C., & Stumpfl, E. F. (1998). Trace elements and precious metals in snow samples from the immediate vicinity of nickel processing plants, Kola Peninsula, northwest Russia. *Environmental Pollution*, *102*, 221–232.
- Johnson-Matthey (2003) The expansion of platinum mining in South Africa. In: *Platinum 2003*. Johnson-Matthey, London, UK. pp 14–17.
- Johnson-Matthey. (2010). *Platinum 2010*. London: Johnson-Matthey.
- Lindell, B. (1997). *DECOS and NEG basis for an occupational standard. Platinum*. Solna: Arbetslivsinstitutet. ISBN 91–7045–420–5.
- Maboeta, M. S., Claassens, S., Van Rensburg, L., & Van Rensburg, P. J. (2006). The effects of platinum mining on the environment from a soil microbial perspective. *Water, Air, and Soil Pollution*, *175*, 149–161.
- Moldovan, M., Veschambre, S., Amouroux, D., Benech, B., & Donard, O. F. X. (2007). Platinum, palladium, and rhodium in fresh snow from the Aspe Valley (Pyrenees Mountains, France). *Environmental Science and Technology*, *41*, 66–73.
- Niskavaara, H., Kontas, E., & Reimann, C. (2004). Regional distribution and sources of Au, Pd and Pt in moss and O-, B- and C-horizon podzol samples in the European Arctic. *Geochemistry-Exploration Environment Analysis*, *4*, 143–159.
- Nriagu, J. O., & Pacyna, J. M. (1988). Quantitative assessment of worldwide contamination of air, water and soils by trace metals. *Nature*, *333*, 134–139.
- Parent, M., Vanhoe, H., Moens, L., & Dams, R. (1997). Investigation of HfO₂ interference in the determination of platinum in a catalytic converter (cordierite) by inductively coupled plasma mass spectrometry. *Talanta*, *44*, 221–230.
- Peucker-Ehrenbrink, B., & Jahn, B. M. (2001). Rhenium-osmium isotope systematics and platinum group element concentrations: loess and the upper continental crust. *Geochemistry, Geophysics, Geosystems*, *2*(10), 1061. doi:10.1029/2001GC000172.
- Rauch, S., Hemond, H. F., Barbante, C., Owari, M., Morrison, G. M., Peucker-Ehrenbrink, B., & Wass, U. (2005). Importance of automobile exhaust catalyst emissions for the deposition of platinum, palladium, and rhodium in the Northern Hemisphere. *Environmental Science and Technology*, *39*, 8156–8162.
- Rauch, S., & Fatoki, O. S. (2010). Platinum and lead in South African road dust. In S. Rauch, G. M. Morrison, & A. Monzon (Eds.), *Highway and urban environment* (pp. 161–166). Heidelberg: Springer.
- Rauch, S., Lu, M., & Morrison, G. M. (2001). Heterogeneity of platinum group metals in airborne particles. *Environmental Science and Technology*, *35*, 595–599.
- Rauch, S., & Morrison, G. M. (2008). The environmental relevance of platinum group elements. *Elements*, *4*, 259–263.
- Ravindra, K., Bencs, L., & Van Grieken, R. (2004). Platinum group elements in the environment and their health risk. *The Science of the Total Environment*, *318*, 1–43.
- Schäfer, J., Eckhardt, J. D., Berner, Z. A., & Stüben, D. (1999). Time-dependent increase of traffic-emitted platinum-group elements (PGE) in different environmental compartments. *Environmental Science and Technology*, *33*, 3166–3170.
- Steyn, S. (2005). *The C region of North West Province, South Africa. Doctoral thesis*. South Africa: University of Pretoria.
- Vermaak, C. F. (1997). A brief overview of South Africa's mineral industry: world context and changing local circumstances. *Mineralium Deposita*, *32*, 312–322.
- Wilhelm, H. J., Zhang, H., Chen, F. L., Elsenbroek, J. H., Lombard, M., & deBruin, D. (1997). Geochemical exploration for platinum-group elements in the Bushveld Complex, South Africa. *Mineralium Deposita*, *32*, 349–361.
- Zimmermann, S., & Sures, B. (2004). Significance of platinum group metals emitted from automobile exhaust gas converters for the biosphere. *Environmental Science and Pollution Research*, *11*, 194–199.