

Relationships between vegetation zonation and environmental factors in newly formed tidal marshes of the Yangtze River estuary

Yanlong He · Xiuzhen Li · Christopher Craft ·
Zhigang Ma · Yongguang Sun

Received: 1 February 2011 / Accepted: 17 May 2011 / Published online: 1 June 2011
© Springer Science+Business Media B.V. 2011

Abstract The Yangtze River delta is characterized by rapidly accreting sediments that form tidal flats that are quickly colonized by emergent vegetation including *Scirpus mariqueter* and the invasive species *Spartina alterniflora*. We measured soil surface elevation, water table depth, soil salinity, water content and compaction in the tidal flat, the *Scirpus* and *Spartina* zones and their borders to identify relationships between environmental factors and colonization by *Scirpus* and *Spartina*. With increasing elevation from tidal flat to *Spartina*, inundation frequency and duration, moisture and depth to water table decreased whereas soil salinity, temperature and compaction increased. High soil moisture and groundwater and low salinity were the characteristics of the tidal flat and its border with *Scirpus*. The *Spartina* zone and its border with *Scirpus* were characterized by greater salinity and elevation relative to the other zones. Our findings suggest that soil salinity controls patterns of plant zonation in the newly formed tidal salt marshes whereas elevation is of secondary importance. Our results suggest that patterns of vegetation zonation in tidal marshes of the Yangtze River delta are controlled

by environmental factors, especially (low) salinity that favors colonization by *Scirpus* in the lower elevations of the marsh.

Keywords Tidal marshes · Salinity · *Spartina alterniflora* · *Scirpus mariqueter* · Yangtze River estuary

Introduction

Since the 20th century, ecologists have studied the relationship between environment factors and the distribution of vegetation communities of salt marshes (Simas et al. 2001; Caçador et al. 2007). Salt marshes have attracted extensive attention from plant ecologists because of their low species diversity, simple community structure and striking zonation (Levine et al. 1998).

The distribution of salt marsh vegetation is related to energy and material inputs (i.e. tidal action, soil nutrients), and to the constraints of physical factors (such as flooding, anoxia, salinity, etc.). Tides play an important role in the development of estuarine salt marshes with zonal distribution of plant communities along the gradient of elevation from open water to uplands (Vince and Snow 1984). These zones are determined by differences in elevation that affect inundation and salinity. Most studies to date have focused on the role of abiotic factors in plant zonal

Y. He · X. Li (✉) · Z. Ma · Y. Sun
State Key Laboratory of Estuarine and Coastal Research,
East China Normal University, Shanghai 200062, China
e-mail: xzli@sklec.ecnu.edu.cn

C. Craft
School of Public and Environmental Affairs,
Indiana University, Bloomington, IN 47405, USA

pattern, especially salinity and flooding (Pennings et al. 2005; Silvestri et al. 2005).

The mechanisms controlling vegetation zonation in coastal salt marshes may be universal though the importance of individual factors may vary geographically (Pennings et al. 2005; Castillo et al. 2008). One region that is understudied is the tidal marshes of East Asia, including the Yangtze and Yellow River deltas. The eastern headlands of Chongming Island is an important salt marsh of the Yangtze River delta. It contains abundant resources of zoobenthos and vegetation and is a stopover for migrating waterfowls across the Asia–Pacific region (Huang et al. 1993).

In recent years, invasion of salt marsh communities by introduced species has become an important issue of tidal marsh research in the region and elsewhere (Daehler and Strong 1996; Zedler and Kereher 2004). *Spartina alterniflora*, a salt marsh species native to the east coast of the US, was introduced to China for sediment stabilization in 1979 and spread to Chongming Island by 1995. Because of its adaptability to high salinity and anoxia, *Spartina* began to colonize the tidal flats and began to displace native species, *Scirpus mariqueter* and *Phragmites australis*. It is considered a threat to benthos and birds which use *Scirpus* as their primary habitat (Yong and Zhang 1992). To date, no studies have focused on the spatial patterns of plant

communities, including the invasive species *Spartina*, and environmental factors in rapidly accreting tidal flats and newly colonizing tidal marshes of the Yangtze River delta.


We measured soil salinity, surface elevation, soil moisture, groundwater, soil compaction, soil temperature and inundation in the mudflat and the stands of *Scirpus*, *Spartina* and the transition zones to understand the role of environment factors in *Spartina* invasion in the Yangtze River delta. Our goal was to identify (1) the environmental factors that structure vegetation zonation in the Yangtze Estuary tidal marshes, and (2) the conditions that favor *Spartina* invasion into this habitat.

Materials and methods

Study site

The study site is located at the eastern end of Chongming Island (121°45'E, 31°30'N) in the Yangtze River estuary (Fig. 1). The wetland was established as a national nature reserve in 1992 and was designated as a Ramsar Site in 2001. It has a subtropical monsoon climate, with mean annual sunshine hours of 2138 h, frostless period of 229 days, and average annual

Fig. 1 Location of the study sites in the Yangtze River delta, eastern headlands of Chongming Island






Fig. 2 Diurnal tide curves, by month at the Sheshan tidal gauge station, Chongming Island

temperature of 15.3°C. July and August are the warmest months with monthly average temperature of 26.8°C. The coldest months are January and February with a mean temperature of 3.0°C. Mean annual precipitation ranges from 900 to 1,050 mm, most of which falls from April to September.

The study site has an irregular semidiurnal tide with mean tidal range of 2.43 and 3.08 m (Fig. 2). The soil texture is loamy with finer materials at the higher elevation of the marsh (Salt content of the soil ranges from 0.2 to 0.6%).

The native vegetation consists of *Scirpus mariqueter* and *Phragmites australis*. *Scirpus* is an herbaceous perennial species and reproduces both sexually and asexually though, like many salt marsh species, it spreads mostly by clonal propagation. Since 1995, when *Spartina* was first reported in marshes of the eastern headlands, *Spartina* gradually

colonized and spread into habitats of the native species. Today, a mosaic pattern of the invasive and native species exists in the marshes with sharp boundaries delineating the different communities.

Experimental design

We established three transects, each perpendicular to the levee and each separated by 200 m. Along each transect, three smaller subtransects were established to transit through the tidal flat, the *Scirpus* zone and the *Spartina* zone to sample four vegetated and one unvegetated habitats; tidal flat, tidal flat-*Scirpus* edge, *Scirpus* zone, *Spartina*-*Scirpus* edge zone and *Spartina* zone. At each location, three replicate measurements of soil compaction, salinity, temperature and water content, water table depth and elevation were made. Measurements were made four times between

May and October, 2008, during period of plant emergence (May), growth (July), flowering (September) and senescence (October). All measurements were made during neap tide phase.

Soil moisture, salinity and temperature were measured 10 cm below the surface using a Hydra soil moisture sensor (including Hydra Data Reader and Hydra Probe II Soil Moisture Sensor (SDI-12/RS485)) (Precision: Moisture, $\pm 0.5\%$ vol; Salinity, $\pm 2\%$; temperature, $\pm 0.6^\circ\text{C}$); Stevens Water Monitoring Systems Inc., Australia). Soil compaction was measured using a TJSD-750 soil compaction device (Precision: ± 0.05 kg; Hangzhou Tuopu Ltd. China). Surface elevation was measured using GPS-RTK (Z-Xtreme) (Precision: < 1 cm for static GPS, < 5 cm for kinematic GPS, cm-dm accuracies for baselines < 5 km for kinematic; AshtechTM, USA). The elevations were tied into the nearby Dongtan benchmark site ($121^\circ 57'$, $31^\circ 30'$), part of the Chinese geodetic level. We augered a hole to depth of 1.5 m to measure water table depth. Once the water table stabilized in the hole, we measured the water table depth from the soil surface.

We use the following formula to calculate the flooding depth of different vegetation zones:

$$I = r - h$$

where I represents flooding depth, r represents the changing range of the tide in this region each day (determined from the tide table of the eastern headlands in Chongming island, the Sheshan tide-gauge station ($122^\circ 14'$, $31^\circ 25'$), adjacent to our study site)

and h represents the mean elevation of each vegetation zone.

Statistical analysis

We used analysis of variance (ANOVA) to test for differences in surface elevation and soil properties among the vegetation zones (SPSS 1999). Means were compared using the least significant difference (LSD) test. Principal components analysis (PCA) was used to explore associations between the environmental properties and vegetation zones. We used PC-ORD 5.0 (McCune and Mefford 1999) for PCA using centered and standardized species-environmental variables. According to Jogman et al. (1995), the data were square root-transformed prior to analysis. All tests of significance were conducted at P of 0.05.

Results

Variation of flooding properties among different zones

The tidal cycle of the eastern headlands is characterized by irregular semidiurnal tides (Fig. 2). Combining the tide gauge data with our elevation data, we estimated monthly changes of the flooding duration and depth for the five zones (Fig. 3). One-way ANOVA analysis showed that the depth and duration of flooding in the tidal flat and the tidal flat-*Scirpus*




Fig. 3 Diurnal variation in inundation depth average every month, by vegetation zone, at the study site

edge were significantly ($P < 0.05$) greater than in the other zones and the daily maximum flooding depth was about 60 cm.

Mean monthly flooding depth in the *Scirpus* zone was greater than in the *Spartina* zone. Flooding duration in the *Scirpus* zone also was longer than in the *Spartina* zone. Both flooding depth and duration were less than the two vegetated zones relative to the un-vegetated mudflat (Fig. 4). The tidal flat was flooded every day and, on most days, it was flooded to a depth of 50 cm or more (Fig. 4). The depth of flooding was shallower in the *Scirpus* zone than that in the tidal flat, around 50 cm. And the depth of flooding was less than 40 cm in the *Spartina* zone.

Plots of flooding frequency versus inundation depth (Fig. 4) showed that the *Spartina* zone experienced more than 100 days with no flooding. In this zone, the maximum flooding depth was 160 cm. In the *Scirpus* zone, the number of days with no flooding was similar to the *Spartina* zone though the maximum depth of flooding was greater, about 180 cm (Fig. 4). The tidal flat was inundated longer and to a greater depth than the vegetated zones. In this zone, depth of flooding was 50 cm or more for more than

100 days and the maximum flooding depth was more than 200 cm. The *Spartina-Scirpus* edge had the lowest duration and depth of inundation of the zones sampled (Fig. 4).

Environmental factors and vegetation zones

Surface elevation and soil properties varied among the different vegetation zones ($P < 0.05$) as well. Soil surface elevation was greater in the vegetated zones than that in the tidal flat (Fig. 5a). Surface elevation averaged 2.74 m in the tidal flat, 2.95 m in the *Scirpus* zone and 3.15 m in the *Spartina* zone. Salinity exhibited a similar trend; it was the lowest in the tidal flat, moderate in the *Scirpus* zone and the greatest in the *Spartina* zone (Fig. 5b). Soil water content displayed the opposite trend as it was the greatest in the tidal flat and decreased landward (Fig. 5c). Water table depth, soil compaction and temperature followed the same pattern as elevation, increasing from the tidal flat to the vegetated zones (Fig. 5d, e, f). The *Spartina-Scirpus* edge had the highest elevation, and greatest depth to water table, soil compaction and temperature.





Fig. 4 Inundation depth versus frequency in the tidal flat (TF), tidal flat-*Scirpus* edge (ESM), *Scirpus maritimum* (SM), *Scirpus-Spartina* edge (ESS) and *Spartina alterniflora* (SA) zones. The density probability curve simulates the trend of

frequency (days) versus depth of inundation. The inundation depth and frequency in the *Spartina* zone are significantly lower than those in *Scirpus* and tidal flat zone ($P < 0.05$)

Fig. 5 **a** Mean surface elevation, **b** soil salinity, **c** soil water content, **d** water table depth, **e** soil compaction and **f** temperature (\pm standard error) in the tidal flat (TF), tidal flat-*Scirpus* edge (ESM), *Scirpus marigueter* (SM), *Scirpus-Spartina* edge (ESS) and *Spartina alterniflora* (Sa) zones. Means separated by the same letter are not significantly different ($P < 0.05$) according to the least significant difference (LSD) test



Principle components analysis

Principal components analysis revealed gradients associated with soil salinity, water content and water table depth on the first axis and soil compaction on the second axis (Fig. 6). The first two axes explain

82% of the variation in the data. Tidal flat sites were clustered in areas of high soil water content and water table depth, and low salinity (Fig. 6). *Spartina* sites were clustered in areas of high soil salinity and compaction. *Scirpus* sites were intermediate with respect to soil moisture and salinity (Fig. 6) whereas




Fig. 6 Principal component analysis of environmental factors (vectors) versus vegetation zones (points), tidal flat, tidal flat-*Scirpus* edge, *Scirpus mariquetter*, *Scirpus-Spartina* edge and *Spartina alterniflora*

Scirpus-Spartina zone sampling points were clustered in areas of high soil compaction.

Discussion

Expansion of *Spartina* and the corresponding decline of *Scirpus* coverage in tidal marshes of the eastern headlands in recent years may be attributed to several factors: (1) Reduced freshwater discharge by the Yangtze River since 2000 (Zhang R et al. 2006) and enhanced saltwater intrusion from the rising sea levels have led to an increase in surface soil salinity in the salt marshes of the region (Day et al. 1995) that made it difficult for *Scirpus* to colonize newly formed mudflats and (2) Seaward colonization of tidal flats by salt marsh vegetation caused by sediment deposition that increased surface elevation and salinity. Thus, the increase in salinity in recently colonized *Spartina* zones, in part, may have facilitated the invasion by *Spartina*. Furthermore, *Scirpus* reproduction and growth is more intimately tied to freshwater pulsing than *Spartina*. Li et al. (2010) reported that salinity in the Yangtze estuary is the lowest from May to October which is consistent with the life cycle of *Scirpus* that is completed during this

period. For example, emergence of *Scirpus* seedling occurs in May, followed by growth in summer, then by plant senescence in October. *Spartina*, in contrast, continues to grow throughout the year by producing clonal ramets (Livingstone and Patriquin 1981).

Laboratory studies also support *Spartina*'s greater tolerance to salinity. For example, *Scirpus* dies at surface water salinities of 1.6‰ or greater, whereas *Spartina* tolerates salinities up to 3.2‰ (Chen et al. 2005). Landin (1991) found that the optimum salinity for the growth of *Spartina* was 1–2‰.

Elevation is also important in structuring patterns of plant zonation in tidal marshes (Sánchez et al. 1996; Boorman et al. 2001; Bockelmann et al. 2002) as it is linked to surface inundation, salinity, soil aeration and other soil properties (Snow and Vince 1984; Adam 1990). In addition to soil salinity, water table depth, compaction and temperature all increased with elevation in our study (Fig. 5). Also, as surface elevation increased, flooding frequency decreased (Fig. 3). Although we observed differences in inundation patterns between *Scirpus* and *Spartina*, there was no significant difference in elevation between the two zones (Fig. 5). In addition to larger scale variation in environmental properties along our elevation gradient, there probably are smaller gradients of soil salinity, water content and ground water depth caused by differences in microtopography and other factors (Pennings et al. 2005), including plant species composition.

Principal components analysis also showed that vegetation zonation was strongly linked to soil moisture, salinity, compaction and water table depth (Fig. 6). Longer and deeper inundation and low salinity were characteristics of the tidal flat and tidal flat-*Scirpus* edge. In contrast, high soil salinity and compaction were characteristics of the *Spartina* and *Spartina-Scirpus* edge zone. In addition to salinity and soil moisture, we found that soil temperature differed among *Spartina* and *Scirpus* zones, with higher temperature in the *Spartina* zone. We hypothesize that higher temperature may be attributed to reduced duration and depth of inundation and soil moisture in the *Spartina* zone.

Differences in plant reproduction strategies also may play an important role in the distribution of tidal marsh vegetation (van Zandt et al. 2003). Clonal reproduction is the main mechanism of reproduction by *Spartina* in the Yangtze River delta with

colonization rates of up to 32 cm in one year (Zhang D et al. 2006). Zhi et al. (2007) also found that, on newly colonized sites, *Spartina* produces about 53,748 seeds per square meter, of which 70–80% germinate.

In contrast, *Scirpus* spreads via rhizomes at about one half the rates of *Spartina* (15 cm one year). Like *Spartina*, *Scirpus* also produces considerable numbers of seeds (15–18 per ramet with 2,580 ramets per square meter) that have similar germination rates of 73.8%. Thus, it appears that *Spartina*'s success may be due more to its rate of clonal spread than its seed production and germination.

In summary, environmental factors, especially salinity, structure plant communities of tidal marshes of the Yangtze River delta. Furthermore, increased salinity appears to favor the spread of *Spartina* into habitats historically occupied by *Scirpus*. Other environmental factors, soil moisture and compaction, as well as biological factors, especially rhizome growth also contributes to *Spartina*'s ability to colonize tidal flat habitat in the delta.

Acknowledgments We thank Wenyong Guo, Zaijun Xin and Yue Jia for their assistance with the collection of field data. Our research was supported by the 111 Project (B08022); Ministry of Education, China; Programme Strategic Scientific Alliances between China and the Netherlands (2008DFB90240); FP7.2009-1 (Contract 244104—THESEUS).

References

- Adam P (1990) Saltmarsh ecology. Cambridge University Press, Cambridge, p 461
- Bockelmann AC, Bakker JP, Neuhaus R, Lage J (2002) The relation between vegetation zonation, elevation and inundation frequency in a Wadden Sea salt marsh. *Aquat Bot* 73:211–221
- Boorman LA, Hazelden J, Boorman M (2001) The effect of rates of sedimentation and tidal submersion regimes on the growth of salt marsh plants. *Cont Shelf Res* 21: 2155–2165
- Caçador I, Tibério S, Cabral HN (2007) Species zonation in Corroios salt marsh in the Tagus estuary (Portugal) and its dynamics in the past fifty years. *Hydrobiologia* 587: 205–211
- Castillo JM, Mateos-Naranjo E, Nieva FJ, Figueroa E (2008) Plant zonation at salt marshes of the endangered cordgrass *Spartina maritima* invaded by *Spartina densiflora*. *Hydrobiologia* 614:363–371
- Chen ZY, Li B, Chen JK (2005) The effect of salinity and elevation to the growth of *Spartina alterniflora*. *J Yangtze Univ (Nat sci)* 2(2):6–9 (in Chinese with English abstract)
- Daehler CC, Strong DR (1996) Status, prediction and prevention of introduced cordgrass *Spartina* spp. invasions in Pacific estuaries, USA. *Biol Conserv* 78(1):51–58
- Day JW, Pont D, Hensel PF, Ibañez C (1995) Impacts of sea-level rise on deltas in the Gulf of Mexico and the Mediterranean: the importance of pulsing events to sustainability. *Estuaries* 18(4):636–647
- Huang ZY, Sun ZH, Yu K (1993) Shanghai birds' resources and their habitat. Fudan University Press, Shanghai, p 175 (in Chinese)
- Jogman RHG, ter Braak CJF, van Tongeren OFR (1995) Data analysis in community and landscape ecology. Pudoc, Wageningen, The Netherlands
- Landin M C (1991) Growth habits and other considerations of smooth cordgrass *Spartina alterniflora* Loisel [A]. In: Mumford TF Jr, Peyton P, Sayce JR et al. (eds) *Spartina* workshop record [C], Washington Sea Grant Program, University of Washington, Seattle, pp 115–201
- Levine JM, Brewer JS, Bertness MD (1998) Nutrients, competition and plant zonation in a New England salt marsh. *J Ecol* 86:285–292
- Li Q, An CG, Ma Q, Xu LL, Zhao YL (2010) Species composition and diversity of zooplankton in tidal creeks of the Chongming Dongtan intertidal flat. *Biodiversity Science* 18(1):67–75 (in Chinese with English abstract)
- Livingstone DC, Patriquin DG (1981) Belowground growth of *Spartina alterniflora* Loisel, habit, functional biomass and non-structural carbohydrates. *Estuar Coast Shelf Sci* 12(5):579–587
- McCune B, Mefford MJ (1999) PC-ORD. Multivariate analysis of ecological data, Version 4. MjM Software Design, Gleneden Beach, OR, USA, p 237
- Pennings SC, Grant MB, Bertness MB (2005) Plant zonation in low-latitude salt marshes: disentangling the roles of flooding, salinity and competition. *J Ecol* 93:159–167
- Sánchez JM, Izco J, Medrano M (1996) Relationships between vegetation zonation and altitude in a salt-marsh system in northwest Spain. *J Veg Sci* 7:695–702
- Silvestri S, Defina A, Marco Marani M (2005) Tidal regime, salinity and salt marsh plant zonation. *Estuar Coast Shelf Sci* 62:119–130
- Simas T, Nunes JP, Ferreira JG (2001) Effects of global climate change on coastal salt marshes. *Ecol Model* 139: 1–15
- Snow AA, Vince SW (1984) Plant zonation in an Alaskan salt marsh II: an experimental study of the role of edaphic conditions. *J Ecol* 72:669–684
- SPSS (1999) SPSS 10.0.1 for Windows. SPSS Inc, Chicago, IL
- van Zandt PA, Tobler MA, Mouton E, Hasenstein KH, Mopper S (2003) Positive and negative consequences of salinity stress for the growth and reproduction of the clonal plant, *Iris hexagona*. *J Ecol* 91:837–846
- Vince SW, Snow AA (1984) Plant zonation in an Alaskan salt marsh I: distribution, abundance and environmental factors. *J Ecol* 72:651–667
- Yong XK, Zhang LQ (1992) Studies on reproduction ecology of *Scirpus Mariqueter* population. *J East China Norm Univ (Nat Sci)* 4:94–99 (in Chinese with English abstract)
- Zedler JB, Kereher S (2004) Causes and consequences of invasive plants in wetlands: opportunities, opportunists, and outcomes. *Crit Rev Plant Sci* 23:431–452

- Zhang D, Yang MM, Li JX, Chen XY (2006) Vegetative dispersal ability of *Spartina alterniflora* in eastern end of Chongming Island. J East China Norm Univ (Nat Sci) 2:130–135 (in Chinese with English abstract)
- Zhang R, Wang YP, Pan SM (2006) Analyses with Wavelet and Hilbert-Huang transform on monthly water discharges at Datong Station, Yangtze River. J Nanjing Univ (Nat Sci) 42(4):423–434 (In Chinese with English abstract)
- Zhi YB, Li HL, An SQ (2007) Inter-specific competition: *Spartina alterniflora* is replacing *Spartina anglica* in coastal China. Estuar Coast Shelf Sci 74:437–448