

M. Barančková · J. Krojerová-Prokešová
I. V. Voloshina · A. I. Myslenkov · Y. Kawata
T. Oshida · J. Lamka · P. Koubek

The origin and genetic variability of the Czech sika deer population

Received: 8 June 2011 / Accepted: 6 March 2012 / Published online: 9 October 2012
© The Ecological Society of Japan 2012

Abstract Sika deer (*Cervus nippon*), native to Asia, formed two well-established free-living populations in the Czech Republic over the last century and continue to spread. Sika are also maintained in a large number of enclosures; these continue to introduce new individuals from the places of its origin as well as from other European countries. Despite extensive research into the morphology and ethology of the Czech sika deer, conducted over the last three decades, no study using genetic methods has been done. This study aimed to determine the genetic variability and the geographic origin of the Czech sika deer population. Two mitochondrial markers, the cytochrome *b* and the control region were analyzed in this study. Analysis of the two markers confirmed that the founder individuals of the Czech population originated from both native island (Japanese Islands) and native mainland (Far East Russia) popu-

lations. Results showed that the genetic variability of the Czech sika deer population is lower than the variability of the native Japanese population, but higher than that of the sampled part of the native Russian population. Also, the genetic variability was found to be higher within the samples from enclosures.

Keywords *Cervus nippon* · Cytochrome *b* gene · Control region · Introduced species

Electronic supplementary material The online version of this article (doi:10.1007/s11284-012-0992-y) contains supplementary material, which is available to authorized users.

M. Barančková (✉) · J. Krojerová-Prokešová · P. Koubek
Institute of Vertebrate Biology, Academy of Sciences of the Czech Republic, v.v.i., Květná 8, Brno 603 65, Czech Republic
E-mail: barancekova@ivb.cz
Tel.: +420-54-3422543
Fax: +420-54-3211346

I. V. Voloshina · A. I. Myslenkov
Lazovsky State Nature Reserve, Centralnaya St. 56, Lazo,
Primorsky Krai 692980, Russia

Y. Kawata · T. Oshida
Obihiro University of Agriculture and Veterinary Medicine,
Inada-cho, Obihiro, Hokkaido 080-8555, Japan

J. Lamka
Faculty of Pharmacy, Charles University Prague,
500 05 Hradec Králové, Czech Republic

P. Koubek
Department of Forest Protection and Game Management,
Faculty of Forestry and Wood Sciences, Czech University
of Life Sciences Prague, Kamýcká 1176,
165 21 Prague 6, Suchbát, Czech Republic

Introduction

The sika deer (*Cervus nippon* Temminck 1838) belongs to the subfamily Cervinae. It originated on the mainland of southeastern Asia and subsequently spread to Taiwan and the Japanese Archipelago (McCullough 2009a). Over the last two centuries, sika deer have been introduced into many parts of the world and they have become a stable element of the fauna in various European countries (Bartoš 2009; McCullough et al. 2009).

Sika originally spread from Japan through the southern Ussuri, Korea, Manchuria, and Eastern China to Vietnam. However, sika in their native range have greatly suffered due to habitat changes (Yuasa et al. 2007) and uncontrolled hunting, which led to the extinction of local wild populations in Vietnam, Taiwan, and South Korea (McCullough 2009a, b). Based on morphological characters, sika in the world were originally classified into 13 subspecies (Whitehead 1993). Six of these subspecies are found in Japan (Ohtaishi 1986): *Cervus nippon yesoensis* (Hokkaido Island), *C. n. centralis* (Honshu and Tsushima Island), *C. n. nippon* (Kyushu Island, Shikoku Island, and Goto Island), *C. n. mageshimae* (Mageshima and Tanegashima Island), *C. n. yakushimae* (Yakushima and Kuchinoerabu Island), and *C. n. keramae* (Ryukyu Island). Small, free-living populations of sika can still be found in mainland Asia and Taiwan (McCullough 2009b). The historical range of Russian sika (*C. n. hortulorum*) was primarily in Primorsky Krai extending to the borders of China (Aramilev 2009). Up until the early 19th

century, five morphological subspecies were abundant throughout mainland China (Sheng and Ohtaishi 1993). At present, wild sika are extinct over most of their original mainland range, with only three subspecies, *C. n. hortulorum*, *C. n. sichuanicus*, and *C. n. kopschi*, surviving in northeastern, southern, and western China (Guo and Zheng 2000). Wild sika are extinct in South Korea (Wo and Smith 1999), and there is only a small amount of information on their current distribution in North Korea (McCullough 2009b). In Vietnam, only captive populations survive, with a rather low, but not dangerously low, genetic diversity, suggesting a bottleneck (Thévenon et al. 2004). The last free-living Taiwan sika (*Cervus nippon taiouanus*) was killed in 1969, and its current population exists thanks to the release of animals bred in captivity (McCullough 2009a).

However, the classification based on morphology is not completely supported by genetic studies. Several studies reported the existence of two mitochondrial lineages in Japan the northern and southern Japanese groups (Nagata et al. 1995, 1999; Tamate and Tsuchiya 1995; Tamate et al. 1998), which does not reflect the morphological classification. The divergence of these lineages probably occurred before the colonization of Japan from China (Kawamura 2009). Furthermore, analyses showed that the southern Japanese lineage is more closely related to the mainland subspecies *C. n. hortulorum* than to the northern Japanese lineage (Nagata 2009). The three surviving Chinese subspecies appear to be valid based on DNA evidence; however, due to the lack of DNA samples, the status of the two extinct subspecies (*C. n. grassianus* and *C. n. mandarinus*) cannot be verified (Wilson 2000). Also, DNA studies confirmed the status of Vietnamese *C. n. pseudaxis* as a separate subspecies (Wilson 2000).

The introduction of sika to Europe began about 150 years ago. Altogether, sika were introduced to 35 different European countries at the end of the 19th century and the beginning of the 20th century (Bartoš 2009). Outside Asia, free-ranging populations are established in Australasia (New Zealand); North America (Kentucky, Maryland, North Carolina, Texas, Virginia); Europe (Austria, Czech Republic, Denmark, France, Germany Ireland, Poland, United Kingdom). Most of these introductions date from the last years of the 19th century (1890s) through to 1930s, although some have continued until more recently. The most authoritative reviews of distribution, history of introductions, and current status will be found in McCullough (2009b), Feldhamer and Demarais (2009), Banwell (2009), Bartoš (2009), and Swanson and Putman (2009) (see also Eick (1995a, b) for Continental Europe and Ratcliffe (1987) for United Kingdom).

In the Czech Republic, sika were first released into the Bor enclosure near Poděbrady in 1891 (Kokeš 1970). The geographical origin and subspecies status of most of the introduced individuals can no longer be ascertained as the archives of the main supplier, the Hagenbeck Company from Hamburg (Eick 1995a, b), were destroyed (Bartoš 2009). The animals reportedly originated from

Japan, eastern China, Korea, and Russia. Later, animals from England and Austria were also imported (Kokeš 1970). Up until World War II the sika were kept in enclosures; however, after the war some of these enclosures were destroyed or the animals were released on purpose, and the sika escaped (Vavruněk and Wolf 1977). These escaped sika were the basis of the West Bohemian and Moravian (Bouzovsko) populations (Doležal 1960; Babička et al. 1977). Since then, sika have colonized a large area of the Czech Republic (Vavruněk and Wolf 1977). Currently, there are two well-established populations inhabiting both the West Bohemia and the Bouzovsko area, and their populations are reported to be spreading—hunters repeatedly report the presence of sika males in previously sika-free localities. The recommendations for management of this exotic species are strictly set. However, due to very imprecise methods for the estimation of deer densities combined with the unwillingness of hunters to stop sika spreading, the situation in reality is greatly different. At present, the population size of sika in the Czech Republic, as provided by hunters for the year 2009, is 9,031 individuals (http://eagri.cz/public/web/file/67788/sumar_2009.pdf). However, real population numbers are much higher than the figures estimated and reported on a yearly basis.

Sika may have considerable ecological impacts due to their potential to cause significant damage to forestry, agriculture, and habitat structure (Chadwick et al. 1996; Abernethy 1998; Putman and Moore 1998; Putman 2000; Kelly 2002; Lowe 1994). There is also the additional concern of hybridization between introduced sika and native deer species. Sika subspecies are not only able to hybridize intra-specifically, they can interbreed with other deer species, such as red deer, hog deer, and axis (Bartoš et al. 1981). Hybridization occurs in regions of natural contact of sika with another deer species, for example the Ussuri River on the Russia–China border (Heptner et al. 1961), and sika have also been intentionally crossbred with other deer species for research purposes (e.g., Harrington 1973, 1982) or to improve the quality of sika trophies (Eick 1995b). This fact has been known for a long time and the first hybrids in Europe were recorded in 1884 in Ireland (Powerscourt 1884). Despite this, the possibility of interspecific hybridization has been ignored and the introduced animals have often been kept together with other deer species. Hybridization of sika has been recorded in Great Britain (Lowe and Gardiner 1975; Abernethy 1994; Goodman et al. 1999; Diaz et al. 2006; Pemberton et al. 2006; Pérez-Espona et al. 2009; Senn and Pemberton 2009; Swanson and Putman 2009), Germany (Herzog 1987; Gehle and Herzog 1998), Ireland (Harrington 1973; McDevitt et al. 2009), and in the Czech Republic (Bartoš and Žirovnický 1981; Bartoš and Vitek 1993) by morphological and karyological analyses.

One of the first applications of molecular phylogenetic approaches was to simultaneously determine both the phylogenetic and the spatial relationships between different mtDNA haplotypes or sequences. Avise (2000)

termed the joint use of phylogenetic techniques and geographic distributions phylogeography. The rapid mutation rate of the mitochondrial genome, the basis for phylogeographic research, makes it particularly suitable for studies of intraspecific variation (Avise 1994). Two regions of mitochondrial DNA, the cytochrome *b* gene and the control region, have also been extensively used in studies of various deer species (e.g., sika—Nagata et al. 1995; Tamate and Tsuchiya 1992, 1995; red deer—Polzahn and Strobeck 2002; Ludt et al. 2004; Eld's deer—Balakrishnan et al. 2003). The genetic variability and the phylogenetic and geographic relationships of both native (e.g., Nagata et al. 1995; Tamate and Tsuchiya 1995) and introduced (Cook et al. 1999; Pitra et al. 2005) sika populations have been extensively studied.

Even though over the last three decades several studies researched the ecology (Heroldová 1990), ethology (e.g., Bartoš and Žirovnický 1981, 1982; Bartoš et al. 1998), and morphology (Bartoš and Žirovnický 1981; Bartoš and Vitek 1993) of the Czech sika population, until now there have been no studies of the genetic variability and origin of this population. The aim of this study was to obtain the first genetic data about the Czech sika population by investigating (1) the genetic variability of the sika population using mitochondrial DNA markers for the cytochrome *b* gene and the control region and (2) the origins of Czech sika and their phylogeographic relationships with native Asian populations.

Materials and methods

Tissue samples (muscles, ear pieces, skin) for the analyses were collected from introduced Czech (236 samples), native Japanese (102 samples), and Russian (111 samples) populations, and stored in 96 % ethanol. In the Czech Republic, the samples were collected from free-living populations by hunters during the hunting seasons and by J. Lamka during veterinary examinations of the individuals kept in enclosures during the years 2007–2010 (Table 1). The Russian samples from Primorsky Krai were collected by hunters, mostly from cadavers found during winter in the years 2007–2010, while the Japanese samples primarily came from meat samples sent to Obihiro University of Agriculture and Veterinary Medicine in the years 2009–2010 (Table 1). All of the samples collected were tested for genetic polymorphisms in the cytochrome *b* (*cyt b*) gene and in the control region (CR). A further nine sequences for the *cyt b* gene and nine sequences for the CR, representing haplotypes of various sika subspecies from Asia, were downloaded from GenBank database (see the Supplementary material for the accession numbers).

The differences between introduced Czech and native Asian populations were analyzed as well as the differences between the free-living populations and populations from enclosures in the Czech Republic. The decision to analyze the differences between free and kept populations was based on the fact that the population of

Table 1 Tissue samples collected in introduced and native populations

Country	Locality	Sample size
Czech Republic	Bílá Lhota	1
	Bouzovsko	20
	Bydlo	1
	Častolovice	9
	Český les	2
	Doupov	47
	Horšovský Týn	23
	Hrotovice	3
	Lány	49
	Lidmovice	15
	Manětínsko	21
	Měřín	4
	Opočno	8
	Pastviny	2
	Podrážnice	10
	Rovensko	2
	Rouchovany	2
	Uherčice	4
	Úsobí	1
	Veselsko	1
	Vidlice	2
	Vysočina	5
	Private enclosures	4
Russia	Lazovsky District	97
	Terneysky District	14
Japan	Ehime	1
	Hokkaido	34
	Chiba	18
	Kochi	7
	Kyoto	8
	Nagasaki	10
	Osaka	12
	Yamaguchi	12

Table 2 The *cyt b* haplotype and nucleotide diversity of free-living sika deer and sika deer in enclosures in the Czech Republic

	<i>N</i>	<i>N_H</i>	<i>H_D</i>	<i>P_i</i> (JC)	<i>k</i>	<i>S</i>
Enclosures	144	6	0.666	0.02025	22.599	72
Free-living	95	4	0.461	0.01498	16.682	42

N number of samples, *N_H* number of haplotypes, *H_D* haplotype diversity, *P_i* (JC) nucleotide diversity Jukes–Cantor-corrected, *k* average number of nucleotide differences, *S* number of polymorphic sites

sika in enclosures continues to get “fresh genes”—all introductions are to enclosures and not to the wild—and that the population in enclosures includes also subspecies from mainland Asia, which is not present in the free-living population.

DNA extraction, PCR, and sequencing

The DNA was isolated using a Qiagen DNeasy Blood & Tissue Kit (Hilden, Germany) and a Genomed Jetquick Tissue DNA Spin Kit (Löhne, Germany), following the manufacturer's instructions. The PCR amplification of *cyt b* was carried out using the primers L14724 and H15915 (Irwin et al. 1991), and the amplification of the

CR using primers L15926 and H00651 (Kocher et al. 1989). The PCR protocol for *cyt b* was as follows: (1) 94 °C, 3 min; (2) 35 × (94 °C, 40 s; 50 °C, 40 s; 65 °C, 90 s); (3) 65 °C, 5 min, and for CR: (1) 94 °C, 3 min; (2) 35 × (94 °C, 30 s; 53 °C, 30 s; 65 °C, 1 min); (3) 65 °C, 5 min. The amplified DNA fragments were visualized by electrophoresis on a 2 % agarose gel stained with ethidium bromide or GoldView™. The DNA sequencing was performed on an ABI 3730 Genetic Analyzer (Applied Biosystems, Carlsbad, CA, USA). Complete sequences of the *cyt b* and the CR were edited and aligned using Sequencher 4.6 (Gene Codes Corporation, Ann Arbor, MI, USA).

Genetic diversity analysis and statistics

The Akaike information criterion (AIC; Akaike 1974), implemented in the MrModeltest 2.3 program (Nylander 2004), was used to select the best-fitting nucleotide substitution models for PAUP*4 (Swofford 2003) and MrBayes for both the *cyt b* and the CR datasets. Phylogenetic reconstructions were performed using maximum-likelihood criterion (ML; Felsenstein 1981) implemented in PHYML software (Guindon and Gascuel 2003). The robustness of the trees was assessed by bootstrap re-sampling (1,000 random replications for both NJ and ML analyses; Felsenstein 1985). Bayesian estimation (BE) of phylogeny (Yang and Rannala 1997) was performed in the program MrBayes 3.1 (Ronquist and Huelsenbeck 2003). Metropolis-coupled Markov chain Monte Carlo (MCMC) sampling was performed with five chains run for 1,000,000 iterations, using estimated model parameters as the starting values. Bayesian posterior probabilities were obtained from the 50 % majority rule consensus of trees sampled every 20 generations, discarding the trees obtained before the chains a reached stationary distribution. The *cyt b* gene sequence of *Odocoileus virginianus* and CR sequence of *Capreolus pygargus* were used as outgroups to root the phylogenetic trees (see the Supplementary material).

A median-joining network (MJ; Bandelt et al. 1999) was used as another way of visualizing relationships between the haplotypes. The MJ network was constructed using freely accessible Network software (<http://www.fluxus-engineering.com>).

The estimation of genetic diversity was conducted in DnaSP v5 (Librado and Rozas 2009). Genetic diversity indices consisted of haplotype diversity (H_D), nucleotide diversity (P_i ; Jukes-Cantor-corrected) and number of polymorphic sites (S).

Results

A total of 449 samples were collected from 33 localities (Table 1) and analyzed. The analyses yielded 449 complete *cyt b* sequences, 1140 bp, and 447 CR sequences, 975–1143 bp. All of the haplotype sequences were

deposited in GenBank (accession numbers for *cyt b*: JF893469–JF893492, for CR: JF893497–893538).

Sequence divergence of the *cyt b* region

A total of 24 distinct cytochrome *b* haplotypes was identified among the 449 analyzed samples. The haplotype diversity of the whole data set was $H_D = 0.861$. The most frequent haplotype J67D was found in 22.94 % of the samples, and three other haplotypes (J1N, J4N, J35D) were found in more than 10 % of the samples each. While none of the Japanese haplotypes were found in the Czech sika population, both Russian haplotypes were also shared by the Czech sika. The number of haplotypes found on the Czech localities varied from one to four (Fig. 1). The most frequent Czech haplotype was J1N, which was found in 30.38 % of the samples; haplotypes J35D and J4N were present in more than 20 % of the samples. The number and distribution of haplotypes at localities sampled in Japan was similar to that in the Czech Republic and varied from one to five haplotypes (Fig. 2a). Only three of these haplotypes reached more than 10 % representation in the data set, with the most frequent being J132NJ (33.33 %). The haplotype variability in the native population in Russia was very low with only two *cyt b* haplotypes found at both sampled localities (Fig. 2b). More than 80 % of the Russian samples belonged to haplotype J67D.

Out of the 1140 bp in the *cyt b* sequences, up to 78 were variable. The base composition in the sequences was 31 % of A, 28 % of C, 13 % of G and 28 % of T. The highest number of haplotypes and the highest haplotype diversity were found in Japanese samples, whereas the Russian population was the least diversified (see the Supplementary material).

Using Akaike information criterion, the best-fitting model chosen for maximum likelihood (ML) and Bayesian estimation (BE) was a general time reversible model (GTR + I + G; Tavaré 1986), with the gamma shape parameter $\Gamma = 0.6963$ and the proportion of invariable sites $I = 0.486$. Both ML and BE methods, yielded phylogenetic trees with an almost identical topology. The topology of the *cyt b* trees indicated the presence of three haplotype groups: *nippon1*, *nippon2*, and *hortul* (Fig. 3a), with the highest diversity found in the haplogroup *nippon1*. The divergence of the three haplogroups is well supported by both used methods (ML = 83; BE = 0.96) (Fig. 3a). The k2p-distance between the two *nippon* groups was estimated to be between 2.9 and 3.5 %, and the p-k2p-distances between *nippon1*/*hortul* and *nippon2*/*hortul* were 2.9–3.7 % and 3.3–4.2 %, respectively (see the Supplementary material). The haplogroups *nippon1* and *nippon2* contained haplotypes found in both Czech and Japanese sika populations. The most frequent haplotype within *nippon1* was J4N (35.14 %), within *nippon2* it was haplotype J1N (70.59 %), and within the group *hortul* two of three haplogroups were present in more than 40 % of




Fig. 1 Distribution and variability of *cyt b* haplotypes at the localities sampled in the Czech Republic




Fig. 2 Distribution and variability of *cyt b* haplotypes at the localities sampled in **a** Japan and **b** Russia




Fig. 3 Maximum likelihood trees for cytochrome *b* sequences. Bootstrap support of ML analysis is indicated *below the branches*, posterior probabilities of BE *above the branches*; **a** whole sample set, **b** Czech samples

the samples analyzed (D35 = 43.37 %; J67D = 52.55 %). The haplotypes from the *nippon1* group belong to the morphological subspecies with larger bodies, found in northern Japan (*C. n. yesoensis*, *C. n. centralis*), while the haplogroup *nippon2* includes the subspecies with smaller bodies that are found in southern Japan (*C. n. mageshimae*, *C. n. nippon*, *C. n. keramae*, *C. n. pulchellus*). The haplogroup *hortul* includes haplotypes found in the samples from Russian and Czech specimens. These haplotypes belong to the mainland subspecies *C. n. hortulorum*. The largest average number of nucleotide

differences was found between the haplogroups *nippon1* and *hortul* (see the Supplementary material). The most distinctive difference found in the topology of two *cyt b* phylogenetic trees was found in the position of haplogroup *nippon1*.

A haplotype network (Fig. 4) constructed using the full dataset revealed the same structure as the phylogenetic trees reconstructed by ML and BE methods. Each of the three main haplogroups was diverged from the neighboring haplogroups by at least ten mutation steps and contained one to two common haplotypes (Fig. 4).




Fig. 4 Median-joining network of sika deer from the Czech Republic, Japan, and Russia, based on the *cyt b* sequences. Numbers of mutations between the haplotypes are indicated on the

branches; the median vectors are indicated by *black dots*. The most common haplotypes are indicated in relevant *circles*



Fig. 5 Distribution and variability of CR haplotypes at the localities sampled in the Czech population

Phylogenetic reconstruction performed by ML and BE methods exclusively on the Czech samples resulted in trees with a very similar topology to the trees constructed for the complete dataset, and confirmed the presence of three haplogroups (*nippon1*, *nippon2*, *hortul*) in the Czech sika population (Fig. 3b). The haplotype diversity in the Czech sika population was higher in the samples collected from the enclosures. From the total number of eight haplotypes found in the Czech population, six haplotypes were found in the enclosures and four in the free-living populations (Table 2). Both of these populations shared two haplotypes—J1N and J4N, four other haplotypes (J35D, J49D, J67D and J173D) were found only in the enclosures, and the last two haplotypes (J93N and J203N) were only present in the samples from free-living animals. The most frequent haplotypes found in the enclosures were J35D (46.53 %) and J4N (32.64 %),

and those found most often in the free-living population were J1N (69.47 %) and J93N (24.21 %). All samples within the haplogroup *hortul* originated in the enclosures, those from *nippon* haplogroups can be found in free as well as in enclosures. While the haplotypes of sika in enclosures contained 36 exclusive polymorphic mutations, only six exclusive polymorphic mutations were found in the free-living population. In total, 36 mutations shared between both populations. The average number of nucleotide differences between the free-living sika and sika in the enclosures was 36.114.

Sequence divergence of the CR region

The analyses of the CR yielded results very similar to those of the *cyt b*. Thus, we present here only the basic




Fig. 6 Distribution and variability of CR haplotypes at the localities sampled in **a** Japan and **b** Russia

results, more detailed information can be found in the Supplementary material.

Altogether, 42 CR haplotypes were identified in the 447 samples analyzed. The length of the analyzed CR sequences varied from 975 to 1,143 bases (including one to five tandem repeat units (23 bp), found in the sequences between 285 and 443 bp). After removing the tandem repetitions, a fragment with the length of 898 bp (including gaps) remained and was used for further analyses, thus decreasing the number of analyzed haplotypes to 32. The haplotype diversity of the whole dataset was $H_D = 0.868$. The Japanese and Czech populations shared no common haplotype, while the Czech and Primorian populations shared two haplotypes. The number of haplotypes in the Czech localities varied from one to four (Fig. 5). A maximum of three haplotypes were found in the Russian localities (Fig. 6 a), while the number of haplotypes in the Japanese localities varied from one to seven (Fig. 6b).

Following the Akaike information criterion, the most suitable model for maximum likelihood and Bayesian estimation was the Hasegawa, Kishino, and Yano + invariant sites + gamma model (HKY + I + G; Hasegawa et al. 1985), with the gamma shape parameter $\Gamma = 0.5844$ and the proportion of invariable sites $I = 6522$. The analyses performed on the whole data set as well as exclusively on the Czech samples yielded phylogenetic trees with a similar topology. The topology of the CR trees indicated the presence of the same three

well-defined haplotype groups that were also found in the *cyt b* tree: *nippon1*, *nippon2*, *hortul* (Fig. 7a, b). Their existence is once again also supported by the haplotype network reconstruction (Fig. 8). Haplotypes belonging to the *nippon2* and *hortul* haplogroups all contained only one repetition, while in those from the *nippon1* haplogroup one to five tandem repeats were found (Table 3).

As was also the case of *cyt b*, the CR haplotype diversity in the Czech sika population was higher in the samples collected from the enclosures (Table 4). Both populations shared two haplotypes—J1N and J4N, and samples from haplogroup *hortul* were only found in the enclosures. The average number of nucleotide differences between the free-living sika and sika from the enclosures was 39.358, and both populations were found to share 44 mutations.

Discussion

As expected, due to the higher mutation rate (Avice 1994), the number of haplotypes found within the CR sequences was higher than within the cytochrome *b* sequences. However, the overall haplotype diversity of the whole dataset was almost identical for both markers analyzed. The separate analysis on the Czech, Japanese, and Russian samples showed that these relatively high values of haplotype diversity were influenced by the inclusion of the highly diverse Japanese samples. The




Fig. 7 Maximum likelihood trees for CR sequences. Bootstrap support of ML analysis is indicated *below the branches*, posterior probabilities of BE *above the branches*; **a** whole sample set, **b** Czech samples

haplotype diversity of the Czech sika population was lower than that of the Japanese population but higher than the diversity of the Russian population. However, the values of diversity within the Czech population were influenced by the presence of both types of haplotypes, those closely related to the Japanese population as well as those from the Russian populations, as these two native populations were not found to share common haplotypes under natural conditions. Despite this, the genetic diversity of Czech sika is low, as is typical for a recently introduced species. We also separately analyzed the haplotype diversity of Czech samples collected from the free-living population and samples collected from animals in enclosures. The free-living population contained only “Japanese-like haplotypes” (*nippon1* and

nippon2) and its diversity was markedly lower than that of the whole Czech dataset. The low genetic diversity of the free-living population may have been caused by a founder effect, resulting from the low genetic variability of the first introduced sika combined with the relatively small number of animals that escaped from enclosures and founded the free-living population. The cytochrome *b* and CR haplotype diversities of the free-living Czech population were half of those of the native Japanese population, suggesting that the introduced individuals might have originated from restricted area(s) in Japan.

The haplotypes found within the samples from enclosures contained both *nippon* and *hortul* haplotypes, thus the diversity of sika in enclosures was higher than that of the free-living population. This may also have




Fig. 8 Median-joining network of sika deer from the Czech Republic, Japan, and Russia, based on the CR sequences. Numbers of mutations between the haplotypes are indicated on

the branches; the median vectors are indicated by *black dots*. The most common haplotypes are indicated in relevant *circles*

Table 3 Number of tandem repeats found within the Czech, Japanese, and Russian samples

	Number of tandem repeats			
	1	3	4	5
Czech Republic				
Free-living	×		×	×
Enclosure	×			×
Japan				
Hokkaido			×	
Honshu	×	×	×	×
Kyushu	×			
Shikoku	×	×	×	
Russia				
Lazovsky District	×			
Terneysky District	×			

Table 4 CR haplotype and nucleotide diversity of free-living sika deer and sika deer in enclosures in the Czech Republic

	N	N_H	H_D	P_i (JC)	k	S
Enclosures	146	7 (9)	0.675	0.03267	27.774	83
Free-living	95	3 (4)	0.446	0.02135	18.268	45

N number of samples, N_H number of haplotypes, H_D haplotype diversity, P_i (JC) nucleotide diversity Jukes–Cantor-corrected, k average number of nucleotide differences, S number of polymorphic sites

been partly influenced by the continuous import of individuals from Japan and Russia, as well as from other countries with introduced sika (Vach et al. 2010). Despite these new introductions, the genetic variability in the enclosures is still relatively low and the number of *hortul* haplotypes in the Czech population is almost as low as it is in the native Russian population, which suffered a significant decrease in the 1940s (Voloshina and Myslenkov 2009). According to Kokeš (unpublished data), the first confirmed introduction of *C. n. hortulorum* dates back to 1903, and it was followed by further introductions of animals from Vala Co. in Přerov Czech republic and Hagenbeck Co., Germany (Kokeš, unpublished data). Many of the introductions were unsuccessful; the fate of other introduced animals is unknown. Thus, it can be assumed that the number of

founder animals of the Czech population was rather small, resulting in the low haplotype variability in the current enclosed as well as free-living population.

Despite the well-documented dates of the first sika introductions to the Czech Republic, the first introduction in 1891 to the Bor enclosure and the second introduction in 1897 to the Lipí enclosure (Kokeš, unpublished data), little is known about the origins of the introduced animals (Kokeš 1970). The small amount of remaining data indicates that the introduced animals originated from both mainland (Far East Russia) and island (Japanese Islands) populations (Kokeš 1970). In order to determine as precisely as possible the geographical origin of this allochthonous deer species in the Czech Republic, we acquired samples from both places of sika origin—Far East Russia and the Japanese Island. Two mtDNA markers, the cytochrome *b* gene and the control region, were used to determine the phylogeographical origin of the free-living and enclosed Czech sika. The topology of both *cyt b* and CR phylogenetic trees in our study showed the existence of three diverged haplogroups within the whole dataset, as well as within

the Czech population. Two of these groups, *nippon1* and *nippon2*, were found to contain samples belonging to Japanese subspecies, whereas the third group, *hortul*, was found to include Russian haplotypes.

It is generally thought that mammals migrated to the Japanese islands from mainland Asia across land bridges that were repeatedly formed between Japan and the Asian continent (Kawamura 1991). In the case of sika, the molecular analyses suggest that it colonized Japan twice. This double colonization is supported by the existence of two different mitochondrial lineages in Japan—the northern and the southern groups as reported by several studies (Nagata et al. 1995, 1999; Tamate and Tsuchiya 1995; Tamate et al. 1998). Nagata et al. (1995) studied the genetic variability of three populations of Japanese sika by sequencing the mitochondrial 12S rRNA and cytochrome *b*. Their results clearly supported the existence of two different populations of sika in northern and southern Japan. Further evidence of repeated colonization could be the closer relationship between the southern Japanese lineage and the Chinese subspecies *C. n. hortulorum* than with the northern Japanese lineage (Nagata 2009), suggesting that the divergence of both lineages probably occurred before the colonization of Japan from China (Kawamura 2009). The existence of the two mitochondrial lineages was also confirmed in the sika populations introduced to Germany and Austria (Pitra et al. 2005). Also, our results confirm the presence of these two lineages, denoted in the phylogenetic trees as *nippon1* and *nippon2*, and these can also be seen not just in the cytochrome *b* and D-loop phylogenetic trees reconstructed from the whole dataset but also in the trees reconstructed solely from the Czech samples. This suggests that the introduced sika individuals brought to the Czech Republic at the end of the 19th century and at the beginning of the 20th century may have originated from a much larger area than expected and might have belonged to two different genetic lineages. Part of the Czech sika population seems to be related to smaller bodied sika (*C. n. nippon*, *C. n. keramae*, *C. n. mageshimae*, *C. n. pulchellus*) from southern Japan, while the other part is closer to the larger bodied sika (*C. n. centralis*, *C. n. yesoensis*) from northern Japan. The haplotypes from the *hortul* group were only found within enclosures; not even one sample from the free-living animals belonged to this group. The reason for this could be that only a few animals from the *hortul* group escaped the enclosures (according to unpublished data of Kokeš, individuals of *C. n. hortulorum* bred in an enclosure in Vlkava, in Bohemia, became a part of the native fauna after its liquidation). Furthermore, another reason could be intraspecific hybridization. The existence of hybridization between sika subspecies in the Czech Republic was confirmed when 32 samples from enclosures that were thought to be *C. n. hortulorum* actually contained haplotypes belonging to Japanese subspecies (Krojerová-Prokešová et al., in prep.). Of course, this could be caused by mistakes in identification caused by the hybridization among sika subspecies, and

hybridization between sika and red deer. A detailed analysis of microsatellite markers on the samples used in the study could provide further data to support, and maybe even improve, the data on the geographical origin of the Czech sika obtained from the mtDNA analyses. It should also help to evaluate the degree of interspecific (sika subspecies) and intraspecific (sika deer × red deer) introgression within the Czech sika and red deer populations.

Acknowledgments We thank H. Pimenova and S. Bondarchuk from the Sikhote-Alin Reserve, the hunters from the Primorsky District and the Terneysky District, M. Asada from the Chiba Biodiversity Centre, and the hunters from the Czech Republic for collection of the samples. We thank our colleague P. Vallo for his laboratory help at the beginning of the study. We also thank both reviewers for their suggestions and comments on previous versions of this paper. This study was financially supported by Grant No. 524/09/1569 from the Czech Science Foundation and by institutional support RVO:68081766.

References

- Abernethy K (1994) The establishment of a hybrid zone between red and sika deer (genus *Cervus*). *Mol Ecol* 3:551–562. doi:10.1111/j.1365-294X.1994.tb00086.x
- Abernethy K (1998) Sika deer in Scotland. Deer commission for Scotland. Chapman & Hall, London
- Akaike H (1974) A new look at the statistical model identification. *IEEE Trans Autom Control* 19:716–723. doi:10.1109/TAC.1974.1100705.MR0423716
- Aramilev VV (2009) Sika deer in Russia. In: McCullough DR, Kaji K, Takatsuki S (eds) Sika deer. Biology and management of native and introduced populations. Springer, Berlin Heidelberg New York, pp 475–500
- Avise JC (1994) Molecular markers, natural history and evolution. Chapman & Hall, New York
- Avise JC (2000) Phylogeography: The history and formation of species. Harvard University Press, Cambridge
- Babička C, Drábek M, Štika J, Ženožička J (1977) Findings about sika deer breeding (in Czech). *Myslivost* 11:270–271
- Balakrishnan Ch-N, Monfort SL, Gaur A, Singh L, Sorenson MD (2003) Phylogeography and conservation genetics of Eld's deer (*Cervus eldi*). *Mol Ecol* 12:1–10. doi:10.1046/j.1365-294X.2003.01751.x
- Bandelt H-J, Forster P, Röhl A (1999) Median-joining network for inferring intraspecific phylogenies. *Mol Biol Evol* 16:37–48
- Banwell DB (2009) The sika in New Zealand. In: McCullough DR, Kaji K, Takatsuki S (eds) Sika deer. Biology and management of native and introduced populations. Springer, Berlin Heidelberg New York, pp 643–656
- Bartoš L (2009) Sika deer in Continental Europe. In: McCullough DR, Kaji K, Takatsuki S (eds) Sika deer. Biology and management of native and introduced populations. Springer, Berlin Heidelberg New York, pp 573–594
- Bartoš L, Vitek M (1993) Cluster analysis of red and sika deer phenotypes. In: Ohtaishi N, Sheng H-I (eds) Deer of China: biology and management. Elsevier, Amsterdam, pp 15–21
- Bartoš L, Žirovnický J (1981) Hybridization between red and sika deer. II. Phenotype analysis. *Zool Anz* 207:20–36
- Bartoš L, Žirovnický J (1982) Hybridization between red and sika deer. III. Interspecific behaviour. *Zool Anz* 208:30–36
- Bartoš L, Hyánek J, Žirovnický J (1981) Hybridization between red and sika deer. I. Craniological analysis. *Zool Anz* 207:260–270
- Bartoš L, Herrmann H, Šiler J, Losos S, Mikeš J (1998) Variation of mating systems of introduced sika deer. *Rev Ecol Terre Vie* 53:337–345

- Chadwick AH, Ratcliffe PR, Abernethy K (1996) Sika deer in Scotland: density, population size, habitat use and fertility—some comparisons with red deer. *Scott For* 50:8–16
- Cook CE, Wang Y, Sensabaugh G (1999) A mitochondrial control region and cytochrome *b* phylogeny of sika deer (*Cervus nippon*) and report of tandem repeats in the control region. *Mol Phylogenet Evol* 12:47–56. doi:10.1006/mpev.1998.0593
- Diaz A, Hughes S, Putman R, Mogg R, Bond JM (2006) A genetic study of sika (*Cervus nippon*) in the New Forest and in the Purbeck region, southern England. Is there evidence of past hybridization with red deer (*Cervus elaphus*)? *J Zool* 270:227–235. doi:10.1111/j.1469-7998.2006.00130.x
- Doležal V (1960) From Manětínsko (in Czech). *Myslivost* 4:174
- Eick E (1995a) Germany. In: Eick E, König R, Willett J (eds) *Sika, Cervus nippon* Temminck, 1838, vol 1, 2nd edn. International Sika Society, Möhnese, Germany, pp 2.2-D/1–13
- Eick E (1995b) A history of naturalisation. In: Eick E, König R, Willett J (eds) *Sika, Cervus nippon* Temminck, 1838, vol 1, 2nd edn. International Sika Society, Möhnese, Germany pp 9.1–9.14
- Feldhamer GA, Demarais S (2009) Free-ranging and confined sika deer in North America: current status, biology, and management. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 615–641
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368–376. doi:10.1007/BF01734359
- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783–791
- Gehle T, Herzog S (1998) Is there evidence for hybridisation between red deer and sika deer in Germany? In: Zomborszky Z (ed) 4th International Deer Biology Congress, June 30–July 4, 1998, Final program—Abstracts. Pannon Agricultural University, Faculty of Animal Science, Kaposvar, Hungary, pp 121–123
- Goodman SJ, Barton NH, Swanson G, Abernethy K, Pemberton JM (1999) Introgression through rare hybridization: a genetic study of a hybrid zone between red and sika deer (genus *Cervus*) in Argyll, Scotland. *Genetics* 152:355–371
- Guindon S, Gascuel O (2003) A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. *Syst Biol* 52:696–704. doi:10.1080/10635150390235520
- Guo Y-S, Zheng H-Z (2000) On the geological distribution, taxonomic status of species and evolutionary history of sika deer in China (in Chinese with English summary). *Acta Theriol Sinica* 20:168–179
- Harrington R (1973) Hybridization among deer and its implication for conservation. *Irish For J* 30:64–78
- Harrington R (1982) The hybridization of red deer (*Cervus elaphus* L. 1758) and Japanese sika deer (*C. nippon* Temminck 1838). In: O’Gorman F, Rochford J (eds) *Transactions XIVth International Congress of Game Biologists*. Irish Wildlife Publications for the Organising Committee of the XIVth Congress, Dublin, Ireland, pp 559–571
- Hasegawa M, Kishino H, Yano T (1985) Dating of the human-ape splitting by a molecular clock of mitochondrial DNA. *J Mol Evol* 22:160–174. doi:10.1007/BF02101694
- Heptner VG, Nasimovitch AA, Banikov AG (1961) *Mammals of the Soviet Union. Part One. Even-toed and odd-toed ungulates* (in Russian). State University Press, Moscow
- Heroldová M (1990) The diet of sika deer (*Cervus nippon*) outside the growing-season, with a snow cover. *Folia Zool* 39:197–206
- Herzog S (1987) Mechanisms of karyotype evolution in *Cervus nippon* Temminck. *Caryologia* 40:347–353. doi:10.1016/0303-2647(90)90008-O
- Irwin DM, Kocher TD, Wilson AC (1991) Evolution of cytochrome *b* in mammals. *J Mol Evol* 32:128–144. doi:10.1007/BF02515385
- Kawamura Y (1991) Quaternary mammalian faunas in the Japanese islands (in Japanese with English abstract). *Quat Res* 30:213–220
- Kawamura Y (2009) Fossil record of Sika deer in Japan. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 11–26
- Kelly DL (2002) The re-generation of *Quercus petraea* (sessile oak) in southwest Ireland: a 25-year experimental study. *Forest Ecol Manag* 166:207–226. doi:10.1016/S0378-1127(01)00670-3
- Kocher TD, Thomas WK, Meyer A (1989) Dynamics of mitochondrial DNA evolution in animals: amplification and sequencing with conserved primers. *PNAS* 86:6196–6200
- Kokeš O (1970) Asian deer in the Czechoslovakia (in Czech). *Ochrana Fauny* 4:158–161
- Librado P, Rozas J (2009) DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinformatics* 25:1451–1452. doi:10.1093/bioinformatics/btp187
- Lowe R (1994) Deer management: developing the requirements for the establishment of diverse coniferous and broadleaf forests. Unpublished Report. Coilte, Bray, Co. Wicklow
- Lowe VPW, Gardiner AS (1975) Hybridisation between red deer and sika deer, with reference to stocks in north-west England. *J Zool* 177:553–566
- Ludt ChJ, Schroeder W, Rottmann O, Kuehn R (2004) Mitochondrial DNA phylogeography of red deer (*Cervus elaphus*). *Mol Phylogenet Evol* 31:1064–1083. doi:10.1016/j.ympev.2003.10.003
- McCullough DR (2009a) Sika deer in Korea and Vietnam. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 541–548
- McCullough DR (2009b) Sika deer in Taiwan. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 549–560
- McCullough DR, Kaji K, Takatsuki S (eds) (2009) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York
- McDevitt AD, Edwards CJ, O’Toole P, O’Sullivan P, O’Reilly C, Carden RF (2009) Genetic structure of, and hybridisation between, red (*Cervus elaphus*) and sika (*Cervus nippon*) deer in Ireland. *Mamm Biol* 74:263–273. doi:10.1016/j.mambio.2009.03.015
- Nagata J (2009) Two genetically distinct lineages of the Japanese Sika deer based on mitochondrial control regions. In: McCullough et al. (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 27–42
- Nagata J, Masuda R, Yoshida MC (1995) Nucleotide sequences of the cytochrome *b* and 12S rRNA in the Japanese sika deer *Cervus nippon*. *J Mamm Soc Jpn* 20:1–8
- Nagata J, Masuda R, Tamate HB, Hamasaki S, Ochiai K, Asada M, Tatsuzawa S, Suda K, Tado H, Yoshida MC (1999) Two genetically distinct lineages of the sika deer, *Cervus nippon*, in Japanese islands: comparison of mitochondrial D-loop region sequences. *Mol Phylogenet Evol* 13:511–519. doi:10.1006/mpev.1999.0668
- Nylander JAA (2004) MrModeltest v2. Program distributed by the author. Evolutionary Biology Centre, Uppsala University, Uppsala
- Ohtaishi N (1986) Preliminary memorandum of classification, distribution and geographic variation on sika deer (in Japanese). *Mammal Sci* 53:13–17
- Pemberton J, Swanson G, Barton N, Livingstone S, Senn H (2006) Hybridisation between red and sika deer in Scotland. *Deer* 13:22–26
- Pérez-Espona S, Pemberton JM, Putman R (2009) Red and sika deer in the British Isles, current management issues and management policy. *Mammal Biol* 74:247–262. doi:10.1016/j.mambio.2009.01.003
- Pitra Ch, Rehbein S, Lutz W (2005) Tracing the genetic roots of the sika deer *Cervus nippon* naturalized in Germany and Austria. *Eur J Wildl Res* 51:237–241. doi:10.1007/s10344-005-0107-y
- Polzheim R, Strobeck C (2002) A phylogenetic comparison of red deer and wapiti using mitochondrial DNA. *Mol Phylogenet Evol* 22:342–356. doi:10.1006/mpev.2001.1065

- Powerscourt V (1884) On the acclimatization of the Japanese deer at Powerscourt. *P Zool Soc Lond*: 207–209
- Putman RJ (2000) Sika deer. Joint publication by The Mammal Society, London, and the British Deer Society, Hampshire
- Putman RJ, Moore NP (1998) Impact of deer in lowland Britain on agriculture, forestry and conservation habitats. *Mamm Rev* 28:141–163. doi:10.1046/j.1365-2907.1998.00031.x
- Ratcliffe PR (1987) Distribution and current status of sika deer (*Cervus nippon*) in Great Britain. *Mamm Rev* 17:37–58
- Ronquist FR, Huelsenbeck JP (2003) MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574
- Senn HV, Pemberton JM (2009) Variable extent of hybridization between invasive sika (*Cervus nippon*) and native red deer (*C. elaphus*) in a small geographic area. *Mol Ecol* 18:862–876. doi:10.1111/j.1365-294X.2008.04051.x
- Sheng H-L, Ohtaishi N (1993) The status of deer in China. In: Ohtaishi N, Sheng H-L (eds) *Deer of China: biology and management*. Elsevier, Amsterdam, pp 1–11
- Swanson GM, Putman R (2009) Sika deer in the British Isles. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 595–614
- Swofford DL (2003) PAUP*. Phylogenetic Analysis using parsimony (*and other methods). Version 4. Sinauer Associates, Sunderland, Massachusetts
- Tamate HB, Tsuchiya T (1992) Molecular phylogeny of Japanese sika deer (*Cervus nippon*). *Zool Sci* 12:1302. doi:10.1006/mpev.1998.0593
- Tamate HB, Tsuchiya T (1995) Mitochondrial DNA polymorphism in subspecies of the Japanese sika deer, *Cervus nippon*. *J Hered* 86:211–215
- Tamate HB, Tatsuzawa S, Suda K, Izawa M, Doi T, Sunagawa K, Miyahira F, Tado H (1998) Mitochondrial DNA variations in local populations of the Japanese sika deer, *Cervus nippon*. *J Mammal* 78:1396–1403
- Tavaré S (1986) Some probabilistic and statistical problems in the analysis of DNA sequences. *Lect Math Life Sci (Am Math Soc)* 17:57–86
- Thévenon S, Thuy LT, Ly LV, Maudet F, Bonnet A, Jarne P, Maillard J-C (2004) Microsatellite analysis of genetic diversity of the Vietnamese sika deer (*Cervus nippon pseudaxis*). *J Hered* 95:11–18. doi:10.1093/jhered/esh001
- Vach M et al. (2010) Development of game management and hunting in the Czech Republic. Publisher Silvestris PB tisk, Příbram (In Czech)
- Vavruněk J, Wolf R (1977) Breeding of red deer in West-Bohemian region. *Textbook Sci For Inst VŠZ Prague* 20:97–115 (in Czech)
- Voloshina IV, Myslenkov AI (2009) Sika deer distribution changes at the north extent of their range in the Sikhote-Alin Mountains of the Russian Far East. In: McCullough DR, Kaji K, Takatsuki S (eds) *Sika deer. Biology and management of native and introduced populations*. Springer, Berlin Heidelberg New York, pp 501–520
- Whitehead GK (1993) *The Encyclopaedia of deer*. Swan Hill Press, Shrewsbury
- Wilson RL (2000) An investigation into the phylogeography of sika deer (*Cervus nippon*) using microsatellite markers. M.Sc. thesis, University of Edinburgh, Scotland, UK
- Wo C, Smith KG (1999) History and current status of mammals of the Korean Peninsula. *Mammal Rev* 29:3–33. doi:10.1046/j.1365-2907.1999.00034.x
- Yang Z, Rannala B (1997) Bayesian phylogenetic inference using DNA sequences: a Markov chain Monte Carlo method. *Mol Biol Evol* 14:717–724
- Yuasa T, Nagata J, Hamasaki S, Tsuruga H, Furubayashi K (2007) The impact of habitat fragmentation on genetic structure of the Japanese sika deer (*Cervus nippon*) in southern Kantoh, revealed by mitochondrial DNA sequences. *Ecol Res* 22:97–106. doi:10.1007/s11284-006-0190-x