

Identification of a major QTL for time of initial vegetative budbreak in apple (*Malus x domestica* Borkh.)

Maria M. van Dyk · Mogamat Khashief Soeker ·
Iwan F. Labuschagne · David Jasper G. Rees

Received: 18 June 2009 / Revised: 7 December 2009 / Accepted: 11 December 2009 / Published online: 19 January 2010
© Springer-Verlag 2010

Abstract In the Western Cape region of South Africa, dormancy release and the onset of growth does not occur normally in apple (*Malus x domestica* Borkh.) trees during spring due to the mild winter conditions experienced and fluctuations in temperatures experienced during and between winters. In this region, the application of chemicals to induce the release of dormancy forms part of standard orchard management. Increasing awareness of the environmental impact of chemical sprays and global warming has led to the demand for new apple cultivars better adapted to local climatic conditions. We report the construction of framework genetic maps in two F1 crosses using the low chilling cultivar ‘Anna’ as common male parent and the higher chill requiring cultivars ‘Golden Delicious’ and ‘Sharpe’s Early’ as female parents. The maps were constructed using 320 simple sequence repeats, including

116 new markers developed from expressed sequence tags. These maps were used to identify quantitative trait loci (QTL) for time of initial vegetative budbreak (IVB), a dormancy related characteristic. Time of IVB was assessed four times over a 6-year period in ‘Golden Delicious’ x ‘Anna’ seedlings kept in seedling bags under shade in the nursery. The trait was assessed for 3 years on adult full-sib trees derived from a cross between ‘Sharpe’s Early’ and ‘Anna’ as well as for 3 years on replicates of these seedlings obtained by clonal propagation onto rootstocks. A single major QTL for time of IVB was identified on linkage group (LG) 9. This QTL remained consistent in different genetic backgrounds and at different developmental stages. The QTL may co-localize with a QTL for leaf break identified on LG 3 by Conner et al. (1998), a LG that was, after the implementation of transferable microsatellite markers, shown to be homologous to the LG now known to be LG 9 (Kenis and Keulemans 2004). These results contribute towards a better understanding regarding the genetic control of IVB in apple and will also be used to elucidate the genetic basis of other dormancy related traits such as time of initial reproductive budbreak and number of vegetative and reproductive budbreak.

Communicated by E. Dirlwanger

M. M. van Dyk (✉) · M. K. Soeker · D. J. G. Rees
Department of Biotechnology, University of the Western Cape,
Private Bag X17,
Bellville 7535, South Africa
e-mail: daleen.vandyk@up.ac.za

I. F. Labuschagne
Agricultural Research Council (ARC) Infruitec-Nietvoorbij,
Private Bag X5026,
Stellenbosch 7599, South Africa

I. F. Labuschagne
Colors Fruit (SA) (Pty) Ltd,
3rd Floor Newlink Centre, New Street,
Paarl 7646, South Africa

M. M. van Dyk
Department of Genetics, University of Pretoria,
Lynwood Road,
Pretoria 0002, South Africa

Keywords Microsatellites · Marker-assisted selection ·
Dormancy

Introduction

The domesticated apple (*Malus x domestica* Borkh.) has been distributed into diverse climatic conditions worldwide for commercial production of fruit. Apple trees need exposure to cold temperatures, referred to as chill unit (CU) accumulation during winter, in order for budbreak to

occur promptly and uniformly after winter (Cook and Jacobs 2000). In warmer production areas, such as the Western Cape region of South Africa, the application of dormancy breaking chemicals, forming part of standard orchard management, enable successful production of high chilling requiring apple cultivars in suboptimal environmental conditions. Failure to apply dormancy-breaking chemicals can result in prolonged dormancy symptoms, which include extended rest, less synchronized breaking of buds, and reduced branching (Labuschagné et al. 2002b). An increasing awareness of both global temperature increase and the negative effects associated with the use of chemical sprays (for both pest and disease resistance and growth regulation) has resulted in the need to breed cultivars better adapted to current and future environmental conditions.

The breeding of new cultivars using conventional breeding methods is a time-consuming process, especially in perennial tree species with a long juvenile phase such as apple. Markers linked to genes involved in apple disease resistance for a variety of pests and pathogens have been identified (Gardiner et al. 2007) and are already in use in breeding programs (Kellerhals et al. 2008; Tartarini and Sansavini 2003; Tartarini et al. 2000), through the implementation of marker-assisted breeding (MAB) and selection (MAS) that enables the selection of favorable genotypes at a very early seedling stage. The genetic determinants of dormancy-related characteristics, such as time of initial vegetative budbreak (IVB), are still poorly understood, and this hampers the genetic improvement of such characters using MAB. Dormancy characteristics can be controlled by factors residing within the bud itself, referred to as endodormancy, by factors in the plant but outside of the bud (paradormancy) and control by environmental factors (ectodormancy; Khan 1997; Lang et al. 1985). Although our study focused on time of IVB, a character related to endodormancy (Bradshaw and Stettler 1995), various other characteristics can be associated with dormancy, such as position and number of budbreak and budbreak duration.

Unravelling of the genetic basis of complex traits such as dormancy can be undertaken through the construction of a genetic linkage map followed by quantitative trait loci (QTL) identification (Falconer and Mackay 1996; Young 1996). A first attempt towards understanding the genetic control of ‘leaf break’ in apples through the identification of QTLs, was performed by Conner et al. (1998) using a population of 172 trees derived from a cross between ‘Wijcik McIntosh’ and NY 75441-58. Eight genomic regions on seven linkage groups (LGs) could be associated with time of budbreak. The genetic linkage map constructed during their investigation, however did not include transferable simple sequence repeat (SSR) markers, resulting in their inability to align this map with the now more

commonly used LG numbering for apple genetic linkage maps (Maliepaard et al. 1998). Further investigation resulted in alignment of three LGs from these two maps, including one (LG 3) that was homologous to LG 9 of Maliepaard et al. (1998) and carried a QTL for leaf break (Kenis and Keulemans 2004). More recently Segura et al. (2007) used 123 seedlings derived from a cross between ‘Starkrimson’ and ‘Granny Smith’ to identify two QTLs for time of budbreak. The first on LG 8, corresponded to that identified on the corresponding LG 7 by Conner et al. (1998) (see Kenis and Keulemans 2004). The second QTL for time of budbreak identified by Segura et al. (2007) was on LG 6.

In the present study, genetic linkage maps were constructed for two mapping pedigrees with the low chilling requiring cultivar ‘Anna’ as common male parent. ‘Anna’ is one of only a few cultivars worldwide characterized by a low chilling requirement (CR) and with ‘Dorsett Golden’ was reported as varieties needing less than 300 h of chilling in Southern California (<http://ucce.ucdavis.edu/files/filelibrary/5764/33384.pdf>) and North and North Central Florida (Andersen and Crocker 2000). Both published SSR markers (Celton et al. 2009; Guilford et al. 1997; Hemmat et al. 2003, 1997, Liebhard et al. 2002; Silfverberg-Dilworth et al. 2006; Yamamoto et al. 2002a, b) and 116 new SSR markers, developed from expressed sequence tags (ESTs), were used for the construction of the genetic linkage maps used to identify a major QTL for time of IVB on LG 9.

Materials and methods

Plant material

Two F1 progenies, derived from crosses between the low chilling ‘Anna’ (common male parent) and the higher chill requiring ‘Golden Delicious’ (population A) and ‘Sharpe’s Early’ (population B), containing 87 and 92 individuals, respectively, were used. Seedlings from population A were kept in seedling bags under shade netting in Groot Drakenstein (Western Cape, South Africa; 33°50’36” S 18°58’39” E). Seedlings in this population were cut back and re-grown to single shoots on a seasonal basis and no chemical treatment was used to induce budbreak. Seedlings from population B were planted in an orchard in Vyeboom (Western Cape, South Africa; 34°4’15” S 19°4’47” E) characterized by low winter chilling. Resulting trees were in their fifth growing season at the onset of this investigation. Seven clonal replicates from seedlings in population B and the two parental cultivars were grafted onto rootstocks (M793) and planted in seven randomized blocks in an adjacent orchard (34°8’21” S 19°0’44” E). Both sites are

characterized by warmer winters and fluctuating chilling accumulation between winters. At these sites CU accumulation varies between 500 and 1,000 CU annually. Chill units were calculated according to a modified Utah model found to be more suitable for local chilling conditions where negative CU values are not carried from one day to the next (Linsley-Noakes et al. 1994). Orchard management of adult and juvenile clonal trees from population B were typical of commercial practice, except that no pruning and tree growth manipulations, such as dormancy breaking chemicals, were applied.

Phenotypic assessment

The time of IVB was scored as the day on which the first green leaves emerged from the vegetative buds (day 1 being January 1; Labuschagné et al. 2002a, b). Phenotypic trait assessments were performed four times over a period of 6 years (1999, 2000, 2002, and 2004) on the 87 seedlings from population A. Trait assessment of population B was first performed during a 3-year period, from 1996 to 1998, on 60 adult trees, initially in their fifth growing season, followed by trait assessment on the seven clonally replicated juvenile trees of all 92 siblings from 1998 to 2000. The data obtained from population B has been used in previous studies (Labuschagné et al. 2002a, b) during which broad sense heritability of IVB was estimated between 0.62 and 0.92 in clonal trials on young seedlings and between 0.57 and 0.83 for adult seedling trees. We calculated Pearson's correlation coefficients to determine the relationship between different years of phenotypic trait assessment.

DNA extraction

Extraction of seedling and parental cultivar DNA were performed using the hexadecyl trimethyl ammonium bromide (CTAB) method described by Doyle and Doyle (1990) with the addition of polyvinylpyrrolidone (Kim et al. 1997) in order to bind secondary plant products such as polyphenolics.

SSR marker development and implementation

Unigene sets obtained from the large public EST database (>240,000; <http://www.ncbi.nlm.nih.gov>; Naik et al. 2006; Newcomb et al. 2006) for *Malus*, were searched for SSRs using the Tandem Repeats Finder algorithm (Benson 1999). SSRs were selected based on length of the repeat unit, number of repeats (>10 for di-, >7 for tri-, >5 for tetra- and penta-, and >3 for hexanucleotide repeats) and length of sequences flanking SSR regions. Conserved sequences flanking 196 selected SSRs (100 di-, 60 tri-, 25 tetra-, 5 penta-, and 6 hexanucleotide repeats) were used to design

primers resulting in amplicons ranging between 100 and 450 bp in length. Newly developed SSR markers were tested on the three parental cultivars, 'Anna', 'Sharpe's Early', and 'Golden Delicious'.

Markers for map construction were selected based on map position as well as heterozygosity observed during previous studies. They included 238 previously published SSR markers (Celton et al. 2009; Guilford et al. 1997; Hemmat et al. 2003, 1997; Liebhard et al. 2002; Silfverberg-Dilworth et al. 2006; Yamamoto et al. 2002a, b), marker AG11 (unpublished data, A. Patocchi (ETH-Zürich, CH)) and marker *Md-EXP7* (Costa et al. 2008) and were initially screened for polymorphism over the three parental cultivars, 'Anna', 'Sharpe's Early', and 'Golden Delicious'.

SSR analysis

All SSR markers implemented in mapping populations were fluorescently labeled and up to 16 markers were multiplexed using both size and fluorescent dye (6-FAM™, VIC™, NED™, and PET™) differences. Polymerase chain reaction (PCR) reactions were performed using the Qiagen multiplexing kit (QIAGEN Ltd., West Sussex, RH10 9NQ) according to the manufacturer's instructions. Resulting PCR products were prepared for capillary electrophoresis by adding 1 µl of a 1:10 diluted PCR product to 10 µl Hi-Di formamide containing 0.15 µl GeneScan™-500 LIZ™ size standard (Applied Biosystems). Genotyping was performed using the ABI Prism 310 and 3130 (16-capillary array system) Genetic Analyzers (Applied Biosystems, Foster City, CA, USA). Data collection and analysis were performed using GeneMapper®4 software (Applied Biosystems, Foster City, CA, USA).

Genetic linkage map construction

For both progenies, parental genetic linkage maps and integrated genetic linkage maps were constructed using JoinMap®4 (Van Ooijen 2006). A logarithm of the odds (LOD score) of 4 was used to define LGs and genetic distances between markers were calculated using the Kosambi mapping function. On the basis of previously mapped SSRs, LGs were numbered in accordance with the 17 LGs obtained by Maliepaard et al. (1998).

QTL analysis

QTL analysis was performed using MapQTL®5 (Van Ooijen 2004) using the average phenotypic value for the 4 years of phenotypic trait assessment performed on population A and the two 3-year periods of trait assessment performed on adult and juvenile trees from population B. Analyses were also performed separately for each year of

phenotypic trait assessment and in the case of clonal replicates, the mean value per genotype was used. Regions with potential QTL effects were identified using interval mapping with a step size of 1 cM. QTLs were declared significant if the maximum LOD, obtained after multiple rounds of MQM mapping, exceeded the genome wide (GW) LOD threshold (calculated with an error rate of 0.05 over 1,000 permutations). QTLs were characterized by the maximum LOD score and the percentage of phenotypic variation explained. For each QTL the differences in mean time of IVB associated with the different genotypic classes, *ac*, *ad*, *bc*, and *bd*, derived for an *ab* x *cd* cross, are reported. QTLs were graphically displayed as bars next to the LGs on which they were identified, with bars corresponding to a 95% confidence interval (LOD score drop of 0.5) and dotted lines corresponding to a 90% confidence interval (LOD score drop of 1).

Results

Phenotypic trait assessment

Bi-modal distribution patterns were observed during most years of phenotypic trait assessment (Fig. 1). The distribution patterns indicate budbreak was occurring earlier during consecutive years as trees matured. Significant levels of correlation were found between the different years during which phenotypic trait assessment were conducted (Table 1). High broad sense heritability values ($h^2=0.69$) for IVB were calculated by Labuschagné et al. (2002a).

SSR marker development and implementation

The amplification success of newly developed SSR markers was 86% (168 SSRs from a total of 196). From these, a total of 116 new SSR markers were polymorphic in at least one of the three parental cultivars used and were mapped in one or both mapping populations (Table 2). Of the 240 previously published markers, including 238 SSR markers (Celton et al. 2009; Guilford et al. 1997; Hemmat et al. 2003, 1997; Liebhard et al. 2002; Silfverberg-Dilworth et al. 2006; Yamamoto et al. 2002a, b), marker AG11 (unpublished data, A. Patocchi (ETH-Zürich, CH)) and marker *Md-EXP7* (Costa et al. 2008), 232 markers yielded amplification products of which 204 markers were heterozygous in one or more of the three cultivars tested.

Designing new SSR markers so that the resulting amplicons vary in size, enabled effective multiplexing of up to 16 markers in one PCR reaction, greatly reducing the cost involved in the screening of mapping populations. Markers used within each multiplex are very flexible when using the QIAGEN multiplexing kit (QIAGEN Ltd., West

Sussex, RH10 9NQ) that provides optimal reaction conditions that increases specificity and minimizes the effect of primer-dimers and non-specific artifacts often associated with multiplex PCR reactions. The ease with which different multiplexes could be assembled enabled easy assembly of new multiplexes containing highly informative markers for each specific mapping pedigree.

Genetic linkage map construction

The four parental maps constructed (Fig. 2) enabled the positioning of 286 SSR markers on 17 LGs corresponding to the number of chromosomes in the apple haploid genome. The number of SSR markers per LG range from ten SSR markers on LG 3 to 28 SSR markers on LG 10, with an average of 17 SSR markers per LG. The positioning of the 116 newly developed SSR markers (Table 2) range from two SSR markers on LG 1 to 15 SSR markers on LG 10.

Genetic linkage map construction allowed the positioning of five previously published but unmapped markers (Liebhard et al. 2002). CH01b09b was mapped to LG 4, CH01e09b was mapped to LG 10 and CH02h11b was mapped to LG 12 in both mapping populations. CH01e12₁ was mapped to LG 8 and CH05c02 was mapped to LG 11 in the ‘Golden Delicious’ x ‘Anna’ mapping population. Three markers were mapped to different LGs when compared to their location on previously published maps: (1) CH03e03 was mapped to LG 5 compared to LG 3 (Liebhard et al. 2002), most likely due to the amplification of a different locus as observed fragment sizes are slightly larger than published (a fragment size of 216 bp was observed in ‘Prima’ compared to the published 186 bp), (2) Hi23g12 was mapped to LG 15 compared to LG 8 (Silfverberg-Dilworth et al. 2006) confirming results obtained by Patocchi et al. (2009); (3) CH05d04 was mapped to LG 5 compared to LG 12 (Liebhard et al. 2002), also most likely due to the amplification of a different locus as observed fragment sizes are slightly smaller than published (fragments of 154 and 175 bp were observed in ‘Prima’ compared to the published 176 and 186 bp. The marker CH05g07 (Liebhard et al. 2002) was found to amplify two loci, both mapping to LG 12. A locus amplified by the marker Hi03a03 (Silfverberg-Dilworth et al. 2006) was confirmed to map onto LG 6 in both mapping populations used while a second locus amplified by the same marker was found to map to LG 14 in the ‘Anna’ x ‘Sharpe’s Early’ mapping population, confirming structural homology between LG 6 and LG14 (Celton et al. 2009).

Population A

Of the 285 SSR markers screened on 87 seedlings from the ‘Golden Delicious’ x ‘Anna’ mapping pedigree, 260 markers




Fig. 1 Histogram showing the distribution of time of initial vegetative budbreak (IVB) observed in 1 ‘Golden Delicious’ x ‘Anna’ seedlings during 4 years of phenotypic trait assessment a 1999, b 2000, c 2002, and d 2004; 2 ‘Sharpe’s Early’ x ‘Anna’ juvenile trees during 3 years

of phenotypic trait assessment a 1998, b 1999, and c 2000; and 3 ‘Sharpe’s Early’ x ‘Anna’ adult trees during 3 years of phenotypic trait assessment a 1996, b 1997, and c 1998

were positioned on the integrated F1 genetic linkage map (map coverage: 1,376.7 cM). Genetic linkage maps constructed for the parental cultivars ‘Golden Delicious’ (map coverage: 1,124.5 cM) and ‘Anna’ (map coverage: 1,292.6 cM) consisted of 163 (including 72 new SSRs) and 170 (including 71 new SSRs) markers, respectively. Parental maps were aligned using 92 SSR markers in common (Fig. 2).

Population B

The ‘Sharpe’s Early’ x ‘Anna’ genetic map was constructed using 230 SSRs genotyped over the 92 F1 seedlings. The integrated F1 genetic linkage map (map coverage: 1,242.6 cM) consisted of 207 mapped SSR markers. Genetic linkage maps constructed for the parental cultivars

Table 1 Pearson’s correlation coefficients indicating phenotypic association ($P < 0.0001$) between different years for time of initial vegetative budbreak (IVB)

Mapping population	Association between different years of phenotypic trait assessment					
	Years 1+2	Years 1+3	Years 2+3	Years 1+4	Years 2+4	Years 3+4
‘Golden Delicious’ x ‘Anna’	0.68	0.69	0.68	0.70	0.78	0.67
‘Sharpe’s Early’ x ‘Anna’ (Adult trees)	0.96	0.94	0.96			
‘Sharpe’s Early’ x ‘Anna’ (Young seedlings) ^a	0.81	0.80	0.90			

^a Clonal trial

Table 2 Summary of 116 new SSR markers, accession number, repeat motif, primers, resulting fragment sizes and genetic linkage map position

Marker	Repeat motif	Forward primer	Reverse primer	Segregating alleles scored		Genetic linkage maps			
				'Golden Delicious' x 'Anna'	'Sharpe's Early'	F1	Parental		
				'Golden Delicious' x 'Anna'	'Sharpe's Early' x 'Anna'	'Golden Delicious' x 'Anna'	'Sharpe's Early' x 'Anna'	'Golden Delicious' x 'Anna'	'Sharpe's Early' x 'Anna'
S Ams CO865608	TC (13)	CAACAAGTGTGCCTCTGTGG	AGCAAGCAACAGATCAAGCC	160-168	160-164	1	1	1	1
S Ams DR995748	TC (16.5)	TACACCAAGCGCCACACCG	TGGCGAGCAGCATGAGCG	314-334	334	1	1	1	1
S Ams CN495924	TC (14.5)	CTCTCAATGAGTCCCTGC	AAACCCTGTTCATCTTCC	148-175	150-173	2	2	2	2
S Ams CN581002	TC (10)	TGGAGGAAAGGAGAAGCAG	CTTGGAAAGCTTCTGTACAGC	253-267	241-253	2	2	2	2
S Ams CO904847	GTT (11.3)	GTTGGGTGTGTTTTTGTATGG	AGCTAAAGGAGAGCTACACC	190-193	182	2	2	2	2
S Ams CN944528	GAC (11.7)	GACGACGGAAAAGAAAGACG	ATTACGCTGTTCAGAGAGC	204	204-214	2	2	2	2
S Ams EB106592	TCC (8.7)	CTTTGGAAGCCAAACGAACC	AGAGGAGCTTGTGTGAGG	236	233-236	2	2	2	2
S Ams EE663746	GA (15.5)	TGGCAATACCCGTTTGACC	CCATCAATACAAAGCCACC	305-307	305-317	3	3	3	3
S Ams AU301301	CT (14.5)	GGCATAGCAATGCTTGAAGG	GAATAGCACAAGGAGGTTGC	228-234	223-241	3	3	3	3
S Ams CN944444	AAG (8.7)	TAGTGCAAGTACTGGGCCC	CATCGATAGATAGGACGGC	371-374	371-378	3	3	3	3
S Ams EB132187	GGA (8.3)	TCTCCCTCACTCGAGCTTG	GTTGCAGGAAAGGAGTGTCC	243-250	253	4	4	4	4
S Ams CV128959	TC (11.5)	AAATAGTGTGGAAGACGGGG	CAATATACTAATGATCCTTCC	240	232-242	4	4	4	4
S Ams CN579721	CT (14)	GATCCAAATCTCAAACCCCTCC	GTTGAAAGCGTGGTTGGCC	246-259	248-259	4	4	4	4
S Ams EB153928	CT (25)	CTCAAATCCCAAGAGATTATCC	GTCCTCGAATCGTCTCTCC	348-350	350-357	4	4	4	4
S Ams CO052033	CT (11.5)	TTGCCAATCCGCAATTCGCC	TGAGGTTCCCGCCCTTGC	118	118-196	5	5	5	5
S Ams CO756306	AAAT (5.8)	GTAATATATCACCAACCCCGC	ACACAGAACTGTGATCATCG	180-184	180	5	5	5	5
S Ams CO416051	AG (16)	CCTCACTAAACGCATTTGCAC	CGGTACGATGAGGATCATCC	120-133	120-130	5	5	5	5
S Ams CN922831	TC (13)	TTTAGAATTCGGAGAGGATAGG	CTGCTTGGAAATCCTCGAGC	293	290-293	5	5	5	5
S Ams CN887525	TTTA (7.8)	TAGTAGCTACACACTTTTCC	GCAATGCTTGAAGCTCCAG	207	207-214	5	5	5	5
S Ams CN544835	AG (17.5)	AGGAGAGCTTTCTGCAATTC	AGCGTATCCCCAGCTGC	301-303	303-305	5	5	5	5
S Ams CN444942	CT (16.5)	GCCTCAAAAGTCTCTCCAGC	TACGGACTCTCTTTGGGGC	265-273	275	6	6	6	6
S Ams EG661303	AT (25)	GGCATGTGAATATGGTGGAGC	CCAATCAATGTCTTGTCTACC	330-351	327	6	6	6	6
S Ams EB127535	GA (30)	AACACACACACACCACCATTCG	TAGGAAAGTCGACGATGTCG	326-330	322-330	6	6	6	6
S Ams CV657225	TAT (10.3)	TCCCTGTCAATCGAATGATGC	GCAAAACCCAAATCAGAAAGGAC	193	193-198	6	6	6	6
S Ams DT041144	AG (15)	AAATGCTGCAGTGAAGGCC	GAATTCATCTAAACAGAGAGC	349-351	349	6	6	6	6
S Ams CN927330	ACC (7.3)	TTAAACTGCCAAATTTGCACGG	GTTGGGTATTTGCATGTTGG	438-443	431-438	7	7	7	7
S Ams CN903950	AGA (14.3)	TTTCCCTTTTGGCCAGTGCA	GTTTGGGCCCTCGATGATGG	306-319	297-319	7	7	7	7
S Ams CO756781	CT (19.5)	ATAAGTTTAGGCTCATCTGCC	AAACCCATCCCACTTAAAGCC	355-361	333-361	7	7	7	7
S Ams CO901343	(CT) 15.5	CACCTCTCCCTCATCATGTC	CGACAAGGAGACTGAGAGG	208-222	208	7	7	7	7
S Ams CN488733	TC (13)	CACAACCAATCCCAACATC	CAGCCGGAGCATCTACC	127-131	131-142	7	7	7	7
S Ams EB127208	AG (14.5)	ATTCTCTCAACCCCTATCC	CACAGTGTGTTAAAGCTGG	479-491	491	7	7	7	7

S_AmsABI62040	TC (39.5)	GGAGTGCATTAGCTCTCTCC	TCCTTGAATCTCAACTCTAGG	266	266-272	7	7	8	8
S_AmsCV883434	TC (23)	CGAAACTGGTCGAAGAACCCT	AAACTACACAGACGAAATGG	331-335	350-354	8	8	8	8
S_AmsEB151277	TC (29)	TCCTCAATCTCTCTCAATAACC	GCGTTCTAGAGAGAGAAAAGG	179-197	197-202	8	8	8	8
S_AmsCN891581	TCC (8)	CCAAAACTCCACGACCGC	CCAGAGCTTTAGGACTCG	294-297	294-null	8	8	8	8
S_AmsEB176883	TGCT (8.5)	AAAAGCTGCTTGCTTGATTCG	ACCAATCAGCTGGGTTCTCG	330-338	322	9	9	9	9
S_AmsCX025465	GAC (10.7)	TGCTAGAGCTGGGTTCTCC	TCGCAGACTGCTCGCTGC	232-238	232-238	9	9	9	9
S_AmsCO903298	TC (14)	TTGAGAAACAATGCTGCCTC	TGCCACAGTTGGAAGGTGG	344	344-350	9	9	9	9
S_AmsEB149750	TC (19)	ATCAAAGGTGTGAGTGTGTGC	AAGCTTGCATCTCTAGGTCC	258-263	255	9	9	9	9
S_AmsCO900452	GA (12.5)	CAAGGCATCTCCCTCATTTGG	TACTACAGTTCCGATCAAAAGC	291-314	293-295	9	9	9	9
S_AmsEG631184	GA (10)	CTTATGGACCTGCAAAATGG	AGACTCTGTACAFACATCTCC	447-464	447	9	9	9	9
S_AmsDR992457	AGC (13.7)	TCTCCAAGTGGACGAATFCAG	TCCTCAGTGAAGACAAAACC	360-370	356-368	9	9	9	9
S_AmsCO898678	CT (16)	CCCAAGTGCACCAACATACAG	AGCTTCTGGCAGCAAGTGC	242	238-244	9	9	9	9
S_AmsDR999029	TC (14)	CGCCCTCACCTCAITTCAGTC	TCAAATGAACTTTCAGTCCG	440	440-443	9	9	9	9
S_AmsCO865207	GA (13.5)	TGCACCAAATAAGCCGATCC	CAAGAAATGCAACCAGTCGA	134-138	120	9	9	9	9
S_AmsCN444550	TGGAT (5)	AGCATCAAGCCAAATCTTAAAGC	GTATGCTCTTCTTCTCATGG	346-351	341	10	10	10	10
S_AmsEB132791a	CT (17.5)	CACTACAGAACTCTCTATCC	GTGGATGGAACCCGAAACC	312	312-316	10	10	10	10
S_AmsEB132791b				344-350	340-350	10	10	10	10
S_AmsCO755814	CT (21)	AACATCAAGACAGAGAAGAGC	CGTCTTCTTCAACAACCTCCG	263	257-263	10	10	10	10
S_AmsCN996777	CACCT (5)	TGACAACTATGATCGAAGTGG	TTT CATATCACATGACGTGGC	270-275	275	10	10	10	10
S_AmsCN865016	CAT (14)	TTCTTACACCCCTCAATCC	AAAGCCCTGGGATTGGC	340-345	334-340	10	10	10	10
S_AmsU50187	GA (17.5)	ACCTGAGAGAGCTCCAAAAGC	GTGCCCCAGTCAAATACG	160-null	149-162	10	10	10	10
S_AmsEB153442	CT (23.5)	GGTTCACAAAGCCAACTTTG	ATGGTTCGATCGGTTTAAATGC	366-373	373	10	10	10	10
S_AmsDR990381	TCT (9.7)	AAACACTACTGTGCTGGTGG	AGTCCACTTACTACTCTCC	287-300	300	10	10	10	10
S_AmsDR996792	CT (15)	AGGCTTCTCTCTTCTTCTCC	GGACCATTGTGGTGGAGC	378-399	397	10	10	10	10
S_AmsCO751676	TC (15)	TGTGGCTCTGGATGGTTCC	TACGAGTCCATCCGTAATGC	233	218-228	10	10	10	10
S_AmsCN879152	ATC (7.3)	CGTTGGAGATGATCAGTACG	ACCTACAATAGTAGTGGAGAC	256-null	243-256	10	10	10	10
S_AmsCN489062	GA (13)	ACAACTTGGTTACCGGACAC	GAACAAGTATAGGTCGCTGG	296-300	284-314	10	10	10	10
S_AmsDR994153	AG (14.5)	CACGAGGCGAAAACCGATC	AGTCTCTCAGAACCCTGAGC	465-472	463-465	10	10	10	10
S_AmsEB149851	AGA (10.3)	GAA CAG AGG GAA GCA GAC G	AGA AGT GGC AAC CAT GTT GC	187-190	190-202	10	10	10	10
S_AmsCN877882	CTAGT (6.8)	AAC TTGCTGAGAGAGTAATGG	CAACCAAAGGCCCTGAAAGC	485	495-500	11	11	11	11
S_AmsEB128431	TAA (17)	ACGTAGTGATACCCGGATTCC	AGAGTAGCTAGAGATAITCC	335	342-null	11	11	11	11
S_AmsDR994274	ACC (12.3)	CCACCCACAAAACATACACC	TGCTGTTGTGGTGTGATGG	228	221-228	11	11	11	11
S_AmsDR993043	TC (13.5)	CACGAGGGTAAAGCTCCCC	TTGGGGTATTGCTCTGACG	298-314	279-304	11	11	11	11
S_AmsCN942929	GTTT (5)	ACGCTAGGAGAGAGGAACG	GAGCATCCCGTATAAATCCG	519-524	524-529	11	11	11	11
S_AmsCN580620	CGG (7)	TGCGGTCAAACGATGCTTCC	AAGTACAAGCCCGCAAAGG	380	377-380	12	12	12	12
S_AmsEB139609	AG (32)	ACCATATACATCTCTCTCTGC	TTCAAGAGCTGTTGTTGTTGG	322-334	313-358	12	12	12	12
S_AmsCN943613	CTT (7.3)	TAGCAGAAAACACAGATGG	GAAAGCCCAATTGGAGC	165-174	174	12	12	12	12
S_AmsDR995002	GAT (8)	ATCTGATGGTGCATCGGTAG	TTAGGGTCTTCTTGTACCGC	329-332	332	12	12	12	12
S_AmsCN492206	TTG (10.7)	ACATACTGGAGTCTGGGAGC	CAATACGCTAGTGAAGACGC	398	398-471	13	13	13	13
S_AmsCO052555	AT (12.5)	GAA GTTCTATCAAGTCTTGC	GCTTCTGCACAATGGCTGG	232-234	236	13	13	13	13
S_AmsCN445562	TC (23.5)	CACAAAACCAACCGTCTAACCC	GCTCTTGATCATAGGCGTGG	139-154	150-154	13	13	13	13

Table 2 (continued)

Marker	Repeat motif	Forward primer	Reverse primer	Segregating alleles scored		Genetic linkage maps					
				‘Anna’	‘Golden Delicious’	‘Sharpe’s Early’	F1		Parental		
							‘Golden Delicious’ x ‘Anna’	‘Sharpe’s Early’ x ‘Anna’	‘Golden Delicious’ x ‘Anna’	‘Sharpe’s Early’ x ‘Anna’	
S Ams CO416477	CT(14)	CCACACAACACAAACCAACC	GAGCAATTGATCCTCATCGT	218	218-224	13		13			
S Ams CO068842	TC (22)	TGGTTGGAGATGTTCCATGG	ACCACTAGATTATCTTCTGC	455-null	401-447	13		13			
S Ams EB154452	GATC (5)	CACCTAACTCACGTTTCTCC	AGGCAGAAAGGCAAGAGAGG	169-174	174-184	14		14		14	14
S Ams CN880881	CCA (10)	AFAGCTCATACCGCTTCTCC	GTGACGAAAACCAAGAACCC	427-429	406-408	14		14		14	14
S Ams CN491038	TC (19)	GCCTGTCTCGTTGATCGG	AGTGTCTCACCCCTCTTGC	498-510	510	14		14		14	14
S Ams CN581649a	CAT (13.3)	AGCCCTGATCTTCCCTTAGC	GACAATCTTCTGAAAAGTCTGG	343-351	351-354	14		14		14	14
S Ams EB144379	GGCGGT (4.5)	AGCTGATGGCCAGA ACTGC	GAGGGTCCAAGTTACAAAGG	418	412-418	14		14		14	14
S Ams CN494928	ATC (14)	AATTATATCCGTCGACTCCA	TTACTGCTACCTGATGATCC	226	209-219	14		14		14	14
S Ams EB114233	GA (11.5)	GCATCCGCCATTGTTGTCG	TGGATTGAGACTGAGAGAGG	221-227	217-223	14		14		14	14
S Ams EB147331	CT (26)	CCTA ACTCTGACTCAGTTGC	AGTGTCTGCTGGAGCTTCC	257	261-266	15		15		15	15
S Ams CN944665	TATG (10.8)	GTCTCTGCTTGC TTAATTCAG	AGGCCAATCTGACTATAG	320	224-320	15		15		15	15
S Ams CN490349	AGG (8.7)	GTACTATCAGCAGAAA ACTGG	GATTTGAGCACAACTACGG	200	200-206	15		15		15	15
S Ams CN445253	CTG (8)	TGCAAGAAATCATCCACTTCC	TTGACCTGTGAGGACTCC	478-494	491	15		15		15	15
S Ams CO900034	AAG (10.3)	AAAGTCCGTTTTGGGCTGAG	GCTCTGCTGCCAATTCC	361-367	353-367	15		15		15	15
S Ams CO051709	CTC.AAG (3.5)	CTGTGCCGTCACTATATGC	ACCA AAGAGGGAAGAGACG	193	193-200	15		15		15	15
S Ams CN580637	TC(16.5)	ACAACAGCTGACGAACAAGC	CTACTCTCGAAGTACGGCC	418	406-418	15		15		15	15
S Ams CO415353	AG(14)	ATGAACAGTCAAGACTATGC	AACGAAGCAAAAGGAAGACGG	329-333	329-333	15		15		15	15
S Ams CN947446	CTT(8.3)	CCGTTACAGCTATCCAAAACC	ATAATGGCCATTCTGTTACAGC	178-181	181-184	15		15		15	15
S Ams EB126773	CT(23.5)	GTTTGTGTTTGAACAA CGACC	GTGGTTGTTGAGGTCGTGG	447-453	441-447	15		15		15	15
S Ams DT042298	GT(12)	AGCATGTTGTGGGAAAGCCC	GCATACTCTATAFAAAAGTCCG	227-229	225-227	15		15		15	15
S Ams DR997862	TCTG (7.8)	CACAATCATAITCCCGCAGC	TCTTCTCCGATGAGCAAGC	275-280	275-283	15		15		15	15
S Ams CN581649b	CAT (13.3)	AGCCCTGATCTTCCCTTAGC	GACAATCTTCTGAAAAGTCTGG	332-338	332-347	15		15		15	15
S Ams CO868594	CT (19)	CACCTCTTCAAA CAACA CACC	GGCGGAGGTAGTTTATCC	412-414	418-436	15		15		15	15
S Ams CO905375	AG (23.5)	AGTCTCTGTTTTGCTCGTTC	GAACGCCGGGTCCCTGC	407	407-427	15		15		15	15
S Ams CO755991	TC (16.5)	AATCTCTGCTGCAAAACC	GGCACTGAGCGCACTTGG	154	150-154	15		15		15	15
S Ams CN930386	AGA (13)	TTGGGTTTGTGCTGAAAACC	TGACCGGACTGTTTACAGG	94-111	94	16		16		16	16
S Ams CV084260	AG (22)	CAAAGCAAAAACAGAGGATTTG	GGAGCGCATGAAATTA CTGC	226-256	262	16		16		16	16
S Ams CN900718	CAG (7.7)	AGCATCTGAACTACCAATACC	ACCGATATAGTCTGTTGC	278	268-278	16		16		16	16
S Ams EB154700	AG (24)	TTTTTGGGATTTGGGTGCG	GTTGCTGAGAGTGATGATGG	229-236	229-234	16		16		16	16
S Ams CO066563	GA (11)	ACAAAAGAAACAGTGAAGACTC	TACTTGTCTGCATAGTTTGG	422-431	425	16		16		16	16
S Ams EB135348	CCA (11)	ATCCCTAACCCAGGATGG	AGCATGTGAAAATCGTATACC	330-333	330	16		16		16	16

SAmCN881550	CAG (14.3)	ATCCAAACAACCCCAATTGCG	AGTCGATGTTGAACGGCTCCA	346-348	356	16	16
SAmCN868149	AT (10)	TTGCTGCTGTGTGTTTGC	GTCCTGTCGAAATCTTAAAGG	246-252	252	16	16
SAmSDT001786	GA (17.5)	TTTCTGTCTGTGAAAITGCG	GTTAACTGAGCTCTGGTATTCC	143-147	141	16	16
SAmCN943252	TTC (9.7)	TCCCACTGACACTATCAC	TGCAGGAAATGAGAAATGCGC	194	194-197	16	16
SAmEB106034	AAG (11.7)	AGAAAGAAGCCCAATCCCAGC	TTCACCTTCGTCGGCAITGG	191-194	194	16	16
SAmCN910302	TCTG (6.8)	TTTTCAAGGCATCACTGTCC	AICAGGATTTCAACAGCGC	466-484	484	16	16
SAmEE663640	GA (12.5)	AGTGTAGCAACCAAAACGGTG	TTAFTTCTCGTCGGCAAGG	486	483-488	17	17
SAmAU301254	TC (15)	TCCCGGAAATTTTCAACGC	AACGCTAGGATTTGGTCCG	233-246	233	17	17
SAmCO414947	AG (12)	TTTGATTTGGACTGCAATGG	TTAGCAGCTGCTTCAAGTGTG	346-350	341-354	17	17
SAmCN492417	TC (10)	TACCAATGTTTTAGCACCATGG	GGCCAAATAGGTCACAGACG	122	122-126	17	17
SAmCN490324	AG (16)	ATAGAGAGGTAGAGGACTGG	TTCCGCCAGTGTAAACATTGG	230-232	223-232	17	17
SAmCN938125	TTC (13.7)	GCCTTCATCCCCCTTGA	GGTGATATAGGAATCTTGGAG	338-345	345-352	17	17
SAmCN910036	CTT (13.7)	GAGAAACCGTTTGATTACAGC	CTCCATCCCAATCACACC	235-241	232-235	17	17
SAmCN855917	AAT (15.3)	CTCTTCTTCTCCCTTCTCC	GATGAGATCCAAATCCGTAGT	149-174	159-174	17	17
SAmCN929037	TA (13.5)	AGTTGACTACTCTCTCCCGC	GTGGTTCTCACGGTACACG	218-225	218-239	17	17

‘Sharpe’s Early’ (map coverage: 1,012.9 cM) and ‘Anna’ (map coverage: 1,050.6 cM) consisted of 127 (including 41 new SSRs) and 126 (including 45 new SSRs) markers respectively. Parental maps were aligned using 79 SSR markers in common (Fig. 2). The parental map constructed for ‘Anna’ has 94 SSR markers in common with the parental map for ‘Anna’ constructed for population A.

QTL detection and mapping

A single major QTL for time of IVB was detected on LG 9 (Fig. 3). Analyses performed on the average time of IVB for the different populations and developmental stages showed that this QTL exceeded the GW LOD threshold during phenotypic trait assessment performed on adult trees from population B. LOD scores obtained for the analyses performed on averages from population A and juvenile trees from population B were just below the GW LOD thresholds. Separate QTL analysis for the different years of phenotypic trait assessment performed on seedlings from population A resulted in GW LOD thresholds being reached during trait assessment performed on seedlings in their fourth (2002) and sixth (2004) year (Table 3). GW LOD thresholds were exceeded during all 3 years phenotypic trait assessment has been performed on adult trees from population B (Table 3). Separate QTL analysis for the three different years of phenotypic trait assessment performed on juvenile trees from population B resulted in GW LOD thresholds not being reached during the first three juvenile years (Table 3). One-way analysis of variance indicated significant association ($P < 0.0001$) between specific NZmsCN943946 alleles inherited from the parental cultivar ‘Anna’ and time of (IVB). This association was true during all years of phenotypic trait assessment on ‘Golden Delicious’ x ‘Anna’ ($30.22 < F > 91.73$) and ‘Sharpe’s Early’ x ‘Anna’ adult ($34.39 < F > 49.9$) and juvenile ($30.6 < F > 69.27$) trees.

Differences in time of IVB associated with the four genotypic classes, *ac*, *ad*, *bc*, and *bd*, derived from an *ab* x *cd* cross, indicate that the phenotypic variation can be associated with alleles inherited from the common male parent ‘Anna’. This QTL explains between 4.8% and 40.1% of the phenotypic variation observed in population A and between 11.9% and 44.6% of the phenotypic variation observed in population B.

Discussion

The genetic linkage maps constructed are composed entirely of SSR markers and since a very large proportion of these markers are derived from EST sequences (more



Fig. 2 Parental genetic linkage maps of ‘Golden Delicious’ (GD) and ‘Anna’ from population A and ‘Sharpe’s Early’ (SE) and ‘Anna*’ from population B. Numbering of LGs are according to Maliepaard et

al. (1998). Newly developed SSRs are prefixed by SAmS and are indicated in *bold, italic, and underlined*

GDxAn.LG9

SExAn.LG9




Fig. 3 Position of the QTL for time of IVB detected on LG 9 of the consensus ‘Golden Delicious’ x ‘Anna’ (GDxAn) map and the ‘Sharpe’s Early’ x ‘Anna’ (SExAn) map. QTLs are represented by boxes where the length of the box corresponds to a 5% confidence interval and extended lines to a 10% confidence interval. Boxes

representing average time of IVB are filled and boxes representing time of IVB for separate years are open. Boxes indicating QTL detected on juvenile trees are green and those indicating QTL detected on adult trees are red

replace markers with low polymorphism now included in the reference set, due to lack of more polymorphic SSR markers in certain regions (Silfverberg-Dilworth et al. 2006).

The time of IVB showed a wide bi-modal distribution in the seedlings derived from both mapping populations.

Although bi-modality could be explained by seedlings having a difference in their rapidity of response to favorable conditions after their CR was satisfied (Labuschagné et al. 2003), the distribution of time of IVB can be explained by the fact that the trait is controlled by a major QTL together

Table 3 Parameters associated with the QTL for time of initial vegetative budbreak (IVB) identified on LG 9 of the consensus map used for population A and population B, using multiple QTL mapping (MQM)

Year	LOD ^a	mu_ac{00} ^b	mu_bc{00} ^b	mu_ad{00} ^b	mu_bd{00} ^b	% Expl. ^c
Population A: ‘Golden Delicious’ x ‘Anna’						
Average	6.07 (6.8)	271	271	247	252	36.7
1999	3.1 (5.4)	309	311	285	292	25.4
2000	3.91 (8.9)	307	312	277	302	4.8
2002	5.7 (4.4)	282	279	269	271	23.6
2004	7 (4.8)	254	254	226	235	40.1
Population B: ‘Sharpe’s Early’ x ‘Anna’—adult trees						
Average	8.65 (7.0)	252	262	219	219	41.6
1996	9.52 (8.6)	258	267	221	222	44.6
1997	8.04 (6.1)	254	262	224	222	39
1998	6.83 (5.3)	245	256	212	212	38.2
Population B: ‘Sharpe’s Early’ x ‘Anna’—juvenile trees						
Average	4.68 (4.9)	246	246	226	225	17.6
1997	2.56 (4.5)	260	256	245	242	11.9
1998	4.44 (4.6)	254	255	235	232	20.9
1999	4.49 (5.8)	251	253	230	226	17.9

^a Maximum LOD score with considered threshold in parentheses

^b Estimated mean of the distribution of time of IVB associated with each genotypic class with alleles “a” and “b” inherited from the parental cultivars ‘Golden Delicious’ and ‘Sharpe’s Early’, respectively, and alleles “c” and “d” inherited from the cultivar ‘Anna’

^c Percentage of the variance explained by the QTL

with some minor QTLs. High heritability estimates, although specific to the experimental conditions in which they have been calculated, were calculated for time of IVB by both Labuschagné et al. (2002a) ($h^2=0.69$) and Segura et al. (2007) ($h^2=0.58$), indicating that the trait has a strong genetic influence and that it can be selected for using marker-assisted selection. Heritability is not always related to the power of QTL detection (Segura et al. 2007), as the latter is also influenced by population size and the number of QTLs affecting the trait. The small number of individuals included in phenotypic trait assessment (87 from ‘Golden Delicious’ x ‘Anna’ and 60 and 92 for adults and juveniles from the ‘Sharpe’s Early’ x ‘Anna’ mapping pedigrees) and the amount of variation observed among seedlings from the same mapping population, allowed for the detection of only one QTL with large effect. The fact that this QTL explains up to 40.1% and 44.6% of the phenotypic variation observed in populations A and B, respectively, indicates that there are further QTLs affecting time of IVB. These may include several QTLs with smaller effect that are statistically not detectable due the restricted population sizes used and the phenotypic variation observed in the seedlings. During initial interval mapping (van Dyk et al. 2009), the involvement of several minor QTLs were suggested. Implementation of more markers leading to better genome coverage and the ability to perform MQM analysis enabled the identification of a QTL with large effect in the current study.

Genetic linkage maps constructed for both mapping populations enabled the efficient detection of a major QTL affecting the time of IVB on LG 9 (Table 3). This QTL may co-localize with one of eight QTLs involved in leaf break that was identified by Conner et al. (1998). The QTL identified on LG 3 of the genetic linkage map produced by Conner et al. (1998) was, after the implementation of transferable microsatellite markers, shown to be homologous to the LG now known to be LG 9 (Kenis and Keulemans 2004). In the present study, the QTL on LG 9 can be associated with specific allele inheritance from the common parent ‘Anna’. Performing QTL analyses on an integrated parental map when working with an outbreeder, as was done during this study, enables the determination of both the effect of alleles inherited from a single parent and the interaction between alleles inherited from both parents. Results (Table 3) indicated a clear difference in average time of IVB between seedlings that inherited allele “c” from ‘Anna’ (average “ac” and “bc”) compared to seedlings that inherited allele “d” from ‘Anna’ (average “ad” and “bd”). No clear difference could be detected between seedlings that inherited different alleles from the other parental cultivar involved in each mapping pedigree or seedlings with a specific combination of parental alleles.

The power of QTL detection (LOD score) increased during consecutive years of phenotypic trait assessment being performed on seedlings from population A and juvenile trees from population B. This suggests that although the QTL can be associated with time of IVB in young seedlings, the association between the QTL and the trait becomes stronger as the tree matures. The QTL was found to be significant (LOD score exceeding GW LOD thresholds) in all 3 years during which phenotypic trait assessment was performed on adult trees from population B (Table 3). Although significant GW LOD thresholds are not met in juvenile trees from population B, the association between the QTL and time of IVB can be seen from obtained phenotypic means associated with each of the genotypic classes (Table 3). Budbreak occurring earlier as trees mature has not been reported before. Preliminary results suggest no correlation between the earlier time of vegetative budbreak, associated with seedling age in two apple populations studied, and the CU accumulated during different years. These results suggest that the CR, which is the major determinant of time of budbreak (Bradshaw and Stettler 1995), has been met and that the time of vegetative budbreak is also influenced by factors associated with tree age. These results need to be confirmed in future studies, including several years of phenotypic trait assessment performed during different developmental stages and on different populations.

Markers linked to the QTL identified will be used in a validation test on a larger progeny sharing common parentage. The QTL region will be saturated with markers selected for their positioning on the genetic linkage map as a result of selective (bin) mapping on a subset of individuals (van Dyk and Rees 2009). The ideal will be the identification of markers flanking the QTL that can be used for the implementation of MAS in breeding for cultivars that are better adapted to local climatic conditions.

Acknowledgments We would like to thank the Deciduous Fruit Producers Trust (DFPT), the Department of Trade and Industry (DTI-THRIP), and the National Research Foundation (NRF) for financial support. Thank you to Dr. Jean-Marc Celton and Dr. W.E. van de Weg for useful discussion regarding mapping. We are especially grateful to Prof. B. D. Wingfield from the University of Pretoria for the use of facilities.

References

- Andersen PC, Crocker TE (2000) Low chill apple cultivars for North and North Central Florida, University of Florida Extension. Institute for Food and Agricultural Sciences
- Benson G (1999) Tandem repeats finder: a program to analyze DNA sequences. *Nucleic Acids Res* 27:573–580
- Bradshaw BDJ, Stettler RF (1995) Molecular genetics of growth and development in *Populus*. IV. Mapping QTLs with large effects on

- growth, form and phenology traits in a forest tree. *Genetics* 139:963–973
- Celton J-M, Tustin DS, Chagne D, Gardiner SE (2009) Construction of a dense genetic linkage map for apple rootstocks using SSRs developed from *Malus* ESTs and *Pyrus* genomic sequences. *Tree Genet Genom* 5:93–107
- Conner PJ, Brown SK, Weeden NF (1998) Molecular-marker analysis of quantitative traits for growth and development in juvenile apple trees. *Theor Appl Genet* 96:1027–1035
- Cook NC, Jacobs G (2000) Progression of apple (*Malus x domestica* Borkh.) bud dormancy in two mild winter climates. *J Hort Sci Bio* 75:233–236
- Costa F, Van de Weg WE, Stella S, Dondini L, Pratesi D, Musacchi S, Sansavini S (2008) Map position and functional allelic diversity of *Md-Exp7*, a new putative expansin gene associated with fruit softening in apple (*Malus x domestica* Borkh.) and pear (*Pyrus communis*). *Tree Genet Genom* 4:575–586
- Doyle JJ, Doyle JL (1990) Isolation of plant DNA from fresh tissue. *Focus* 12:13–15
- Falconer DS, Mackay TFC (1996) Introduction to quantitative genetics, 4th edn. Pearson, England
- Gardiner S, Bus VGM, Rusholme RL, Chagne D, Rikkerink EHA (2007) Apple. In: Kole C (ed) *Genome mapping and molecular breeding in plants. Fruits and Nuts*, vol. 4. Springer, Berlin, pp 1–62
- Guilford P, Prakash S, Zhu JM, Rikkerink E, Gardiner S, Bassett H, Forster R (1997) Microsatellites in *Malus x domestica* (apple): abundance, polymorphism and cultivar identification. *Theor Appl Genet* 94:249–254
- Hemmat M, Weeden NF, Brown SK (2003) Mapping and evaluation of *Malus x domestica* microsatellites in apple and pear. *J Am Soc Hortic Sci* 128:515–520
- Hemmat M, Weeden NF, Conner PJ, Brown SK (1997) A DNA marker for columnar growth habit in apple contains a simple sequence repeat. *J Am Soc Hortic Sci* 122:347–349
- Kellerhals M, Patocchi A, Duffy B, Frey J (2008) Presented at the Ecofruit—13th International conference on cultivation technique and phytopathological problems in organic fruit growing. Weinsberg, Germany
- Kenis K, Keulemans J (2004) QTL analysis of growth characteristics in apple. *Eucarpia symposium on fruit breeding and genetics, Angers, France*, vol 663. *Acta Horticulturae* 663:369–374
- Khan AA (1997) Quantification of plant dormancy: introduction to the workshop. *Hort Science* 32:608–614
- Kim CS, Lee CH, Shin JS, Chung YS, Hyung NI (1997) A simple and rapid method for isolation of high quality genomic DNA from fruit trees and conifers using PVP. *Nucleic Acids Res* 25:1085–1086
- Labuschagné IF, Louw JH, Schmidt K, Sadie A (2002a) Genetic variation in chilling requirement in apple progeny. *J Am Soc Hortic Sci* 127:663–672
- Labuschagné IF, Louw JH, Schmidt K, Sadie A (2002b) Genotypic variation in prolonged dormancy symptoms in apple families. *Hort Science* 37:157–163
- Labuschagné IF, Louw JH, Schmidt K, Sadie A (2003) Budbreak number in apple seedlings as selection criterion for improved adaptability to mild winter climates. *Hort Science* 38:1186–1190
- Lang GA, Early JD, Arroyave NJ, Darnell RL, Martin GC, Stutte GW (1985) Dormancy: towards a reduced, universal terminology. *Hort Science* 20:809–812
- Liebhart R, Gianfranceschi L, Koller B, Ryder CD, Tarchini R, Van de Weg E, Gessler C (2002) Development and characterisation of 140 new microsatellites in apple (*Malus x domestica* Borkh.). *Mol Breeding* 10:217–241
- Linsley-Noakes GC, Allan P, Matthee G (1994) Modification of rest completion prediction models for improved accuracy in South African stone fruit orchards. *J South African Hort Sci* 4:13–15
- Maliepaard C, Alston FH, Van Arkel G, Brown LM, Chevreau E, Dunemann F, Evans KM, Gardiner S, Guilford P, Vand Heusden AW, Janse J, Laurens F, Lynn JR, Manganaris AG, Den Nijs APM, Periam N, Roche REP, Ryder C, Sansavini S, Schmidt H, Tartarini S, Verhaegh JJ, Vrieling-van Ginkel M, King GJ (1998) Aligning male and female linkage maps of apple (*Malus pumila* Mill.) using multi-allelic markers. *Theor Appl Genet* 97:60–73
- Naik S, Hampson C, Gasic K, Bakkeren G, Korban SS (2006) Development and linkage mapping of E-STS and RGA markers for functional gene homologues in apple. *Genome* 49:959–968
- Newcomb RD, Crowhurst RN, Gleave AP, Rikkerink EHA, Allan AC, Beuning LL, Bowen JH, Gera E, Jamieson KR, Janssen BJ, Laing WA, McArtney S, Nain B, Ross GS, Snowden KC, Souleyre EJJ, Walton EF, Yauk Y-K (2006) Analysis of expressed sequence tags from apple (*Malus x domestica*). *Plant Physiol* 141:1–20
- Patocchi A, Fernandez-Fernandez F, Evans K, Gobbin D, Rezzonico F, Boudichevskaia A, Dunemann F, Stankiewicz-Kosyl M, Mathis-Jeanneteau F, Durel CE, Gianfranceschi L, Costa F, Toller C, Cova V, Mott D, Komjane M, Barbaro E, Kodde L, Rikkerink E, Gessler C, Van de Weg WE (2009) Development and test of 21 multiplex PCRs composed of SSRs spanning most of the apple genome. *Tree Genet Genom* 5:211–223
- Segura V, Denance C, Durel CE, Costes E (2007) Wide range QTL analysis for complex architectural traits in a 1-year-old apple progeny. *Genome* 50:159–171
- Silfverberg-Dilworth E, Matasci CL, Van de Weg WE, Van Kaaunen MPW, Walser M, Kodde LP, Soglio V, Gianfranceschi L, Durel CE, Costa F, Yamamoto T, Koller B, Gessler C, Patocchi A (2006) Microsatellite markers spanning the apple (*Malus x domestica* Borkh.) genome. *Tree Genet Genom* 2:202–224
- Tartarini S, Sansavini S (2003) The use of molecular markers in pome fruit breeding. *Acta Hort* 622:129–132
- Tartarini S, Sansavini S, Vinatzer BA, Gennari F, Domizi C (2000) Efficiency of marker assisted selection (MAS) for the *Vf* scab resistance gene. *Acta Hort* 538:549–553
- van Dyk MM, Rees DJG (2009) Bin Mapping of EST-SSRs in Apple (*Malus x domestica* Borkh.). *Acta Hort* 814:681–687
- van Dyk MM, Labuschagné IF, Rees DJG (2009) Genetic linkage map construction and identification of qtls affecting time of initial vegetative budbreak in apple (*Malus x domestica* Borkh.). *Acta Hort* 814:586–590
- Van Ooijen JW (2004) MapQTL®5, software for the mapping of quantitative trait loci in experimental populations. Kyazma B. V., Wageningen, Netherlands
- Van Ooijen JW (2006) JoinMap®4. Software for the calculation of genetic linkage maps in experimental populations. Kyazma B. V., Wageningen, Netherlands
- Yamamoto T, Kimura T, Sawamura Y, Manabe T, Kotobuki K, Hayashi T, Ban Y, Matsuta N (2002a) Simple sequence repeats for genetic analysis in pear. *Euphytica* 124:129–137
- Yamamoto T, Kimura T, Shoda M, Ban Y, Hayashi T, Matsuta N (2002b) Development of microsatellite markers in the Japanese pear (*Pyrus pyrifolia* Nakai). *Mol Ecol Notes* 2:14–16
- Young ND (1996) QTL mapping and quantitative disease resistance in plants. *Annu Rev Phytopathol* 34:479–501