




Maturation effect on physical capacities and anabolic hormones in under-16 elite footballers: a cross-sectional study

Ebrahim Eskandarifard¹ · Rui Silva² · Hadi Nobari^{3,4}  · Filipe Manuel Clemente^{2,5} · Jorge Pérez-Gómez³ · António José Figueiredo⁶

Received: 8 March 2021 / Accepted: 11 June 2021 / Published online: 20 December 2021
© The Author(s) 2021

Abstract

Purpose The purposes of this study were to describe the fitness and hormonal levels according to playing time (PT) (i.e., PT during season less (PT1) or more (PT2) than 50% of the total time) and maturation level (ML) (i.e., normal (ML1) and early maturity levels (ML2)), and to analyze the differences between groups for the measures of aerobic capacity, anaerobic power, power performance, and hormonal concentrations.

Methods Twenty-four youth footballers of a U16 team participated in this study. Anthropometric measures, maturity status, growth hormone, insulin-like growth factor (IGF-1), maximal oxygen uptake, fatigue index, and countermovement jump were collected.

Results Significant differences were found between both PT and ML groups for maturational status, aerobic capacity, power performance, and IGF1 concentrations. The interaction of PT and ML revealed significant differences for maturity offset and power performance. When using the skeletal age as a covariant, the previously significant differences found were reduced only to the fatigue index measure.

Conclusions The response variables analyzed in the present study seem to be influenced by PT and ML. This must be considered when planning training, and coaches must be sensible to these effects as they may assume a preponderant role in PT.

Keywords Soccer · Motor development · Performance · Growth

Introduction

Football (soccer) has evolved into a more physically, technical, and tactical demanding sport over the past few years [1]. Indeed, this fact has been corroborated by studies using data collected via modern devices, such as global positioning systems, enabling to reveal that elite football teams are dealing with greater match demands [2]. As such, this information can be used by coaches and scouts for talent identification and player selection [3]. Considering the approaches for talent identification and player selection, it has been documented that coaches tend to use unidimensional settings, where the levels of strength, speed, and morphological characteristics of young football athletes are prioritized [4].

During youth, especially from 10 to 15 years, it is expected that children present significant variations in maturation status [5, 6]. In fact, great differences in maturation status (biological age) are commonly seen within children of the same age, based on the date of birth (chronological age) [7]. Football academies usually organize youth teams

✉ Hadi Nobari
hadi.nobari1@gmail.com

✉ Jorge Pérez-Gómez
jorgepg100@gmail.com

¹ Department of Exercise Physiology, Faculty of Sport Sciences, University of Isfahan, 81746-7344 Isfahan, Iran

² Escola Superior Desporto e Lazer, Instituto Politécnico de Viana do Castelo, Rua Escola Industrial e Comercial de Nun'Álvares, 4900-347 Viana do Castelo, Portugal

³ HEME Research Group, Faculty of Sport Sciences, University of Extremadura, 10003 Cáceres, Spain

⁴ Sports Scientist, Sepahan Football Club, Isfahan, Iran

⁵ Delegação da Covilhã, Instituto de Telecomunicações, 1049-001 Lisboa, Portugal

⁶ Faculty of Sport Sciences and Physical Education, University of Coimbra, Coimbra, Portugal

by grouping athletes according to their chronological age, and consequently, these practices result in greater within-subjects variation in maturity status [8]. From a practical perspective, late-maturing footballers must deal with an unfair fight, considering that early maturing players are commonly taller, heavier, and outperform their late-maturing colleagues, as well as being preferred by coaches for selection [9]. For those reasons, it is of paramount importance to estimate the maturation level (ML) of young footballers. Although this can be done through the Fels method [10], which is the most valid and reliable method to calculate the skeletal age, it requires equipment that may not be reachable for some sport contexts. However, coaches have simple and inexpensive tools to assess the maturation status of athletes at their disposal. It is possible to predict how far from the peak height velocity (PHV) children are, or if they already reached it, through anthropometric measures and a simple calculation [11].

Aerobic capacity has been considered an important performance determinant of adult elite footballers, where values of maximal oxygen consumption (VO_{2max}) can go up to $69 \text{ ml kg}^{-1} \text{ min}^{-1}$ [12, 13]. And it is commonly assessed via incremental tests to exhaustion, such as the yo-yo intermittent recovery (YYIR) levels 1 and 2 [14]. On the other hand, adolescent footballers have demonstrated an average level of VO_{2max} of $48 \text{ ml kg}^{-1} \text{ min}^{-1}$, which can be slightly different depending on the level of competition (elite or non-elite) and the maturity of the players [15–17]. In the same line, anaerobic power has an important role in football due to the repeated short high-intensity actions that are required during training and matches [17, 18]. This highlights the importance of assessing the repeated sprint ability (RSA) and their related fatiguing effects (fatigue index), in adolescent footballers [19, 20]. Lower limb explosive power is also one of the key-determinants for achieving success in elite football performance, and it is usually assessed using vertical jumps such as the countermovement jump (CMJ) [15]. Given the abovementioned physical determinants, it is fundamental to highlight the between-player variations and status of those physical aspects in critical stages of human maturation, such as puberty. In fact, it has been documented that early maturing players have greater physical characteristics and capabilities than late maturers [7]. Also, considering the typical hormonal changes during puberty, associations between the activity of some hormones, such as growth hormone (GH) and insulin-like growth factor (IGF1) with physical performance have been made [21, 22]. Therefore, coaches tend to overlook late maturing players, having a strong preference for selecting early maturing players as they augment the chances to win matches [23].

Although it has been previously documented that in lower competitive levels (U9-U10), early maturing players revealed greater playing time (PT) than mature late, in

higher competitive levels (> U11) this was not clear [24]. Notwithstanding the attempt to reveal the influence of ML on PT, there is still limited evidence regarding these patterns [24]. Also, to the best of our knowledge, there is a lack of evidence supporting the influence of PT and maturational levels on physical and physiological levels between early and normal maturing players.

As football sessions and other training stimuli may influence the concentration levels of GH and IGF1 during adolescence, and knowing that chronic training load exposure may elicit increased GH concentrations [22, 25], it is hypothesized that players with more PT may present greater physical and physiological levels than players with lesser PT. Given the fact that male athletes advanced in maturity levels possess greater strength capabilities, it is expected that they present greater jump performance. While, it is also hypothesized that normal maturing players have the better aerobic capacity, as they seem to perform better in intermittent tasks [7], such as the YYIR1 test.

Considering that in a youth football team there are great development differences between-players in the same team [4], it is expected that this factor is manifested by greater physical qualities differences in the same team. Given that, it is of utmost importance to analyze the within-team maturational status for determining players' physical capacity. Understanding the influence that PT and maturational levels have on the physical and physiological development of early and normal mature athletes may be an optimal strategy to sensitize coaches to plan training considering these factors. From a practical perspective, this is of paramount importance to make football practice sessions fair and motivating for youth. For these reasons, the purposes of this study were: (i) to describe the fitness status, and hormonal indices according to maturation and competitive level (PT); and (ii) to analyze the comparisons between early and normal mature players (within each competitive level) for the measures of minutes playing, VO_{2max} , seven repeated sprint test (7RST), fatigue index, CMJ, GH, and IGF-1 level.

Materials and methods

Participants

Twenty-four youth footballers (Mean \pm standard deviation (SD)); chronological age 15.6 ± 0.2 years; height 176.4 ± 5.3 cm; body mass 60.6 ± 8.3 kg; skeletal age 16.6 ± 0.8 years; maturity offset 2.0 ± 0.4 years; minutes of playing 1383.4 ± 260.7 min) participated in this study. Players were playing in the Regional Premier League of U16. Positions and number of players in each position were: 2 goalkeepers, 9 defenders, 5 central midfielders, 4 wingers, and 4 attackers. The inclusion criteria were: (1) at

least playing football for 3 years; (2) they banned for using any kind of food supplement which could affect on growth; and (3) do not participate in any extra training outside this study. The exclusion criteria were: (1) participating in less than 80% of training and games (formal and no formal); (2) absence in any examination or tests of this study; (3) late maturity status. This study had an Ethics code from the University of Isfahan, also we followed all ethics rights recommendations of the Helsinki Declaration. Before starting the study all players and parents were aware of the benefits and risks of this study and signed a permission letter.

Experimental approach to the problem

This study was performed as a quasi-experimental and cohort research which was conducted on a cross-sectional basis, the results will be practical. The participants were divided into two groups in terms of PT. The first group played less than 50% of the season games (PT1), and the second group played more than 50% of the season games (PT2). Furthermore, other grouping was used based on the maturity level of the players, which were considered as two categories of normal maturity (ML1) and early maturity (ML2) [7]. For grouping players, we used the subtraction of skeletal age and chronological age, if it was more than + 1 we called them early mature, and if it was between - 1 and + 1 we called them on time. All minutes of formal and informal games were included in the variable of minutes playing by the players. At the end of the season after three days’ rest the assessment was measured during four days and in the morning, t in the first day, we did blood sampling, X-Ray, and anthropometric measurement for identifying the maturity status of the player. Furthermore, physical fitness tests used in this study were CMJ (second day), the 7RST (third day), and the YYIR1 Test (fourth days). All of these tests were performed in the morning, under the same weather conditions (21–23 °C temperature and 50% humidity) [26]. Timetable of this study is in Table 1. Participants completed an average of 188–200 training sessions during the season for 8 months, including 43 matches (21 official + 12 non-officials) and no long-term injury reports (7 players’ minor sports injuries) during the course.

Measurements

The tests in this study were performed in 4 days, at the same temperature (21–23 °C and 50% humidity) [26], participants stopped physical activity 3 days before the tests.

Blood sampling On the first day, blood samples were taken from the vein of the front arm, after 12 h of fasting. To minimize measurement error, all blood samples were taken at 8 a.m. by a skilled expert. To measure serum IGF-1 and GH levels, the Chemiluminescence method and IMMULITE system were used. Blood samples were centrifuged and kept at - 70°C until analyzing.

Skeletal age The EOS Imaging device was used for posterior-anterior imaging of the hand-wrist graph, which is a new device in radiography, where the radiation dose is much lower than previous devices [27–29], and it has also better picture quality [30, 31]. Two-dimensional graph of the EOS Imaging system was used for the determination of skeletal age. This assessment was done by an experienced person, he used the Fels method. The Fels method is one of the three methods for assessing skeletal age from radiography, which is known as the most valid method [10].

Anthropometric measurements Finally, on the first day, anthropometric characteristics were measured [32], the participants stood without shoes, with their shoulders upright and in a normal position, then their standing height was measured. The sitting height was calculated when they sat on a 50-cm bench, with their shoulders and buttocks directly attached to the stadiometer. To measure standing and sitting height, the German SECA device was used with an accuracy of 0.05 m, and to measure the weight of the German SECA 813 device with an accuracy of 0.1 kg. On the other hand, anthropometric measurements were used to maturity offset and age at PHV [11]: Maturity offset = $-9.236 + 0.0002708 (\text{leg length} \times \text{sitting height}) - 0.001663 (\text{age} \times \text{leg length}) + 0.007216 (\text{age} \times \text{sitting height}) + 0.02292 (\text{Weight by Height ratio})$, $R = 0.94$, $R^2 = 0.891$, and $SEE = 0.592$) and for leg length = $\text{Standing Height (cm)} - \text{Sitting height (cm)}$ [11].

Table 1 Control and assessment period during season

Year	2018					2019				Total
	August	September	October	November	December	January	February	March	End of March	
Month										
Phase	First preparation phase (8 week)		Regional games			Best of south region	Second preparation phase (6 week)	Best of Iran (National)	Assessment	
Official games no			4	4	4	5		4	4 days	21
Non official games no	2	4					6			12

Physiological tests On the second day of the test, after 15 min of a standardized warm-up, which included jogging, stretching, explosive exercises, similar to the test, under the supervision of a conditioning trainer, players performed the CMJ test without swinging the hand, with the hand on the lateral part of the above pelvis for measurements of lower body explosive power [33]. Following the 90 degrees of flexion of the knee, they performed their jump with a maximum power by command of the tester. Two jumps were performed with 5 min' recovery between them, then the best performance was recorded. An electronic pad, the Newest Power Timer company made in Finland was used for testing.

On the third day, the players participated in the 7RST, which included seven 34.5-m sprints with a 25-s recovery time between them, it assessed anaerobic power [34]. The protocol was started, with a warm-up that included jogging, stretching, and short sprints. After a 15-min warm-up, players were instructed how they should have performed. For identifying the fatigue index we subtracted the worse performance to best performance [35]. To assess the time of each of the players' performances we used Newest Power Timer device made by Finland company and the 25 s, recovery time was calculated by a stopwatch. The player was notified at the end of each recovery period for the next run, and players were encouraged during the performance.

On the last day of measurement, the aerobic capacity of the players was assessed by the YYIR1 test. The test started with 40 m running where players run a 20-m back and forth, with 10 m of walking recovery, which is a 5-m back and forth. In this test, with the sound of each beep, the players must be at the exact point of the test, and each level of the test increased, 0.5 km/h progressively. Players continued the test until the players could not reach the exact point two times, the last level that the player was able to perform was registered as his record. The following formula was used to determine the VO_{2max} : $VO_{2max} \text{ (ml kg}^{-1} \text{ min}^{-1}) = IR1 \text{ distance (m)} \times 0.0084 + 36.4$ [12].

Football training In scientific literature, there is a term called the special sport training, and because this research is in the field of football, so we examined the duration of football training, which indicates the time that the player has been engaged in football from the beginning of his life in months. For this purpose, the history of each player was checked from the archive and database of the club, and after that, for more accuracy and precision, the available results were reviewed again with each player and recorded as "Football training" in "months". The average is in Table 1.

Content of each training session The training program for this age group was as follows: 5 training sessions of 90 min per week, including 10 min of warm-up, 20 min of physical training, 10 min of technical training, 20 min of tactical

training, 25 min of training game (including playing in the small-sided game) and at the end there was a recovery for 5 min. The time of the matches in this age group was 90 min per game, which was held on the official field according to the rules of the Football Federation of the country for this age group. Training goals in the age group include goals in the small-sided game (development of ball possession, ball transition in speed and rapid organization of zonal defense, retreat and recovery), tactical goals (using defensive and offensive principles quickly), technical goals (focus on passing and controls skills, as well as ball control in small and large spaces) and physical fitness goals (development of aerobic power, linear speed, and explosive power) and were applied in the exercises of each session.

Statistical method

Mean and SD of variables were used for descriptive statistics. After examining the normality of the test data (Smirnov Kolmogorov) and homogeneity of variances (Leven test), the Multivariate Analysis of Variance (MANOVA) test, to contrast athletes of early and normal maturity, in both groups of players, PT1 and PT2 were used. Furthermore, Multivariate analyses of covariance (MANCOVA) for controlling contending factor apply two time; first, minutes of playing and second, ML as a covariate, also partial eta-square (ηp^2) were used to show the magnitude effect size (trivial < 0.2 , small ≥ 0.2 ; medium ≥ 0.5 ; large ≥ 0.8 and above) all variables were examined at a significance level of 0.05 using the SPSS 22.

Results

Table 2 presents the descriptive characteristics of this study sample in Mean \pm SD in both ML and PT groups. We summarized some variables like total season playing time: 1383.4 ± 260.7 min; height: 176.4 ± 5.3 cm; VO_{2max} : 49.5 ± 3.5 ml.kg⁻¹.min⁻¹; IGF-1 level: 392.0 ± 107.7 ng/dl; skeletal age: 16.6 ± 0.8 , that was used to divided players in the two groups.

The results of MANOVA are presented in Table 3. Significant differences between PT1 and PT2 were found for minutes of playing ($F = 45.1$, $p < 0.01$, $\eta p^2 = 0.7$), age at PHV ($F = 8.1$, $p < 0.01$, $\eta p^2 = 0.3$), maturity offset ($F = 11.2$, $p < 0.01$, $\eta p^2 = 0.4$), standing height ($F = 6.0$, $p < 0.01$, $\eta p^2 = 0.2$), sitting height ($F = 8.5$, $p < 0.01$, $\eta p^2 = 0.3$), VO_{2max} ($F = 9.8$, $p < 0.01$, $\eta p^2 = 0.3$), CMJ ($F = 9.2$, $p < 0.01$, $\eta p^2 = 0.3$) and IGF-1 level ($F = 17.0$, $p < 0.01$, $\eta p^2 = 0.5$). Also, in maturation level groups there were significant differences between early and on time mature players for skeletal age ($F = 29.0$, $p < 0.01$, $\eta p^2 = 0.6$), weight ($F = 7.5$, $p < 0.01$, $\eta p^2 = 0.3$), VO_{2max} ($F = 5.2$, $p < 0.05$, $\eta p^2 = 0.2$),

Table 2 Descriptive statistics of playing time and maturity groups in inter-variability of U16 youth Football players

Variables	Total <i>n</i> = 24	Playing time groups		Maturation level groups	
		PT1 <i>n</i> = 12	PT2 <i>n</i> = 12	ML1 <i>n</i> = 10	ML2 <i>n</i> = 14
Minutes playing (min)	1383.4 ± 260.7	1166.8 ± 170.1	1600 ± 104.5	1509.0 ± 256.6	1293.7 ± 232.2
Chronological age (year)	15.6 ± 0.2	15.66 ± 0.2	15.57 ± 0.3	15.6 ± 0.2	15.6 ± 0.3
Skeletal age (year)	16.6 ± 0.8	17.0 ± 0.8	16.3 ± 0.8	15.9 ± 0.6	17.2 ± 0.5
Football training (month)	79.9 ± 19.5	77.8 ± 19.7	82.0 ± 19.9	80.3 ± 15.2	79.6 ± 22.6
Age at PHV (year)	13.6 ± 0.3	13.4 ± 0.4	13.8 ± 0.1	13.7 ± 0.3	13.6 ± 0.3
Maturity offset (year)	2.0 ± 0.4	2.2 ± 0.4	1.7 ± 0.3	1.9 ± 0.3	2.0 ± 0.5
Standing height (cm)	176.4 ± 5.3	179.2 ± 5.4	173.7 ± 3.8	175.5 ± 4.5	177.1 ± 6.0
Sitting height (cm)	93.1 ± 2.6	94.5 ± 2.8	91.6 ± 1.3	92.8 ± 2.6	93.2 ± 2.7
Weight (kg)	60.6 ± 8.3	63.6 ± 10.1	57.7 ± 4.6	55.7 ± 5.7	64.2 ± 8.1
VO _{2max} (ml kg ⁻¹ min ⁻¹)	49.5 ± 3.5	47.3 ± 1.8	51.7 ± 3.4	51.7 ± 3.2	47.9 ± 2.8
7RST (s)	6.7 ± 0.2	6.7 ± 0.3	6.7 ± 0.2	6.6 ± 0.2	6.8 ± 0.3
Best of 7RST (s)	6.4 ± 0.2	6.5 ± 0.2	6.4 ± 0.3	6.4 ± 0.2	6.5 ± 0.2
Worse of 7RST (s)	7.0 ± 0.3	7.0 ± 0.4	7.0 ± 0.3	6.9 ± 0.3	7.1 ± 0.3
Fatigue Index (s)	0.5 ± 0.3	0.5 ± 0.2	0.6 ± 0.3	0.5 ± 0.2	0.7 ± 0.3
CMJ (cm)	40.8 ± 4.4	38.0 ± 3.9	43.6 ± 3.0	43.2 ± 2.4	39.1 ± 4.8
GH level (ng/dl)	1.6 ± 1.9	1.1 ± 1.6	2.3 ± 2.2	2.1 ± 1.9	1.3 ± 1.9
IGF1 level (ng/dl)	392.0 ± 107.7	317.8 ± 66.4	466.3 ± 88.5	459.0 ± 93.3	343.5 ± 91.8

PT1 group of player who play less than 50 percent of total season, PT2 group of player who play more than 50 percent of total season, ML1 the group of player who were on normal maturation, ML2 the group of player who were early maturation, PHV Peak Height Velocity, VO_{2max} maximal oxygen uptake, 7RST 7 Repeated Sprint Test, CMJ Countermovement Jump, GH Growth Hormone, IGF1 Insulin-like Growth factor-1

Table 3 Results of MANOVA to test the effect of Playing Time and maturity groups in inter-variability of U16 youth Football players

Variables	Effects of playing time			Effects of maturation level			Effects of playing × maturation		
	<i>F</i>	<i>p</i>	η ²	<i>F</i>	<i>p</i>	η ²	<i>F</i>	<i>p</i>	η ²
Minutes playing (min)	45.1	<0.001*	0.7	1.2	0.283	0.1	0.7	0.413	<0.001
Chronological age (year)	0.8	0.382	<0.001	0.1	0.330	0.1	1.8	0.199	0.1
Skeletal age (year)	0.5	0.493	<0.001	29.0	<0.001*	0.6	0.5	0.476	<0.001
Football training (month)	0.4	0.516	<0.001	0.0	0.843	<0.001	0.7	0.430	<0.001
Age at PHV (year)	8.1	0.010*	0.3	0.3	0.590	<0.001	1.5	0.242	0.1
Maturity offset (year)	11.2	0.003*	0.4	<0.001	0.942	<0.001	4.7	0.043*	0.2
Standing height (cm)	6.0	0.024*	0.2	<0.001	0.972	<0.001	1.0	0.327	0.1
Sitting height (cm)	8.5	0.009*	0.3	0.2	0.639	0.0	1.9	0.180	0.1
Weight (kg)	0.5	0.483	<0.001	7.5	0.012*	0.3	4.4	0.049*	0.2
VO _{2max} (ml kg ⁻¹ min ⁻¹)	9.8	0.005*	0.3	5.2	0.034*	0.2	<0.001	0.927	<0.001
7RST (s)	0.1	0.713	<0.001	2.5	0.129	0.1	0.8	0.370	<0.001
Best of 7RST (s)	0.1	0.779	<0.001	1.2	0.275	0.1	0.9	0.362	<0.001
Worse of 7RST (s)	1.7	0.206	0.1	2.1	0.100	0.1	0.9	0.355	<0.001
Fatigue Index (s)	1.6	0.220	0.1	0.6	0.454	<0.001	<0.001	0.858	<0.001
CMJ (cm)	9.2	0.007*	0.3	4.4	0.048*	0.2	4.5	0.046*	0.2
GH level (ng/dl)	1.6	0.223	0.1	0.2	0.647	<0.001	<0.001	0.845	<0.001
IGF1 level (ng/dl)	17.0	0.001*	0.5	4.9	0.039*	0.2	0.1	0.333	0.5

PHV Peak Height Velocity, VO_{2max} maximal oxygen uptake, 7RST 7 Repeated Sprint Test, CMJ Counter Movement Jump, GH Growth Hormone, IGF1 Insulin-like Growth Factor-1

*Significant difference at (*p* ≤ 0.05)

CMJ ($F = 4.4, p < 0.05, \eta p^2 = 0.2$) and IGF-1 level ($F = 4.9, p < 0.05, \eta p^2 = 0.2$). The interaction of groups of PT and their maturation ($PT \times ML$) showed that maturity offset ($F = 4.7, p < 0.05, \eta p^2 = 0.2$), weight ($F = 4.4, p < 0.05, \eta p^2 = 0.2$) and CMJ ($F = 4.5, p < 0.05, \eta p^2 = 0.2$) were significantly different.

A MANCOVA was performed to test the effect of PT and ML in inter-variability of U16 youth footballers, and the results are shown in Table 4. When controlling ML, skeletal age as a covariate, just the parameter of fatigue index ($F = 47, p < 0.05, \eta p^2 = 0.2$) was significant.

Discussion

The purposes of the present study were to describe the fitness and hormonal status according to ML and PT, and to analyze the differences between ML1 and ML2 within PT1 and PT2 groups, for the measures of aerobic capacity, anaerobic power, power performance, and hormonal concentrations. Results revealed significant differences between both PT and ML groups for maturational status, aerobic capacity, power performance and for IGF1 concentrations. Also, the interaction of PT and ML revealed significant differences for maturity offset and power performance. However, after controlling the ML, using the skeletal age as a covariant, the latter significant differences were reduced only to the fatigue index measure.

The first aim of the present study was to describe the fitness and hormonal levels of both PT1, PT2, ML1, and ML2 groups. Results revealed the greater overall physical performance and greater hormonal mean values for PT2 and ML12? groups, than PT1 and ML21? groups. The

characterization of fitness and hormonal measures of young footballers players have previously been made [21, 36, 37]. For instance, in a recent study, it was revealed that footballers had greater changes during 2 seasons in physical and hormonal parameters than non-athletic adolescents from the same chronological age as in the present study [21]. Also, the same study revealed that GH concentrations of 2.2 ng/dl were similar to our study. In contrast, the IGF1 mean concentration was lower in our study compared to that found in the study of Hammami et al. [21], which showed IGF1 concentrations of 870 ng/dl. In fact, IGF1 values of our sample were similar to the non-athletic population of the above-mentioned study. However, those differences may be due to the divergent methodologies used, as well as the type, duration, and frequency of training that was not controlled in the present study.

Regarding the second aim of the present study, a MANOVA was executed to test the effects of PT and ML on between-group differences of the analyzed response variables. Results revealed that PT influenced the VO_{2max} , CMJ, and IGF1 measures, with the PT2 group presenting the greatest values. Other studies revealed that long exposures to football training results in enhanced physical capacities [21, 38]. Indeed, Nobari et al. [16] showed that accumulated training loads influenced the significant differences found at different assessment timings for aerobic capacity, anaerobic power, fatigue index, and maximal speed. Also, despite acute training doses may cause a suppression of IGF1 levels, it seems that chronic exposures to training elicit increases in both GH and IGF1 hormones, that have a great impact on growth and physical performance [22]. For those reasons, it is suggested that players with more PT during a full season show greater physical performance and anabolic hormonal

Table 4 Results of MANCOVA to control playing time and maturation level of U16 youth football players

Variables	Playing time as covariate			Skeletal age as covariate		
	F	p	ηp^2	F	p	ηp^2
Football training (month)	0.9	0.355	<0.001	0.1	0.831	<0.001
Standing height (cm)	0.7	0.414	<0.001	0.5	0.489	<0.001
Sitting height (cm)	0.4	0.551	<0.001	0.1	0.787	<0.001
Weight (kg)	4.4	0.051	0.2	0.9	0.358	0.1
VO_{2max} (ml kg ⁻¹ min ⁻¹)	1.8	0.199	0.1	0.6	0.439	<0.001
7RST (s)	0.5	0.494	<0.001	0.2	0.659	<0.001
Best of 7RST (s)	1.1	0.315	0.1	1.8	0.200	0.1
Worse of 7RST (s)	0.4	0.548	<0.001	0.7	0.425	<0.001
Fatigue Index (s)	<0.001	0.987	<0.001	4.7	0.043*	0.2
CMJ (cm)	<0.001	0.855	<0.001	0.9	0.356	0.1
GH level (ng/dl)	1.3	0.263	0.1	<0.001	0.962	<0.001
IGF1 level (ng/dl)	1.5	0.232	0.1	1.1	0.317	0.1

PHV Peak Height Velocity, VO_{2max} maximal oxygen uptake, 7RST 7 Repeated Sprint Test, CMJ Counter Movement Jump, GH Growth Hormone, IGF1 Insulin-like Growth Factor-1

*Significant difference at ($p \leq 0.05$)

concentrations. Given that, special attention should be given to players with less PT, thus, it would be of paramount importance for coaches to promote a between-player balanced training stimulus.

Although there is a lack of studies investigating the effects of PT on the analyzed dependent variables in the present study, other studies attempted to test the influence of physical and hormonal measures, and maturational status on PT [24, 39]. In fact, Deprez et al. [39] showed that jump performance influenced approximately 17% of the variance in future PT. Also, Goto et al. [24] revealed that early mature players had more PT than their late-maturing counterparts in lower age categories. This was not observed in the present study, as the ML2 group, early maturing players showed slightly lesser minutes of play, but not representing significant differences. However, our sample is from higher age categories (U-16), and the above-mentioned study [24], highlighted that at higher age categories the differences between early and late maturing players were not clear. Given that, more studies should be conducted for ensuring greater information about these types of influences of PT on physical, physiological, and hormonal measures, and vice-versa.

Furthermore, ML showed effects on player's weight, VO_{2max} , CMJ, and IGF1 measures, being the ML1 group the lighter, and with the greatest values of VO_{2max} , CMJ, and IGF1. Thus, after removing the influence of skeletal age, used as a covariate, from the analysis, the significant differences were reduced to fatigue index only. Indeed it has been documented that the ML of footballers are strongly associated with greater physical performance [7, 40]. Greater development of aerobic capacity is shown at the PHV, and greater maximal aerobic power, strength, and power development might occur slightly later [38, 41, 42]. Also, as children advance in maturity status during puberty, it is expected a greater activity of operating hormones such as the GH and IGF1 concentrations, which facilitates human growth [22, 37]. Thus, when footballers are at critical stages of puberty, it is expected that they become more heavy and taller, confirming our results considering the effects of ML1 on player's weight. Coaches should be aware of these effects as they may assume a preponderant role when considering the selection and PT.

The present study has some limitations. One of the main limitations is the small size of the sample, which include only one U16 football team. However, this is a common limitation considering elite sports contexts. Another limitation is related to the fact that we did not consider any external load and heart rate-based measures [43–45]. It would be of interest that future studies focused on the influence that PT and ML might have on different heart rate and external variables. Despite those limitations, the present study showed that both PT and maturation status may influence the level

of anthropometric measures, aerobic capacity, power performance, and IGF1 concentrations. Given that, coaches must provide balanced PT opportunities within a youth team, and individualize training to promote greater physical, physiological and hormonal adaptations to the whole team players.

Conclusions

In short, significant differences between both PT and ML groups for maturational status, aerobic capacity, power performance, and for IGF1 concentrations were found. PT and ML together revealed significant differences for maturity offset and power performance. Considering the present study findings, it is suggested that the above-mentioned measures seem to be influenced by PT and ML, as after controlling the skeletal age as a covariant, the latter significant differences were reduced only to the fatigue index measure.

Funding Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethics approval This study had an Ethics code from the University of Isfahan, also we followed all ethics rights recommendations of the Helsinki Declaration. Before starting the study all players and parents were aware of the benefits and risks of this study and signed a permission letter (ie., consent form).

Informed consent Informed consent was obtained from all subject and their parents.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Sarmento H, Marcelino R, Anguera MT, Campaniço J, Matos N, Leitão JC (2014) Match analysis in football: a systematic review. *J Sports Sci* 32(20):1831–43. <http://www.ncbi.nlm.nih.gov/pub-med/24787442>

2. Dellal A, Lago-Peñas C, Rey E, Chamari K, Orhant E (2015) The effects of a congested fixture period on physical performance, technical activity and injury rate during matches in a professional soccer team. *Br J Sports Med* 49(6):390–394. <https://doi.org/10.1136/bjsports-2012-091290>
3. Barnes C, Archer D, Hogg B, Bush M, Bradley P (2014) The evolution of physical and technical performance parameters in the English premier league. *Int J Sports Med* 35(13):1095–1100. <https://doi.org/10.1055/s-0034-1375695>
4. Sarmiento H, Anguera MT, Pereira A, Araújo D (2018) Talent identification and development in male football: a systematic review. *Sport Med* 48(4):907–931. <https://doi.org/10.1007/s40279-017-0851-7>
5. Till K, Cogley S, O'Hara J, Chapman C, Cooke C (2013) A longitudinal evaluation of anthropometric and fitness characteristics in junior rugby league players considering playing position and selection level. *J Sci Med Sport* 16(5):438–443
6. Nobari H, Aquino R, Clemente FM, Khalafi M, Adsuar JC, Pérez-Gómez J (2020) Description of acute and chronic load, training monotony and strain over a season and its relationships with well-being status: a study in elite under-16 soccer players. *Physiol Behav* 225:113117. <https://linkinghub.elsevier.com/retrieve/pii/S0031938420304315>
7. Malina RM, Rogol AD, Cumming SP, Coelho-E-Silva MJ, Figueiredo AJ (2015) Biological maturation of youth athletes: assessment and implications. *Br J Sports Med* 49(13):852–859
8. Figueiredo AJ, Coelho-E-Silva MJ, Sarmiento H, Moya J, Malina RM (2020) Adolescent characteristics of youth soccer players: do they vary with playing status in young adulthood? *Res Sport Med* 28(1):72–83. <https://doi.org/10.1080/15438627.2019.1586704>
9. Vaeyens R, Lenoir M, Williams AM, Philippaerts RM (2008) Talent identification and development programmes in sport: current models and future directions. *Sport Med* 38(9):703–714
10. Roche AF, Thissen D, Chumlea W (1988) Assessing the skeletal maturity of the hand-wrist: fels method. Thomas
11. Mirwald RL, Baxter-Jones ADG, Bailey DA, Beunen GP (2002) An assessment of maturity from anthropometric measurements. *Med Sci Sport Exerc* 34(4):689–694
12. Bangsbo J, Iaiia FM, Krustup P (2008) The Yo-Yo intermittent recovery test. *Sport Med* 38(1):37–51. <https://doi.org/10.2165/00007256-200838010-00004>
13. Tønnessen E, Hem E, Leirstein S, Haugen T, Seiler S (2013) Maximal aerobic power characteristics of male professional soccer players, 1989–2012. *Int J Sports Physiol Perform* 8(3):323–329
14. Schmitz B, Pfeifer C, Kreitz K, Borowski M, Faldum A, Brand SM (2018) The Yo-Yo intermittent tests: a systematic review and structured compendium of test results. *Front Physiol* 9(JUL):1–16
15. Slimani M, Nikolaidis PT (2018) Anthropometric and physiological characteristics of male soccer players according to their competitive level, playing position and age group: a systematic review. *J Sports Med Phys Fitness* 2019:141–163
16. Nobari H, Silva AF, Clemente FM, Siahkhouhian M (2020) Analysis of fitness status variations of under-16 soccer players over a season and their relationships with maturational status and training load. *Front Physiol*. <https://doi.org/10.3389/fphys.2020.597697>
17. Nobari H, Tubagi Polito LF, Clemente FM, Pérez-Gómez J, Ahmadi M, Garcia-Gordillo MÁ, et al (2020) Relationships between training workload parameters with variations in anaerobic power and change of direction status in elite youth soccer players. *Int J Environ Res Public Health* 17(21):7934. <https://www.mdpi.com/1660-4601/17/21/7934>
18. Nikolaidis P (2011) Anaerobic Power across Adolescence in Soccer Players. *Hum Mov* 12(4):342–347. <http://www.degruyter.com/view/j/humo.2011.12.issue-4/v10038-011-0039-1/v10038-011-0039-1.xml>
19. Buchheit M, Mendez-Villanueva A, Simpson BM, Bourdon PC (2010) Repeated-sprint sequences during youth soccer matches. *Int J Sports Med* 31(10):709–716. <https://doi.org/10.1055/s-0030-1261897>
20. Duarte JP, Coelho-E-Silva MJ, Costa D, Martinho D, Luz LGO, Rebelo-Gonçalves R et al (2019) Repeated sprint ability in youth soccer players: independent and combined effects of relative age and biological maturity. *J Hum Kinet* 67(1):209–221
21. Hammami MA, Ben Abderrahman A, Rhibi F, Nebigh A, Coppalle S, Ravé G et al (2018) Somatotype hormone levels and physical fitness in elite young soccer players over a two-year monitoring period. *J Sport Sci Med* 17(3):455–464
22. Eliakim A, Nemet D (2010) Exercise training, physical fitness and the growth hormone-insulin-like growth factor-1 axis and cytokine balance. *Med Sport Sci* 55:128–140
23. Henriksen K, Stambulova N, Roessler KK (2010) Holistic approach to athletic talent development environments: a successful sailing milieu. *Psychol Sport Exerc* 11(3):212–222
24. Goto H, Morris JG, Nevill ME (2019) Influence of biological maturity on the match performance of 8- to 16-year-old, elite, male, youth soccer players. *J Strength Cond Res* 33(11):3078–84. <http://journals.lww.com/00124278-201911000-00023>
25. Hadzović A, Nakas-Ićindić E, Kucukalić-Selimović E, Avdagić N, Zacririć A (2004) The level of physical activity and the growth hormone (GH) response to acute physical exercise. *Bosn J Basic Med Sci* 4(3):47–49
26. Lippincott W, Wilkins (2013) ACSM's guidelines for exercise testing and prescription. Medicine ACoS
27. Ilharreborde B, Ferrero E, Alison M, Mazda K (2016) EOS micro-dose protocol for the radiological follow-up of adolescent idiopathic scoliosis. *Eur Spine J* 25(2):526–531
28. Luo TD, Stans AA, Schueler BA, Larson AN (2015) Cumulative radiation exposure with EOS imaging compared with standard spine radiographs. *Spine Deform* 3(2):144–150
29. Hui SCN, Pialasse JP, Wong JYH, Lam T, Ng BKW, Cheng JCY et al (2016) Radiation dose of digital radiography (DR) versus micro-dose x-ray (EOS) on patients with adolescent idiopathic scoliosis: 2016 SOSORT- IRSSD “John Sevastic Award” Winner in Imaging Research. *Scoliosis Spinal Disord* 11(1):1–8
30. Wade R, Yang H, McKenna C, Faria R, Gummerson N, Woolacott N (2013) A systematic review of the clinical effectiveness of EOS 2D/3D X-ray imaging system. *Eur Spine J* 22(2):296–304
31. Deschenes S, Charron G, Beaudoin G, Labelle H, Miron M, Parent S (2016) Diagnostic imaging of spinal deformities: reducing patients' radiation dose. *Spine (Phila Pa 1976)* 35(9):989–994
32. Arazi H, Mirzaei B, Nobari H (2015) Anthropometric profile, body composition and somatotyping of national Iranian cross-country runners. *Turkish J Sport Exerc* 17(2):35
33. Bangsbo J, Mohr M (2012) Fitness testing in football. *BangsboSport*
34. Haugen TA, Tønnessen E, Seiler S (2012) Speed and counter-movement-jump characteristics of elite female soccer players, 1995–2010. *Int J Sports Physiol Perform* 7(4):340–349. <http://www.ncbi.nlm.nih.gov/pubmed/22645175>
35. Kaplan T (2010) Examination of repeated sprinting ability and fatigue index of soccer players according to their positions. *J Strength Cond Res Natl Strength Cond Assoc* 24(6):1495–1501
36. Perroni F, Fittipaldi S, Falcioni L, Ghizzoni L, Borriore P, Vetrano M et al (2019) Effect of pre-season training phase on anthropometric, hormonal and fitness parameters in young soccer players. Boullosa D, editor. *PLoS ONE* 14(11):e0225471. <https://doi.org/10.1371/journal.pone.0225471>
37. Nobari H, Kargarfard M, Minasian V, Cholewa JM, Pérez-Gómez J (2021) The effects of 14-week betaine supplementation on

- endocrine markers, body composition and anthropometrics in professional youth soccer players: a double blind, randomized, placebo-controlled trial. *J Int Soc Sports Nutr* 18(1):20. <https://doi.org/10.1186/s12970-021-00417-5>
38. Vääntinen T, Blomqvist M, Nyman K, Häkkinen K (2011) Changes in body composition, hormonal status, and physical fitness in 11-, 13-, and 15-year-old Finnish regional youth soccer players during a two-year follow-up. *J Strength Cond Res* 25(12):3342–3351. <http://journals.lww.com/00124278-201112000-00016>
39. Deprez DN, Franssen J, Lenoir M, Philippaerts RM, Vaeyens R (2015) A retrospective study on anthropometrical, physical fitness, and motor coordination characteristics that influence dropout, contract status, and first-team playing time in high-level soccer players aged eight to eighteen years. *J Strength Cond Res* 29(6):1692–1704. <http://journals.lww.com/00124278-201506000-00031>
40. Lovell R, Franssen J, Ryan R, Massard T, Cross R, Eggers T, et al (2019) Biological maturation and match running performance: a national football (soccer) federation perspective. *J Sci Med Sport* 22(10):1139–1145. <https://linkinghub.elsevier.com/retrieve/pii/S1440244018309848>
41. Philippaerts RM, Vaeyens R, Janssens M, Van Renterghem B, Matthyss D, Craen R et al (2006) The relationship between peak height velocity and physical performance in youth soccer players. *J Sports Sci* 24(3):221–230. <https://doi.org/10.1080/02640410500189371>
42. Nobari H, Alves AR, Clemente FM, Pérez-Gómez J, Clark CCT, Granacher U et al (2021) Associations between variations in accumulated workload and physiological variables in young male soccer players over the course of a season. *Front Physiol* 12:233. <https://doi.org/10.3389/fphys.2021.638180/full>
43. Clemente FM, Silva R, Chen Y-S, Aquino R, Praça GM, Castellano J, et al (2021) Accelerometry-workload indices concerning different levels of participation during congested fixture periods in professional soccer: a pilot study conducted over a full season. *Int J Environ Res Public Health* 18(3):1137. <https://www.mdpi.com/1660-4601/18/3/1137>
44. Nobari H, Praça GM, Clemente FM, Pérez-Gómez J, Carlos Vivas J, Ahmadi M (2020) Comparisons of new body load and metabolic power average workload indices between starters and non-starters: a full-season study in professional soccer players. *Proc Inst Mech Eng Part P J Sport Eng Technol*. <https://doi.org/10.1177/1754337120974873>
45. Nobari H, Oliveira R, Brito JP, Pérez-Gómez J, Clemente FM, Ardigò LP (2021) Comparison of running distance variables and body load in competitions based on their results: a full-season study of professional soccer players. *Int J Environ Res Public Health* 18(4):2077. <https://www.mdpi.com/1660-4601/18/4/2077>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.