



## Correction to: Influence of *Leifsonia* sp. on U(VI) removal efficiency and the Fe–U precipitates by zero-valent iron

Shuibo Xie<sup>1</sup> · Xue Xiao<sup>2</sup> · Wenfa Tan<sup>2,3</sup>  · Junwen Lv<sup>2</sup> · Qinwen Deng<sup>3</sup> · Qi Fang<sup>2</sup>

Published online: 21 January 2020

© Springer-Verlag GmbH Germany, part of Springer Nature 2020

**Correction to: Environmental Science and Pollution Research**

<https://doi.org/10.1007/s11356-019-07306-0>

The original publication of this paper contains a mistake.

The correct figure 7 is presented in this paper.

The original article was corrected.

---

The online version of the original article can be found at <https://doi.org/10.1007/s11356-019-07306-0>


---

✉ Wenfa Tan  
[nhwftan@163.com](mailto:nhwftan@163.com)

<sup>1</sup> Key Discipline Laboratory for National Defense of Biotechnology in Uranium Mining and Hydrometallurgy, University of South China, Hengyang 421001, China

<sup>2</sup> School of Resource Environment and Safety Engineering, University of South China, Hengyang 421001, China

<sup>3</sup> Hengyang Key Laboratory of Soil Pollution Control and Remediation, University of South China, Hengyang 421001, China



**Fig. 7** The characterization of SEM images and EDS analysis of ZVI (**a,d**), *Leifsonia* sp. (**b,e**) and ZVI + *Leifsonia* sp. (**c,f**). U(VI)=10 mg/L, pH=5, t=48 h, ZVI dosage=1.0 g/L. Illustration is an enlarged image of the load