



Associations of bisphenol exposure with the risk of gestational diabetes mellitus: a nested case–control study in Guangxi, China

Peng Tang¹ · Jun Liang¹ · Qian Liao¹ · Huishen Huang¹ · Xiaojing Guo¹ · Mengrui Lin² · Bihu Liu¹ · Bincai Wei¹ · Xiaoyun Zeng¹ · Shun Liu³ · Dongping Huang² · Xiaoqiang Qiu¹

Received: 31 August 2021 / Accepted: 23 November 2021 / Published online: 27 November 2021
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

A growing number of epidemiologic studies have estimated the associations between endocrine-disrupting chemicals and gestational diabetes mellitus (GDM). However, reports on the association between bisphenol A (BPA) substitutes and GDM are limited. This investigation aimed to explore the associations of maternal serum BPA, bisphenol B (BPB), bisphenol F (BPF), bisphenol S (BPS), and tetrabromobisphenol A (TBBPA) with the risk of GDM. A nested case–control study was performed among 500 pregnant women. In conditional logistic regression models, the OR for BPS was significantly increased in the medium exposure groups (OR = 1.77; 95% CI: 1.01, 3.13) compared with the reference group, while BPA (OR: 0.38, 95%CI: 0.29, 0.50) and TBBPA (OR: 0.67, 95%CI: 0.54, 0.85) were negatively associated with the risk of GDM. In the Bayesian kernel machine regression (BKMR) analysis, the joint effect of bisphenols was positively associated with the risk of GDM. BPS showed positively relationship, while BPA and TBBPA showed negatively relationship, respectively. The quantile g-computation revealed a statistically significant and negative joint effect of the five bisphenols on the risk of GDM (OR: 0.57; 95% CI: 0.46, 0.72) with BPA (70.2%), TBBPA (21.3%), and BPB (8.5%) had positive contribution to the overall effect. These findings suggested that BPS had a positive effect on the risk of GDM, while BPA and TBBPA had negative effect on the risk of GDM. Moreover, exposure to the mixture of the five bisphenols was negatively associated with the risk of GDM.

Keywords Bisphenol · Gestational diabetes mellitus · Nested case–control study · Joint effect models

Introduction

Gestational diabetes mellitus (GDM), which is a common complication during pregnancy, is any degree of glucose intolerance with onset or first recognition during pregnancy (American-Diabetes-Association 2013). GDM can lead to short-term and long-term health effects on mothers and their offspring, such as a higher risk of obesity and diabetes later in life (Burlina et al. 2019). Thus, finding the potential risk factors for GDM is important. In addition to traditional risk factors, such as maternal age, obesity, ethnicity, family history of diabetes, and high-carbohydrate diets (Ehrlich et al. 2016; Farahvar et al. 2019), some recent studies have shown that endocrine-disrupting chemicals (EDCs) may play an important role in the incidence and development of GDM (Ehrlich et al. 2016; Filardi et al. 2020; Liu et al. 2018b; Xu et al. 2020).

Bisphenol A (BPA) is an EDC that is widely used in the production of plastic products (Michałowicz 2014) and has

Responsible Editor: Lotfi Aleya

Peng Tang and Jun Liang contributed equally to this work

-
- ✉ Shun Liu
liushun@gxmu.edu.cn
 - ✉ Dongping Huang
dongpinghuang@gxmu.edu.cn
 - ✉ Xiaoqiang Qiu
xqiu9999@163.com

- ¹ Department of Epidemiology and Health Statistics, School of Public Health, Guangxi Medical University, No. 22 Shuangyong Road, Nanning 530021, Guangxi, China
- ² Department of Sanitary Chemistry, School of Public Health, Guangxi Medical University, No. 22 Shuangyong Road, Nanning 530021, Guangxi, China
- ³ Department of Maternal, Child and Adolescent Health, School of Public Health, Guangxi Medical University, No. 22 Shuangyong Road, Nanning 530021, Guangxi, China

been found in various types of human biological samples (Rochester 2013). The effects of BPA on human health have been extensively studied, with its estrogenic activity being one of its best-known outcomes. BPA can interact with estrogen receptors because of its phenolic structure, possibly resulting in estrogenic actions (Konieczna et al. 2015). Some animal studies have suggested that BPA exposure may disrupt glucose homeostasis (Batista et al. 2012; Ding et al. 2014). Previous epidemiological studies have determined a strong association between BPA and type 2 diabetes mellitus (T2DM) (Ranci re et al. 2019; Sun et al. 2014). However, findings related to the relationship between BPA and GDM have been inconsistent (Fisher et al. 2018; Hou et al. 2021; Robledo et al. 2013; Shapiro et al. 2015; Wang et al. 2017; Yang et al. 2021; Zhang et al. 2019).

Given the growing concern about the ecological and long-term health implications of bisphenols, new BPA-related chemicals have been considered to be safer alternatives for industrial applications (Owczarek et al. 2018). Considering their structural similarity to BPA, BPA analogs that are used more widely and frequently in daily life may present endocrine-disrupting effects similar to those of BPA (Rochester and Bolden 2015). However, the potential effects of these BPA substitutes on glucose homeostasis are still unclear. Therefore, we conducted a nested case–control study to explore the potential disrupting effects of exposure to bisphenol on glucose metabolism.

We quantified the serum concentrations of five bisphenols, namely, BPA, bisphenol B (BPB), bisphenol F (BPF), bisphenol S (BPS), and tetrabromobisphenol A (TBBPA). To our knowledge, few studies have assessed the associations of bisphenol mixtures with the risk of GDM. In addition to estimating the contributions of individual elements in bisphenol mixtures, exploring the effect of the overall mixture is necessary to understand the impact of bisphenols on the risk of GDM. We used two new statistical approaches known as Bayesian kernel machine regression (BKMR) (Bobb et al. 2018, 2015) and quantile-based g-computation (Keil et al. 2020) to investigate the effect of the overall mixture.

Methods

Study population

The Guangxi Zhuang Birth Cohort (GZBC) is a prospective and ongoing birth cohort study that has been conducted in county-level hospitals of six major counties in Guangxi Province of China since June 2015. The baseline design, inclusion, and exclusion criteria of this cohort have been reported (Liang et al. 2020). The present study is a subset of the GZBC. In this study, women with a history of diabetes mellitus and insufficient serum sample size were excluded.

A total of 100 pregnant women diagnosed with GDM were selected. For each subject with GDM, four healthy women without GDM were selected as pair-matched controls (maternal age and fetal sex). All subjects had detailed questionnaire data and medical records. All women gave their informed consent, and the study was approved by the Ethical Committee of Guangxi Medical University (no. 20140305–001). After the informed consent form was signed, a standardized structured questionnaire was administered to each participant through an in-person interview to collect information.

GDM diagnosis

An oral glucose tolerance test (OGTT; 75 g) was used to screen for GDM at 24–28 weeks of gestation. GDM was diagnosed according to the Diagnostic Criteria for Gestational Diabetes Mellitus (WS311-2011) released by the Ministry of Health of China, which coincided with the recommendations by the International Association of Diabetes and Pregnancy Study Groups. Pregnant women who met one or more of the following criteria were diagnosed with GDM: (1) fasting blood glucose (FBG) was more than 5.1 mmol/L; (2) 1-h blood glucose > 10.0 mmol/L; or (3) 2-h blood glucose > 8.5 mmol/L. The subjects were tested in the morning after an overnight fast of at least 8 h. The first blood sample was taken to measure the FBG before 9 a.m.

Exposure assessment

Spot serum samples collected during the first trimester were stored in polypropylene containers at –80 °C until further analysis. Serum bisphenol concentrations were quantified using an ultrahigh liquid performance chromatography–tandem mass spectrometer (UPLC-MS, Waters, USA) with isotope-labeled internal standards as previously described (Liang et al. 2020). Briefly, 500 µL of the serum was spiked with three isotope-labeled internal standards (i.e., BPA-D₁₆, BPS-¹³C₁₂, and TBBPA-¹³C₁₂), sodium dihydrogen phosphate dihydrate buffer (pH 5.4), and β-glucuronidase/sulfatase. The mixtures were hydrolyzed and incubated overnight in the dark at 37 °C. After enzymatic hydrolysis, the solution was extracted twice with 2 mL of the solvent [n-hexane:acetone (7:3, v/v)] and 2 mL of methyl tert-butyl ether. The supernatants were combined, evaporated until dry at 40 °C, redissolved in 100 µL of methanol:0.1% ammonia solution (50:50, v/v), and filtered for instrumental analysis. Chromatographic separation was achieved by using an Acquity UPLC BEH C18 (1.7 mm, 2.1 × 100 mm, Waters, USA) analytical column with a mobile phase gradient of ammonia solution and acetonitrile. The bisphenols were detected by using negative-ion electrospray ionization mass spectrometry and multiple reaction monitoring mode.

Procedure blank, solvent blank, and calibration standard samples were measured in each batch sample. As previously reported, the limits of detection (LODs) of BPA, BPB, BPF, BPS, and TBBPA were 0.193, 0.232, 0.507, 0.046, and 0.454 ng/mL, respectively (Liang et al. 2020). The determined concentration below the LOD was calculated as the LOD divided by the square root of 2. The resulting quotients were then used in the subsequent data analyses.

Covariates

For each participant, a face-to-face interview was conducted at the hospital by professionally trained interviewers using a standardized and structured questionnaire to collect information, including demographics (e.g., age, ethnicity, employment status, and self-reported prepregnancy weight) and lifestyles (e.g., drinking and smoking before pregnancy and passive smoking during pregnancy). Moreover, maternal information (e.g., height, parity, gravidity, and pregnancy complications) and birth characteristics (e.g., sex, gestational age, and anthropometric measures) were obtained from the medical data.

Statistical analysis

Summary statistics were calculated and reported as the means \pm standard deviation for continuous variables and n (%) for categorical variables in both the GDM and non-GDM groups. Continuous variables were compared using Mann–Whitney U tests, while categorical variables were compared using χ^2 tests. A natural log transformation [$\ln(X)$] was applied to the bisphenols to normalize their distributions. This step stabilized the variances for parametric model assumptions and reduced the influence of extreme values. The distributions of bisphenol concentrations were presented as percentiles and geometric means. Spearman's rank correlation analysis was used to explore correlations among the serum bisphenol concentrations in the study population.

We categorized participants into tertiles based on the distribution of serum BPA, BPB, BPS, and TBBPA concentrations among the whole population. Given that a high proportion of the samples was below the LOD, BPFs with concentrations $<$ LOD were classified into the low-exposure group (Referent), while those with detected concentrations were divided into the middle- (LOD median) and high-exposure groups (\geq median) based on the median of the detected concentration levels. Conditional logistic regression analysis was employed to the association between individual bisphenol exposure and the risk of GDM, and individual serum bisphenol concentrations were modeled as continuous and categorical variables. Covariates were selected based on either their biological plausibility (regardless of statistical

significance) or the association with GDM in the bivariate analysis ($P < 0.10$). The crude model was a basic unadjusted model. In the adjusted models, covariates included prepregnancy body mass index (BMI), area of residence, passive smoking during pregnancy, gravidity, parity, and regular exercise. The linear P values were derived by modeling the median value of each category of bisphenol as a continuous variable in the statistical model.

Given the potential effect of prepregnancy BMI and the difference in fetal sex on the risk of GDM, stratified analyses were performed using prepregnancy BMI (18.5–22.9, ≥ 23.0 kg/m²) and infant sex to evaluate the potential effect modification on the association between bisphenols and the risk of GDM. The BMI cutoff point of 23.0 kg/m² was selected for overweight subjects, according to a reported optimal cutoff value of BMI for urban Chinese female adults (Zeng et al. 2014).

Considering the possible nonlinearity and nonadditive effects among mixed bisphenols, BKMR (Bobb et al. 2018, 2015) was used to assess the joint effect of all bisphenols on the risk of GDM, the impact of an individual bisphenol as part of a bisphenol mixture, the nonlinear dose–response effect of these bisphenols and the risk of GDM, and the possible interaction among different bisphenols. Predictors with a posterior inclusion probability (PIP) greater than or equal to 0.5 were considered to be meaningful (Lebeaux et al. 2020). Quantile-based g -computation (Keil et al. 2020) was used to corroborate the overall effect to compare with results from the BKMR models. This computation is an adaptation of a mixture modeling method used in environmental epidemiology known as weighted quantile sum regression (Carrico et al. 2015) (WQSR). Compared with the traditional WQSR, quantile g -computation does not require the directional homogeneity of effect estimates. The covariates adjusted in the BKMR and quantile g -computation were the same as those in the single-bisphenol conditional logistic regression analysis. Serum bisphenol levels were \ln -transformed in the BKMR and quantile g -computation.

All data were analyzed using R (version 3.6.1; R Foundation for Statistical Computing), and a two-sided $P < 0.05$ was considered to be statistically significant.

Results

The demographic characteristics of the participants are summarized in Table 1. Compared with non-GDM women, pregnant women with GDM had a higher rate (33.0% vs. 22.2%) of obesity (BMI ≥ 23 kg/m²). In addition, pregnant women with GDM were more likely to be primigravida (49.0% vs. 38.3%). No statistically significant differences between groups were observed in terms of maternal age, area

Table 1 Population characteristics of a nested case–control study based on GDM status in Guangxi, China from 2015 to 2019

Characteristic	Cases (n = 100)	Controls (n = 400)	P value
Age(years)	30.62 ± 6.46	30.6 ± 6.41	0.99
Prepregnancy BMI (kg/m ²)			0.03
< 18.5	11 (11.0%)	79 (19.8%)	
18.5–22.99	56 (56.0%)	232 (58.0%)	
≥ 23	33 (33.0%)	89 (22.2%)	
Area of residence			0.76
Urban	9 (9.0%)	40 (10.0%)	
Rural	91 (91.0%)	360 (90.0%)	
Gravidity			0.29
Nulliparous	24 (24.0%)	77 (19.3%)	
Multiparous	76 (76%)	323 (80.7%)	
Parity			0.05
Primigravida	49 (49.0%)	153 (38.3%)	
Multigravida	51 (51.0%)	247 (61.7%)	
Exercise regularly			0.23
Yes	32 (32.0%)	154 (38.5%)	
No	68 (68.0%)	246 (61.5%)	
Passive smoking during pregnancy			0.11
Yes	52 (52.0%)	172 (43.0%)	
No	48 (48.0%)	228 (57.0%)	
Fetal sex			0.82
Male	55 (55.0%)	225 (56.2%)	
Female	45 (45.0%)	175 (43.8%)	

Abbreviations: *BMI* body mass index; *GDM* gestational diabetes mellitus

P values estimates were derived from Mann–Whitney U tests for continuous variables expressed as mean ± SD, and Pearson chi-squared test for categorical variables expressed as *n* (%)

of residence, passive smoking during pregnancy, gravidity, regular exercise, or infant sex.

Maternal serum bisphenol levels are shown in Table 2. BPA had the highest detection rate (> LOD) (99.6%), followed by BPB (88.8%), BPS (82.2%), TBBPA (71.4%), and BPF (67.2%). The geometric mean BPA and TBBPA levels in the participants with GDM were significantly lower than those in the controls (*P* < 0.001 and *P* = 0.003, respectively). BPA also showed the highest geometric mean concentration (4.726 ng/mL), followed by TBBPA (0.783 ng/mL), BPF

(0.461 ng/mL), BPB (0.243 ng/mL), and BPS (0.094 ng/mL). Most of the bisphenols were significantly correlated with each other (Spearman correlation coefficients = − 0.27 to 0.32, Fig. S1).

The individual effect of each bisphenol on the risk of GDM is shown in Table 3. After adjusting for covariates, BPA (OR: 0.38, 95%CI: 0.29, 0.50) and TBBPA (OR: 0.67, 95%CI: 0.54, 0.85) were negatively associated with the risk of GDM, respectively. Moreover, the concentrations of bisphenols were classified into categorical variables. The

Table 2 Distributions of serum bisphenols concentrations (ng/mL) in study participants

Bisphenols	Detection rate (%)	GM (ng/mL)	Cases (GM, ng/mL)	Controls (GM, ng/mL)	<i>P</i> value	Percentiles (ng/mL)			
						P25	P50	P75	P95
BPA	99.6	4.726	2.016	5.848	< 0.001	2.463	5.552	10.883	24.201
BPB	88.8	0.243	0.241	0.243	0.298	0.233	0.236	0.269	0.360
BPF	67.2	0.461	0.465	0.46	0.373	< LOD	0.605	0.609	0.625
BPS	82.2	0.094	0.103	0.092	0.075	0.050	0.097	0.107	0.832
TBBPA	71.4	0.783	0.525	0.866	0.003	< LOD	0.511	1.643	7.890

Abbreviations: *LOD* limit of detection; *GM* geometric mean

Bisphenol concentrations were expressed as percentiles and GM. *P* values were derived from Mann–Whitney U tests, to distinguish differences between the two groups

Table 3 The association between concentrations of bisphenols and the risk of gestational diabetes mellitus (GDM)

Bisphenols	Cases/controls	Crude OR (95%CI) ^a	Adjusted OR (95%CI) ^b
BPA			
Log [BPA]	100/400	0.40 (0.31, 0.51)	0.38 (0.29, 0.50)
Low	64/101	Reference	Reference
Medium	30/135	0.28 (0.16, 0.49)	0.29 (0.16, 0.52)
High	6/164	0.04 (0.02, 0.11)	0.04 (0.01, 0.10)
<i>P</i> for trend		<0.01	<0.01
BPB			
Log [BPB]	100/400	0.93 (0.49, 1.79)	0.84 (0.42, 1.66)
Low	33/132	Reference	Reference
Medium	41/132	1.23 (0.73, 2.08)	1.14 (0.66, 1.97)
High	26/136	0.78 (0.45, 1.38)	0.67 (0.37, 1.22)
<i>P</i> for trend		0.160	0.076
BPS			
Log [BPS]	100/400	1.09 (0.91, 1.30)	1.06 (0.88, 1.28)
Low	15/132	Reference	Reference
Medium	45/132	1.71 (0.99, 2.96)	1.77 (1.01, 3.13)
High	40/136	1.68 (0.96, 2.94)	1.68 (0.95, 2.99)
<i>P</i> for trend		0.047	0.049
BPF			
Log [BPF]	100/400	1.04 (0.67, 1.62)	1.07 (0.67, 1.71)
Low	31/133	Reference	Reference
Medium	31/127	1.05 (0.61, 1.79)	1.08 (0.62, 1.89)
High	38/140	1.16 (0.69, 1.97)	1.18 (0.68, 2.05)
<i>P</i> for trend		0.671	0.611
TBBPA			
Log [TBBPA]	100/400	0.67 (0.53, 0.83)	0.67 (0.54, 0.85)
Low	36/129	Reference	Reference
Medium	45/120	1.41 (0.85, 2.33)	1.32 (0.78, 2.21)
High	19/151	0.42 (0.22, 0.78)	0.42 (0.22, 0.79)
<i>P</i> for trend		<0.01	<0.01

Abbreviations: *OR* odds ratio; *CI* confidence interval; *BMI* body mass index

^aCrude model had no adjustment

^bAdjusted factors included by prepregnancy BMI, area of residence, passive smoking during pregnancy, gravidity, parity, and exercise regularly

associations of BPA and TBBPA with the risk of GDM remained significant and negative. The OR for BPS was significantly increased in the medium-concentration groups (OR = 1.77; 95% CI: 1.01, 3.13) compared with the low-concentration groups.

We also explored whether the associations between bisphenols and the risk of GDM might be modified by fetal sex (Table S1). Significant results were only observed among women with female fetuses, and the OR for BPS was significantly increased in the high-concentration groups (OR = 2.74; 95% CI: 1.07, 7.47) compared with the low-concentration groups. We also determined whether the bisphenol-GDM associations were modified by prepregnancy BMI (Table S2). The OR for BPS was only significantly increased in the high-concentration groups (OR = 2.74; 95% CI: 1.07,

7.47) compared with the low-concentration groups among women with BMI ≥ 23.0 kg/m², while the per unit increase in ln-transformed TBBPA was only associated with the risk of GDM among women with normal weight.

We visualized the BKMR model. The overall association between the bisphenol mixture and the risk of GDM is shown in Fig. 1A. The risk of GDM decreased with the increase in exposure. The change in the ln-transformed BPS concentration from the 25th to the 75th percentile was associated with a significant increase in the risk of GDM when other bisphenols were fixed at the 25th, 50th, and 75th percentiles. A significant and negative association was observed between the risk of GDM and BPA when all the other bisphenols were fixed at the 25th, 50th, and 75th percentiles. TBBPA displayed a significant and negative effect when all




Fig. 1 Associations between serum bisphenol and GDM risk among the study population by Bayesian kernel machine regression model. **A** The cumulative effect of the serum bisphenols (estimates and 95% credible intervals). The overall effect of mixture shows the estimated exposure–response associations when bisphenols are at a particular percentile (X-axis) compared to when all bisphenols are at 50th percentile. **B** The single-exposure effect (estimates and 95% credible intervals) when all the other bisphenols were fixed at 25th, 50th, and 75th percentiles. **C** Univariate exposure response functions and 95%

confidence intervals between Ln-transformed bisphenol levels (ng/mL) and risk of GDM for each bisphenol when the other bisphenols fixed at the median. **D** Bivariate exposure response functions for each bisphenol on risk of GDM. Exposure response for exposure1 was presented when exposure2 fixed at either the 25th, 50th, or 75th percentile and the rest of bisphenols is fixed at the median. Adjusted factors included prepregnancy BMI, area of residence, passive smoking during pregnancy, gravidity, parity, and regular exercise

the other bisphenols were fixed at the 50th and 75th percentiles (Fig. 1B). The potential nonlinear exposure–response relationships were examined using the BKMR when the levels of the other metals were held at the corresponding median concentration. A suggestion of nonlinear relationships was observed between the risk of GDM and BPA and BPS (Fig. 1C). Furthermore, the BKMR models suggested evidence of interactions between BPA and TBBPA on the risk of GDM (Fig. 1D). The PIPs using BKMR models are summarized in Table S3.

In quantile g-computation analysis, bisphenol mixtures were related to decrease the risk of GDM (OR = 0.57; 95% CI: 0.46, 0.72). BPA (70.2%), TBBPA (21.3%), and BPF (8.5%) had positive contribution to the overall effect, while BPS (71.1%) and BPF (28.9%) had positive weights. In general, the quantile g-computation and BKMR models were in agreement that the overall effect estimates of the ln-transformed bisphenol mixtures were associations in the same direction (Figs. 1 and 2). Both mixture models




Fig. 2 Association between bisphenol composite levels and the risk of GDM using quantile-based g-computation model. The model runs in a negative direction with respect to GDM risk. Adjusted logistic regression models reveal that the quantile g-computation index predominantly consisted of contributions from BPA and TBBPA associated with decreased odds of GDM (OR=0.57, 95% CI: 0.46, 0.72). Model was adjusted by prepregnancy body mass index (BMI), area of residence, passive smoking during pregnancy, gravidity, parity, and regular exercise

suggested that ln-transformed bisphenols had a negative association with the risk of GDM.

Discussion

In the present study, we assessed independent and joint associations between coexposure to five bisphenols (BPA, BPB, BPF, BPS, and TBBPA) and the risk of GDM. BPA and TBBPA were associated with a decreased risk of GDM, while BPS increased the risk of GDM. To the best of our knowledge, this study is the first report to explore the associations of BPB and TBBPA with the risk of GDM. We also determined a significant and negative joint effect of the mixture of the five bisphenols on the risk of GDM, but BPA and TBBPA appeared to be more important than the other three bisphenols in the mixture. This result was consistent across BKMR, quantile g-computation, and multivariable conditional logistic regression models.

The maternal BPA level in the present study was higher than that previously reported (Fisher et al. 2018; Minatoya et al. 2018; Shekhar et al. 2017; Zhang et al. 2013) but lower than that found by another study in China (Yang et al. 2021). BPS levels in this study were lower than those in some areas in China (Jin et al. 2018; Li et al. 2020). In addition, the determined concentrations of BPF were higher than those reported for Czech (median: 0.28 ng/mL) (Kolatorova Sosvorova et al. 2017) and Polish subjects (median: 0.115 ng/mL) (Owczarek et al. 2018). BPB was previously detected at a low frequency (Cobellis et al. 2009; Jin et al. 2018; Li et al. 2020), in contrast to our study which showed a high detection frequency (88.8%) as well as higher concentrations.

The reported TBBPA concentrations in human serum from Belgium (0.08 ng/mL), Norway (0.005 ng/mL), and China (<LOD) were lower than the observed TBBPA median concentration in this study. The differences might be related to the area-specific usage and production of bisphenols, dietary habits, bisphenol contamination in food or in the aquatic environment, and different national regulations for bisphenols.

Previous epidemiological studies have also examined the associations between individual maternal bisphenol levels and the risk of GDM but have mainly focused on BPA. Inconsistent results were obtained. Data from these earlier studies are presented in Table S4 (Fisher et al. 2018; Hou et al. 2021; Robledo et al. 2013; Shapiro et al. 2015; Wang et al. 2017; Yang et al. 2021; Zhang et al. 2019). These reports did not find a positive association between BPA and the risk of GDM. In a Chinese prospective cohort study, maternal urine BPA exposure was associated with a decreased risk of GDM (Wang et al. 2017). Another prospective study from China reported that BPS might be a potential risk factor for GDM. To the best of our knowledge, the present study is the first to show a significant and positive association between BPS levels and the risk of GDM but a negative association between TBBPA levels and the risk of GDM. We also determined that serum BPA was associated with a decreased risk of GDM.

The results showed that serum BPA was associated with a decreased risk of GDM. A study from China also reported that exposure to BPA was associated with a decreased risk of GDM (Wang et al. 2017). Similar findings were obtained in Mexico, where BPA levels were lower in women with GDM (Martínez-Ibarra et al. 2019). A previous small-sample study reported that pregnant women with high BPA levels have low ratios of GDM (Robledo et al. 2013). Some studies have shown that 17 β -estradiol (E2) and xenoestrogens modulate estrogen receptors (ER) in pancreatic β -cells to prevent T2DM (Liu et al. 2009, Tiano and Mauvais-Jarvis 2012, Tiano et al. 2011). Glucose tolerance could be improved by E2 at low concentrations (10^{-8} M) via ER α /ER β and the G-protein coupled ER in rodent models of T2DM (Tiano et al. 2011). Given that BPA has a similar molecular structure to E2, BPA may bind to ERs and act with similar biological effects. Low doses of 1 nM and 10 nM BPA were found to significantly increase insulin content, and 1 nM BPA was equally as effective as E2 when intact islets were cultivated in the presence of BPA at concentrations from 0 to 1000 nM (Alonso-Magdalena et al. 2005; Soriano et al. 2012). Some in vivo experiments have also shown that low-dose BPA in mice led to a rapid increase in plasma insulin and a significant decrease in glycemia (Batista et al. 2012; Nadal et al. 2009, 2004). In the current study, TBBPA was also associated with a decreased risk of GDM. Given its structural resemblance to BPA, TBBPA may have an endocrine effect

similar to that of BPA (Covaci et al. 2009). The estrogenic effect of BPA and TBBPA has been confirmed (Kitamura et al. 2005). An *in vitro* experiment reported that exposure to TBBPA significantly increased the expression levels of insulin (Jiang et al. 2019). However, other studies (Bellavia et al. 2018; Chiu et al. 2017) presented contrasting results in which BPA levels were positively associated with blood glucose levels among pregnant women.

We also identified a positive association between BPS and the risk of GDM. A study from China also reported that BPS was associated with increased fasting plasma glucose (FPG) levels among pregnant women (Zhang et al. 2019). Another study in the French population showed a positive association between exposure to BPS and the incidence of T2DM (Rancière et al. 2019). The disrupting effects of BPS on glucose metabolism were also reported in animal studies (Meng et al. 2018). An experimental study showed that exposure to 1 µg/L and 10 µg/L BPS reduced plasma insulin levels in male zebrafish. This phenomenon altered fasting glycemia, impacted glucose homeostasis, and downregulated the expression of preproinsulin and glucagon genes in visceral tissue after exposure to BPS (Zhao et al. 2018). 17-Hydroxyprogesterone (17-OHP), an intermediate steroid in the biosynthetic pathway that converts cholesterol to cortisol, binds to and stimulates the transcriptional activity of glucocorticoid receptor (GR) (Lu et al. 2020). Previous studies showed a positive association between 17-OHP and the rate of GDM (Nelson et al. 2017; Rebarber et al. 2007; Waters et al. 2009). BPS was less estrogenic and antiandrogenic than other analogs, but BPS showed the largest efficacy on 17-OHP, and BPA had little effect on 17-OHP (Rosenmai et al. 2014). Exposure to BPS may increase the risk of GDM by affecting 17-OHP. Thus, contrary to the results of BPA and TBBPA, BPS may be associated with an increased risk of GDM.

To identify the joint effects and the possible interactions among bisphenols, we used the BKMR method. The results showed a negative joint effect of the mixture of BPA, BPB, BPF, BPS, and TBBPA, and this phenomenon was consistent with that from the quantile *g*-computation models. This approach can be used to analyze not only the overall effect of a mixture but also the effects of each component and dose–response relationships when other metabolites are fixed at a particular percentile (Valeri et al. 2017). The BKMR and quantile *g*-computation models suggested a negative association of BPA and TBBPA with the risk of GDM and a positive association between BPS and this risk. The results were consistent with the traditional model. We also identified the potential interaction between BPA and TBBPA, perhaps because of their similar endocrine effects (Covaci et al. 2009). We determined that BPA played the most critical role in the decreased risk of GDM in the context of various bisphenols. The other bisphenols showed a

low contribution to the joint effect of the five bisphenols on the risk of GDM. We believe that the consistency across multiple modeling types strengthened our results.

In the stratified analysis, we only observed the effect of BPS on the risk of GDM among women with a female fetus. This result was consistent with previous research (Zhang et al. 2019). The effect of fetal sex may be related to the different levels of sex hormones in maternal circulation. A previous study also reported the sex-specific effects of BPS (Wan et al. 2018). In addition, we only observed the effects of BPS among overweight/obese women and those of TBBPA among women with normal weight in the stratified analysis by prepregnancy BMI. These phenomena may be due to the high risk of GDM in overweight/obese women (Chu et al. 2007). A synergistic effect may exist between BPS and overweight/obesity on the risk of GDM in pregnant women, while an antagonistic effect may exist between TBBPA and overweight/obesity.

This study has several strengths. Our prospective study design with a large sample size and mixture modeling approaches enabled us to evaluate individual and joint effects of exposures on the development of GDM. We used two different mixture modeling approaches, namely, BKMR and quantile *g*-computation, which were appropriate for identifying which bisphenol was the most predictive of the risk of GDM and calculating the overall effect of the mixture. Both approaches largely corroborated each other and generally resembled the results of the more traditional multivariable modeling. These characteristics enhanced our internal validity. Moreover, we explored the association between bisphenol A substitutes and the risk of GDM, thus providing a reference for evaluating the health effects of bisphenols.

Nevertheless, several limitations of this study are noteworthy. First, the time interval between the baseline measurement of bisphenols and the period of GDM diagnosis was close due to the short duration of the entire pregnancy. Second, diet is an important source of bisphenols (Liu et al. 2018a), and this information was not available in the cohort we investigated. Future studies should further investigate these important sources of bisphenols and whether those factors operate as confounders or effect modifiers of the observed associations. In addition, some other known or unknown risk factors for GDM were not considered completely, and this limitation may influence the results of the study.

Conclusion

We assessed the individual, overall, and joint exposure effects of maternal bisphenols on the maternal risk of GDM by using multiple robust statistical techniques. The results showed an inverse association between BPA and TBBPA

levels and the risk of GDM. We provided the first evidence that BPS levels were positively associated with the risk of GDM. Exposure to the mixture of five bisphenols was negatively associated with the risk of GDM, and BPA displayed the most significant effect on the risk of GDM. Our findings may have important public health implications and provide a reference for evaluating the health effects of bisphenols. Further epidemiologic and experimental studies are needed to confirm these observations and to investigate the underlying molecular mechanisms.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11356-021-17794-8>.

Acknowledgements The authors really appreciate all the participants who collaborated with this study and donated serum samples as well as all staff for assisting in collecting the samples and data.

Author contribution Peng Tang: methodology, formal analysis, writing (original draft), visualization. Jun Liang: methodology, investigation, data curation, validation. Qian Liao: methodology, validation, investigation. Huishen Huang: validation, investigation. Xiaojing Guo: validation, investigation. Mengrui Lin: investigation. Bihu Liu: investigation. Bincai Wei: investigation. Xiaoyun Zeng: writing (review and editing), supervision. Shun Liu: writing (review and editing), supervision. Dongping Huang: methodology, writing (review and editing), supervision, funding acquisition. Xiaoqiang Qiu: project administration, resources, data curation, writing (review and editing), supervision, funding acquisition.

Funding This research was supported by Guangxi Key Research Program (AB17195012) and the National Natural Science Foundation of China (81860587).

Data availability The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval All participants were informed at recruitment and they signed an informed consent. Ethics approval was provided by Guangxi Medical University, China (no. 20140305–001).

Consent to participate Not applicable.

Consent for publication Not applicable.

Conflict of interest The authors declare no competing interests.

References

- Alonso-Magdalena P, Laribi O, Ropero AB, Fuentes E, Ripoll C, Soria B, Nadal A (2005) Low doses of bisphenol A and diethylstilbestrol impair Ca²⁺ signals in pancreatic alpha-cells through a nonclassical membrane estrogen receptor within intact islets of Langerhans. *Environ Health Perspect* 113:969–977
- American-Diabetes-Association (2013) Diagnosis and classification of diabetes mellitus. *Diabetes Care* 36(Suppl 1):S67–74
- Batista TM, Alonso-Magdalena P, Vieira E, Amaral ME, Cederroth CR, Nef S, Quesada I, Carneiro EM, Nadal A (2012) Short-term treatment with bisphenol-A leads to metabolic abnormalities in adult male mice. *PLoS One* 7:e33814
- Bellavia A, Cantonwine DE, Meeker JD, Hauser R, Seely EW, McElrath TF, James-Todd T (2018) Pregnancy urinary bisphenol-A concentrations and glucose levels across BMI categories. *Environ Int* 113:35–41
- Bobb JF, Valeri L, Claus Henn B, Christiani DC, Wright RO, Mazumdar M, Godleski JJ, Coull BA (2015) Bayesian kernel machine regression for estimating the health effects of multi-pollutant mixtures. *Biostatistics* 16:493–508
- Bobb JF, Claus Henn B, Valeri L, Coull BA (2018) Statistical software for analyzing the health effects of multiple concurrent exposures via Bayesian kernel machine regression. *Environ Health* 17:67
- Burlina S, Dalfrà MG, Lapolla A (2019) Short- and long-term consequences for offspring exposed to maternal diabetes: a review. *J Matern Fetal Neonatal Med* 32:687–694
- Carrico C, Gennings C, Wheeler DC, Factor-Litvak P (2015) Characterization of weighted quantile sum regression for highly correlated data in a risk analysis setting. *J Agric Biol Environ Stat* 20:100–120
- Chiu YH, Minguez-Alarcón L, Ford JB, Keller M, Seely EW, Messerlian C, Petrozza J, Williams PL, Ye X, Calafat AM, Hauser R, James-Todd T (2017) Trimester-specific urinary bisphenol A concentrations and blood glucose levels among pregnant women from a fertility clinic. *J Clin Endocrinol Metab* 102:1350–1357
- Chu SY, Callaghan WM, Kim SY, Schmid CH, Lau J, England LJ, Dietz PM (2007) Maternal obesity and risk of gestational diabetes mellitus. *Diabetes Care* 30:2070–2076
- Cobellis L, Colacurci N, Trabucco E, Carpentiero C, Grumetto L (2009) Measurement of bisphenol A and bisphenol B levels in human blood sera from healthy and endometriotic women. *Biomed Chromatogr* 23:1186–1190
- Covaci A, Voorspoels S, Abdallah MA, Geens T, Harrad S, Law RJ (2009) Analytical and environmental aspects of the flame retardant tetrabromobisphenol-A and its derivatives. *J Chromatogr A* 1216:346–363
- Ding S, Fan Y, Zhao N, Yang H, Ye X, He D, Jin X, Liu J, Tian C, Li H, Xu S, Ying C (2014) High-fat diet aggravates glucose homeostasis disorder caused by chronic exposure to bisphenol A. *J Endocrinol* 221:167–179
- Ehrlich S, Lambers D, Baccarelli A, Khoury J, Macaluso M, Ho SM (2016) Endocrine disruptors: a potential risk factor for gestational diabetes mellitus. *Am J Perinatol* 33:1313–1318
- Farahvar S, Walfisch A, Sheiner E (2019) Gestational diabetes risk factors and long-term consequences for both mother and offspring: a literature review. *Expert Rev Endocrinol Metab* 14:63–74
- Filardi T, Panimolle F, Lenzi A, Morano S (2020) Bisphenol A and phthalates in diet: an emerging link with pregnancy complications. *Nutrients* 12:525
- Fisher BG, Frederiksen H, Andersson AM, Juul A, Thankamony A, Ong KK, Dunger DB, Hughes IA, Acerini CL (2018) Serum phthalate and triclosan levels have opposing associations with risk factors for gestational diabetes mellitus. *Front Endocrinol (lausanne)* 9:99
- Hou Y, Li S, Xia L, Yang Q, Zhang L, Zhang X, Liu H, Huo R, Cao G, Huang C, Tian X, Sun L, Cao D, Zhang M, Zhang Q, Tang N (2021) Associations of urinary phenolic environmental estrogens exposure with blood glucose levels and gestational diabetes mellitus in Chinese pregnant women. *Sci Total Environ* 754:142085
- Jiang S, Miao J, Wang X, Liu P, Pan L (2019) Inhibition of growth in juvenile Manila clam *Ruditapes philippinarum*: potential adverse outcome pathway of TBBPA. *Chemosphere* 224:588–596

- Jin H, Zhu J, Chen Z, Hong Y, Cai Z (2018) Occurrence and partitioning of bisphenol analogues in adults' blood from China. *Environ Sci Technol* 52:812–820
- Keil AP, Buckley JP, O'Brien KM, Ferguson KK, Zhao S, White AJ (2020) A quantile-based g-computation approach to addressing the effects of exposure mixtures. *Environ Health Perspect* 128:47004
- Kitamura S, Suzuki T, Sanoh S, Kohta R, Jinno N, Sugihara K, Yoshihara S, Fujimoto N, Watanabe H, Ohta S (2005) Comparative study of the endocrine-disrupting activity of bisphenol A and 19 related compounds. *Toxicol Sci* 84:249–259
- Kolatorova Sosvorova L, Chlupacova T, Vitku J, Vlk M, Heracek J, Starka L, Saman D, Simkova M, Hampl R (2017) Determination of selected bisphenols, parabens and estrogens in human plasma using LC-MS/MS. *Talanta* 174:21–28
- Konieczna A, Rutkowska A, Rachoń D (2015) Health risk of exposure to Bisphenol A (BPA). *Rocz Panstw Zakl Hig* 66:5–11
- Lebeaux RM, Doherty BT, Gallagher LG, Zoeller RT, Hoofnagle AN, Calafat AM, Karagas MR, Yolton K, Chen A, Lanphear BP, Braun JM, Romano ME (2020) Maternal serum perfluoroalkyl substance mixtures and thyroid hormone concentrations in maternal and cord sera: The HOME Study. *Environ Res* 185:109395
- Li A, Zhuang T, Shi W, Liang Y, Liao C, Song M, Jiang G (2020) Serum concentration of bisphenol analogues in pregnant women in China. *Sci Total Environ* 707:136100
- Liang J, Liu S, Liu T, Yang C, Wu Y, Jennifer Tan HJ, Wei B, Ma X, Feng B, Jiang Q, Huang D, Qiu X (2020) Association of prenatal exposure to bisphenols and birth size in Zhuang ethnic newborns. *Chemosphere* 252:126422
- Liu J, Wattar N, Field CJ, Dinu I, Dewey D, Martin JW (2018a) Exposure and dietary sources of bisphenol A (BPA) and BPA-alternatives among mothers in the APRON cohort study. *Environ Int* 119:319–326
- Liu S, Le May C, Wong WP, Ward RD, Clegg DJ, Marcelli M, Korach KS, Mauvais-Jarvis F (2009) Importance of extranuclear estrogen receptor-alpha and membrane G protein-coupled estrogen receptor in pancreatic islet survival. *Diabetes* 58:2292–2302
- Liu X, Zhang L, Li J, Meng G, Chi M, Li T, Zhao Y, Wu Y (2018b) A nested case-control study of the association between exposure to polybrominated diphenyl ethers and the risk of gestational diabetes mellitus. *Environ Int* 119:232–238
- Lu Y, Wang E, Chen Y, Zhou B, Zhao J, Xiang L, Qian Y, Jiang J, Zhao L, Xiong X, Lu Z, Wu D, Liu B, Yan J, Zhang R, Zhang H, Hu C, Li X (2020) Obesity-induced excess of 17-hydroxyprogesterone promotes hyperglycemia through activation of glucocorticoid receptor. *J Clin Invest* 130:3791–3804
- Martínez-Ibarra A, Martínez-Razo LD, Vázquez-Martínez ER, Martínez-Cruz N, Flores-Ramírez R, García-Gómez E, López-López M, Ortega-González C, Camacho-Arroyo I, Cerbón M (2019) Unhealthy levels of phthalates and bisphenol A in Mexican pregnant women with gestational diabetes and its association to altered expression of miRNAs involved with metabolic disease. *Int J Mol Sci* 20:3343
- Meng Z, Wang D, Yan S, Li R, Yan J, Teng M, Zhou Z, Zhu W (2018) Effects of perinatal exposure to BPA and its alternatives (BPS, BPF and BPAF) on hepatic lipid and glucose homeostasis in female mice adolescent offspring. *Chemosphere* 212:297–306
- Michałowicz J (2014) Bisphenol A—sources, toxicity and biotransformation. *Environ Toxicol Pharmacol* 37:738–758
- Minatoya M, Itoh S, Yamazaki K, Araki A, Miyashita C, Tamura N, Yamamoto J, Onoda Y, Ogasawara K, Matsumura T, Kishi R (2018) Prenatal exposure to bisphenol A and phthalates and behavioral problems in children at preschool age: the Hokkaido Study on Environment and Children's Health. *Environ Health Prev Med* 23:43
- Nadal A, Ropero AB, Fuentes E, Soria B, Ripoll C (2004) Estrogen and xenoestrogen actions on endocrine pancreas: from ion channel modulation to activation of nuclear function. *Steroids* 69:531–536
- Nadal A, Alonso-Magdalena P, Soriano S, Quesada I, Ropero AB (2009) The pancreatic beta-cell as a target of estrogens and xenoestrogens: implications for blood glucose homeostasis and diabetes. *Mol Cell Endocrinol* 304:63–68
- Nelson DB, McIntire DD, McDonald J, Gard J, Turrichi P, Leveno KJ (2017) 17-alpha Hydroxyprogesterone caproate did not reduce the rate of recurrent preterm birth in a prospective cohort study. *Am J Obstet Gynecol* 216:600.e1–600.e9
- Owczarek K, Kubica P, Kudlak B, Rutkowska A, Konieczna A, Rachoń D, Namieśnik J, Wasik A (2018) Determination of trace levels of eleven bisphenol A analogues in human blood serum by high performance liquid chromatography-tandem mass spectrometry. *Sci Total Environ* 628–629:1362–1368
- Rancièrè F, Botton J, Slama R, Lacroix MZ, Debrauwer L, Charles MA, Roussel R, Balkau B, Magliano DJ (2019) Exposure to bisphenol A and bisphenol S and incident type 2 diabetes: a case-cohort study in the french cohort D. E. S. I. R. *Environ Health Perspect* 127:107013
- Rebarber A, Istwan NB, Russo-Stieglitz K, Cleary-Goldman J, Rhea DJ, Stanziano GJ, Saltzman DH (2007) Increased incidence of gestational diabetes in women receiving prophylactic 17alpha-hydroxyprogesterone caproate for prevention of recurrent preterm delivery. *Diabetes Care* 30:2277–2280
- Robledo C, Peck JD, Stoner JA, Carabin H, Cowan L, Koch HM, Goodman JR (2013) Is bisphenol-A exposure during pregnancy associated with blood glucose levels or diagnosis of gestational diabetes? *J Toxicol Environ Health A* 76:865–873
- Rochester JR (2013) Bisphenol A and human health: a review of the literature. *Reprod Toxicol* 42:132–155
- Rochester JR, Bolden AL (2015) Bisphenol S and F: A systematic review and comparison of the hormonal activity of bisphenol a substitutes. *Environ Health Perspect* 123:643–650
- Rosenmai AK, Dybdahl M, Pedersen M, Alice van Vugt-Lussenburg BM, Wedebye EB, Taxvig C, Vinggaard AM (2014) Are structural analogues to bisphenol a safe alternatives? *Toxicol Sci* 139:35–47
- Shapiro GD, Dodds L, Arbuckle TE, Ashley-Martin J, Fraser W, Fisher M, Taback S, Keely E, Bouchard MF, Monnier P, Dallaire R, Morisset A, Ettinger AS (2015) Exposure to phthalates, bisphenol A and metals in pregnancy and the association with impaired glucose tolerance and gestational diabetes mellitus: The MIREC study. *Environ Int* 83:63–71
- Shekhar S, Sood S, Showkat S, Lite C, Chandrasekhar A, Vairamani M, Barathi S, Santosh W (2017) Detection of phenolic endocrine disrupting chemicals (EDCs) from maternal blood plasma and amniotic fluid in Indian population. *Gen Comp Endocrinol* 241:100–107
- Soriano S, Alonso-Magdalena P, García-Arévalo M, Novials A, Muhammed SJ, Salehi A, Gustafsson JA, Quesada I, Nadal A (2012) Rapid insulinotropic action of low doses of bisphenol-A on mouse and human islets of Langerhans: role of estrogen receptor β . *PLoS One* 7:e31109
- Sun Q, Cornelis MC, Townsend MK, Tobias DK, Eliassen AH, Franke AA, Hauser R, Hu FB (2014) Association of urinary concentrations of bisphenol A and phthalate metabolites with risk of type 2 diabetes: a prospective investigation in the Nurses' Health Study (NHS) and NHSII cohorts. *Environ Health Perspect* 122:616–623
- Tiano J, Mauvais-Jarvis F (2012) Selective estrogen receptor modulation in pancreatic β -cells and the prevention of type 2 diabetes. *Islets* 4:173–176
- Tiano JP, Delghingaro-Augusto V, Le May C, Liu S, Kaw MK, Khuder SS, Latour MG, Bhatt SA, Korach KS, Najjar SM, Prentki M, Mauvais-Jarvis F (2011) Estrogen receptor activation reduces

- lipid synthesis in pancreatic islets and prevents β cell failure in rodent models of type 2 diabetes. *J Clin Invest* 121:3331–3342
- Valeri L, Mazumdar MM, Bobb JF, Claus Henn B, Rodrigues E, Sharif OIA, Kile ML, Quamruzzaman Q, Afroz S, Golam M, Amarasiriwardena C, Bellinger DC, Christiani DC, Coull BA, Wright RO (2017) The joint effect of prenatal exposure to metal mixtures on neurodevelopmental outcomes at 20–40 months of age: evidence from rural Bangladesh. *Environ Health Perspect* 125:067015
- Wan Y, Huo W, Xu S, Zheng T, Zhang B, Li Y, Zhou A, Zhang Y, Hu J, Zhu Y, Chen Z, Lu S, Wu C, Jiang M, Jiang Y, Liu H, Yang X, Xia W (2018) Relationship between maternal exposure to bisphenol S and pregnancy duration. *Environ Pollut* 238:717–724
- Wang X, Wang X, Chen Q, Luo ZC, Zhao S, Wang W, Zhang HJ, Zhang J, Ouyang F (2017) Urinary bisphenol a concentration and gestational diabetes mellitus in Chinese women. *Epidemiology* 28(Suppl 1):S41–s47
- Waters TP, Schultz BAH, Mercer BM, Catalano PM (2009) Effect of 17alpha-hydroxyprogesterone caproate on glucose intolerance in pregnancy. *Obstet Gynecol* 114:45–49
- Xu H, Zhou Q, Zhang J, Chen X, Zhao H, Lu H, Ma B, Wang Z, Wu C, Ying C, Xiong Y, Zhou Z, Li X (2020) Exposure to elevated per- and polyfluoroalkyl substances in early pregnancy is related to increased risk of gestational diabetes mellitus: A nested case-control study in Shanghai, China. *Environ Int* 143:105952
- Yang J, Wang H, Du H, Xu L, Liu S, Yi J, Chen Y, Jiang Q, He G (2021) Serum Bisphenol A, glucose homeostasis, and gestational diabetes mellitus in Chinese pregnant women: a prospective study. *Environ Sci Pollut Res Int* 28:12546–12554
- Zeng Q, He Y, Dong S, Zhao X, Chen Z, Song Z, Chang G, Yang F, Wang Y (2014) Optimal cut-off values of BMI, waist circumference and waist:height ratio for defining obesity in Chinese adults. *Br J Nutr* 112:1735–1744
- Zhang T, Sun H, Kannan K (2013) Blood and urinary bisphenol A concentrations in children, adults, and pregnant women from china: partitioning between blood and urine and maternal and fetal cord blood. *Environ Sci Technol* 47:4686–4694
- Zhang W, Xia W, Liu W, Li X, Hu J, Zhang B, Xu S, Zhou Y, Li J, Cai Z, Li Y (2019) Exposure to bisphenol a substitutes and gestational diabetes mellitus: a prospective cohort study in China. *Front Endocrinol (lausanne)* 10:262
- Zhao F, Jiang G, Wei P, Wang H, Ru S (2018) Bisphenol S exposure impairs glucose homeostasis in male zebrafish (*Danio rerio*). *Ecotoxicol Environ Saf* 147:794–802

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.