

## Erratum to: Widespread endocrine activity in river sediments in Hesse, Germany, assessed by a combination of *in vitro* and *in vivo* bioassays

Simone Galluba · Jörg Oehlmann


Published online: 11 January 2012  
© Springer-Verlag 2011

Erratum to: J Soils Sediments  
DOI 10.1007/s11368-011-0454-0

Table 2: The values highlighted in light grey were missing in this table.

Table 2 Contamination levels at sampling sites expressed as sums of quality limit value exceedances

Sampling site	Metals	PCB	PAH	OTC	Total
9	2.32	0	6.05	0	8.37
10	0	0	4.15	0	4.15
15	0	0	3.95	0	3.95
16	0	0	3.05	0	3.05
18	1.06	0	40.8	1.30	43.2
20	10.3	0	12.8	0	23.1
21	0	0	3.10	0	3.10
22	0	0	0	0	0
23	3.36	2.45	19.8	1.35	26.9
24	7.07	2.75	7.65	0	17.5
25	2.90	1.15	19.4	0	23.5
26	2.62	0	10.5	0	13.1
28	3.00	1.05	33.4	0	37.4
31	15.3	2.40	5.40	0	23.1
32	1.09	0	0	0	1.09
33	1.21	0	4.40	0	5.61
34	1.35	0	13.6	0	15.0
36	3.57	0	3.10	0	6.67
39	14.0	17.7	6.75	0	38.4
41	5.95	0	0	1.11	7.06
43	2.26	0	5.60	0	7.86
44	3.95	0	6.95	12.5	23.4
47	2.59	0	3.40	0	5.99
48	2.20	0	5.10	0	7.30



Contamination levels (represented by shades of grey) are in steps of 20%, with 100% representing the highest contamination of a substance class (for details see Table 1). For sampling sites not listed here, no chemical analytical data were available.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s11368-011-0454-0>.

S. Galluba (✉) · J. Oehlmann  
Department of Aquatic Ecotoxicology,  
Goethe University Frankfurt am Main,  
Max-von-Laue-Str. 13,  
60438 Frankfurt am Main, Germany  
e-mail: [galluba@bio.uni-frankfurt.de](mailto:galluba@bio.uni-frankfurt.de)