



# Cytotoxic activity screening of Bangladeshi medicinal plant extracts

Raushanara Akter · Shaikh J. Uddin ·  
I. Darren Grice · Evelin Tiralongo

Received: 26 March 2013 / Accepted: 10 June 2013 / Published online: 12 July 2013  
© The Japanese Society of Pharmacognosy and Springer Japan 2013

**Abstract** The cytotoxic activity of 23 crude methanol extracts from 19 Bangladeshi medicinal plants was investigated against healthy mouse fibroblasts (NIH3T3), healthy monkey kidney (VERO) and four human cancer cell lines (gastric, AGS; colon, HT-29; and breast, MCF-7 and MDA-MB-231) using MTT assay. High cytotoxicity across all cell lines tested was exhibited by *Aegiceras corniculatum* (fruit) and *Hymenodictyon excelsum* (bark) extracts (IC<sub>50</sub> values ranging from 0.0005 to 0.9980 and 0.08 to 0.44 mg/mL, respectively). Fourteen extracts from 11 plant species, namely *Clitoria ternatea* (flower and leaf), *Dillenia indica* (leaf), *Diospyros peregrina* (leaf), *Dipterocarpus turbinatus* (bark and leaf), *Ecbolium viride* (leaf), *Glinus oppositifolius* (whole plant), *Gnaphalium luteoalbum* (leaf), *Jasminum sambac* (leaf), *Lannea coromandelica* (bark and leaf), *Mussaenda glabrata* (leaf) and *Saraca asoca* (leaf), were also significantly cytotoxic (IC<sub>50</sub> < 1.0 mg/mL) against at least one of the cancer cell lines tested. More selectively, *Avicennia alba* (leaf), *C. ternatea* (flower and leaf),

*Caesalpinia pulcherrima* (leaf), *E. viride* (leaf) and *G. oppositifolius* (whole plant) showed cytotoxicity only against both of the breast cancer cell lines (MCF-7 and MDA-MB-231). In contrast, *C. ternatea* (flower and leaf) exhibited high cytotoxic activity against MDA-MB-231 (IC<sub>50</sub> values of 0.11 and 0.49 mg/mL, respectively), whereas *E. viride* and *G. oppositifolius* whole plant extracts exhibited high activity against MCF-7 cells (IC<sub>50</sub> values of 0.06 and 0.15 mg/mL, respectively). The cytotoxic activity test results for 9 of the plant species correlate with their traditional use as anticancer agents, thus making them interesting sources for further drug development.

**Keywords** Bangladeshi medicinal plants · Traditional use · Cytotoxic activity · Anticancer · MTT assay

## Introduction

Identification of medicinal plants with significant cytotoxic potential useful for the development of cancer therapeutics has gained increasing importance in the last decade, and research in this field is expanding [1]. More than 1000 plants species have been identified with significant anti-cancer potential [2]. About 80 % of the population in developing countries relies on traditional plant-based medicines for their primary health care needs [3]. In Bangladesh, it is estimated that more than 500 species of medicinal plants exist. These indigenous medicinal plants have been extensively used in the preparation of Unani, Ayurvedic and homeopathic medicines in Bangladesh and have been serving as important raw materials in many modern medicinal preparations [4]. Pharmacological activities of 64 plants have been studied and reported in the last two decades in Bangladesh [5], with some of these

**Electronic supplementary material** The online version of this article (doi:10.1007/s11418-013-0789-5) contains supplementary material, which is available to authorized users.

R. Akter · S. J. Uddin · E. Tiralongo (✉)  
School of Pharmacy and Griffith Health Institute,  
Griffith University, Gold Coast Campus, Gold Coast,  
Queensland 4222, Australia  
e-mail: e.tiralongo@griffith.edu.au

S. J. Uddin  
Pharmacy Discipline, Khulna University, Khulna 9208,  
Bangladesh

I. D. Grice  
Institute for Glycomics and School of Medical Science,  
Griffith University, Gold Coast Campus, Gold Coast,  
Queensland 4222, Australia

investigated for anticancer activity [6], as well as cytotoxic activity [7, 8]. However, most of these plants have not thoroughly been evaluated for cytotoxic activity. Thorough scientific evaluation of the pharmacological properties of these plants used in different traditional formulations carries enormous potential and promise for the twenty first century [4, 5]. Based on ethno-medical information, the present study reports on the evaluation of cytotoxic activity

of selected Bangladeshi medicinal plant extracts. A total of 19 medicinal plant species, namely *Aegiceras corniculatum*, *Argyreia nervosa*, *Avicennia alba*, *Caesalpinia pulcherrima*, *Clerodendrum viscosum*, *Clitoria ternatea*, *Dillenia indica*, *Dipterocarpus turbinatus*, *Diospyros peregrina*, *Ecbolium viride*, *Glinus oppositifolius*, *Glycosmis pentaphylla*, *Gnaphalium luteoalbum*, *Hymenodictyon excelsum*, *Jasminum sambac*, *Lannea coromandelica*,

**Table 1** List of the selected Bangladeshi medicinal plants with their traditional uses

Plant species	Family	Local name	Voucher no.	Traditional uses
<i>Aegiceras corniculatum</i>	Myrsinaceae	Kholisha	DACB 31584	B, L and S: fish poison, asthma, diabetes, <i>inflammation</i> and rheumatism [9, 23]
<i>Argyreia nervosa</i>	Convolvulaceae	Bichtarak	DACB 37954	S: hypotensive, hallucinogenic; AP: stomach trouble, small pox, syphilis, diarrhoea, dysentery; L: wound healing, local stimulant, skin diseases; R: rhematic affections, diseases of nervous system, diuretic, gonorrhoea, urinary diseases and <i>chronic ulcers</i> [4, 9]
<i>Avicennia alba</i>	Avicenniaceae	Morcha baen	DACB 40556	Treatment of infertility, skin diseases, <i>tumours, ulcers</i> , (resin) [23]
<i>Caesalpinia pulcherrima</i>	Caesalpinaceae	Krishnachura	DACB 32020	L: purgative, abortifacient, <i>anticancer</i> , fungi toxic, liver disorders; F: cough, asthma, bronchitis, malaria fever; R: cholera, infantile convulsions; W: diarrhoea, dysentery, skin diseases [4, 9]
<i>Clerodendrum viscosum</i>	Verbenaceae	Bhant	DACB 37953	WP: hypotensive, L: anthelmintic, emetic, antiperiodic in malaria, cough, asthma, <i>tumour</i> , skin disease, snakebite, R: antifungal [4]
<i>Clitoria ternatea</i>	Papilionaceae	Aparajita	DACB 32021	R: demulcent, aperients, laxative, diuretic; S: cathartic, ascites, sore throat, <i>tumours</i> , dropsy and skin diseases; AP: colic, gonorrhoea and skin diseases [4, 9]
<i>Dillenia indica</i>	Dilleniaceae	Chalta	DACB 32019	Fr: expectorant, laxative, tonic, abdominal pain; L: astringent; B: astringent; S: antifungal and <i>antibacterial</i> [9]
<i>Diospyros peregrina</i>	Ebenaceae	Gab	DACB 30323	B: astringent, dysentery, biliousness; Fr: astringent, sore throat, wounds, <i>ulcers</i> , cough, dyspnoea; S: diarrhoea, dysentery; SB: antiprotozoal, antiviral, diuretic and <i>anticancer</i> [4]
<i>Dipterocarpus turbinatus</i>	Dipterocarpaceae	Garjan	DACB 32026	Gonorrhoea, gleet, rheumatism, <i>ulcer</i> , ringworm and skin diseases [9]
<i>Ecbolium viride</i>	Acanthaceae	Nilkanta	DACB 32018	L: <i>antitumour</i> ; R: <i>tumour</i> , jaundice, rheumatism; WP: gout, dysuria, cardiovascular diseases [9]
<i>Glinus oppositifolius</i>	Molluginaceae	Gima	DACB 32014	WP: CNS depressant, diuretic, stomachic, aperients, antiseptic and skin diseases [9]
<i>Glycosmis pentaphylla</i>	Rutaceae	Daton	DACB 37931	L: fever, liver complaints, coughs, rheumatism, anaemia, jaundice, eczema and other skin infections; R: fever; Fr: dysentery [9]
<i>Gnaphalium luteoalbum</i>	Compositae	Boro karma	DACB 37955	L: astringent, diuretic, haemostatic, T: counter-irritant [4]
<i>Hymenodictyon excelsum</i>	Rubiaceae	Bhui kadam	DACB 32013	B: astringent, febrifuge, antiperiodic, hypotensive, <i>antimicrobial</i> , diarrhoea; W: herpes [9]
<i>Jasminum sambac</i>	Oleaceae	Beli phul	DACB 31262	AP: CNS depressant, hypotensive; L: <i>indolent ulcer</i> , <i>breast tumours</i> ; R: emmenagogue [4]
<i>Lannea coromandelica</i>	Anacardiaceae	Jeol/jiga	DACB 35242	B: astringent, leprous and <i>obstinate ulcer</i> , mouth sores; L: local swellings and pain of the body [9]
<i>Mussaenda glabrata</i>	Rubiaceae	Patralekha	DACB 32023	R: white leprosy; L: jaundice; F: diuretic; asthma, fever, dropsy, chest pain and <i>ulcer</i> [9]
<i>Myrica nagi</i>	Myricaceae	Kaiphall	DACB 32029	B: astringent, carminative, antiseptic, fever, cough asthma, cholera, chronic dysentery, anaemia, piles, chronic bronchitis, <i>ulcer</i> and <i>tumours</i> [4]
<i>Saraca asoca</i>	Caesalpinaceae	Ashok	DACB 32007	B: <i>antitumour</i> , menorrhagia, bleeding haemorrhoids, haemorrhagic dysentery, colic, <i>ulcers</i> ; L: blood purification, F: haemorrhagic dysentery, biliousness, syphilis, uterine tonic [4, 9]

AP aerial parts, B bark, F flowers, Fr fruits, L leaves, R roots, RB root bark, S seeds, SB stem bark, T tomentum, W wood, WP whole plant

*Mussaenda glabrata*, *Myrica nagi* and *Saraca asoca*, were evaluated for their cytotoxic activity. Most of these plants grow all over Bangladesh, two plants (*A. corniculatum*, *A. alba*), which are mangrove plants, grow specifically in the coastal region, such as the Sundarban, and some (*D. turbinatus*, *M. glabrata* and *M. nagi*) grow specifically in the hilly area of Sylhet and Chittagong. Traditionally, these plants have been used as antitumour/anticancer, anti-infective and anti-inflammatory agents as well as for the treatment of other diseases (Table 1). Different parts of 9 plants, namely *A. alba*, *C. pulcherrima*, *C. ternatea*, *C. viscosum*, *D. peregrina*, *E. viride*, *J. sambac*, *M. nagi* and *S. asoca*, have a reputation of being used traditionally as anticancer medications [3, 4, 9].

Cytotoxic activity screening of 19 Bangladeshi medicinal plants against healthy mouse fibroblast (NIH3T3), healthy monkey kidney (VERO), gastric cancer (AGS), colon cancer (HT-29), breast cancer (oestrogen-dependent, MCF-7; non-oestrogen-dependent, MDA-MB-231) cells are reported here for the first time to confirm the traditional use of some plants as anticancer agents and to additionally identify new plants with significant anticancer potential.

## Materials and methods

### Plant collection and identification

Plants were collected from the coastal Sundarban tidal forest as well as from other parts of Bangladesh during March 2006 to May 2007. The plant materials were subjected to shade drying and identified by the Bangladesh National Herbarium, Mirpur, Dhaka, where a voucher specimen for individual plant species was deposited for future reference.

### Extraction of plant materials

The shade-dried, powdered plant materials were extracted by soaking in methanol for 24 h. The extracts were then filtered and the solvent evaporated (rotary evaporator), followed by freeze drying.

### Phytochemical profile

The phytochemical profiles of the plant extracts were obtained by analytical HPLC (Luna C18, 250 × 4.6 mm column) using 10–90 % methanol in water as the mobile phase with UV detection at 210 and 280 nm (see supplementary data).

## Cytotoxic screening

### Cell culture

Two normal cell lines, namely mouse fibroblast (NIH3T3, ATCC CRL-1658) and healthy monkey kidney cells (VERO, ATCC CCL-81), and four human cancer cell lines, namely gastric (AGS, ATCC CRL-1739), colon (HT-29, ATCC HTB-38), non-oestrogen-dependent breast (MDA-MB-231, ATCC HTB-26) and oestrogen-dependent breast (MCF-7, ATCC: HTB-22) cancer cells, were used for cytotoxicity screening of the selected Bangladeshi medicinal plant extracts. All cell lines were purchased from ATCC, Manassas, VA 20108, USA. Cell lines were cultured in Advanced DMEM supplemented with 10 % inactivated NBS and 5 mM L-glutamine, and grown at 37 °C in a humidified atmosphere of 5 % CO<sub>2</sub> in air.

### MTT colorimetric assay

The MTT colorimetric assay was performed to evaluate the cytotoxicity of the selected Bangladeshi plant extracts according to the method described and validated by Uddin et al. [8]. Briefly, the cells were seeded in 96-well plates at a density of  $1.0 \times 10^4$ – $2.0 \times 10^4$  cells/well. Following 24 h incubation and attachment, the cells were treated with different concentrations of plant extract for 48 h. Washing and incubation with MTT solution for 2 h was followed by cells being lysed with dimethyl sulfoxide (DMSO). The absorbance was measured after 45 min using a microplate reader (Wallac 1420 Multilabel counter, PerkinElmer) at a wavelength of 560 nm. DMSO (2 %), was used to dissolve the extracts. It showed less than 20 % cell growth inhibition and served as the negative control, whereas 20 % DMSO (>80 % cell growth inhibition) and cycloheximide served as positive controls. The results are generated from two independent experiments; each experiment was performed in triplicate. The IC<sub>50</sub> values were calculated with Probit analysis software (LdP Line software, USA).

## Results and discussion

Many Bangladeshi medicinal plants are traditionally known to have cytotoxic and antitumour properties, with some having a folkloric reputation of being used in the treatment of different types of cancer [4, 8]. Our study reports on the investigation into the cytotoxic activity of 19 Bangladeshi medicinal plant species which have not been investigated for such activity previously. Cytotoxic activities of the tested extracts are summarized in Table 2.

**Table 2** Cytotoxic activity (IC<sub>50</sub>) of selected Bangladeshi medicinal plant extracts

Plant species	Plant (part)	Yield %	Cytotoxic activity (IC <sub>50</sub> ) <sup>a</sup> (mg/mL)					
			VERO	NIH3T3	AGS	HT-29	MCF-7	MDA-MB-231
<i>Aegiceras corniculatum</i>	Fruit	5.7	<b>0.150</b>	<b>0.097</b>	<b>0.0005</b>	<b>0.998</b>	<b>0.091</b>	<b>0.461</b>
<i>Argyrea nervosa</i>	Leaf	5.0	>2.5	>2.5	2.20	>2.5	>2.5	>2.5
<i>Avicennia alba</i>	Leaf	4.7	>2.5	>2.5	>2.5	>2.5	1.17	1.34
<i>Caesalpinia pulcherrima</i>	Leaf	8.3	>2.5	>2.5	>2.5	>2.5	2.40	1.15
<i>Clerodendrum viscosum</i>	Leaf	4.3	>2.5	2.03	2.23	<b>0.88</b>	<b>0.05</b>	1.69
<i>Clitoria ternatea</i>	Flower	12.4	>2.5	>2.5	>2.5	>2.5	1.14	<b>0.11</b>
<i>Clitoria ternatea</i>	Leaf	11.3	>2.5	>2.5	>2.5	>2.5	1.70	<b>0.49</b>
<i>Dillenia indica</i>	Leaf	8.9	>2.5	>2.5	1.18	>2.5	<b>0.34</b>	<b>0.54</b>
<i>Diospyros peregrina</i>	Leaf	5.9	>2.5	>2.5	1.58	>2.5	<b>0.007</b>	<b>0.33</b>
<i>Dipterocarpus turbinatus</i>	Bark	1.7	>2.5	>2.5	1.81	>2.5	1.68	<b>0.27</b>
<i>Dipterocarpus turbinatus</i>	Leaf	7.1	>2.5	>2.5	1.50	1.50	>2.5	<b>0.008</b>
<i>Ecbolium viride</i>	Leaf	13.3	>2.5	>2.5	>2.5	>2.5	<b>0.06</b>	1.40
<i>Glinus oppositifolius</i>	Whole plant	9.1	>2.5	>2.5	>2.5	>2.5	<b>0.15</b>	1.30
<i>Glycosmis pentaphylla</i>	Leaf	5.9	>2.5	>2.5	1.48	>2.5	>2.5	1.10
<i>Gnaphalium luteoalbum</i>	Leaf	1.9	>2.5	>2.5	<b>0.98</b>	>2.5	<b>0.34</b>	>2.5
<i>Hymenodictyon excelsum</i>	Bark	3.1	<b>0.23</b>	<b>0.07</b>	<b>0.09</b>	<b>0.16</b>	<b>0.08</b>	<b>0.44</b>
<i>Hymenodictyon excelsum</i>	Wood	4.5	>2.5	<b>0.18</b>	<b>0.58</b>	1.12	<b>0.72</b>	1.94
<i>Jasminum sambac</i>	Leaf	2.3	>2.5	>2.5	1.25	>2.5	<b>0.007</b>	>2.5
<i>Lannea coromandelica</i>	Bark	9.1	>2.5	>2.5	<b>0.09</b>	>2.5	<b>0.27</b>	<b>0.16</b>
<i>Lannea coromandelica</i>	Leaf	8.1	>2.5	>2.5	<b>0.67</b>	<b>0.52</b>	1.61	<b>0.71</b>
<i>Mussaenda glabrata</i>	Leaf	6.4	>2.5	>2.5	1.15	>2.5	1.33	0.15
<i>Myrica nagi</i>	Leaf	19.8	1.77	>2.5	<b>0.02</b>	>2.5	1.72	>2.5
<i>Saraca asoca</i>	Leaf	4.7	>2.5	>2.5	2.22	>2.5	>2.5	<b>0.40</b>
Cycloheximide	Positive control		ND	0.0003	0.0010	0.0036	0.061	0.0004

Highly cytotoxic results (IC<sub>50</sub> < 1 mg/mL) are displayed in bold

ND not determined

<sup>a</sup> IC<sub>50</sub> (inhibition of cell growth 50 %) calculated by Probit analysis (LdP Line software, USA), each experiment performed in triplicate

### High non-selective cytotoxicity

The most toxic extracts among all extracts tested were the those from *A. corniculatum* (fruit) and *H. excelsum* (bark) which showed high cytotoxicity across all cell lines tested (IC<sub>50</sub> values ranging from 0.0005 to 0.9980 and 0.07 to 0.44 mg/mL, respectively). Notably, high cytotoxicity (IC<sub>50</sub> = 0.5 µg/mL) against AGS was shown by *A. corniculatum* fruit which is 2 times higher than the cytotoxicity (IC<sub>50</sub> = 1.0 µg/mL) exhibited by the positive control, cycloheximide. We have previously reported on the cytotoxicity of *A. corniculatum* bark extract against NIH3T3, HT29, AGS and MDA-MB-435S cell lines [8], but no previous reports were found on the cytotoxicity evaluation on the fruits of this plant. Cytotoxic activity study of *A. corniculatum* bark [8] showed potent cytotoxicity against NIH3T3, HT-29, MDA-MB-435S (IC<sub>50</sub> values of 0.02, 0.33 and 0.66 mg/mL, respectively) but no cytotoxicity was exhibited against AGS cancer cells. In contrast, in our

study, highly significant cytotoxicity was shown by *A. corniculatum* fruit extract against AGS cancer cells (IC<sub>50</sub> = 0.5 µg/mL) and this extract also showed significant cytotoxic potential against VERO, NIH3T3, AGS, HT-29, MCF-7 and MDA-MB-231 cancer cells (IC<sub>50</sub> values of 0.150, 0.097, 0.0005, 0.998, 0.091, 0.461 mg/mL, respectively). Pentacyclic triterpenes such as maslinic acid, oleanolic acid and lupeol have been isolated previously from *A. corniculatum* bark [10–12]. Maslinic acid and lupeol have been reported to exert anticancer activity without effecting non-neoplastic cell lines via the inhibition of NF-κB activity [13, 14]. Another report indicates the apoptosis-inducing effect of oleanolic acid on colon cancer cells (HT-29) [15]. In addition, resveratrol, a natural phytoalexin isolated from this plant [12] as well as grapes, affects the cell cycle of cancer cells through inhibition of protein kinase C and D activities [16]. Resveratrol also induces apoptosis and decreases the activity of the transcription factors NF-κB and AP-1 [16]. Some of these

compounds could be responsible for the significant cytotoxic activity displayed by *A. corniculatum* fruit extract; however, it is still unknown if and to what extent these constituents are also present in the fruits of this plant.

Coumarin and its glycosides, such as scopoletin, aesculin and hymexelsin, have been identified and isolated from the bark of *H. excelsum* [17, 18]. A variety of coumarins are reported to have profound anticancer and anti-proliferative potential, thus displaying inhibitory effects against several tumour cells lines in vitro and in vivo [19–22]. Thus, the high cytotoxicity of bark extract of *H. excelsum* could be due, at least in part, to the presence of coumarins. Furthermore, *H. excelsum* wood extract displayed high non-selective antiproliferative activity ( $IC_{50} < 1.0$  mg/mL) against NIH3T3, AGS and MCF-7 cancer cells in our study. At this stage, however, it is not clear which constituents are present in the wood extract and may be responsible for the detected activity.

#### High selective cytotoxicity

Fourteen methanol extracts from 11 plants, namely *C. ternatea* (flower and leaf extracts), *D. indica* (leaf), *D. peregrina* (leaf), *D. turbinatus* (bark and leaf), *E. viride* (leaf), *G. luteoalbum* (leaf), *G. oppositifolius* (whole plant), *J. sambac* (leaf), *L. coromandelica* (bark and leaf), *M. glabrata* (leaf) and *S. asoca* (leaf) showed high selective cytotoxic potential ( $IC_{50} < 1.0$  mg/mL) against a minimum of one cancer cell line tested in this study. Among these 11 medicinal plants, 6 (*A. alba*, *D. peregrina*, *E. viride*, *J. sambac*, *C. ternatea* and *S. asoca*) have traditionally been used to treat cancer [4, 23, 24].

#### High selective cytotoxicity against breast cancer cells

It is worth noting that *A. alba* (leaf), *C. ternatea* (flower and leaf), *C. pulcherrima* (leaf), *E. viride* (leaf) and *G. oppositifolius* (whole plant) showed selective cytotoxicity only against both of the breast cancer cell lines (MCF-7 and MDA-MB-231). *C. ternatea* flower and leaf extracts exhibited high cytotoxic activity ( $IC_{50} < 1.0$  mg/mL) against MDA-MB-231, in addition *E. viride* leaf and *G. oppositifolius* whole plant extracts also showed high activity but against MCF-7 cancer cells ( $IC_{50}$  values of 0.06 and 0.15 mg/mL, respectively). It is also notable that the cytotoxicity of *E. viride* leaf extract was similar to that of the positive control, cycloheximide against the MCF-7 cancer cells ( $IC_{50}$  values of 0.060 and 0.061 mg/mL, respectively). To date no reports exist on cytotoxicity studies for *A. alba*, *C. ternatea*, *E. viride* and *G. oppositifolius* extracts. Although the cytotoxic activity of aerial parts and wood extracts of *C. pulcherrima* [25, 26] as well as anticancer activity of isolated compounds from this plant

(diterpenoid, 12-demethyl neocaesalpin F) against two cancer lines, HL-60 and HeLa, have been reported [27].

Extracts of *G. luteoalbum* and *J. sambac* leaf exhibited significant selective cytotoxicity only against MCF-7 (oestrogen-dependent breast cancer cells) ( $IC_{50}$  values of 0.34 and 0.007 mg/mL, respectively), whereas *D. turbinatus* and *S. asoca* leaf extracts showed potent cytotoxicity only against MDA-MB-231 (non-oestrogen-dependent breast cancer cells), ( $IC_{50}$  values of 0.008 and 0.40 mg/mL, respectively). Noticeably, the cytotoxicity exhibited by *D. peregrina* and *J. sambac* leaf extracts ( $IC_{50}$  values of 0.007 mg/mL for both) against MCF-7 cells was about 9 times higher than the cytotoxicity displayed by cycloheximide ( $IC_{50} = 0.061$  mg/mL). Interestingly, it is reported that *J. sambac* leaves have been traditionally used to treat breast cancer [4], whereas the flowers have reported antileukaemic activity against K562, P3HR1, Raji and U937 leukaemia cells [28]. Reports also state in vitro antiproliferative activity of *S. asoca* crude extract towards human tumour cell lines, including human erythromyeloid K562, B-lymphoid Raji, T-lymphoid Jurkat and erythroleukaemic HEL leukaemia. In vivo chemopreventive properties of *S. asoca* flower flavonoids on second-stage skin carcinogenesis are also reported [29, 30]. Neither cytotoxic nor anticancer activity studies of *D. turbinatus* and *G. luteoalbum* have been reported to date.

Different plant parts of *M. nagi*, *M. glabrata*, *J. sambac*, *D. peregrina*, *A. nervosa*, *S. asoca*, *L. coromandelica*, and *D. turbinatus* have been used traditionally as ulcer healing agents [4, 9]. Interestingly, we found high cytotoxic activity against gastric cancer cells (AGS) for *M. nagi* leaf extract ( $IC_{50} = 0.02$  mg/mL) and for *L. coromandelica* bark and leaf extracts ( $IC_{50}$  values of 0.090 and 0.67 mg/mL, respectively). Moreover, some gastric cancer cell cytotoxicity was detected for different plant part extracts of the remaining plants mentioned above; *M. glabrata* leaf ( $IC_{50} = 1.15$  mg/mL), *J. sambac* leaf ( $IC_{50} = 1.25$  mg/mL), *D. peregrina* leaf ( $IC_{50} = 1.58$  mg/mL), *D. turbinatus* bark and leaf ( $IC_{50}$  values of 1.81 and 1.50 mg/mL, respectively), *A. nervosa* leaf ( $IC_{50} = 2.20$  mg/mL) and *S. asoca* leaf ( $IC_{50} = 2.22$  mg/mL). Importantly, cytotoxic activity screening against gastric cancer cells of these plants had not been reported previously.

Moreover, in our study, 2 different parts of 4 plants such as flowers and leaves of *C. ternatea*, bark and leaves of *D. turbinatus*, bark and wood of *H. excelsum* and bark and leaves of *L. coromandelica* were screened for their cytotoxic activities to identify their effectiveness and selectivity against the cancer cells tested in our study. No significant difference in cytotoxic activity and selectivity was found in the case of flowers and leaf extracts of *C. ternatea*. But in the case of the bark and leaf extracts of *D. turbinatus*, the bark and wood extracts of *H. excelsum*, and the bark and

leaf extracts of *L. coromandelica* significant differences in the cytotoxicity as well as in selectivity against MCF-7, VERO and HT-29 cancer cells were exhibited.

Furthermore, *H. excelsum* wood exhibited no cytotoxicity ( $IC_{50} > 2.5$  mg/mL) to VERO, but displayed high cytotoxicity ( $IC_{50} = 0.18$  mg/mL) against NIH3T3 indicating a different cytotoxic potential against various healthy cell lines.

Of note, *A. nervosa* leaf extract showed very low cytotoxicity ( $IC_{50} > 2.0$  mg/mL) against all cell lines tested in this study.

Phyto-constituents such as polyphenols, flavonoids and catechins have long been recognised as having potential anticancer, anti-inflammatory, antioxidant and antimicrobial properties [31–33]. Previous research reports exist on the isolation of flavonoids from different parts of *A. nervosa*, *A. corniculatum*, *C. pulcherrima*, *C. viscosum*, *D. indica*, *D. peregrina*, *G. luteoalbum*, *G. oppositifolius*, *J. sambac*, *L. coromandelica* and *M. nagi* [12, 34–44]. It is likely that these constituents are also associated with the anticancer activity observed in this study [8, 26].

## Conclusion

The 9 plant species showing significant cytotoxic activity in this study have all been used traditionally as antitumour/anticancer agents. Seven (*A. alba*, *C. pulcherrima*, *D. peregrina*, *E. viride*, *J. sambac*, *C. ternatea* and *S. asoca*) of these plants showed high selective cytotoxic properties against at least one of the tested cancer cells, but not against the two healthy cell lines. The remaining 2 species (*A. corniculatum* fruit and *H. excelsum* bark) showed the highest, but non-selective cytotoxicity overall. These results lend support for the traditional use of the ‘active’ plants as anticancer agents. Further work will focus on isolation and characterisation of the cytotoxic and other bioactive constituents.

**Acknowledgments** Authors are thankful to Griffith University, Australia for providing a PhD scholarship to Raushanara Akter.

## References

- Al-Kalaldeh JZ, Abu-Dahab R, Afifi FU (2010) Volatile oil composition and antiproliferative activity of *Laurus nobilis*, *Origanum syriacum*, *Origanum vulgare*, and *Salvia triloba* against human breast adenocarcinoma cells. *Nutr Res* 30:271–278
- Mukherjee AK, Basu S, Sarkar N, Ghosh AC (2001) Advances in cancer therapy with plant based natural products. *Curr Med Chem* 8:1467–1486
- FAO (2004) Trade in medicinal plants. Economic and Social Department, Food, and Agriculture Organization of the United Nations, Rome, pp 2–3
- Ghani A (2003) Medicinal plants of Bangladesh with chemical constituents and uses, 2nd edn. Asiatic Society of Bangladesh, Dhaka
- Rahman S, Hasnat A, Hasan CM, Rashid MA, Ilias M (2001) Pharmacological evaluation of Bangladeshi medicinal plants—a review. *Pharm Biol* 39:1–6
- Costa-Lotuf LV, Khan Mahmud TH, Ather A, Wilke Diego V, Jimenez Paula C, Pessoa C, de Moraes Maria EA, de Moraes MO (2005) Studies of the anticancer potential of plants used in Bangladeshi folk medicine. *J Ethnopharmacol* 99:21–30
- George S, Bhalerao SV, Lidstone EA, Ahmad IS, Abbasi A, Cunningham BT, Watkin KL (2010) Cytotoxicity screening of Bangladeshi medicinal plant extracts on pancreatic cancer cells. *BMC Complement Altern Med* 10:52
- Uddin SJ, Grice ID, Tiralongo E (2011) Cytotoxic effects of Bangladeshi medicinal plant extracts. *eCAM* 2011:1–7. doi: 10.1093/ecam/nep111
- Bandaranayake WM (1998) Traditional and medicinal uses of mangroves. *Mangroves Salt Marshes* 2:133–148
- Gowri PM, Rao AHCVHA, Saidulu R, Shusma S, Madhu A, Raju TV (2012) Unusual isomeric corniculatolides from Mangrove, *Aegiceras corniculatum*. *J Nat Prod* 75:275–279
- Xu J-j, Long S-j (2009) Study on chemical constituents from *Aegiceras corniculatum*. *Huaxi Yaoxue Zazhi* 24:120–123
- Wang JD, Dong ML, Zhang W, Shen X, Guo YW (2006) Chemical components of Mangrove plant *Aegiceras corniculatum*. *Zhongguo Tianran Yaowu* 4:275–277
- Lee TK, Poon RTP, Wo JY, Ma S, Guan XY, Myers JN, Altevogt P, Yuen APW (2007) Lupeol suppresses cisplatin-induced nuclear factor-Kb activation in head and neck squamous cell carcinoma and inhibits local invasion and nodal metastasis in an orthotopic nude mouse model. *Cancer Res* 67:8800–8809
- Li C, Yang Z, Zhai C, Qiu W, Li D, Yi Z (2010) Maslinic acid potentiates the anti-tumor activity of tumor necrosis factor A by inhibiting NF-κB signaling pathway. *Mol Cancer* 9:73
- Juan ME, Planas JM, Ruiz-Gutierrez V, Daniel H, Wenzel U (2008) Antiproliferative and apoptosis-inducing effects of maslinic and oleanolic acids, two pentacyclic triterpenes from olives, on HT-29 colon cancer cells. *Br J Nutr* 100:36–43
- Androutsopoulos VP, Ruparella KC, Papakyriakou A, Filippakis H, Tsatsakis AM, Spandidos DA (2011) Anticancer effects of the metabolic products of the resveratrol analogue, Dmu-212: structural requirements for potency. *Eur J Med Chem* 46:2586–2595
- Rao PS, Asheervadam Y, Khaleelullah M, Rao NS, Murray RDH (1988) Hymexelsin, an apiose-containing scopoletin glycoside from the stem bark of *Hymenodictyon excelsum*. *J Nat Prod* 51:959–961
- Gibson CS, Simonsen JL (1918) The constituents of the bark of the *Hymenodictyon excelsum*. *J Chem Soc* 114:151–152
- Ruiz-Marcial C, Reyes CR, Estrada E, Reyes-Esparza J, Garrido Farina G, Rodriguez-Fragoso L (2007) Antiproliferative, cytotoxic and antitumour activity of coumarins isolated from *Calophyllum brasiliense*. *J Pharm Pharmacol* 59:719–725
- Kawase M, Sakagami H, Motohashi N, Hauer H, Chatterjee SS, Spengler G, Vigyikanne AV, Molnar A, Molnar J (2005) Coumarin derivatives with tumor-specific cytotoxicity and multidrug resistance reversal activity. *In Vivo* 19:705–711
- Ishihara M, Yokote Y, Sakagami H (2006) Quantitative structure-cytotoxicity relationship analysis of coumarin and its derivatives by semiempirical molecular orbital method. *Anticancer Res* 26:2883–2886
- Yang H, Protiva P, Gil RR, Jiang B, Baggett S, Basile MJ, Reynertson KA, Weinstein IB, Kennelly EJ (2005) Antioxidant and cytotoxic isoprenylated coumarins from *Mammea americana*. *Planta Med* 71:852–860

23. Bandaranayake WM (2002) Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetlands Ecol Manage* 10:421–452
24. Devmurari V, Shivanand P, Goyani MB, Vaghani S, Jivani NP (2009) A review: *Carissa congesta*: phytochemical constituents, traditional use and pharmacological properties. *Pharmacogn Rev* 3:375–377
25. Chanda S, Baravalia Y (2011) Brine shrimp cytotoxicity of *Caesalpinia pulcherrima* aerial parts, antimicrobial activity and characterisation of isolated active fractions. *Nat Prod Res* 25:1955–1964
26. Pawar CR, Mutha RE, Landge AD, Jadhav RB, Surana SJ (2009) Antioxidant and cytotoxic activities of *Caesalpinia pulcherrima* wood. *Indian J Biochem Biophys* 46:198–200
27. Das B, Srinivas Y, Sudhakar C, Mahender I, Laxminarayana K, Reddy PR, Raju TV, Jakka NM, Rao JV (2010) New diterpenoids from *Caesalpinia* species and their cytotoxic activity. *Bioorg Med Chem Lett* 20:2847–2850
28. Chiang L-C, Cheng H-Y, Liu M-C, Chiang W, Lin C-C (2004) *In vitro* evaluation of antileukemic activity of 17 commonly used fruits and vegetables in Taiwan. *Lebensm-Wiss Technol* 37: 539–544
29. Dabur R, Gupta A, Mandal TK, Singh DD, Bajpai V, Gurav AM, Lavekar GS (2007) Antimicrobial activity of some Indian medicinal plants. *Afr J Tradit Complem* 4:313–318
30. Cibin TR, Devi DG, Abraham A (2012) Chemoprevention of two-stage skin cancer in vivo by *Saraca asoca*. *Integr Cancer Ther* 11:279–286
31. Ren WY, Qiao ZH, Wang HW, Zhu L, Zhang L (2003) Flavonoids: promising anticancer agents. *Med Res Rev* 23:519–534
32. Gonzalez-Mejia ME, Voss OH, Murnan EJ, Doseff AI (2010) Apigenin-induced apoptosis of leukemia cells is mediated by a bimodal and differentially regulated residue-specific phosphorylation of heat-shock protein-27. *Cell Death Dis* 1:64
33. Park KD, Lee SG, Kim SU, Kim SH, Sun WS, Cho SJ, Jeong DH (2004) Anticancer activity of 3-O-acyl and alkyl(-)-epicatechin derivatives. *Bioorg Med Chem Lett* 14:5189–5192
34. Prakash S, Gautam S, Ashok K (2011) Phytochemical screening of selected medicinal plants used in skin diseases. *Int J Pharm Pharm Sci* 3:1402–1406
35. Das B, Thirupathi P, Ravikanth B, Kumar RA, Sarma AVS, Basha SJ (2009) Isolation, synthesis, and bioactivity of homio-soflavonoids from *Caesalpinia pulcherrima*. *Chem Pharm Bull* 57:1139–1141
36. Srinivas KVNS, Koteswara RY, Mahender I, Das B, Rama Krishna KVS, Hara Kishore K, Murty USN (2003) Flavanoids from *Caesalpinia pulcherrima*. *Phytochem* 63:789–793
37. Roy R, Pathak NKR, Biswas M, Pandey VB (1994) Flavonoids of *Clerodendron infortunatum*. *Orient J Chem* 10:169–170
38. Pavanasisavan G, Sultanbawa MUS (1975) Flavonoids of some Dilleniaceae species. *Phytochem* 14:1127–1128
39. Sahu R, Dewanjee S, Dua TK, Gangopadhyay M, Das AK, Dey SP (2012) Dereplication coupled with in vitro antioxidant assay of two flavonoid glycosides from *Diospyros peregrina* fruit. *Nat Prod Res* 26:454–459
40. Morimoto M, Kumeda S, Komai K (2000) Insect antifeedant flavonoids from *Gnaphalium affine* D. Don. *J Agric Food Chem* 48:1888–1891
41. Singh BP, Singh RP, Jha OP (1982) Flavonoids of some Aizoaceae and Molluginaceae of Bhagalpur. *Biol Bull India* 4:157–163
42. Liu H, Ni W, Yuan M, Chen C (2004) Chemical constituents of *Jasminum sambac*. *Yunnan Zhiwu Yanjiu* 26:687–690
43. Yun X-j, Shu H-m, Ji M-h, Chen G-y, Chen M-l (2012) Studies on the chemical constituents from the bark of *Lannea coromandelica*. *Huaxue Yanjiu Yu Yingyong* 24:610–613
44. Yang W, C-m Tang, Li X, Zhou Y, Wang L, Li L (2011) Study on the chemical constituents of *Myrica esculenta*. *Yunnan Daxue Xuebao Ziran Kexueban* 33:453–457