

Impact of Prenatal Immune System Disturbances on Brain Development

Amrita Madhusudan · Prisca Vogel · Irene Knuesel

Received: 10 January 2012 / Accepted: 27 April 2012 / Published online: 13 May 2012
© Springer Science+Business Media, LLC 2012

Abstract As research into various aging-associated neurodegenerative disorders reveals their immense pathophysiological complexity, the focus is currently shifting from studying changes in an advanced disease state to investigations involving pre-symptomatic periods, possible aberrations in early life, and even abnormalities in brain development. Recent studies on the etiology of schizophrenia and autism spectrum disorders revealed a profound impact of neurodevelopmental disturbances on disease predisposition, onset and progression. Here, we discuss how a prenatal immune challenge can affect the developing brain—with a selective focus on the impact on microglia, the brain’s immune cells—and the implications for brain aging and its associated risk of developing Alzheimer’s disease.

Keywords Polyriboinosinic-polyribocytidilic acid · PolyI:C · Late gestation · Infection · Neuronal health · Brain aging

Microglia in development

The immune system has developed to ensure cells of the innate immune system reside in all organs of the body. The brain, far from being a completely immune-privileged site, as thought earlier, possesses its own immune cells (Ransohoff and Perry 2009). These are the microglia, and are the only cells in the central nervous system (CNS) that do not originate from the neuroectoderm (Allen and Barres 2009), but belong to the

myeloid lineage. After some debate about how these cells populate the CNS—either via entering the developing brain from peripheral circulation as monocytes followed by differentiation (Perry et al. 1985), or multiplying from intraparenchymal precursors that existed before the development of the vasculature (Alliot et al. 1999), recent studies provided supporting evidence that microglia derive from macrophages that originate from the yolk-sac and then colonize the brain (Herbomel et al. 2001; Ginhoux et al. 2010). Still, what percentage of microglia in the adult pool are remnants from these developmental precursors is unknown. While the predominant view is that microglia are long-lived cells that are not often replenished from the periphery, different studies have shown that, under certain conditions, peripheral bone-marrow derived mononuclear phagocytes (BMDPs) can integrate into the CNS (Priller et al. 2001; Massengale et al. 2005; Prinz and Mildner 2011). Moreover, differential expression of certain markers, for instance, *Hoxb8*, seem to indicate that at least two different microglial populations exist in the adult mouse brain (Chen et al. 2011), which may have different implications for their function.

Once the microglia colonize the brain, several studies (Pont-Lezica et al. 2011) have shown that contrary to being randomly and evenly distributed throughout the brain, these cells appear to concentrate specifically at certain locations, either being associated with apoptotic neurons and radial glial cells, for example, or being excluded from others, like the lens and photoreceptor layer of the retina. The distribution of microglial cells in the brain show region-specific differences with highest density in the hippocampus, basal ganglia, olfactory telencephalon and substantia nigra, followed by cerebral cortex, thalamus and hypothalamus, and lowest density in cerebellum brain stem areas, and myelinated fibre tracts (Lawson et al. 1990; Lawson et al. 1993). Moreover, there even seems to be a variation in where the majority of the morphologically distinct microglia generally reside. Those with the shortest processes

A. Madhusudan · P. Vogel · I. Knuesel (✉)
Institute of Pharmacology and Toxicology, University of Zurich,
Winterthurerstrasse 190,
8057 Zurich, Switzerland
e-mail: knuesel@pharma.uzh.ch

are found in the circumventricular regions, while the most ramified microglia are found in the grey matter (Lawson et al. 1990). This region-specific density, which varies around five-fold, as well as the differences in morphology, appears to be closely linked to the vast number of functions performed by microglia.

During development, they play significant roles in the formation of a healthy working environment in the CNS, and accomplish this in various ways. First, microglia are found in close association with cells undergoing developmental cell death, and have been shown to be capable of phagocytosing apoptotic neurons (Peri and Nusslein-Volhard 2008). They have also been shown to control which cells undergo apoptosis, via NGF signaling or triggering oxidative stress (Frade and Barde 1998; Wakselman et al. 2008). Large microglial populations have also been discovered free of apoptotic cells, and these are thought to have other functions. Those in close association with the developing vasculature have been suggested to play a key role in angiogenesis, as evidenced by the underdeveloped vascular system in mice lacking microglia (Rymo et al. 2011). Others are associated with developing axon tracts throughout the brain (Herbomel et al. 2001), at the right time to influence neurite development and axon remodeling, as well as synaptic pruning in development (Paolicelli et al. 2011). Microglia have been shown to contain large inclusions that often resemble axon terminals and even spines (Tremblay 2012), and could contribute to experience-dependent remodeling of synapses and developing circuits. In line, members of the complement system, often derived from microglia, have also been shown to be involved in synapse elimination (Stevens et al. 2007). Recent research also showed that mice lacking CX3CR1 (a microglia-specific chemokine receptor) show significant impairments in synaptic development as compared to wild-type littermates (for recent review, Ransohoff and Perry 2009).

Microglia functions in the adult brain

The close and tightly regulated interaction between microglia and neurons during brain development is maintained in the adult brain. This has been highlighted by several studies recently summarized by Tremblay and colleagues (2011), indicating that microglia are actively involved in the remodeling of the perisynaptic environment and fine-tuning of neuronal activity, likely also involving feedback signals provided by the neurons to modulate the state of activity in microglia (Tremblay et al. 2011). This bidirectional communication is achieved through interactions between cell-surface receptors and cell-surface-bound ligands (i.e. CD200-CD200 receptors), or at a distance after the generation of diffusible ligands, as well as via neurotransmission-associated inhibition involving dopamine and noradrenaline and their respective receptors

(for recent review, Ransohoff and Cardona 2010). In turn, through the release of a multitude of soluble factors including neurotrophic factors chemokines, cytokines, microglia not only perform classical glia functions, but also accomplish a crucial immune function by surveillance of the brain for damage infection, and engulfing infectious agents, dead cells and debris (Streit 2006; Streit and Xue 2009).

At least two distinct configurations have been observed—highly branched with long, fine processes, or rounded and amoeboid. The former were considered to be “resting microglia” that occur in a healthy, undisturbed CNS. This term later proved to be a misnomer, since these cells were shown to be highly dynamic, forming networks with their processes, while their soma remain more or less stationary, that probe and test the entire brain in a matter of a few hours (Nimmerjahn et al. 2005; Schlegelmilch et al. 2011; Wrenfeldt et al. 2011). The onus is to designate these microglia more appropriately as “surveying microglia” (Hanisch and Kettenmann 2007; Marin-Teva et al. 2012). The amoeboid cells are indeed more active in the immune sense, with the surveying glia transforming to this phenotype, possessing upregulated antigen presenting molecules, when phagocytosis is required. This rapid conversion to the amoeboid form is termed “activation of microglia” and can be triggered by a variety of stimuli, ranging from infection and trauma to ischemia or any disturbance in the normal functioning of the brain (Kettenmann et al. 2011). The abrupt response to neuronal injury could also be related to a disrupted cell-cell contact and the concomitant loss of neuron-mediated inhibition of microglia (for recent review, Ransohoff and Cardona 2010). Whether these properties also relate to the observed differences in the basal activation state of microglia in white versus grey matter (Carson et al. 2007) remains to be determined. However, it is conceivable that these functional differences are linked to the expression profile of cell surface markers, through which microglia are not only able to control distinct population of neurons but also modulate a variety of cellular and immune functions (for recent review, Lynch 2009). For instance, microglia can promote the infiltration of circulating cells into the CNS (upregulation of CD11b, an integrin family member which pairs with CD18 to form the complement receptor 3), can be non-phagocytic and produce proinflammatory molecules (CD40), or be phagocytic and motile (MHCII), as shown for the highly effective clearance of apoptotic neurons. In the adult brain, microglia are responsible for all apoptotic debris, including the majority of new born neurons (Sierra et al. 2010; Graeber and Streit 2010).

Based on morphology, it is thus possible to differentiate microglia into different states including amoeboid (surveillance) and ramified (activated) forms. A third potential state has recently been hypothesized, by postulating that microglia can devolve from their “surveying” state into an “alerted” one, before becoming fully activated. This resembles the process of priming of macrophages in the periphery,

where activation occurs in two steps—priming, followed by triggering (Dilger and Johnson 2008). A “primed” microglial cell would thus react to a greater degree than a non-primed one. In line with these hypothesis, once activated, microglia tend not to return all the way to their “surveying” state, thereby eliciting elevated responses to the next stimulus (Schwartz et al. 2006). Microglia differ, however, from peripheral immune cells in their preference for humoral responses rather than eliciting the classical cytotoxic responses, which involves the ingestion of antigens, exiting the tissue and entering the draining lymph nodes, where stimulation of naive T cells occurs. This happens perforce due to the nature of the CNS environment—where the neurons cannot regenerate as easily as cells in the periphery, and collateral damage cannot be repaired.

Microglia in disease and aging

Several diseases, not only those of the CNS, appear to impact on microglial activity and function. Infectious diseases of the CNS (Mariani and Kielian 2009), conditions of neuropathic pain (Ji and Suter 2007), even air pollution (Block and Calderon-Garciduenas 2009) can create environments where microglia become highly activated, and after a point, cannot protect the CNS any more. Looking at two neurodegenerative diseases—multiple sclerosis (MS) and traumatic spinal cord injury (SCI)—the role of microglia in repairing the damage is highlighted. In cases of MS or other demyelinating conditions, it has been shown that there exists a subset of microglia at the site of demyelination that can recruit oligodendrocyte precursor cells and phagocytose damaged neurons (Olah et al. 2011). In SCI, cells of the myeloid origin have been shown to play a role in intraspinal trafficking past the injury (Hawthorne and Popovich 2011).

Beyond diseases, aging also represents a critical “primer” of microglia activity, and is also accompanied by microglial dystrophy and degeneration, a feature that appears to be accompanied by diminution of microglial neuroprotective functions and hypothesized to significantly contribute to neurodegenerative changes characteristic of AD (Streit 2004). A combination of microglia senescence on one side and priming-related hyperactivity on the other side is therefore expected to have a detrimental influence on neuronal health. This represents a highly relevant theory regarding AD pathogenesis because it integrates aging as the most important risk factor of this progressive neurodegenerative disease. However, the molecular mechanisms underlying aging-associated microglia dysfunction are largely unknown. Besides a decline in proteasome activity and other aging-associated impairments in protein homeostasis (Stolzing and Grune 2003), limited information is available regarding the lifespan of microglia (Streit 2006). It has been shown that nerve injury induces—

beside recruitment of bone marrow-derived precursors that can infiltrate the brain—a burst in mitotic activity and a wave of proliferation that is followed by apoptotic cell death of microglia (Graeber et al. 1988; Gehrman and Banati 1995; Lassmann et al. 1995). This represents an essential defense and wound healing mechanism shown to be maintained in aged rodents (Conde and Streit 2006) and suggested to be required to keep the number of these endogenous immune defenders of the CNS constant. It is conceivable, therefore, that repetitive exposures to brain injury and inflammation induces replicative senescence, eventual loss of mitotic activity and degeneration of microglia during aging. Evidence for this hypothesis has been provided by *in vitro* findings showing that microglia stimulation by mitogens induces telomere shortening (Flanary and Streit 2004). Recent data also showed that interferon- γ exposure is sufficient to trigger activation-induced microglia cell death (Yun et al. 2011), in line with postmortem investigations of AD patients showing caspase activity in plaque-associated microglia in AD (Yang et al. 1998).

Specifically with respect to AD, the significance between the association of microglia with amyloid plaques has long been debated. Several studies have looked at the possibility of using the phagocytic ability of microglia therapeutically to clear the plaques, but largely, this doesn't seem to have the desired results on cognition. It has further been observed that plaque-associated microglia still possess their processes at the interface, and are capable of phagocytosing A β peptides (Bolmont et al. 2008). However, it has also been demonstrated that microglia show an age-dependent reduction in their ability to clear A β peptides, despite preservation of functional phagocytic activity and even adherence to A β plaques (Floden and Combs 2011), underscoring the importance of aging for microglial health and activity.

Prenatal PolyI:C model of accelerated aging

Interestingly, degeneration of microglia has also been reported in neurological disorders without major neurodegenerative changes, such as schizophrenia, (Wierzbica-Bobrowicz et al. 2004), or associations with aging. Besides genetic risk factors, maternal infection during pregnancy represents one of the currently most extensively studied risk factor in experimental schizophrenia research. This is based on a large body of epidemiological data that has provided compelling evidence for enhanced risk of schizophrenia following prenatal exposure to infection with various viral or bacterial pathogens, or genital and/or reproductive infections (Brown et al. 2006; Patterson 2007; Boksa 2008). This data indicates that the early inflammatory insult not only strongly affects the developing nervous system and predisposes the offspring to psychosis-related behavioural abnormalities in adulthood, but likely also critically influences the

proliferation, maturation and function of the developing immune system. The findings of significantly elevated levels of degenerating microglia seen in frontal and temporal cortices of schizophrenia patients compared to healthy subjects (Wierzbica-Bobrowicz et al. 2004) indeed supports the hypothesis that early prenatal infection-induced expression of pro-inflammatory cytokines and other mediators of the innate immunity might reduce the proliferative capacity of microglia and hence reduce their lifespan.

To investigate the impact of a prenatal infection on brain and immune system development, we established a mouse model that is based on the critical contribution of the gestational stage during which the maternal immune challenge occurs. It builds on findings reported by other groups that demonstrated a wide spectrum of behavioural, neuroanatomical, and neurochemical changes following exposure to viral or bacterial pathogens in rats and mice that mimicked critical phenotypic symptoms of schizophrenia (for recent review, Meyer et al. 2009). Based on the highly comparable phenotype induced by the various types of pathogens we reasoned that the downstream effects, i.e. the production of pro-inflammatory cytokines might be the critical common denominator of the behavioural and neurochemical changes described. To test this, we employed a viral mimic, Polyriboinosinic-polyribocytidilic acid (PolyI:C), a double-stranded RNA shown to bind to Toll-like receptor 3 (TLR3), leading to activation of nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) and production of pro-inflammatory cytokines such as TNF- α , IL-6 and IL-12 (Alexopoulou et al. 2001; Okun et al. 2011). Given the critical processes during the development of the brain and its resident immune cells, we decided to contrast early, mid- and late-gestational stages. We hypothesize that the developmental stage critically determines the phenotype of the adult offspring, allowing us to test whether prenatal infections may—in addition to the widely accepted role in the etiology of neuropsychiatric diseases—also contribute to brain aging and neurodegenerative diseases.

Our first findings indeed confirmed that the time of maternal immune challenge critically determines the patterns of fetal immunological, juvenile neurochemical, as well as adult behavioural abnormalities displayed by the offspring (Meyer et al. 2006). While the PolyI:C challenge during early-gestation (gestational day 6, GD6) resulted in a high incidence of abortion, a viral-like infection during mid-gestation (GD9) confirmed the precipitation of psychosis-related symptoms in the adult offspring, in line with previous findings by other groups (Borrell et al. 2002; Shi et al. 2003; Zuckerman et al. 2003; Zuckerman and Weiner 2005; Ozawa et al. 2006; Romero et al. 2007). A late-gestational (GD17) immune challenge resulted in a behavioural phenotype characterized by cognitive impairments, elevation of fetal IL6, IL10, and TNF α , enhanced apoptosis in the dentate gyrus, as well as a





Fig. 1 Schematic representation of the critical differences resulting from a prenatal immune challenge at different time periods in gestation

decline in Reelin-positive cells in juvenile brains (Fig. 1). We showed for the first time that the fetal brain can directly contribute to the specific changes in inflammatory cytokine protein levels after late-gestational but not mid-gestational immune challenge, suggesting a close correlation between acute immunological modulators and histopathological changes in the juvenile offspring. We also reported that the mid- and late-gestational infections had a differential impact on the developing neurotransmitter systems (Meyer et al. 2008). Using immunohistochemical approaches and unbiased stereological evaluations, we provided first evidence that a prenatal immune challenge around mid-gestation has a stronger impact on the prefrontal dopamine receptor D1 expression as compared to the treatment effect at GD17. These neurochemical alterations were accompanied with behavioural abnormalities with relevance to positive symptom clusters characteristic of schizophrenia. On the other hand, an in utero immune challenge at GD17 preferentially affected the expression levels of the hippocampal NR1 subunit of the NMDA receptor as compared to GD9 and control treatments, which were paralleled by impairments in working memory performance. These studies revealed also that a prenatal immune challenge during late gestation resulted in acceleration of aging, as indicated by the earlier formation of Reelin-positive extracellular amyloid-like deposits, a neuropathological feature detected in the hippocampal formation in several

Fig. 2 Glial cells in prenatal astrocyte cultures undergo morphological changes upon immune stimulation. **a–b**, Immunofluorescence staining of astrocytes (mouse anti-GFAP, 1:5000, Dako, Z334; green) and microglia (rabbit anti-Iba1, 1:4000, Wako Nr. 019–19741; red) in astrocyte cultures prepared from E18 rats that were exposed at day 3 to PolyI:C (200 μ g/ml, P9582, Sigma), vehicle (0.9 % pyrogen-free NaCl, B. Braun, 534534), or left undisturbed. The nuclei were counterstained with DAPI (Thermo Scientific Pierce, Switzerland; blue). Triple-immunofluorescence labelings of the paraformaldehyde-fixed cells (at day 7 in vitro) were visualized by confocal microscopy using sequential acquisition of separate channels. Z-stacks of consecutive optical sections were summed and projected in the z dimension (maximal intensity) and merged using the image analysis software Imapis (Bitplane, Zurich, Switzerland) for visual display. Note that the immune challenge with PolyI:C altered the ramified phenotype found in untreated and NaCl-treated cultures into an amoeboid-like appearance indicative of activated microglia. Scale bar: **a**: 50 μ m, **b**: high magnification image: 15 μ m



species that is accompanied by a reduction in Reelin-expressing neurons (Knuesel et al. 2009; Knuesel 2010). Both neuropathological alterations significantly correlated with hippocampus-dependent episodic-like memory deficits (Knuesel et al. 2009). We found that healthy and cognitively normal subjects were able to sufficiently clear and degrade these deposits, whereas cognitively impaired aged mice failed to do so. Our results showed for the first time that prenatal exposure to infection during late embryonic developmental stages is not only an important factor in the segregation of certain behavioural symptom clusters with relevance for neuropsychiatric disorders but does also represent a critical driving force of aging-related neuropathological processes.

Effects of PolyI:C in vitro and in vivo on microglia

These observations indicated that the effect of PolyI:C on microglia and astrocytes could be highly relevant for such neuropathology. Supporting this view, our studies with primary cultures of these cells revealed that PolyI:C could trigger the morphological changes underlying the transformation of microglia into their activated state, as compared to the ramified morphology of the untreated, or saline treated, surveying microglia, but not changes in astrocyte morphology (Fig. 2; Vogel et al., unpublished data). The implication here is that PolyI:C induces a strong and distinct immune response through the TLR3 found on microglia, but saline, a likely stressor, activates different cellular pathways, leading only to a mild or non-selective immune response.

The mRNA of different TLRs has previously been shown to be enriched in the vicinity of amyloid plaques (Frank et al. 2009), likely reflecting their localization on microglia. Our own observations in aged prenatally immune-challenged mice reveal a striking overlap between microglial and TLR3 immunoreactivity (Vogel et al., unpublished data). Besides establishing a chronic adverse neuronal environment, the prenatal immune challenge might prime microglia and thereby alter their ability to react to different stimuli and create an innate memory allowing a faster and exaggerated response upon a second challenge; changes highly comparable to mechanisms in the adaptive immune system. The strong glial response may then have devastating effects on surrounding neurons resulting in demyelination and apoptosis due to release of ROS.

New perspectives and future outlooks

Having seen how delicately balanced the immune system is in the CNS, it becomes more evident that inflammation at critical points during development can have a serious and long-lasting impact on microglial function, and thus, on how the brain

ages. Studying microglia during development and understanding the different aspects of their priming and activation could provide insights to disorders that set in much later in life, like AD. Everyone's goal of healthy aging seems thus to depend more on proper developmental processes than so far anticipated.

Acknowledgements This study was supported by the Swiss National Science Foundation, Grant Nr 310030–132629 (IK), and National Center for Competence in Research (NCCR project 2, Alzheimer's Disease, IK). The authors thank Prof. Jean-Marc Fritschy for his advice, support and inputs regarding this project. The authors also wish to thank all the people of the Animal Services of the Institute of Pharmacology and Toxicology for animal husbandry and care.

References

- Alexopoulou L, Holt AC, Medzhitov R, Flavell RA (2001) Recognition of double-stranded RNA and activation of NF-kappaB by toll-like receptor 3. *Nature* 413:732–738
- Allen NJ, Barres BA (2009) Neuroscience: glia—more than just brain glue. *Nature* 457:675–677
- Alliot F, Godin I, Pessac B (1999) Microglia derive from progenitors, originating from the yolk sac, and which proliferate in the brain. *Brain Res Dev Brain Res* 117:145–152
- Block ML, Calderon-Garciduenas L (2009) Air pollution: mechanisms of neuroinflammation and CNS disease. *Trends Neurosci* 32:506–516
- Boksa P (2008) Maternal infection during pregnancy and schizophrenia. *J Psychiatry Neurosci* 33:183–185
- Bolmont T, Haiss F, Eicke D, Radde R, Mathis CA, Klunk WE, Kohsaka S, Jucker M, Calhoun ME (2008) Dynamics of the microglial/amyloid interaction indicate a role in plaque maintenance. *J Neurosci* 28:4283–4292
- Borrell J, Vela JM, Arevalo-Martin A, Molina-Holgado E, Guaza C (2002) Prenatal immune challenge disrupts sensorimotor gating in adult rats. Implications for the etiopathogenesis of schizophrenia. *Neuropsychopharmacology* 26:204–215
- Brown AS, Schaefer CA, Quesenberry CP Jr, Shen L, Susser ES (2006) No evidence of relation between maternal exposure to herpes simplex virus type 2 and risk of schizophrenia? *Am J Psychiatry* 163:2178–2180
- Carson MJ, Bilousova TV, Puntambekar SS, Melchior B, Doose JM, Ethell IM (2007) A rose by any other name? The potential consequences of microglial heterogeneity during CNS health and disease. *Neurotherapeutics* 4:571–579
- Chen SK, Tvrdik P, Peden E, Cho S, Wu S, Spangrude G, Capecchi MR (2011) Hematopoietic origin of pathological grooming in Hoxb8 mutant mice. *Cell* 141:775–785
- Conde JR, Streit WJ (2006) Effect of aging on the microglial response to peripheral nerve injury. *Neurobiol Aging* 27:1451–1461
- Dilger RN, Johnson RW (2008) Aging, microglial cell priming, and the discordant central inflammatory response to signals from the peripheral immune system. *J Leukoc Biol* 84:932–939
- Flanary BE, Streit WJ (2004) Progressive telomere shortening occurs in cultured rat microglia, but not astrocytes. *Glia* 45:75–88
- Floden AM, Combs CK (2011) Microglia demonstrate age-dependent interaction with amyloid-beta fibrils. *J Alzheimers Dis* 25:279–293
- Frade JM, Barde YA (1998) Microglia-derived nerve growth factor causes cell death in the developing retina. *Neuron* 20:35–41
- Frank S, Copanaki E, Burbach GJ, Muller UC, Deller T (2009) Differential regulation of toll-like receptor mRNAs in amyloid

- plaque-associated brain tissue of aged APP23 transgenic mice. *Neurosci Lett* 453:41–44
- Gehrmann J, Banati RB (1995) Microglial turnover in the injured CNS: activated microglia undergo delayed DNA fragmentation following peripheral nerve injury. *J Neuropathol Exp Neurol* 54:680–688
- Ginhoux F, Greter M, Leboeuf M, Nandi S, See P, Gokhan S, Mehler MF, Conway SJ, Ng LG, Stanley ER, Samokhvalov IM, Merad M (2010) Fate mapping analysis reveals that adult microglia derive from primitive macrophages. *Science* 330:841–845
- Graeber MB, Streit WJ (2010) Microglia biology and pathology. *Acta Neuropathol* 119:89–105
- Graeber MB, Tetzlaff W, Streit WJ, Kreutzberg GW (1988) Microglial cells but not astrocytes undergo mitosis following rat facial nerve axotomy. *Neurosci Lett* 85:317–321
- Hanisch UK, Kettenmann H (2007) Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 10:1387–1394
- Hawthorne AL, Popovich PG (2011) Emerging concepts in myeloid cell biology after spinal cord injury. *Neurotherapeutics* 8:252–261
- Herbomel P, Thisse B, Thisse C (2001) Zebrafish early macrophages colonize cephalic mesenchyme and developing brain, retina, and epidermis through a M-CSF receptor-dependent invasive process. *Dev Biol* 238:274–288
- Ji RR, Suter MR (2007) p38 MAPK, microglial signaling, and neuropathic pain. *Mol Pain* 3:33
- Kettenmann H, Hanisch UK, Noda M, Verkhratsky A (2011) Physiology of microglia. *Physiol Rev* 91:461–553
- Knuesel I (2010) Reelin-mediated signaling in neuropsychiatric and neurodegenerative diseases. *Prog Neurobiol* 91:257–274
- Knuesel I, Nyffeler M, Mormede C, Muhia M, Meyer U, Pietropaolo S, Yee BK, Pryce CR, LaFerla FM, Marighetto A, Feldon J (2009) Age-related accumulation of reelin in amyloid-like deposits. *Neurobiol Aging* 30:697–716
- Lassmann H, Bancher C, Breitschopf H, Wegiel J, Bobinski M, Jellinger KA, Wisniewski HM (1995) Cell death in Alzheimer's disease evaluated by DNA fragmentation in situ. *Acta Neuropathol* 89:35–41
- Lawson LJ, Perry VH, Dri P, Gordon S (1990) Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience* 39:151–170
- Lawson LJ, Matyszak MK, Perry VH (1993) Lessons from microglia in special sites. *Clin Neuropathol* 12:310–313
- Lynch MA (2009) The multifaceted profile of activated microglia. *Mol Neurobiol* 40:139–156
- Mariani MM, Kielian T (2009) Microglia in infectious diseases of the central nervous system. *J Neuroimmune Pharmacol* 4:448–461
- Marin-Teva JL, Cuadros MA, Martin-Oliva D, Navascues J (2012) Microglia and neuronal cell death. *Neuron Glia Biol* 1:1–16
- Massengale M, Wagers AJ, Vogel H, Weissman IL (2005) Hematopoietic cells maintain hematopoietic fates upon entering the brain. *J Exp Med* 201:1579–1589
- Meyer U, Nyffeler M, Engler A, Urwyler A, Schedlowski M, Knuesel I, Yee BK, Feldon J (2006) The time of prenatal immune challenge determines the specificity of inflammation-mediated brain and behavioral pathology. *J Neurosci* 26:4752–4762
- Meyer U, Nyffeler M, Yee BK, Knuesel I, Feldon J (2008) Adult brain and behavioral pathological markers of prenatal immune challenge during early/middle and late fetal development in mice. *Brain Behav Immun* 22:469–486
- Meyer U, Feldon J, Fatemi SH (2009) In-vivo rodent models for the experimental investigation of prenatal immune activation effects in neurodevelopmental brain disorders. *Neurosci Biobehav Rev* 33:1061–1079
- Nimmerjahn A, Kirchhoff F, Helmchen F (2005) Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 308:1314–1318
- Okun E, Griffioen KJ, Mattson MP (2011) Toll-like receptor signaling in neural plasticity and disease. *Trends Neurosci* 34:269–281
- Olah M, Amor S, Brouwer N, Vinet J, Eggen B, Biber K, Boddeke HW (2011) Identification of a microglia phenotype supportive of remyelination. *Glia* 60:306–321
- Ozawa K, Hashimoto K, Kishimoto T, Shimizu E, Ishikura H, Iyo M (2006) Immune activation during pregnancy in mice leads to dopaminergic hyperfunction and cognitive impairment in the offspring: a neurodevelopmental animal model of schizophrenia. *Biol Psychiatry* 59:546–554
- Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira TA, Guiducci E, Dumas L, Ragozzino D, Gross CT (2011) Synaptic pruning by microglia is necessary for normal brain development. *Science* 333:1456–1458
- Patterson PH (2007) Neuroscience. Maternal effects on schizophrenia risk. *Science* 318:576–577
- Peri F, Nusslein-Volhard C (2008) Live imaging of neuronal degradation by microglia reveals a role for v0-ATPase a1 in phagosomal fusion in vivo. *Cell* 133:916–927
- Perry VH, Hume DA, Gordon S (1985) Immunohistochemical localization of macrophages and microglia in the adult and developing mouse brain. *Neuroscience* 15:313–326
- Pont-Lezica L, Bechade C, Belarif-Cantaut Y, Pascual O, Bessis A (2011) Physiological roles of microglia during development. *J Neurochem*
- Priller J, Flugel A, Wehner T, Boentert M, Haas CA, Prinz M, Fernandez-Klett F, Prass K, Bechmann I, de Boer BA, Frotscher M, Kreutzberg GW, Persons DA, Dirnagl U (2001) Targeting gene-modified hematopoietic cells to the central nervous system: use of green fluorescent protein uncovers microglial engraftment. *Nat Med* 7:1356–1361
- Prinz M, Mildner A (2011) Microglia in the CNS: immigrants from another world. *Glia* 59:177–187
- Ransohoff RM, Cardona AE (2010) The myeloid cells of the central nervous system parenchyma. *Nature* 468:253–262
- Ransohoff RM, Perry VH (2009) Microglial physiology: unique stimuli, specialized responses. *Annu Rev Immunol* 27:119–145
- Romero E, Ali C, Molina-Holgado E, Castellano B, Guaza C, Borrell J (2007) Neurobehavioral and immunological consequences of prenatal immune activation in rats. Influence of antipsychotics. *Neuropsychopharmacology* 32:1791–1804
- Rymo SF, Gerhardt H, Wolfhagen Sand F, Lang R, Uv A, Betsholtz C (2011) A two-way communication between microglial cells and angiogenic sprouts regulates angiogenesis in aortic ring cultures. *PLoS One* 6:e15846
- Schlegelmilch T, Henke K, Peri F (2011) Microglia in the developing brain: from immunity to behaviour. *Curr Opin Neurobiol* 21:5–10
- Schwartz M, Butovsky O, Bruck W, Hanisch UK (2006) Microglial phenotype: is the commitment reversible? *Trends Neurosci* 29:68–74
- Shi L, Fatemi SH, Sidwell RW, Patterson PH (2003) Maternal influenza infection causes marked behavioral and pharmacological changes in the offspring. *J Neurosci* 23:297–302
- Sierra A, Encinas JM, Deudero JJ, Chancey JH, Enikolopov G, Overstreet-Wadiche LS, Tsirka SE, Maletic-Savatic M (2010) Microglia shape adult hippocampal neurogenesis through apoptosis-coupled phagocytosis. *Cell Stem Cell* 7:483–495
- Stevens B, Allen NJ, Vazquez LE, Howell GR, Christopherson KS, Nouri N, Micheva KD, Mehalow AK, Huberman AD, Stafford B, Sher A, Litke AM, Lambris JD, Smith SJ, John SW, Barres BA (2007) The classical complement cascade mediates CNS synapse elimination. *Cell* 131:1164–1178

- Stolzing A, Grune T (2003) Impairment of protein homeostasis and decline of proteasome activity in microglial cells from adult wistar rats. *J Neurosci Res* 71:264–271
- Streit WJ (2004) Microglia and Alzheimer's disease pathogenesis. *J Neurosci Res* 77:1–8
- Streit WJ (2006) Microglial senescence: does the brain's immune system have an expiration date? *Trends Neurosci* 29:506–510
- Streit WJ, Xue QS (2009) Life and death of microglia. *J Neuroimmune Pharmacol* 4:371–379
- Tremblay ME (2012) The role of microglia at synapses in the healthy CNS: novel insights from recent imaging studies. *Neuron Glia Biol*:1–10.
- Tremblay ME, Stevens B, Sierra A, Wake H, Bessis A, Nimmerjahn A (2011) The role of microglia in the healthy brain. *J Neurosci* 31:16064–16069
- Wakselman S, Bechade C, Roumier A, Bernard D, Triller A, Bessis A (2008) Developmental neuronal death in hippocampus requires the microglial CD11b integrin and DAP12 immunoreceptor. *J Neurosci* 28:8138–8143
- Wierzba-Bobrowicz T, Lewandowska E, Kosno-Kruszewska E, Lechowicz W, Pasennik E, Schmidt-Sidor B (2004) Degeneration of microglial cells in frontal and temporal lobes of chronic schizophrenics. *Folia Neuropathol* 42:157–165
- Wirenfeldt M, Babcock AA, Vinters HV (2011) Microglia - insights into immune system structure, function, and reactivity in the central nervous system. *Histol Histopathol* 26:519–530
- Yang F, Sun X, Beech W, Teter B, Wu S, Sigel J, Vinters HV, Frautschy SA, Cole GM (1998) Antibody to caspase-cleaved actin detects apoptosis in differentiated neuroblastoma and plaque-associated neurons and microglia in Alzheimer's disease. *Am J Pathol* 152:379–389
- Yun HJ, Yoon JH, Lee JK, Noh KT, Yoon KW, Oh SP, Oh HJ, Chae JS, Hwang SG, Kim EH, Maul GG, Lim DS, Choi EJ (2011) Daxx mediates activation-induced cell death in microglia by triggering MST1 signalling. *EMBO J* 30:2465–2476
- Zuckerman L, Weiner I (2005) Maternal immune activation leads to behavioral and pharmacological changes in the adult offspring. *J Psychiatr Res* 39:311–323
- Zuckerman L, Rehavi M, Nachman R, Weiner I (2003) Immune activation during pregnancy in rats leads to a postpubertal emergence of disrupted latent inhibition, dopaminergic hyperfunction, and altered limbic morphology in the offspring: a novel neurodevelopmental model of schizophrenia. *Neuropsychopharmacology* 28:1778–1789