

A new family of facet defining inequalities for the maximum edge-weighted clique problem

Franklin Djeumou Fomeni¹

Received: 9 November 2015 / Accepted: 17 June 2016 / Published online: 22 June 2016
© Springer-Verlag Berlin Heidelberg 2016

Abstract This paper considers a family of cutting planes, recently developed for mixed 0–1 polynomial programs and shows that they define facets for the maximum edge-weighted clique problem. There exists a polynomial time exact separation algorithm for these inequalities. The result of this paper may contribute to the development of more efficient algorithms for the maximum edge-weighted clique problem that use cutting planes.

Keywords Edge-weighted clique problem · Cutting planes · Separation algorithm · Integer programming · Boolean quadric polytope · Facet defining inequalities

1 Introduction

The maximum edge-weighted clique problem (MEWCP) is a well known combinatorial optimisation problem which consists of finding a maximum weight clique with no more than b nodes in a node- and edge-weighted complete graph. The weight of a clique is defined as the sum of the weights of all its nodes and edges. More formally, the MEWCP is defined as follows. Given a complete undirected graph $G = (N, E)$ with node set N , edge set E , an integer number $b \leq |N| - 1$, weights $w_i \in \mathbb{R}$ associated with each node $i \in N$ and weights $c_e \in \mathbb{R}$ associated with each edge $e \in E$, the MEWCP consists of finding a sub-clique $C = (U, F)$ of G such that the sum of the weights of nodes in U and edges in F is maximised and $|U| \leq b$. It can be formulated as follows:

✉ Franklin Djeumou Fomeni
franklin.djeumou-fomeni@polymtl.ca

¹ GERAD and Department of Mathematics and Industrial Engineering, Polytechnique Montréal, Montreal, QC H3T 3A7, Canada

$$\max \sum_{i \in N} w_i x_i + \sum_{e \in E} c_e y_e \quad (1a)$$

$$s.t. \quad \sum_{i \in N} x_i \leq b \quad (1b)$$

$$y_{ij} \leq x_i \quad \text{for } (i, j) \in E \quad (1c)$$

$$y_{ij} \leq x_j \quad \text{for } (i, j) \in E \quad (1d)$$

$$x_i + x_j \leq y_{ij} + 1 \quad \text{for } (i, j) \in E \quad (1e)$$

$$x_i \in \{0, 1\} \quad \text{for } i \in N \quad (1f)$$

$$y_e \in \{0, 1\} \quad \text{for } e \in E \quad (1g)$$

Note that due to the McCormick inequalities [12] (1c)–(1e) and the constraint (1f), the variables y_e , $e \in E$ can be assumed to be continuous between 0 and 1.

The MEWCP has many applications, especially in certain facility location problems, see [3, 10, 17, 18]. Other important applications of the MEWCP that arise in molecular biology are given in Hunting [6]. The MEWCP is a generalization of the well studied maximum clique problem, which is known to be NP-hard, see [20] for a review of solution approaches for the maximum clique problem. On the other hand, the above formulation of the MEWCP can also be seen as a particular case of the quadratic knapsack problem for which plenty of exact and heuristic methods exist, see [2, 5, 16].

Numerous solution methods have been proposed in the literature for the MEWCP. We refer the reader to Wu and Hao [20] for a recent review of exact and heuristic solution methods for the MEWCP. The most successful algorithms proposed in the literature for the MEWCP use a branch-and-cut framework. The availability of strong valid inequalities is key to the success of these algorithms. Ideally, one would like to use cutting planes that are facet defining and computationally ‘easy’ to generate. Several families of facet defining inequalities are proposed in the literature for this purpose, see for example [7–9, 11, 13, 14, 19].

In this paper, we first consider a family of cutting planes that have recently been developed by Djeumou Fomeni et al. [4] for the general mixed 0–1 polynomial programs, and which can be separated efficiently in polynomial time. Then we prove that under certain conditions, one of the inequalities in this family defines a facet for the MEWCP. This result may contribute to the development of more efficient algorithms for the MEWCP that use cutting planes.

The rest of this paper is organised as follows. In Sect. 2, we review the relevant literature, and in Sect. 3 we provide the proof that the (s, t) inequalities define facets of the MEWCP.

2 Literature review

We refer the reader to [1, 3, 7–9, 11, 13, 14, 19] for more details on other existing facet defining inequalities and solution methods for the MEWCP. For the sake of brevity, we restrict our literature review to the paper of Djeumou Fomeni et al. [4] in which they presented the cutting planes that are discussed in this paper.

2.1 The family of (s, t) inequalities for 0–1 quadratic programs

Given a linear inequality $\alpha^T x \leq \beta$, with $\alpha \in \mathbb{Q}^n, \beta \in \mathbb{Q}$ let us define the corresponding quadratic knapsack polytope as

$$\mathcal{Q} := \text{conv} \left\{ (x, y) \in \{0, 1\}^{n+\binom{n}{2}} : \alpha^T x \leq \beta, y_{ij} = x_i x_j \text{ for } (i, j) \in E \right\}$$

For any $S \subset N$ and any $\alpha \in \mathbb{Q}^n$, we will let $\alpha(S)$ denote $\sum_{i \in S} \alpha_i$, S^+ denote $\{i \in S : \alpha_i > 0\}$ and S^- denote $\{i \in S : \alpha_i < 0\}$.

The method for generating inequalities presented in [4] is based on the following idea. First, we construct a cubic valid inequality, by which we mean a non-linear inequality that involves products of up to three x variables, but no y variables. Then, we weaken the cubic inequality, in order to make it valid for \mathcal{Q} . For example, we can take the inequality $\alpha^T x \leq \beta$, and two binary variables, say x_s and x_t , and form the following three cubic inequalities:

$$(\beta - \alpha^T x)x_s x_t \geq 0 \tag{2}$$

$$(\beta - \alpha^T x)x_s(1 - x_t) \geq 0 \tag{3}$$

$$(\beta - \alpha^T x)(1 - x_s)(1 - x_t) \geq 0. \tag{4}$$

Since quadratic terms of the form $x_i x_j$ can be replaced with y_{ij} , and linear and constant terms can be left unchanged, the only real issue is how to deal with cubic terms, of the form $x_i x_j x_k$. The following lemma addresses this issue:

Lemma 1 *Let x_i, x_j and x_k be three variables, all constrained to lie in the interval $[0, 1]$. Let $y_{ij} = x_i x_j$, and similarly for y_{ik} and y_{jk} . Then we have the following lower bounds on $x_i x_j x_k$:*

$$x_i x_j x_k \geq \max \{0, y_{ij} + y_{ik} - x_i, y_{ij} + y_{jk} - x_j, y_{ik} + y_{jk} - x_k\}, \tag{5}$$

and the following upper bounds:

$$x_i x_j x_k \leq \min \{y_{ij}, y_{ik}, y_{jk}, 1 - x_i - x_j - x_k + y_{ij} + y_{ik} + y_{jk}\}. \tag{6}$$

It is shown in [4] that (5) and (6) provide the best possible under- and over-estimators of the product term $x_i x_j x_k$.

The following theorem characterises the cutting planes that can be derived by weakening the cubic inequalities (2), (3) and (4), respectively. It turns out that they give rise to three huge (exponentially-large) families of valid inequalities for \mathcal{Q} .

Theorem 1 *For any pair $\{s, t\} \subset N$, let S, T and W be disjoint subsets of $N \setminus \{s, t\}$ and let R denote $N \setminus (\{s, t\} \cup S \cup T \cup W)$.*

1. *Then the following (s, t) inequalities are valid for \mathcal{Q} :*

$$\begin{aligned} \sum_{i \in S \cup W} \alpha_i y_{is} + \sum_{i \in T \cup W} \alpha_i y_{it} - \sum_{i \in W} \alpha_i x_i &\leq -\alpha(W^-) + \alpha(S^+ \cup W^-)x_s \\ &+ \alpha(T^+ \cup W^-)x_t + (\beta - \alpha(\{s, t\} \cup S^+ \cup T^+ \cup W^- \cup R^-)) y_{st}. \end{aligned} \tag{7}$$

2. The following mixed (s, t) inequalities are valid for \mathcal{Q} :

$$\sum_{i \in W} \alpha_i x_i + \sum_{i \in TUR} \alpha_i y_{is} - \sum_{i \in TUW} \alpha_i y_{it} \leq \alpha(W^+) + (\beta - \alpha(\{s\} \cup S^- \cup W^+)) x_s - \alpha(W^+ \cup T^-) x_t + (\alpha(\{s\} \cup S^- \cup T^- \cup W^+ \cup R^-) - \beta) y_{st}. \tag{8}$$

3. The following reverse (s, t) inequalities are valid for \mathcal{Q} :

$$\sum_{i \in SUTUR} \alpha_i x_i - \sum_{i \in TUR} \alpha_i y_{is} - \sum_{i \in SUR} \alpha_i y_{it} \leq \beta - \alpha(W^-) + (\alpha(S^+ \cup W^-) - \beta) x_s + (\alpha(T^+ \cup W^-) - \beta) x_t + (\beta - \alpha(S^+ \cup T^+ \cup W^- \cup R^-)) y_{st}. \tag{9}$$

These inequalities can be strengthened further, see [4] for details. Our contribution in this paper consists of proving that under certain conditions, the (s, t) inequalities (7) are facet defining. We also remark that the particular case of the mixed (s, t) inequalities obtained when $S = T = R = \emptyset$ and $\alpha = (1, \dots, 1)$ was previously given in [7] and proved to be facet defining for the MEWCP.

2.2 Separation of the (s, t) inequalities in $\mathcal{O}(n^3)$ time

Since the inequalities presented in Theorem 1 are exponential in number, we need separation algorithms. For a given family of inequalities, the separation algorithm takes a fractional point (x^*, y^*) , solution of the LP relaxation, as input, and outputs a violated inequality in that family, if one exists.

It turns out that the separation problems for the (s, t) inequalities (7), mixed (s, t) inequalities (8) and reverse (s, t) inequalities (9) can each be solved exactly in $\mathcal{O}(n^3)$ time [4]. Indeed, there are $\binom{n}{2}$ choices for the pair $\{s, t\}$. Now assume that s and t are fixed. The (s, t) inequality can be rewritten as:

$$\begin{aligned} & \sum_{i \in S^+} \alpha_i (y_{is} + y_{st} - x_s) + \sum_{i \in T^+} \alpha_i (y_{it} + y_{st} - x_t) + \sum_{i \in W^+} \alpha_i (y_{is} + y_{it} - x_i) \\ & + \sum_{i \in S^-} \alpha_i y_{is} + \sum_{i \in T^-} \alpha_i y_{it} + \sum_{i \in W^-} \alpha_i (1 - x_i - x_s - x_t + y_{is} + y_{it} + y_{st}) \\ & + \sum_{i \in R^-} \alpha_i y_{st} \leq (\beta - \alpha_s - \alpha_t) y_{st}. \end{aligned}$$

Observe that, in this form, the right-hand side is a constant for the given α, β, s and t . Then, to find a most-violated (s, t) inequality, if any exists, it suffices to maximise the left-hand side. This can be done using the following algorithm. Consider each node $i \in N \setminus \{s, t\}$ in turn. If $\alpha_i > 0$, place i in one of the sets S, T, W or R , according to which of the following four quantities is largest: $y_{is}^* + y_{st}^* - x_s^*, y_{it}^* + y_{st}^* - x_t^*, y_{is}^* + y_{it}^* - x_i^*$ and zero. (Ties can be broken arbitrarily.) If $\alpha_i < 0$, place i in S, T, W or R according to which of the following four quantities is smallest: $y_{is}^*, y_{it}^*, 1 - x_i^* - x_s^* - x_t^* + y_{is}^* + y_{it}^* + y_{st}^*$ and y_{st}^* . (Again, ties can be broken arbitrarily.) If

$\alpha_i = 0$, then i can be placed in an arbitrary set, since it has no effect on the violation. Note that, for any i , only a constant number of comparisons is needed. Therefore the algorithm runs in $\mathcal{O}(n)$ time for the given α, β, s and t .

3 Facet proof

In this Section, we provide the proof that under certain conditions, the family of (s, t) inequalities (7) are facet defining for the MEWCP. We can note from the cardinality constraint (1b) that the coefficients $\alpha_i, i = 1, \dots, n$, are all positive and equal to 1, i.e. $S = S^+, T = T^+, W = W^+, R = R^+$ and for each of these sets, the sum of coefficients α is simply equal to its cardinality (for example $\alpha(S) = |S|$). For these reasons, the (s, t) inequality for the MEWCP can be written as follows:


$$\sum_{i \in SUW} y_{is} + \sum_{i \in T UW} y_{it} - \sum_{i \in W} x_i \leq (|S|)x_s + (|T|)x_t + (b - 2 - |S| - |T|) y_{st}. \tag{10}$$

For the rest of this paper, the set \mathcal{Q} corresponds to

$$\mathcal{Q} := \text{conv} \left\{ (x, y) \in \{0, 1\}^{n+\binom{n}{2}} : \sum_{i=1}^n x_i \leq b, y_{ij} = x_i x_j \text{ for } (i, j) \in E \right\}$$

Theorem 2 *Let s, t, S, T and W be defined as in Sect. 2. If S and T are non empty, $|S| \leq b - 2, |T| \leq b - 2, W = \emptyset$ and $|S \cup T| \geq b - 1$, then the (s, t) inequalities (10) define facets of \mathcal{Q} .*

Note that with the settings of Theorem 2, the supporting graph of the (s, t) inequalities (10) is a double star tree as follows.



Proof Under the hypothesis that $W = \emptyset$, the (s, t) inequalities (10) becomes

$$\sum_{i \in S} y_{is} + \sum_{i \in T} y_{it} \leq (|S|)x_s + (|T|)x_t + (b - 2 - |S| - |T|) y_{st}. \tag{11}$$

Let $F = \{(x, y) \in \mathcal{Q} : (11) \text{ holds with equality}\}$, and $a(x, y) \leq a_0$ i.e. let

$$a_1x_1 + a_2x_2 + \dots + a_nx_n + a_{12}y_{12} + a_{13}y_{13} + \dots + a_{n-1,n}y_{n-1,n} \leq a_0$$

be an inequality valid for \mathcal{Q} such that every point $(x, y) \in F$ satisfies $a(x, y) = a_0$. We will use some integer points in \mathcal{Q} that satisfy (11) to equality i.e integer points in F to find the coefficients a and a_0 uniquely up to scalar multiplication.

Let e_i be i^{th} unit vector of size n , e_{ij} the $\binom{n}{2}$ -vector with all components equal to zero except the $(i, j) - th$ component which is equal to 1.

1. The vector $(x, y) = (0, 0) \in F$; this implies that $a_0 = 0$.
2. $(e_i, 0) \in F$ for $i \neq s, t$; this implies that $a_i = 0$ for all $i \neq s, t$. Note that the nodes s and t can be isolated in the set N without ambiguity since $|S| \leq b - 2$ and $|T| \leq b - 2$.
3. $(e_i + e_j, e_{ij}) \in F$ for all $i, j \neq s, t$ and $i \neq j$; it follows that $a_{ij} = 0$ for all $i, j \neq s, t$ and $i \neq j$.
4. We now prove that $a_{it} = 0$ for any node $i \in N \setminus (T \cup \{s, t\})$. Let $i \in N \setminus (T \cup \{s, t\})$, we define:
 - C_{it}^s to be a star tree with node set $T \cup \{i, t\}$ (it is possible to have such a star tree since $T \neq \emptyset$) such that all the edges are incident to t . Since $C_{it}^s \in F$, it follows that

$$a_t + \sum_{k \in T} a_{kt} + a_{it} = 0 \quad (i)$$

- C_t^i to be a star tree with node set $T \cup \{t\}$ such that all the edges are incident to t this is the same as the star tree C_{it}^s without the edge (i, t) . Since $C_t^i \in F$, it follows that

$$a_t + \sum_{k \in T} a_{kt} = 0 \quad (ii)$$

Subtracting (ii) from (i) yields $a_{it} = 0$ for $i \in N \setminus (T \cup \{s, t\})$.

5. Similarly to the above point, $a_{js} = 0$ for $j \in N \setminus (S \cup \{s, t\})$, also using the fact that $S \neq \emptyset$.
6. For $i, j \in S \cup T$, we want to show that: a) $a_{is} = a_{js}$ when $i, j \in S$, b) $a_{it} = a_{jt}$ when $i, j \in T$, and c) $a_{is} = a_{jt}$ when $i \in S$ and $j \in T$. Let $i, j \in S \cup T$ with $i \neq j$ and let $A \subseteq S \cup T \setminus \{i, j\}$ such that $|A| = b - 3$, (since $|S \cup T| \geq b - 1$). Let C_{ist}^j be a double star tree with node set $A \cup \{i, s, t\}$ obtained by linking all the nodes in $A \cap S$ to s , all the nodes in $A \cap T$ to t and connecting the node s to the node t .
 - Since $C_{i,s,t}^j \in F$, it follows that

$$a_s + a_t + \sum_{k \in A \cap S} a_{ks} + \sum_{k \in A \cap T} a_{kt} + a_{is} + a_{it} + a_{st} = 0 \quad (iii).$$

- Since $C_{j,s,t}^i \in F$, it follows that

$$a_s + a_t + \sum_{k \in A \cap S} a_{ks} + \sum_{k \in A \cap T} a_{kt} + a_{js} + a_{jt} + a_{st} = 0 \quad (iv).$$

Subtracting (iii) from (iv) yields $a_{is} + a_{it} = a_{js} + a_{jt}$. So, using steps 4 and 5 we have the following:

- (a) If $i, j \in S$ then $a_{is} = a_{js}$.
 - (b) If $i, j \in T$ then $a_{it} = a_{jt}$.
 - (c) If $i \in S$ and $j \in T$, then $a_{is} = a_{jt}$.
7. Using $a_{it} = a_{jt}$ for $i, j \in T$, as given by b) in the above point, and considering equation (ii), we have $a_t + |T|a_{it} = 0$ for $i \in T$. Therefore, $a_{it} = -\frac{a_t}{|T|}$, (since $T \neq \emptyset$).
 Similarly, $a_s + |S|a_{is} = 0$ for $i \in S$, i.e $a_{is} = -\frac{a_s}{|S|}$, (since $S \neq \emptyset$).
8. Let $i \in S$ and $j \in T$, we define the set A as in step 6 and denote $\omega_s = |A \cap S| + 1$ and $\omega_t = |A \cap T|$. It follows from (iii) that $a_s + a_t + \omega_s a_{is} + \omega_t a_{jt} + a_{st} = 0$ i.e. $a_{st} = -a_s - a_t + \frac{a_s \omega}{|S|} + \frac{a_t \omega_t}{|T|}$ for $i \in S$ and $j \in T$.
9. Finally, considering the above steps, the inequality

$$a_1x_1 + a_2x_2 + \dots + a_nx_n + a_{12}y_{12} + a_{13}y_{13} + \dots + a_{n-1,n}y_{n-1,n} \leq a_0$$

reduces to

$$a_sx_s + a_tx_t + \sum_{i \in S} a_{is}y_{is} + \sum_{i \in T} a_{it}y_{it} + a_{st}y_{st} \leq 0$$

which, using the identities $a_{is} = a_{jt}$, $a_{jt} = -\frac{a_t}{|T|}$ and $a_{is} = -\frac{a_s}{|S|}$ for $i \in S, j \in T$, is equivalent to

$$a_sx_s + a_tx_t - \frac{a_s}{|S|} \sum_{i \in S} y_{is} - \frac{a_t}{|T|} \sum_{i \in T} y_{it} + \left(\frac{a_s \omega_s}{|S|} + \frac{a_t \omega_t}{|T|} - a_s - a_t \right) y_{st} \leq 0.$$

We finally have

$$\frac{a_s}{|S|} \left[|S|x_s + |T|x_t - \sum_{i \in S} y_{is} - \sum_{i \in T} y_{it} - (|S| + |T| - \omega_s - \omega_t)y_{st} \right] \leq 0.$$

Since $(e_s, 0)$ satisfies the inequality $a(x, y) \leq a_0$, i.e $a_s \leq 0$, and given that $\omega_s + \omega_t = b - 2$, this ends the proof.

Acknowledgements Part of the work of the author was done during a stay at GERAD as a postdoctoral researcher with Professor Miguel Anjos. The author is grateful to Professor Adam Letchford for his comments and feedback on this work, and finally thanks the referees for their careful reading of the paper and their helpful feedback.

References

1. Alidaee, B., Glover, F., Kochenberger, G., Wang, H.: Solving the maximum edge weight clique problem via unconstrained quadratic programming. *Eur. J. Oper. Res.* **181**, 592–597 (2007)
2. Caprara, A., Pisinger, D., Toth, P.: Exact solution of the quadratic knapsack problem. *INFORMS J. Comput.* **11**, 125–137 (1998)
3. Dijkhuizen, G., Faigle, U.: A cutting-plane approach to the edge-weighted maximal clique problem. *Eur. J. Oper. Res.* **69**(1), 121–130 (1993)
4. Djeumou Fomeni, F., Kaparis, K., Letchford, A.N.: Cutting planes for first-level RLT relaxations of mixed 0–1 programs. *Math. Progr.* **151**(2), 639–658 (2015)
5. Djeumou Fomeni, F., Letchford, A.N.: A dynamic programming heuristic for the quadratic knapsack problem. *INFORMS J. Comp.* **26**(1), 173–183 (2014)
6. Hunting, M.: Relaxation techniques for discrete optimization problems: theory and algorithms. PhD thesis, University of Twente (1998)
7. Hunting, M., Faigle, U., Kern, W.: A Lagrangean relaxation approach to the edge-weighted clique problem. *Eur. J. Oper. Res.* **131**, 119–131 (2001)
8. Helmberg, C., Rendl, F., Weismantel, R.: A semidefinite programming approach to the quadratic knapsack problem. *J. Comb. Optim.* **4**, 197–215 (2000)
9. Johnson, E.L., Mehrotra, A., Nemhauser, G.L.: Min-cut clustering. *Math. Progr.* **62**, 133–151 (1993)
10. Kubj, M.J.: Programming models for facility dispersion: the p -dispersion and maximum dispersion problem. *Geograph. Anal.* **9**, 315–329 (1987)
11. Macambria, E.M., de Souza, C.C.: The edge-weight clique problem: valid inequalities, facets and polyhedral computations. *Eur. J. Oper. Res.* **123**, 346–371 (2000)
12. McCormick, G.P.: Computability of global solutions to factorable nonconvex programs: part I-convex underestimating problems. *Math. Progr.* **10**, 146–175 (1979)
13. Mehrotra, A.: Cardinality constrained boolean quadratic polytope. *Disc. Appl. Math* **79**, 137–154 (1997)
14. Park, K., Lee, K., Park, S.: An extended formulation approach to the edge-weight maximal clique problem. *Eur. J. Oper. Res.* **95**, 671–682 (1995)
15. Padberg, M.W.: The Boolean quadratic polytope: some characteristics, facets and relatives. *Math. Progr.* **45**, 139–172 (1989)
16. Pisinger, D.: The quadratic knapsack problem: a survey. *Discr. Appl. Math.* **155**, 623–648 (2007)
17. Ravi, S.S., Rosenkrantz, D.J., Tayi, G.K.: Heuristic and special case algorithms for dispersion problems. *Oper. Res.* **42**(2), 299–310 (1994)
18. Späth, H.: Heuristically determining cliques of given cardinality and with minimal cost within weighted complete graphs. *Z. Oper. Res.* **29**(3), 125–131 (1985)
19. Sørensen, M.M.: New facets and a branch-and-cut algorithm for the weighted clique problem. *Eur. J. Oper. Res.* **154**, 57–70 (2004)
20. Wu, Q., Hao, J.K.: A review on algorithms for maximum clique problems. *Eur. J. Oper. Res.* **242**(3), 693–709 (2015)