



Feature selection in SVM via polyhedral k -norm

Manlio Gaudioso¹ · E. Gorgone² · J.-B. Hiriart-Urruty³

Received: 10 November 2018 / Accepted: 12 September 2019 / Published online: 18 September 2019
© Springer-Verlag GmbH Germany, part of Springer Nature 2019

Abstract

We treat the feature selection problem in the support vector machine (SVM) framework by adopting an optimization model based on use of the ℓ_0 pseudo-norm. The objective is to control the number of non-zero components of the normal vector to the separating hyperplane, while maintaining satisfactory classification accuracy. In our model the polyhedral norm $\|\cdot\|_{[k]}$, intermediate between $\|\cdot\|_1$ and $\|\cdot\|_\infty$, plays a significant role, allowing us to come out with a DC (difference of convex) optimization problem that is tackled by means of DCA algorithm. The results of several numerical experiments on benchmark classification datasets are reported.

Keywords Sparse optimization · Cardinality constraint · k -norm · Support vector machine · DC optimization

1 Introduction

A relevant problem in binary classification is to design good quality classifiers by resorting to a minimal number of sample parameters. One of the main motivations is to gather a more clear interpretation of phenomena underlying the class membership distribution of the samples. In the more general setting of Machine Learning, such problem falls in the area of Feature Selection (FS), which has been the object of intensive research in recent decades (see, e.g., the survey [13]).

✉ Manlio Gaudioso
gaudioso@dimes.unical.it

E. Gorgone
egorgone@unica.it

J.-B. Hiriart-Urruty
jbhu@math.univ-toulouse.fr

¹ Dipartimento di Ingegneria Informatica, Modellistica, Elettronica e Sistemistica, Università della Calabria, 87036 Rende, CS, Italy

² Dipartimento di Matematica e Informatica, Università di Cagliari, 09124 Cagliari, Italy

³ Institut de Mathématiques, Université Paul Sabatier, 31062 Toulouse, France

In this paper we focus on FS in binary classification (see [7,28] for FS models in other areas of machine learning, such as unsupervised learning and regression, respectively). In particular, we consider the Support Vector Machine (SVM) framework [5,29], where binary classification is pursued by finding an “optimal” two-class separating hyperplane, either in the original parameter (or “feature”) space or upon appropriate kernel transformation.

Numerical optimization algorithms play a relevant role in SVM area and, more specifically, in FS. The problem is to guarantee a reasonable trade-off between classification accuracy and the number of features actually used. Controlling the latter consists basically in minimizing the number of non-zero components of the normal vector to the separating hyperplane.

The literature offers several contributions. In [1] NP-hardness of the problem has been assessed.

In [3,4] the model adopted is based on considering the step function for each component of the normal vector; discontinuity is handled by introducing two different approximations, the standard sigmoid and a concave exponential, respectively. In particular, by adopting the concave approximation, FS problem is tackled by solving a finite sequence of linear programs. Different approximations are given in [25,32], where, in particular, the concave and separable objective functions, derived by the approximations, are handled via variants of Frank-Wolfe method.

It is interesting to note that in [20] an approximation scheme of the step function is cast into a DC (Difference of Convex) framework, providing thus the opportunity of resorting to the algorithmic machinery for dealing with such class of nonconvex problems. An early survey on properties and relevance of this class of functions is in [16] (see also [26]).

Parallel to treatment of FS via approximation methods, Mixed Integer Programming formulations have been successfully adopted. The idea is to introduce binary variables, one for each component of the normal vector to the supporting hyperplane, that are switched to “1” if and only if the corresponding component is non-zero. From among the proposed approaches, we recall here [2,9,21].

In this paper we tackle FS in the general setting of sparse optimization, where one is faced to the (regularized) problem:

$$\min_{x \in \mathbb{R}^n} f(x) + \|x\|_0 \quad (1.1)$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $\|\cdot\|_0$ is the ℓ_0 pseudo-norm, which counts the number of non-zero component of any vector. Sometimes sparsity of the solution, instead of acting on the objective function, is enforced by introducing a constraint on the ℓ_0 pseudo-norm of the solution, thus defining a cardinality-constraint problem [24].

In many applications, the ℓ_0 pseudo-norm in (1.1) is replaced by the ℓ_1 -norm, which is definitely more tractable from the computational point of view, yet ensuring sparsity, to a certain extent (see [33] for a discussion on a general regularization scheme).

In the seminal paper [31], a class of polyhedral norms (the k -norms), intermediate between $\|\cdot\|_1$ and $\|\cdot\|_\infty$, is introduced to obtain sparse approximation solutions to systems of linear equations. The use of other norms to recover sparsity is described in

[8]. In more recent years the use of k -norms has received much attention and has led to several proposals for dealing with ℓ_0 pseudo-norm cardinality constrained problem [12,15,27,34].

In this paper we cast the classic SVM approach into the sparse optimization framework (1.1). Our work is inspired by [12], the main difference being in the explicit (and not parametric) minimization of the ℓ_0 pseudo-norm. We formulate our SVM- ℓ_0 pseudo-norm problem ($SV M_0$, for short) and we tackle it by means of a penalization approach which allows us to put the problem in DC form. The algorithm adopted is of DCA type [19].

The paper is organized as follows. In Sect. 2 we recall first the standard SVM model and introduce the FS problem as a sparse optimization one. Then we discuss the use of the polyhedral k -norm in sparse optimization, coming out with a DC formulation. On such basis we formulate in Sect. 3 the $SV M_0$ problem. The results of several numerical experiments on benchmark datasets are in Sect. 4. Some conclusions are finally drawn in Sect. 5.

2 Feature selection and ℓ_0 pseudo-norm minimization

The binary classification problem in the SVM setting is usually put in the following form. Given two point-sets $\mathcal{A} \triangleq \{a_1, \dots, a_{m_1}\}$ and $\mathcal{B} \triangleq \{b_1, \dots, b_{m_2}\}$ in \mathbb{R}^n , we look for *linear separation* of the two sets, that is for a hyperplane $\{x|x \in \mathbb{R}^n, x^\top w' = \gamma'\}$, ($w' \in \mathbb{R}^n, \gamma' \in \mathbb{R}$), strictly separating \mathcal{A} and \mathcal{B} , thus ensuring

$$a_i^\top w' < \gamma', \quad i = 1, \dots, m_1 \quad \text{and} \quad b_l^\top w' > \gamma', \quad l = 1, \dots, m_2.$$

It is easy to verify that such a hyperplane exists if and only if there exists a hyperplane $\{x|x \in \mathbb{R}^n, x^\top w = \gamma\}$, ($w \in \mathbb{R}^n, \gamma \in \mathbb{R}$), such that

$$a_i^\top w \leq \gamma - 1, \quad i = 1, \dots, m_1 \quad \text{and} \quad b_l^\top w \geq \gamma + 1, \quad l = 1, \dots, m_2.$$

Gordan’s theorem of the alternative [22] guarantees linear separation if and only if $\text{conv}\mathcal{A} \cap \text{conv}\mathcal{B} = \emptyset$, a property which is not usually known in advance to hold.

Consequently, an *error function* of (w, γ) , which is convex, piecewise linear and nonnegative, is introduced. It assumes zero value if and only if (w, γ) actually defines a (strictly) separating hyperplane and it has the form:

$$e(w, \gamma) = \sum_{i=1}^{m_1} \max\{0, a_i^\top w - \gamma + 1\} + \sum_{l=1}^{m_2} \max\{0, -b_l^\top w + \gamma + 1\} \quad (2.1)$$

The SVM approach consists in solving the following convex problem:

$$\min_{w, \gamma} \|w\| + Ce(w, \gamma) \quad (2.2)$$

where the addition of the norm of w to the error function is aimed at obtaining a maximum-margin separation, C being a positive trade-off parameter [5].

Several variants of problem (2.2) are available in literature, according to possible choices of the norm. The ℓ_2 norm is mostly adopted, while for feature selection purposes the ℓ_1 norm is preferred. We observe that the latter plays also a relevant role in many areas of machine learning through the so called LASSO approach (see, e.g. [14,28]). Weighted sums of the ℓ_1 and ℓ_2 norms have been considered within the elastic net [35] and the doubly regularized SVM [30] approaches.

In the following two sub-sections we address first problem (2.2), casting it into the sparse optimization paradigm (1.1), by means of the ℓ_0 pseudo-norm. Then we discuss a DC formulation of the resulting problem.

2.1 Sparse minimization and k -norms

As previously mentioned, the ℓ_0 pseudo-norm counts the number of non-zero components of any vector. We resume here its basic properties and highlight the relationship with polyhedral k -norm, which allows us to reformulate the sparse optimization problem (1.1).

The usual notation $\|\cdot\|_0$ for indicating the ℓ_0 pseudo-norm is motivated by the observation

$$(\|\cdot\|_p)^p \rightarrow \|\cdot\|_0 \quad \text{when } p \rightarrow 0. \quad (2.3)$$

Relevant properties of function $x \mapsto \|x\|_0$ are:

- (i) it is *lower-semicontinuous*, that is to say

$$\liminf_{k \rightarrow +\infty} \|x_k\|_0 \geq \|x\|_0 \quad \text{whenever } x_k \rightarrow x,$$

a property which is fundamental in view of using *descent* algorithms;

- (ii) it is homogeneous of degree 0, ($\|\lambda x\|_0 = \|x\|_0$, for $\lambda \neq 0$); constancy along rays makes difficult the design of minimization algorithms;
- (iii) the convex hull of $\|\cdot\|_0$ on the ball $\{x \mid \|x\|_\infty \leq r\}$ is exactly the function $\frac{1}{r} \|\cdot\|_1$. This is a mathematical justification for frequent substitution of $\|\cdot\|_0$ by $\|\cdot\|_1$, which in fact ensures, in practical applications, attractive *sparsity* properties of the solution.

In our approach to FS we fix $\|w\|_0$ in formulation (2.2), thus we consider a sparse optimization problem of the type

$$f_0^* = \min_{x \in \mathbb{R}^n} f(x) + \|x\|_0, \quad (2.4)$$

where $f : \mathbb{R}^n \rightarrow \mathbb{R}$, $n \geq 2$, is convex, not necessarily differentiable. We observe, in passing, the significant parallelism between sparse optimization and certain problems in matrix optimization [17,23]. See, in particular, the recent approach to the rank function minimization described in [12].

In tackling problem (2.4) we resort to the vector k -norm of x , indicated by $\|x\|_{[k]}$, which is defined as the sum of k largest components (in modulus) of x , $k = 1, \dots, n$. In fact $\|\cdot\|_{[k]}$ is a *polyhedral norm*, intermediate between $\|\cdot\|_1$ and $\|\cdot\|_\infty$.

The following properties hold:

- (i) $\|x\|_\infty = \|x\|_{[1]} \leq \dots \leq \|x\|_{[k]} \leq \dots \leq \|x\|_{[n]} = \|x\|_1$;
- (ii) $\|x\|_0 \leq k \Rightarrow \|x\|_1 - \|x\|_{[s]} = 0, \quad k \leq s \leq n$.

Moreover, it is easy to prove the equivalence, valid for $1 \leq k \leq n$,

$$\|x\|_0 \leq k \Leftrightarrow \|x\|_1 - \|x\|_{[k]} = 0, \tag{2.5}$$

which allows to replace any constraint of the type $\|x\|_0 \leq k$ with a *difference of norms*, that is a DC constraint.

Taking any point $\bar{x} \in \mathbb{R}^n$ and letting $I_{[k]} \triangleq \{i_1, \dots, i_k\}$ be the index set of k largest in modulus components of \bar{x} , a subgradient $\bar{g}^{[k]}$ of the vector k -norm at \bar{x} can be calculated by setting:

$$\bar{g}_i^{[k]} = \begin{cases} 1 & \text{if } i \in I_{[k]} \text{ and } \bar{x}_i \geq 0 \\ -1 & \text{if } i \in I_{[k]} \text{ and } \bar{x}_i < 0 \\ 0 & \text{otherwise} \end{cases}$$

To tackle problem (2.4), we start from the observation (see [23,34]):

$$\|x\|_{[k]} = \max_{y \in \psi_k} y^\top x,$$

where ψ_k is the subdifferential of $\|\cdot\|_{[k]}$ at point 0,

$$\psi_k = \{y \in \mathbb{R}^n \mid y = u - v, \quad 0 \leq u, \quad v \leq e, \quad (u + v)^\top e = k\},$$

with e being the vector of n ones. Then we formulate the following problem:

$$f_c^* = \min_{x,u,v,z} f(x) + z \tag{2.6}$$

$$e^\top (u + v) = z \tag{2.7}$$

$$(u - v)^\top x \geq \|x\|_1 \tag{2.8}$$

$$0 \leq u, \quad v \leq e, \quad x \in \mathbb{R}^n. \tag{2.9}$$

Remark 2.1 From $\|x\|_1 \geq \|x\|_{[k]}$, $k = 1, \dots, n$ it follows that at any feasible point constraint (2.8) is satisfied by equality.

The following proposition highlights the relationship, at the optimum, between variable x and variables u, v .

Proposition 2.2 *At any optimal (local) solution (x^*, u^*, v^*, z^*) of problem (2.6)–(2.9), the following relations hold for $j = 1, \dots, n$:*

$$\begin{cases} x_j^* > 0 \Leftrightarrow u_j^* = 1 \\ x_j^* < 0 \Leftrightarrow v_j^* = 1 \\ x_j^* = 0 \Leftrightarrow u_j^* = v_j^* = 0 \end{cases} \tag{2.10}$$

Proof Note first that u_j^* and v_j^* cannot be both positive. In fact, in such case the solution obtained by replacing u_j^* and v_j^* by $u_j^* - \delta_j$ and $v_j^* - \delta_j$, with $\delta_j \triangleq \min\{u_j^*, v_j^*\} > 0$, would be still feasible and would reduce the objective function value.

Now observe that, while constraint (2.9) ensures

$$-|x_j| \leq x_j(u_j - v_j) \leq |x_j|, \quad j = 1, \dots, n,$$

satisfaction of constraint (2.8) guarantees $x_j(u_j - v_j) = |x_j|$, $j = 1, \dots, n$. Thus it is proved, in particular, $x_j^* > 0 \Rightarrow u_j^* = 1$ ($x_j^* < 0 \Rightarrow v_j^* = 1$).

The implication $x_j^* = 0 \Rightarrow u_j^* = v_j^* = 0$ can be proved by a simple contradiction argument, taking into account optimality of the solution. The same contradiction argument guarantees that the implications $x_j^* > 0 \Leftrightarrow u_j^* = 1$ ($x_j^* < 0 \Leftrightarrow v_j^* = 1$) hold true, while the last implication $x_j^* = 0 \Leftrightarrow u_j^* = v_j^* = 0$ can be proved by observing that $u_j^* = v_j^* = 0$ and $x_j^* \neq 0$ would lead to violation of constraint (2.8). \square

Remark 2.3 Implications (2.10) ensure $z^* = \|x^*\|_0$. Moreover, letting (x^*, u^*, v^*, z^*) be any local minimum of problem (2.6)–(2.9), then x^* is a local minimum for problem (2.4).

By eliminating the scalar variable z in problem (2.6)–(2.9) we come out with the reformulation

$$f_c^* = \min_{x,u,v} f(x) + e^\top(u + v) \tag{2.11}$$

$$(u - v)^\top x \geq \|x\|_1 \tag{2.12}$$

$$0 \leq u, v \leq e, \quad x \in \mathbb{R}^n, \tag{2.13}$$

which we approach by penalizing the nonlinear nonconvex constraint (2.12) through the scalar penalty parameter $\sigma > 0$. We obtain

$$f_c(\sigma) = \min_{x,u,v} f(x) + e^\top(u + v) + \sigma(\|x\|_1 - (u - v)^\top x) \tag{2.14}$$

$$0 \leq u, v \leq e, \quad x \in \mathbb{R}^n. \tag{2.15}$$

Problem (2.14)–(2.15) is box-constrained and nonconvex, due to presence of a bilinear term in the objective function. We introduce next a possible DC decomposition.

2.2 DC decomposition

To obtain the required decomposition, we observe first that, letting $p^\top \triangleq (x^\top, u^\top, v^\top)$, the term

$$q(p) \triangleq (v - u)^\top x,$$

in (2.14) can be written in DC form as

$$q(p) = p^\top Q_1 p - p^\top Q_2 p,$$

where the symmetric positive semidefinite matrices Q_1 and Q_2 of dimension $(3n, 3n)$ are as follows:

$$Q_1 = \frac{1}{4} \begin{pmatrix} 2I & -I & I \\ -I & I & 0 \\ I & 0 & I \end{pmatrix} \tag{2.16}$$

and

$$Q_2 = \frac{1}{4} \begin{pmatrix} 2I & I & -I \\ I & I & 0 \\ -I & 0 & I \end{pmatrix}, \tag{2.17}$$

with I and 0 being, respectively, the identity matrix and the zero matrix of dimension (n, n) .

Summing up, the objective function (2.14) is decomposed as follows:

$$f(x) + e^\top (u + v) + \sigma (\|x\|_1 - (u - v)^\top x) = h_1(p) - h_2(p),$$

with

$$h_1(p) \triangleq f(x) + e^\top (u + v) + \sigma (\|x\|_1 + p^\top Q_1 p),$$

and

$$h_2(p) \triangleq \sigma p^\top Q_2 p.$$

Before concluding the section, we state a property related to parametric normalization of a convex function via the ℓ_1 norm. It will be useful in explaining the role of parameter σ in the numerical experiments of Sect. 4.

Proposition 2.4 *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and define, for $\sigma > 0$, $f_\sigma(x) = f(x) + \sigma \|x\|_1$. If*

$$\sigma > \min_{g \in \partial f(0)} \|g\|_\infty,$$

then $x^ = 0$ is the unique minimum of f_σ .*

Proof Our aim is to prove

$$f(x) + \sigma \|x\|_1 > f(0) \quad \forall x \in \mathbb{R}^n, \quad x \neq 0. \tag{2.18}$$

According to the definition of $g \in \partial f(0)$,

$$f(x) \geq f(0) + g^\top x, \quad \forall x \in \mathbb{R}^n,$$

whence

$$f(x) + \sigma \|x\|_1 \geq f(0) + \sigma \|x\|_1 + g^\top x, \quad \forall x \in \mathbb{R}^n.$$

Since $g^\top x \geq -\|g\|_\infty \|x\|_1$, we infer from the above

$$f(x) + \sigma \|x\|_1 \geq f(0) + (\sigma - \|g\|_\infty) \|x\|_1, \quad \forall x \in \mathbb{R}^n.$$

To ensure (2.18) it suffices to have $\sigma - \|g\|_\infty > 0$ for one $g \in \partial f(0)$. This is secured with the assumption $\sigma > \min_{g \in \partial f(0)} \|g\|_\infty$. □

In next section, after recalling the standard ℓ_1 norm formulation of the SVM model, we state the ℓ_0 FS model and we adapt to it the previously defined DC decomposition.

3 The ℓ_0 -SVM problem (SVM₀)

We rewrite first the SVM model (2.2), taking into account the definition (2.1) of the classification error and adopting the ℓ_1 norm. We obtain problem $SV M_1$.

$$z^* = \min_{w, \gamma, \xi, \zeta} \|w\|_1 + C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) \tag{3.1}$$

subject to

$$a_i^\top w - \gamma + 1 \leq \xi_i, \quad i = 1, \dots, m_1 \tag{3.2}$$

$$-b_l^\top w + \gamma + 1 \leq \zeta_l, \quad l = 1, \dots, m_2 \tag{3.3}$$

$$\xi_i \geq 0, \quad i = 1, \dots, m_1 \tag{3.4}$$

$$\zeta_l \geq 0, \quad l = 1, \dots, m_2, \tag{3.5}$$

where the nonnegative auxiliary variables $\xi_i, i = 1, \dots, m_1$ and $\zeta_l, l = 1, \dots, m_2$ have been introduced to eliminate nonsmoothness in the definition (2.1).

Moreover, by letting

$$w = w^+ - w^-, \quad w^+, w^- \geq 0,$$

and indicating by e the vector of ones of dimension n , the above problem can be rewritten in a Linear Programming form as follows

$$z^* = \min_{w^+, w^-, \gamma, \xi, \zeta} e^\top (w^+ + w^-) + C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) \tag{3.6}$$

subject to

$$a_i^\top (w^+ - w^-) - \gamma + 1 \leq \xi_i, \quad i = 1, \dots, m_1 \tag{3.7}$$

$$-b_l^\top (w^+ - w^-) + \gamma + 1 \leq \zeta_l, \quad l = 1, \dots, m_2 \tag{3.8}$$

$$\xi_i \geq 0, \quad i = 1, \dots, m_1 \tag{3.9}$$

$$\zeta_l \geq 0, \quad l = 1, \dots, m_2, \quad (3.10)$$

$$w^+ \geq 0, \quad w^- \geq 0 \quad (3.11)$$

Of course (3.1)–(3.5) or (3.6)–(3.11) are equivalent formulations of $SV M_1$.

We remark that choice of $\|\cdot\|_1$ in (2.2), instead of $\|\cdot\|_2$, has a beneficial effect in terms of feature selection (see [3]).

To guarantee, however, a better control on the number of features actually entering the classification process, we replace $\|\cdot\|_1$ with $\|\cdot\|_0$ into the model (3.1)–(3.5).

In the sequel we tailor the sparse optimization approach via k -norm of previous section to Feature Selection in SVM model. We obtain the $SV M_0$ problem

$$z^* = \min_{w^+, w^-, \gamma, \xi, \zeta, u, v} C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) + e^\top (u + v) \quad (3.12)$$

subject to

$$(u - v)^\top (w^+ - w^-) \geq e^\top (w^+ + w^-) \quad (3.13)$$

$$a_i^\top (w^+ - w^-) - \gamma + 1 \leq \xi_i, \quad i = 1, \dots, m_1 \quad (3.14)$$

$$-b_l^\top (w^+ - w^-) + \gamma + 1 \leq \zeta_l, \quad l = 1, \dots, m_2 \quad (3.15)$$

$$\xi_i \geq 0, \quad i = 1, \dots, m_1 \quad (3.16)$$

$$\zeta_l \geq 0, \quad l = 1, \dots, m_2 \quad (3.17)$$

$$w^+ \geq 0, \quad w^- \geq 0 \quad (3.18)$$

$$0 \leq u, v \leq e \quad (3.19)$$

Penalizing, as in (2.14), the (nonlinear) constraint (3.13) we obtain

$$z^* = \min_{w^+, w^-, \gamma, \xi, \zeta, u, v} e^\top (u + v) + C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) + \sigma(e^\top (w^+ + w^-) - (u - v)^\top (w^+ - w^-)) \quad (3.20)$$

subject to

$$a_i^\top (w^+ - w^-) - \gamma + 1 \leq \xi_i, \quad i = 1, \dots, m_1 \quad (3.21)$$

$$-b_l^\top (w^+ - w^-) + \gamma + 1 \leq \zeta_l, \quad l = 1, \dots, m_2 \quad (3.22)$$

$$\xi_i \geq 0, \quad i = 1, \dots, m_1 \quad (3.23)$$

$$\zeta_l \geq 0, \quad l = 1, \dots, m_2, \quad (3.24)$$

$$w^+ \geq 0, \quad w^- \geq 0 \quad (3.25)$$

$$0 \leq u, v \leq e \quad (3.26)$$

Remark 3.1 Definition of an exact nondifferentiable penalty function of the type described in [6] would require the introduction into the objective function of the term

$$\sigma \max (0, e^\top (w^+ + w^-) - (u - v)^\top (w^+ - w^-)),$$

which in our case, according to Remark 2.1, is simply replaced by

$$\sigma(e^\top (w^+ + w^-) - (u - v)^\top (w^+ - w^-)),$$

giving rise to a differentiable exact penalty function

In a way analogous to that followed in previous section, problem (3.20)–(3.26) can be put in DC form. In fact, letting $s^\top \triangleq (w^{+T}, w^{-T}, u^\top, v^\top)$, the function

$$r(s) \triangleq (v - u)^\top (w^+ - w^-)$$

can be rewritten as

$$r(s) = s^\top \hat{Q}_1 s - s^\top \hat{Q}_2 s,$$

where the symmetric positive semidefinite matrices \hat{Q}_1 and \hat{Q}_2 of dimension $(4n, 4n)$ are defined as follows

$$\hat{Q}_1 = \frac{1}{4} \begin{pmatrix} 2I & -2I & -I & I \\ -2I & +2I & I & -I \\ -I & I & I & 0 \\ I & -I & 0 & I \end{pmatrix} \tag{3.27}$$

and

$$\hat{Q}_2 = \frac{1}{4} \begin{pmatrix} 2I & -2I & I & -I \\ -2I & +2I & -I & I \\ I & -I & I & 0 \\ -I & I & 0 & I \end{pmatrix}, \tag{3.28}$$

with I and 0 being, respectively, the identity and the zero matrix of dimension (n, n) .

The objective function is then decomposed in DC form $f_1(w^+, w^-, \gamma, \xi, \zeta, u, v) - f_2(w^+, w^-, u, v)$ with

$$f_1(w^+, w^-, \gamma, \xi, \zeta, u, v) = e^\top (u + v) + C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) + \sigma(e^\top (w^+ + w^-) + s^\top \hat{Q}_1 s),$$

and

$$f_2(w^+, w^-, u, v) = \sigma s^\top \hat{Q}_2 s$$

We apply to problem above the DCA method [19], which tackles the unconstrained minimization of a function $q : \mathbb{R}^n \rightarrow \mathbb{R}$

$$q(x) = q_1(x) - q_2(x),$$

with q_1 and q_2 convex, by solving a sequence of linearized convex problems. In particular, letting $x^{(k)}$ be the estimate of a (local) minimum of g at iteration k , the next iterate $x^{(k+1)}$ is calculated as

$$x^{(k+1)} = \arg \min_{x \in \mathbb{R}^n} \left\{ q_1(x) - [q_2(x^{(k)}) + g^{(k)\top}(x - x^{(k)})] \right\},$$

with $g^{(k)} \in \partial q_2(x^{(k)})$. For other methods to solve DC problems see [10,11,18] and the references therein.

We remark that, in applying the successive linearization method to our case, a convex quadratic minimization problem is solved at each iteration.

3.1 Mixed binary programming formulation MBP₀

Before reporting in next section the results of the implementation of our algorithm to solve SVM_0 , we provide the following mixed binary programming formulation MBP_0 [9] of the ℓ_0 pseudo norm FS, which is based on the introduction of the set of binary variables $y_j, j = 1, \dots, n$.

$$z^* = \min_{w, \gamma, \xi, \zeta, y} \|w\|_1 + C \left(\sum_{i=1}^{m_1} \xi_i + \sum_{l=1}^{m_2} \zeta_l \right) + D \sum_{j=1}^n y_j$$

subject to

$$\begin{aligned} a_i^\top w + \gamma + 1 &\leq \xi_i, & i &= 1, \dots, m_1 \\ -b_l^\top w - \gamma + 1 &\leq \zeta_l, & l &= 1, \dots, m_2 \\ -u_j y_j &\leq w_j \leq u_j y_j, & j &= 1, \dots, n \\ \xi_i &\geq 0, & i &= 1, \dots, m_1 \\ \zeta_l &\geq 0, & l &= 1, \dots, m_2 \\ y_j &\in \{0, 1\}, & j &= 1, \dots, n \end{aligned}$$

where $u_j > 0, j = 1, \dots, n$, is a given bound on the modulus of the j -th component of w (see [9] for a discussion on setting the u_j 's). The binary variable $y_j, j = 1, \dots, n$ is equal to 1 at the optimum if and only if $w_j \neq 0, j = 1, \dots, n$. Consequently the term $\sum_{j=1}^n y_j$ in the objective function represents the ℓ_0 pseudo-norm of w . The positive parameter D provides the trade-off between the ℓ_0 pseudo-norm objective and that of the SVM_1 problem (3.6)–(3.11).

4 Numerical experiments

The core of our numerical experiments has been the comparison of the results provided by SVM_0 with those of MBP_0 . We have also implemented, as a useful reference, the standard SVM_1 model. For better explanation of the results of SVM_0 we have introduced a discussion on tuning of the penalty parameter σ .

We have also provided some additional results comparing our approach with the smoothing methods described in [4,25].

Table 1 Description of the datasets

Datasets of Group 1	m	n	Datasets of Group 2	m	n
Carcinoma (CARC)	36	7457	Breast cancer (BC)	683	10
DLBCL	77	7129	PIMA indians diabetes (PIMA)	768	8
Leukemia (LEK)	72	5327	HEART	270	13
Tumor1 (TUM1)	60	7129	Ionosphere (IONO)	351	34
Tumor2 (TUM2)	50	12,625	Liver disorders (LIVER)	345	5

4.1 SVM₀ versus MBP₀

We have performed our experiments on two groups of five datasets each. They are the same datasets adopted as benchmark for the feature selection method described in [9]. In particular in datasets 1–5 (Group 1), available at <http://www.tech.plym.ac.uk/spmc/>, the number of samples is small with respect to the number of features. The opposite happens for Datasets 6–10 (Group 2), which are available at <https://www.csie.ntu.edu.tw/~cjlin/libsvmtools/datasets/>. Thus for the latter ones a certain class-overlap is expected.

The datasets are listed in Table 1, where $m = m_1 + m_2$ is the total number of samples.

As a possible reference, we report first the results of the $SV M_1$ problem provided by CPLEX solver. A standard tenfold cross validation has been performed. The results are in Table 2, where the columns “Test” and “Train” indicate the average testing and training correctness, respectively, expressed as percentage of samples correctly classified. Columns “ $\|w\|_1$ ” and “Time” report the average ℓ_1 norm of w and the average execution time (in seconds). Finally columns “%ft(0)”–“%ft(–9)” report the average percentage of components of w whose modulus is greater than or equal to 10^0 – 10^{-9} , respectively. Note that, assuming, conventionally, to be equal to “zero” any component w_j of w such that $|w_j| < 10^{-9}$, the percentage of zero-components is $(100 - \%ft(-9))$.

Two different values of parameter C have been adopted for the two dataset groups. In fact we have taken the “best” values obtained through the so called “Model selection” phase performed in [9], where a grid of values ranging in the interval $[10^{-1}, 10^2]$ has been considered.

Before reporting the results of the implementation of our algorithm to solve $SV M_0$, we illustrate in Tables 3 and 4 the results provided by CPLEX implementation of model MBP_0 , described in Sect. 3.1, with maximum running time of 1000 s. Note that on some test problems, the maximum running time has been achieved with no optimality certification. The results we provide in such cases are related to the best solution found.

Some comments are in order. We observe first that the average computation time for solving the Linear Program $SV M_1$ is negligible.

As for classification performance, comparison of Table 2 with Tables 3 and 4 highlights that the use of an explicit feature selection mechanism results in a mild downgrading of the classification correctness. Such phenomenon is compensated by

Table 2 SVM_1 —Cplex implementation—tenfold cross validation

Dataset	Test	Train	$ w _1$	Time	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
$C = 1$								
CARC	97.50	100.00	2.63	0.27	0.00	0.26	0.28	0.28
DLBCL	94.29	100.00	5.07	0.54	0.00	0.42	0.44	0.44
LEUK	96.67	100.00	4.45	0.37	0.00	0.54	0.58	0.58
TUM1	76.67	100.00	7.09	0.52	0.00	0.53	0.55	0.55
TUM2	80.50	100.00	5.12	0.69	0.00	0.23	0.24	0.24
$C = 10$								
BC	97.24	94.00	6.01	0.01	9.00	94.00	94.00	94.00
PIMA	77.62	100.00	8.30	0.02	26.25	100.00	100.00	100.00
HEART	86.83	100.00	7.27	0.01	12.31	100.00	100.00	100.00
IONO	95.46	95.59	43.43	0.01	50.59	95.00	95.59	95.59
LIVER	68.27	100.00	9.33	0.00	80.00	100.00	100.00	100.00

Table 3 MBP_0 —Cplex implementation—tenfold cross validation—Group 1

Dataset	Test	Train	$ w _1$	Time	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
$C = 1, D = 0.01$								
CARC	92.50	100.00	2.65	302.07	0.00	0.16	0.16	0.16
DLBCL	92.86	100.00	5.11	999.98	0.00	0.28	0.28	0.28
LEUK	93.57	100.00	4.48	999.74	0.00	0.35	0.35	0.35
TUM1	76.67	100.00	7.12	998.91	0.00	0.39	0.39	0.39
TUM2	83.00	100.00	5.16	999.89	0.00	0.16	0.16	0.16

Table 4 MBP_0 —Cplex implementation—tenfold cross validation—Group 2

Dataset	Test	Train	$ w _1$	Time	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
$C = 10, D = 10$								
BC	96.42	97.15	5.73	0.06	17.00	68.00	68.00	68.00
PIMA	75.87	77.26	7.75	0.06	26.25	78.75	78.75	78.75
HEART	83.95	86.97	5.16	0.07	9.23	58.46	58.46	58.46
IONO	88.29	94.62	27.06	2.03	36.18	45.29	45.29	45.29
LIVER	63.02	62.95	3.00	0.01	60.00	60.00	60.00	60.00

the reduction in the percentage of the numerically significant features (columns “ft(0)”–“ft(-9)”).

Coming now to our method (referred to, in the sequel, as the “ SVM_0 Algorithm”), we report in Table 5 the results on the two groups of datasets.

We have added the two columns “%Viol.” and “ $e^T(u + v)$ ”. In particular, column “%Viol.” reports the percentage ratio between the average violation of the relaxed constraint (3.13) and the average norm $\|w\|_1$. Column “ $e^T(u + v)$ ” reports the average

Table 5 $SV M_0$ algorithm—tenfold cross validation

Dataset	σ	Test	Train	$ w _1$	Time	%Viol.	$e^\top(u+v)$	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
$C = 1$											
CARC	4	97.50	100.00	6.90	39.40	3.00	2.10	0.03	0.05	0.06	0.06
DLBCL	4	91.43	98.57	12.83	260.48	0.50	3.51	0.05	0.06	0.06	0.06
LEUK	5	98.33	99.83	16.02	188.70	0.00	3.70	0.07	0.07	0.07	0.07
TUM1*	5	76.67	100.00	15.27	50.74	4.70	7.23	0.08	0.14	0.16	0.16
TUM2	4	81.00	96.03	7.40	69.68	7.29	3.05	0.02	0.05	0.06	0.06
$C = 10$											
BC	63	93.17	92.48	1.36	0.59	0.29	1.32	2.00	34.00	34.00	34.00
PIMA	17	75.57	76.19	5.89	3.80	0.00	3.50	26.25	43.75	43.75	43.75
HEART	150	82.32	84.32	2.32	0.09	38.02	2.54	0.00	49.23	50.00	50.00
IONO	17	86.35	89.31	3.14	0.27	0.47	2.56	5.88	13.82	13.82	13.82
LIVER	80	65.60	66.63	4.88	0.19	0.00	2.60	46.00	52.00	52.00	52.00

value of the scalar product $e^\top(u+v)$, which, for small values of the companion parameter “%Viol.,” reasonably approximates $\|w\|_0$.¹

We have run the algorithm on each dataset for different values of the penalty parameter σ , and we indicate in Table 5 the specific value of σ the results refer to.

Some comments follow.

- In terms of classification correctness, the results of the $SV M_0$ Algorithm are comparable with those of $MB P_0$.
- The $SV M_0$ Algorithm provides better results in terms of number of zero-components of w . In fact, the percentage of components conventionally assumed equal to zero, that is $(100 - \%ft(-9))$, is significantly bigger, except that in two cases, in $SV M_0$ Algorithm than in $MB P_0$.
- The computation time is, for both algorithms, negligible on the datasets of Group 2 while it is remarkably smaller for $SV M_0$ Algorithm as far as datasets of Group 1 are concerned.

4.2 Tuning of the penalty parameter

In running the $SV M_0$ Algorithm, the most relevant issue is the appropriate tuning of the penalty parameter $\sigma > 0$, which is, of course, dataset-specific. Two aspects are to be taken into account.

- “Small” values of σ may lead to significant violation of the penalized constraint at the optimum of problem (3.20)–(3.23). Note that in such case, variables u and v may lose their “marker” role highlighted in Proposition (2.2).
- “Large” values of σ may result in trivial solutions ($w = 0$) to the penalized problem (see Proposition 2.4).

¹ Symbol “*” in Dataset column of Table 5 indicates that parameter C has been set to 2.

Table 6 Results on dataset CARC for different values of σ

σ	Test	Train	$ w _1$	Time	% Viol.	$e^\top(u+v)$	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
<i>Dataset: CARC; C = 1</i>										
1	95.00	100.00	2.62	14.45	100	0.00	0.00	0.23	0.25	0.25
2	94.17	100.00	3.50	73.13	44	0.87	0.01	0.14	0.15	0.15
4	97.50	100.00	6.90	39.40	3	2.10	0.03	0.05	0.06	0.06
5	91.67	100.00	6.26	41.41	1	2.07	0.02	0.03	0.03	0.03
6	85.00	98.33	7.23	68.53	0.1	2.18	0.03	0.03	0.03	0.03
7	75.83	96.99	6.08	93.80	0	2.20	0.03	0.03	0.03	0.03
12	65.83	81.40	2.32	81.04	0	0.80	0.01	0.01	0.01	0.01
16	58.33	57.59	0.0	1.81	0	0.00	0.00	0.00	0.00	0.00

Table 7 Results on dataset IONO for different values of σ

σ	Test	Train	$ w _1$	Time	% Viol.	$e^\top(u+v)$	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
<i>Dataset: IONO; C = 10</i>										
1	87.61	94.23	19.39	0.36	52	4.16	18.82	72.94	73.24	73.24
2	87.94	93.53	11.21	0.22	50	4.34	9.41	62.06	63.53	63.53
4	87.96	92.55	7.58	0.26	42	4.31	6.18	47.35	47.35	47.35
10	87.63	89.38	4.23	0.24	19	2.90	5.88	18.53	19.41	19.41
17	86.35	89.31	3.14	0.29	15	2.56	5.88	13.82	13.82	13.82
30	80.80	83.47	1.83	0.23	6	1.77	3.82	7.65	7.94	7.04
50	74.77	74.68	1.00	0.26	3	0.97	2.94	2.94	2.94	2.94
64	64.33	64.24	0.0001	1.84	0	0.0001	0.00	0.00	0.29	0.29
93	64.33	64.24	0.00	0.16	0	0.00	0.00	0.00	0.00	0.00

To illustrate in details the impact on the solution of parameter σ , we analyze the results for increasing values of σ . In particular we focus on the datasets CARC and IONO, from the first and second group, respectively. The results are reported in Tables 6 and 7.

We observe that on both datasets increasing values of σ results, as expected, in deterioration of the classification correctness (under such point of view, significant data are those obtained in the training phase). On the other hand, the violation of the relaxed constraint gets smaller and smaller, until, for too large values of σ , the results become meaningless, as w gets close to zero.

4.3 SVM₀ versus smoothing methods

To provide a better assessment of the proposed method we present some comparisons with results taken from [4,25], where a number of smoothing approaches to ℓ_0 pseudo-norm minimization in SVM framework are described.

Table 8 Comparison with *FSV* algorithm [4]

Dataset	<i>FSV</i>			<i>SVM₀</i>		
	Training correct.	% Nonzero	Time	Training correct.	% Nonzero	Time
PIMA	75.52	66.25	–	76.19	43.75	3.80
Heart	85.30	49.23	–	84.32	50.00	0.09
IONO	88.92	30.58	30.94	89.31	13.82	0.27
LIVER	67.83	75.00	–	66.63	52.00	0.19

Table 9 Description of the datasets

Datasets from [25]	<i>m</i>	<i>n</i>
Colon cancer	62	2000
Nova	1754	16,969

Table 10 *SVM₀* algorithm

Dataset	<i>C</i>	σ	Train	$ w _1$	Time	%Viol.	$e^\top(u+v)$	%ft(0)	%ft(-2)	%ft(-4)	%ft(-9)
Colon cancer	10	40	100.00	9.29	14.42	0.00	9.00	0.25	0.45	0.45	0.45
Nova	100	31	100.00	252.88	137.07	0.00	131.00	0.64	0.77	0.77	0.77

As for comparison with [4] we take in consideration the datasets of Group 2, which have been in fact adopted as test problems in [4]. We do not consider here the Breast Cancer (BC) dataset, since two different variants of such dataset have been used in this paper and in [4]. In Table 8 we compare training correctness and percentage of zero-value features provided by *FSV* algorithm from [4] and by our *SVM₀*. In reporting the results of *SVM₀*, we have assumed equal to zero any component of w smaller than 10^{-9} .

The results are comparable in terms of correctness, while *SVM₀* seems to produce a stronger feature reduction effect. Comparison in terms of computing time does not appear particularly significant, as for *FSV* the available data provided in [4] refer only to IONO dataset.

As for comparison with [25], we have considered the datasets of Table 9, which are both linearly separable.

In Table 10 we report the detailed results obtained by *SVM₀*.

In [25] two different formulations of the sparse minimization problem are discussed; they are based on two smooth concave approximations of the zero pseudo-norm inspired by property (2.3) and are referred to, respectively, as Formulation 1 and 2. The approach adopted is a multi-start one, with 100 runs for randomly selected starting point. The comparison is in Table 11, where for Formulation 1 and 2 are reported both the *average* and the *best* value of number of nonzero features. As for *SVM₀*, the average number of nonzero components in the tenfold cross validation procedure is, as usual, reported.

Table 11 Comparison with [25]

Dataset	Formulation 1		Formulation 2		$SV M_0$	
	# Nonzero feat.	% Time	# Nonzero feat.	Time	# Nonzero feat.	Time
Colon cancer	13.7/7	225	9.4/6	24	9	14.42
Nova	126.7/105	10,448	131.9/114	1003	131	137.7

We note, however, that the approach in [25] is a *global* optimization one, while ours is *local*. This fact makes comparison difficult, mainly from point of view of computation time.

5 Conclusions

We have tackled the Feature Selection problem within SVM binary classification by using the polyhedral k -norm, in the sparse optimization context. The numerical experiments show that the approach is a promising alternative to continuous approximations of the ℓ_0 pseudo-norm and to integer programming-based methods.

As Feature Selection is an important subject also in other areas of Machine Learning (see, e.g., [7,28]), we believe that a possible future development of our work would be the extension of the k -norm approach to regression and clustering problems. Yet another possible research direction would be the use of kernel transformations within our model.

Acknowledgements We are grateful to Francesco Rinaldi for having provided us with the datasets *Colon Cancer* and *Nova* that we have used to compare our results with those in [25].

References

1. Amaldi, E., Kann, V.: On the approximability of minimizing nonzero variables or unsatisfied relations in linear systems. *Theor. Comput. Sci.* **209**(1–2), 237–260 (1998)
2. Bertolazzi, P., Felici, G., Festa, P., Fiscon, G., Weitschek, E.: Integer programming models for feature selection: new extensions and a randomized solution algorithm. *Eur. J. Oper. Res.* **250**(2), 389–399 (2016)
3. Bradley, P.S., Mangasarian, O.L., Street, W.N.: Feature selection via mathematical programming. *INFORMS J. Comput.* **10**(2), 209–217 (1998)
4. Bradley, P.S., Mangasarian, O.L.: Feature selection via concave minimization and support vector machines. In: Shavlik, J., (ed.) *Machine Learning Proceedings of the Fifteenth International Conference (ICML '98)*. Morgan Kaufmann, San Francisco, California, pp. 82–90 (1998)
5. Cristianini, N., Shawe-Taylor, J.: *An Introduction to Support Vector Machines and Other Kernel-Based Learning Methods*. Cambridge University Press, Cambridge (2000)
6. Di Pillo, G., Grippo, L.: Exact penalty functions in constrained optimization. *SIAM J. Control Optim.* **27**(6), 1333–1360 (1989)
7. Dy, J.G., Brodley, C.E., Wrobel, S.: Feature selection for unsupervised learning. *J. Mach. Learn. Res.* **5**, 845–889 (2004)
8. Gasso, G., Rakotomamonjy, A., Canu, S.: Recovering sparse signals with a certain family of nonconvex penalties and DC programming. *IEEE Trans. Signal Process.* **57**(12), 4686–4698 (2009)
9. Gaudioso, M., Gorgone, E., Labbé, M., Rodríguez-Chía, A.M.: Lagrangian relaxation for SVM feature selection. *Comput. Oper. Res.* **87**, 137–145 (2017)

10. Gaudioso, M., Giallombardo, G., Miglionico, G.: Minimizing piecewise-concave functions over polytopes. *Math. Oper. Res.* **43**(2), 580–597 (2018)
11. Gaudioso, M., Giallombardo, G., Miglionico, G., Bagirov, A.M.: Minimizing nonsmooth DC functions via successive DC piecewise-affine approximations. *J. Glob. Optim.* **71**(1), 37–55 (2018)
12. Gotoh, J., Takeda, A., Tono, K.: DC formulations and algorithms for sparse optimization problems. *Math. Program. Ser. B* **169**(1), 141–176 (2018)
13. Guyon, I., Elisseeff, A.: An introduction to variable and feature selection. *J. Mach. Learn. Res.* **3**, 1157–1182 (2003)
14. Hastie, T., Tibshirani, R., Wainwright, M.: *Statistical Learning with Sparsity: The Lasso and Generalizations*. CRC Press, Boca Raton (2015)
15. Hempel, A.B., Goulart, P.J.: A novel method for modelling cardinality and rank constraints. In: 53rd IEEE Conference on Decision and Control, Los Angeles, CA, USA December 15–17, pp. 4322–4327 (2014)
16. Hiriart-Urruty, J.-B.: Generalized Differentiability/Duality and Optimization for Problems Deling with Differences of Convex Functions, *Lecture Notes in Economic and Mathematical Systems*, vol. 256, pp. 37–70. Springer, Berlin (1986)
17. Hiriart-Urruty, J.-B., Ye, D.: Sensitivity analysis of all eigenvalues of a symmetric matrix. *Numer. Math.* **70**(1), 45–72 (1995)
18. Joki, K., Bagirov, A.M., Karmitsa, N., Mäkelä, M.M.: A proximal bundle method for nonsmooth DC optimization utilizing nonconvex cutting planes. *J. Glob. Optim.* **68**, 501–535 (2017)
19. Le Thi, H.A., Dinh, T.P.: The DC (difference of convex functions) programming and DCA revisited with DC models of real world nonconvex optimization problems. *J. Glob. Optim.* **133**, 23–46 (2005)
20. Le Thi, H.A., Le, H.M., Nguyen, V.V., Dinh, T.P.: A DC programming approach for feature selection in support vector machines learning. *Adv. Data Anal. Classif.* **2**, 259–278 (2008)
21. Maldonado, S., Pérez, J., Weber, R., Labbé, M.: Feature selection for support vector machines via mixed integer linear programming. *Inf. Sci.* **279**, 163–175 (2014)
22. Mangasarian, O.L.: *Nonlinear Programming*. McGraw-Hill, New York (1969)
23. Overton, M.L., Womersley, R.S.: Optimality conditions and duality theory for minimizing sums of the largest eigenvalues of symmetric matrices. *Math. Program.* **62**(1–3), 321–357 (1993)
24. Pilanci, M., Wainwright, M.J., El Ghaoui, L.: Sparse learning via Boolean relaxations. *Math. Program. Ser. B* **151**, 63–87 (2015)
25. Rinaldi, F., Schoen, F., Sciandrone, M.: Concave programming for minimizing the zero-norm over polyhedral sets. *Comput. Optim. Appl.* **46**, 467–486 (2010)
26. Strekalovsky, A.S.: Global optimality conditions for nonconvex optimization. *J. Glob. Optim.* **12**, 415–434 (1998)
27. Soubies, E., Blanc-Féraud, L., Aubert, G.: A unified view of exact continuous penalties for ℓ_2 - ℓ_0 minimization. *SIAM J. Optim.* **27**(3), 2034–2060 (2017)
28. Tibshirani, R.: Regression shrinkage and selection via the lasso. *J. R. Stat. Soc. Ser. B* **58**, 267–288 (1996)
29. Vapnik, V.: *The Nature of the Statistical Learning Theory*. Springer, Berlin (1995)
30. Wang, L., Zhu, J., Zou, H.: The doubly regularized support vector machine. *Stat. Sin.* **16**, 589–615 (2006)
31. Watson, G.A.: Linear best approximation using a class of polyhedral norms. *Numer. Algorithms* **2**, 321–336 (1992)
32. Weston, J., Elisseeff, A., Schölkopf, B., Tipping, M.: Use of the zero-norm with linear models and kernel methods. *J. Mach. Learn. Res.* **3**, 1439–1461 (2003)
33. Wright, S.J.: Accelerated block-coordinate relaxation for regularized optimization. *SIAM J. Optim.* **22**(1), 159–186 (2012)
34. Wu, B., Ding, C., Sun, D., Toh, K.-C.: On the Moreau–Yosida regularization of the vector k -norm related functions. *SIAM J. Optim.* **24**(2), 766–794 (2014)
35. Zou, H., Hastie, T.: Regularization and variable selection via the elastic net. *J. R. Stat. Soc. Ser. B* **67**, 301–320 (2005)