



Correction to: Sepsis patients with complication of hypoglycemia and hypoalbuminemia are an early and easy identification of high mortality risk

Makoto Furukawa¹ · Kosaku Kinoshita¹ · Junko Yamaguchi¹ · Satoshi Hori¹ · Atsushi Sakurai¹

Published online: 26 October 2019
© The Author(s) 2019

Correction to: Internal and Emergency Medicine (2019) 14:539–548
<https://doi.org/10.1007/s11739-019-02034-2>

In the original article, there were errors in Unit of serum albumin level and the Cases numbers in Table 4.

Unit of serum albumin level was published incorrectly as 2.8 mg/dl throughout the article. The correct unit should read as 2.8 g/dl.

In Table 4, the numbers appearing in column 2, under “Inter -G + non- hypo-A” and column 6, “Hypo-A” was published incorrectly as 112 and 147. The corrected numbers should read as 102 and 156 respectively.

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1007/s11739-019-02034-2>.

✉ Kosaku Kinoshita
kinoshita.kosaku@nihon-u.ac.jp

Makoto Furukawa
Furukawa.makoto@nihon-u.ac.jp

Junko Yamaguchi
yamaguchi.junko@nihon-u.ac.jp

Satoshi Hori
horisa02@yahoo.co.jp

Atsushi Sakurai
sakurai.atsushi@nihon-u.ac.jp

¹ Division of Emergency and Critical Care Medicine, Department of Acute Medicine, Nihon University School of Medicine, 30-1, Oyaguchi Kami-cho, Itabashi-ku, Tokyo 173-8610, Japan