

Development of a Fish Cell Culture Model to Investigate the Impact of Fish Oil Replacement on Lipid Peroxidation

Melissa K. Gregory · Hamish W. King ·
Peter A. Bain · Robert A. Gibson · Douglas R. Tocher ·
Kathryn A. Schuller

Received: 28 February 2011 / Accepted: 5 April 2011 / Published online: 29 April 2011
© AOCS 2011

Abstract Fish oils are rich in omega-3 long-chain polyunsaturated fatty acids (n-3 LC-PUFA), predominantly 20:5n-3 and 22:6n-3, whereas vegetable oils contain abundant C₁₈-PUFA, predominantly 18:3n-3 or 18:2n-6. We hypothesized that replacement of fish oils with vegetable oils would increase the oxidative stability of fish lipids. Here we have used the long established and easily cultivated FHM cell line derived from the freshwater fish species fathead minnow (*Pimephales promelas*) to test this hypothesis. The FHM cells were readily able to synthesize 20:5n-3 and 24:6n-3 from 18:3n-3 but 22:6n-3 synthesis was negligible. Also, they were readily able to synthesize 20:3n-6 from 18:2n-6 but 20:4n-6 synthesis was negligible. Mitochondrial β -oxidation was greatest for 18:3n-3 and 20:5n-3 and the rates for 16:0, 18:2n-6, 22:6n-3 and 18:1n-9 were significantly lower. Fatty acid incorporation was predominantly into phospholipids (79–97%) with very little incorporation into neutral lipids. Increasing the fatty acid concentration in the growth medium substantially increased the concentrations of 18:3n-3 and 18:2n-6 in the cell phospholipids but this was not the case for 20:5n-3 or 22:6n-3. When they were subjected to oxidative stress, the FHM cells supplemented with either 20:5n-3 or 22:6n-3

(as compared with 18:3n-3 or saturated fatty acids) exhibited significantly higher levels of thiobarbituric reactive substances (TBARS) indicating higher levels of lipid peroxidation. The results are discussed in relation to the effects of fatty acid unsaturation on the oxidative stability of cellular lipids and the implications for sustainable aquaculture.

Keywords Aquaculture · β -oxidation · Cell culture · Fish oil replacement · Lipid peroxidation · Phospholipids · Polyunsaturated fatty acids

Abbreviations

ALA	α -Linolenic acid
ARA	Arachidonic acid
BHT	Butylated hydroxytoluene
CerPCho	Sphingomyelin
DHA	Docosahexaenoic acid
EDTA	Ethylenediamine tetraacetic acid
EPA	Eicosapentaenoic acid
FAF-BSA	Fatty acid free-bovine serum albumin
FAME	Fatty acid methyl esters
FBS	Foetal bovine serum
FHM	Fathead minnow
HP-TLC	High performance-thin layer chromatography
LC-PUFA	Long-chain polyunsaturated fatty acids (carbon chain length \geq C ₂₀ with \geq 3 double bonds)
LNA	Linoleic acid
NR	Neutral red
OLA	Oleic acid
PAM	Palmitic acid
PBS	Phosphate buffered saline
PBSA	Phosphate buffered saline without Ca ²⁺ or Mg ²⁺
PCR	Polymerase chain reaction
PtdCho	Phosphatidylcholine

M. K. Gregory · H. W. King · P. A. Bain · K. A. Schuller (✉)
School of Biological Sciences, Flinders University,
GPO Box 2100, Adelaide, SA 5001, Australia
e-mail: kathy.schuller@flinders.edu.au

R. A. Gibson
School of Agriculture, Food and Wine, University of Adelaide,
Adelaide, Australia

D. R. Tocher
Institute of Aquaculture, University of Stirling,
Stirling FK9 4LA, UK

PtdEtn	Phosphatidylethanolamine
Ptd ₂ Gro	Cardiolipin
PtdIns	Phosphatidylinositol
PtdOH	Phosphatidic acid
PtdSer	Phosphatidylserine
PUFA	Polyunsaturated fatty acids
SDS	Sodium dodecyl sulphate
STA	Stearic acid
TBA	Thiobarbituric acid
TBARS	Thiobarbituric acid reactive substances
TLC	Thin-layer chromatography
TN	Total neutral lipids
T/V	Trypsin/versene

Introduction

Fisheries and aquaculture are major contributors to world food security with $\geq 15\%$ of animal protein for human consumption being derived from these sources in recent years [1]. Traditionally, feeds for farmed fish have contained high proportions of fish oil derived from wild-catch fisheries. However, in the 10 years prior to 2007, wild-catch fisheries production has remained static whereas aquaculture production has almost doubled [1]. Thus, there has been considerable interest in the replacement of fish oils with more sustainable oils [2, 3]. The preferred candidates have been vegetable oils from oilseed plants. However, while these are rich in C₁₈ polyunsaturated fatty acids (PUFA), they are completely lacking in the omega-3 long-chain PUFA (n-3 LC-PUFA) that are abundant in fish oils. These n-3 LC-PUFA give seafood its reputation as a health food. In particular, eicosapentaenoic acid (EPA, 20:5n-3) and docosahexaenoic acid (DHA, 22:6n-3) have been shown to be beneficial in the prevention of cardiovascular disease, rheumatoid arthritis, inflammatory bowel disease, childhood learning and behavior disorders and adult psychiatric and neurodegenerative illnesses [4].

The fatty acid profile of fish flesh generally reflects the fatty acid profile of their diet [5]. Therefore, the dietary substitution of fish oils with vegetable oils reduces the n-3 LC-PUFA content of fish flesh thereby reducing its human health value [6–10]. In freshwater fish and salmonids (e.g. trout and salmon), this may be partially offset by their varying ability to synthesize LC-PUFA from C₁₈ PUFA but in marine fish this is not the case [11–14]. Marine fish appear to be poorly able to synthesize LC-PUFA from C₁₈ PUFA and this may reflect the relative abundance of pre-formed LC-PUFA, particularly 22:6n-3, in the marine food web [5]. Although freshwater fish and salmonids have some capacity to synthesize LC-PUFA from C₁₈ PUFA,

this is insufficient to compensate for a decreased dietary intake of LC-PUFA.

The generally accepted pathway for the synthesis of LC-PUFA from C₁₈ PUFA in fish, as in mammals, is a series of desaturation and elongation steps in the endoplasmic reticulum followed by chain shortening in the peroxisomes (Fig. 1). In particular, the synthesis of 22:6n-3 from 18:3n-3 involves a $\Delta 6$ desaturation step followed by a chain elongation step, a $\Delta 5$ desaturation step and two consecutive chain elongation steps before a second $\Delta 6$ desaturation step and the final chain shortening step. Recent studies indicate that the C₁₈ and C₂₀ elongation steps are catalysed by an ELOVL5-like elongase whereas the C₂₂ elongation step is catalysed by an ELOVL2-like elongase [15, 16]. Studies with fish cell lines indicate that freshwater fish and salmonids have substantial activities of the desaturase and elongase enzymes required to synthesize 20:5n-3 from 18:3n-3 whereas for marine fish there is an apparent limitation at either the C₁₈ elongation or the $\Delta 5$ desaturation step [11–14]. In contrast, synthesis of 22:6n-3 appears to be limited in all fish species.

Fish lipids are highly susceptible to peroxidative deterioration because of the high proportions of LC-PUFA they contain and this explains the shorter shelf life of fish flesh products as compared with terrestrial animal meat products [17]. The C₁₈ PUFA abundant in vegetable oils, 18:2n-6 and 18:3n-3, contain 2 and 3 double bonds, respectively, whereas the LC-PUFA abundant in fish oils, 20:5n-3 and 22:6n-3, contain 5 and 6 double bonds, respectively. Lipid peroxidation occurs when reactive oxygen species (ROS) attack PUFA at their double bonds setting off a chain reaction of hydrogen abstraction and lipid radical formation. This is particularly damaging to cell membranes because of the close proximity of fatty acids in the phospholipid bilayer [17]. Thus, a cell culture model would be useful to precisely define the effects of fish oil replacement at the cellular level.

The abundance of individual fatty acids in cellular lipids, particularly phospholipids, is affected by their uptake and esterification into the different lipid classes as well as by their mitochondrial β -oxidation for energy generation and their possible desaturation and/or elongation to form LC-PUFA [5]. Thus, all of these factors need to be considered when developing a cell culture model. Here we have used the long established and easily cultivated FHM cell line derived from the freshwater fish species fathead minnow (*Pimephales promelas*). In particular we have investigated the capacity of the cell line to synthesize LC-PUFA from C₁₈ PUFA, to metabolize various fatty acids by β -oxidation and to incorporate various fatty acids into cellular lipids. Finally, we have investigated the impact of fatty acid unsaturation on the susceptibility of the cells to lipid peroxidation.




Fig. 1 The generally accepted pathway for the synthesis n-3 and n-6 LC-PUFA from their C₁₈ fatty acid precursors in fish (adapted from Miller et al. [2]) showing the alternative reactions when $\Delta 6$ fatty acyl desaturase activity is limiting (broken lines). $\Delta 5$ $\Delta 5$ fatty acyl

desaturase, $\Delta 6$ $\Delta 6$ fatty acyl desaturase, ARA arachidonic acid, ALA α -linolenic acid, DHA docosahexaenoic acid, ELOVL2 fatty acyl elongase 2, ELOVL5 fatty acyl elongase 5, EPA eicosapentaenoic acid, LNA linoleic acid, short peroxisomal shortening of LC-PUFA

Materials and Methods

Cells, Media and Standard Culture Conditions

A stock culture of the FHM cell line derived from fathead minnow (*Pimephales promelas*) was obtained from the Australian Animal Health Laboratory (CSIRO Livestock Industries, Geelong, Victoria, Australia) and the species of origin was confirmed by polymerase chain reaction (PCR) using the mitochondrial cytochrome oxidase subunit I (*cox1*) primers of Ward et al. [18]. For routine culture, the FHM cells were maintained in 75 cm² flasks containing 20 ml Leibovitz's L-15 medium supplemented with antibiotics (100 IU ml⁻¹ penicillin, 100 μ g ml⁻¹ streptomycin) and 10% (v/v) foetal bovine serum (FBS). To subculture or harvest the cells, the medium was decanted, the cell monolayer was rinsed with phosphate buffered saline without Ca²⁺ or Mg²⁺ (PBSA) and the cells were detached with T/V solution containing 0.05% (w/v) trypsin and 0.02% (w/v) ethylenediamine tetraacetic acid (EDTA) in PBSA. For the experiments with [1-¹⁴C]-labelled fatty acids, the cells were cultured to 80% confluence in 75 cm² flasks containing 20 ml L-15 medium supplemented with 5% (v/v) FBS. The medium was then replaced with 10 ml of fresh medium containing no FBS and the [1-¹⁴C]-labelled fatty acids as complexes with fatty acid free-bovine serum albumin (FAF-BSA). The complexes were prepared as described by Ghioni et al. [19]. For the experiments with unlabelled fatty acids, the cells were cultured to 80% confluence in 75 cm² flasks containing 20 ml L-15 medium supplemented with 10% (v/v) FBS and then they were subcultured (at a split ratio of 1 to 12) into 25 cm² flasks containing 10 ml L-15 medium supplemented with 2% (v/v) FBS and the unlabelled fatty acids as complexes with FAF-BSA. The complexes were prepared as described by Best et al. [20] with a molar ratio of fatty acid to BSA of 4:1. For all experiments, the culture temperature was 25 °C and

the incubation period with either the [1-¹⁴C]-labelled or the unlabelled fatty acids was 24 h.

LC-PUFA Synthesis Assay Using [1-¹⁴C]-labelled Fatty Acids

The FHM cells were incubated for 24 h in 75 cm² flasks containing 1 μ Ci (2 μ M) [1-¹⁴C]-labelled 18:2n-6, 18:3n-3 or 20:5n-3 in 10 ml L-15 medium containing no FBS. At the end of the 24-h incubation, the cells were detached with T/V solution and washed with FAF-BSA to remove any residual labelled fatty acid. Total lipids were extracted from the cell pellets by the addition of 5 ml ice-cold chloroform/methanol (2:1, v/v) containing 0.01% (v/v) butylated hydroxytoluene (BHT) followed by 1 ml 0.88% (w/v) KCl. The method was essentially that of Folch et al. [21] with the modifications described by Tocher et al. [22]. Fatty acid methyl esters (FAME) were prepared by incubating the lipid extract overnight at 50 °C with 1 ml of toluene and 2.5 ml of 1% (v/v) H₂SO₄ in methanol. The method was essentially that of Christie [23]. At the end of the incubation, the FAME were extracted by the addition of 2 ml of 2% (w/v) KHCO₃ followed by 5 ml isohexane/diethyl ether (1:1, v/v) containing 0.01% (v/v) BHT. The mixture was centrifuged at 500g and the upper layer was re-extracted with 5 ml isohexane/diethyl ether (1:1, v/v) containing no BHT. The upper layers were combined and the solvent was evaporated under a stream of N₂. The dried FAME were dissolved in 100 μ l isohexane containing 0.01% (w/v) BHT and separated by thin-layer chromatography (TLC) on 20 \times 20 cm TLC plates which had been impregnated with 2 g AgNO₃ in 20 ml acetonitrile before being dried/activated at 110 °C for 30 min. The TLC plates were developed in toluene/acetonitrile (95:5, v/v). The method was essentially as described by Wilson and Sargent [24]. Autoradiography was performed and the areas of silica containing the individual PUFA were scraped into

scintillation vials and radioactivity determined using a scintillation counter as previously described [25].

β -Oxidation Assay Using [1- 14 C]-labelled Fatty Acids

The FHM cells were incubated for 24 h in 75-cm² flasks containing 0.5 μ Ci (1 μ M) [1- 14 C]-labelled 16:0, 18:1n-9, 18:2n-6, 18:3n-3, 20:5n-3 or 22:6n-3 in 10 ml L-15 medium containing no FBS. At the end of the incubation, 0.5 ml of the medium was removed for scintillation counting and the cells were detached with T/V solution and washed as described above. The washed cell pellets were resuspended in 1 ml PBS and homogenized to disrupt the cells. Acid soluble products were obtained by adding 100 μ l of 6% (w/v) FAF-BSA and 1.0 ml ice-cold 4 M HClO₄ to 500 μ l each of the growth medium and the cell homogenate and then centrifuging to remove the precipitated material. An aliquot of the supernatant (500 μ l) was mixed with 4 ml scintillation fluid and radioactivity determined using a scintillation counter as previously described [25].

Incorporation of [1- 14 C]-labelled Fatty Acids into Various Lipid Classes

The FHM cells were incubated for 24 h in 75 cm² flask containing 1.0 μ Ci (2 μ M) [1- 14 C]-labelled 16:0, 18:1n-9, 18:2n-6, 18:3n-3, 20:5n-3 or 22:6n-3 in 10 ml L-15 medium containing no FBS. At the end of the 24-h incubation, the cells were harvested and total lipid was extracted as described above. To separate the various lipid classes, the total lipid was dissolved in 100 μ l chloroform/methanol (2:1, v/v) and applied to a high performance-thin layer chromatography (HPTLC) plate. The plate was developed in methyl acetate/isopropanol/chloroform/methanol/0.25% (w/v) aqueous KCl (25/25/25/10/9, by volume) as described by Vitiello and Zanetta [26]. The separated lipids were stained with iodine and the corresponding bands of silica were scraped into scintillation vials. Scintillation fluid (2.5 ml) was added and radioactivity determined using a scintillation counter as above.

Incorporation of Unlabelled Fatty Acids into the FHM Cell Phospholipids

The FHM cells were incubated for 24 h in 25-cm² flasks containing 10 ml L-15 medium supplemented with 2% (v/v) FBS and various concentrations of unlabelled 18:3n-3, 18:2n-6, 20:5n-3, 20:4n-6 or 22:6n-3. At the end of the incubation, the cells were harvested and total lipid was extracted from approximately 10⁶ cells according to the method of Folch et al. [21]. The phospholipids were

separated from the neutral lipids by TLC with petroleum ether/glacial acetic acid (3:1, v/v) and then they were transmethylated by incubation with 1% (v/v) H₂SO₄ in methanol for 3 h at 70 °C. The resulting FAME were extracted in heptane and analyzed by gas chromatography using a Hewlett-Packard 6890 gas chromatograph (Hewlett Packard, Palo Alto, CA, USA) fitted with a flame ionisation detector and a BPX-70 50 m capillary column coated with 70% (v/v) cyanopropyl polysilphenylene-siloxane (0.25 mm film thickness and 0.32 mm internal diameter, SGE, Australia). The carrier gas was helium at a flow rate of 2.0 ml min⁻¹ and the split-ratio was 20:1. The injection port temperature was 250 °C and the detector temperature was 300 °C. The column temperature was increased from 140 to 220 °C at a rate of 4 °C min⁻¹ and then held at 220 °C for up to 3 min. The identity of each fatty acid peak in the chromatogram was ascertained by comparison with an authentic lipid standard (Nu-Chek Prep, Inc., MN, USA). The amount of each fatty acid was quantified by comparing its peak area with the peak area of a heptadecaenoic acid (17:0) internal standard added prior to lipid extraction.

Effect of Fatty Acid Unsaturation on Cell Viability and Lipid Peroxidation

The FHM cells were seeded into 6-well plates at a density of 5 × 10⁵ cells per well in L-15 medium containing 2% (v/v) FBS. The cells were allowed to attach overnight and then they were incubated for 24 h with fresh medium containing 2% (v/v) FBS and 20 μ M 16:0, 18:0, 18:3n-3, 20:5n-3 or 22:6n-3 coupled with FAF-BSA in a 4:1 fatty acid to BSA ratio. At the end of the incubation period, the cell monolayer was rinsed twice with PBSA and then lipid peroxidation was induced by treating the cells for 1 h with 500 μ M cumene hydroperoxide plus 100 nM hemin in PBSA. Following this treatment, the cell monolayer was rinsed twice with PBSA and the cells were subjected to either the Neutral Red (NR) cell viability assay [27] or the thiobarbituric acid reactive substances (TBARS) assay for lipid peroxidation [28]. Briefly, the NR cell viability assay involved incubating the cells for 3 h in NR dye diluted in L-15 containing 10% (v/v) FBS and then fixing them in 10% (w/v) CaCl₂:4% (v/v) formaldehyde before solubilizing the incorporated dye with 50% (v/v) ethanol:1% (v/v) acetic acid and reading the absorbance at 550 nm. Viable cells take up the dye whereas non-viable cells do not. For the TBARS assay, the cells were lysed with 1% (w/v) sodium dodecyl sulphate (SDS), the cell lysates were mixed 1:1 with the TBARS reagent and the mixture was incubated for 2 h at 75 °C before removing any insoluble material by centrifugation and reading the fluorescence

using excitation/emission wavelengths of 540/590 nm. The TBARS reagent contained 25 mM thiobarbituric acid (TBA), 15% (w/v) trichloroacetic acid and 0.005% (w/v) BHT dissolved in 1 M HCl.

Statistical Analyses

Statistical analyses were performed using SPSS software. The Levene statistic was calculated to test for homogeneity of variances and means were compared using one-way ANOVA followed by Tukey's post-hoc test. Differences where the *P* value was >0.05 were considered to be significant.

Results

Effect of the FBS Concentration in the Growth Medium on the Fatty Acid Composition of the FHM Cell Phospholipids

The fatty acid composition of the FHM cell phospholipids closely resembled that of the FBS in the growth medium, with the exceptions that 16:0 and 18:0 were less abundant and 18:1n-9 was more abundant (Table 1). Increasing the concentration of FBS in the growth medium from 2 to 20% (v/v) significantly increased the concentrations of 22:5n-3, 22:6n-3 and 20:4n-6 in the cell phospholipids (Fig. 2).

Metabolism of [1-¹⁴C]-labelled Fatty Acids by the FHM Cells

Radioactivity from [1-¹⁴C]18:3n-3 was readily recovered in 20:4n-3, 20:5n-3, 22:5n-3 and 24:6n-3 but there was relatively little recovery in 22:6n-3 (Table 2). Thus, the cells expressed considerable desaturase and elongase activity but only limited peroxisomal chain shortening activity. This was confirmed by supplying the cells with [1-¹⁴C]20:5n-3. A somewhat different pattern was observed when the cells were supplied with [1-¹⁴C]18:2n-6. In this case, the label was incorporated predominantly into 20:3n-6. This indicated considerable $\Delta 6$ desaturase activity to yield 18:3n-6 followed by elongase activity to yield 20:3n-6 but limited $\Delta 5$ desaturase activity to yield 20:4n-6. Thus, the FHM $\Delta 5$ desaturase apparently preferred the n-3 over the n-6 substrate.

The capacity of the FHM cells to metabolise fatty acids by mitochondrial β -oxidation was also investigated (Fig. 3). This showed that 18:3n-3 and 20:5n-3 were more readily oxidized than 16:0, 18:2n-6 or 22:6n-3 which in turn were more readily oxidized than 18:1n-9. This result was consistent with the high retention levels for 18:1n-9 in the cell phospholipids (Table 1).

Table 1 Fatty acid composition of foetal bovine serum (FBS) and the phospholipids of FHM cells maintained in L-15 medium containing 10% (v/v) FBS

Fatty acid	Fatty acid composition (% of total fatty acids)	
	FBS	FHM cells in L-15 + 10% FBS
14:0	0.8	1.3
16:0 (PAM)	23.0	17.8
18:0 (STA)	18.8	9.0
Total saturates ^a	45.3	32.0
16:1n-9	0.9	2.0
16:1n-7	1.3	2.2
18:1n-9 (OLA)	15.3	24.1
18:1n-7	2.0	7.9
Total monounsaturates ^b	23.5	42.4
18:2n-6 (LNA)	7.6	1.8
18:3n-6	0.3	0.4
20:2n-6	0.2	0.4
20:3n-6	3.4	3.0
20:4n-6 (ARA)	9.9	8.4
Total n-6 PUFA ^c	23.3	15.3
18:3n-3 (ALA)	0	0.2
20:5n-3 (EPA)	0.3	0.2
22:5n-3 (DPA)	3.7	1.5
22:6n-3 (DHA)	4.0	2.9
Total n-3 PUFA ^d	8.1	4.9
Total PUFA	31.4	20.2
n-3/n-6	0.35	0.32

Data are expressed as % of total fatty acids. For FBS, the data are the means of 3 separate batches. For the FHM cells, the data are the mean of 3 replicate flasks; ^a also includes 15:0, 17:0, 20:0, 22:0 and 24:0; ^b also includes 20:1, 22:1 and 24:1; ^c also includes 22:4n-6; ^d also includes 18:4n-3 and 20:4n-3

PAM palmitic acid, STA stearic acid, OLA oleic acid, LNA linoleic acid, ARA arachidonic acid, ALA α -linolenic acid, EPA eicosapentaenoic acid, DHA docosahexaenoic acid,

Incorporation of [1-¹⁴C]-labelled Fatty Acids into Lipid Classes in the FHM Cells

Total incorporation into all lipid classes was similar for 18:1n-9, 18:2n-6 and 20:5n-3 but significantly greater for 18:3n-3 and significantly less for 16:0 and 22:6n-3 (Table 3). The lower incorporation rate for 16:0 was consistent with its lower steady state level in the FHM cell phospholipids as compared with FBS (Table 1). This contrasts with the higher steady state level for 18:1n-9 which was apparently due to a lower rate of β -oxidation. For all of the fatty acids tested, the majority of the label (79–97%) was incorporated into phospholipids with very little incorporation into neutral lipids. For 16:0, 18:1n-9




Fig. 2 The effect of increasing FBS concentration in the growth medium on the PUFA content of the FHM cell phospholipids. The growth medium contained 2% (v/v) BSA, 2 μM 18:3n-3 and increasing concentrations of FBS and the cells were cultured for 24 h at 25 $^{\circ}\text{C}$. Each data point represents the mean of 3 flasks and the vertical bars represent the standard error of the mean. For each fatty acid, data points with different letters represent values that are significantly different from one another at the $P = 0.05$ level

and 18:2n-6, most of the label (44–50%) was incorporated into phosphatidylcholine (PtdCho) with phosphatidylethanolamine (PtdEtn) ranked second. For 18:3n-3 and 20:5n-3, the ranking was reversed with the greatest incorporation into PtdEtn followed by PtdCho. For 22:6n-3, the pattern was different again with the greatest incorporation into PtdEtn followed by phosphatidylserine (PtdSer) and then PtdCho. Thus, the lower overall incorporation of 22:6n-3 into total lipids was associated with lower incorporation into PtdCho and higher incorporation into PtdSer. In addition, as the unsaturation of the fatty acids increased, there was a trend away from incorporation into PtdCho.

Effect of Concentration on the Accumulation of Unlabelled Fatty Acids in the FHM Cell Phospholipids

Figure 4 shows the effects of increasing concentrations of 18:3n-3 or 18:2n-6 in the growth medium on their accumulation in the cell phospholipids. In both cases, accumulation in the cell phospholipids paralleled the

Table 2 Metabolism of [$1\text{-}^{14}\text{C}$]-labelled 18:3n-3, 20:5n-3 and 18:2n-6 by desaturation and elongation in FHM cells

Fatty acid	Substrate		
	[$1\text{-}^{14}\text{C}$]18:3n-3	[$1\text{-}^{14}\text{C}$]20:5n-3	[$1\text{-}^{14}\text{C}$]18:2n-6
18:3n-3	14.5 \pm 0.7	–	–
20:3n-3	7.7 \pm 0.3	–	–
18:4n-3/24:5n-3	9.9 \pm 0.5	–	–
20:4n-3	18.3 \pm 0.3	–	–
22:4n-3	1.9 \pm 0.2	–	–
20:5n-3	22.1 \pm 0.8	42.7 \pm 1.6	–
22:5n-3	15.2 \pm 0.5	26.9 \pm 0.7	–
24:5n-3	– ^a	8.3 \pm 1.3	–
22:6n-3	1.0 \pm 0.1	3.2 \pm 0.7	–
24:6n-3	9.4 \pm 0.6	18.9 \pm 0.5	–
18:2n-6	–	–	53.4 \pm 1.4
20:2n-6	–	–	6.7 \pm 0.3
18:3n-6	–	–	7.1 \pm 1.0
20:3n-6	–	–	26.8 \pm 1.0
22:3n-6	–	–	1.6 \pm 0.2
20:4n-6	–	–	4.5 \pm 0.3

^a Indicates not applicable. Data are expressed as percentage of total radioactivity recovered in total lipid and are means \pm SD ($n = 4$)

increase in concentration in the growth medium up to a concentration of 5 μM but above this concentration an incorporation threshold was reached. The accumulation of 18:3n-3 was roughly paralleled by an accumulation of 20:3n-3, the direct elongation product of 18:3n-3. Thus, $\Delta 6$ desaturase activity, which would have produced 18:4n-3, appeared to be limiting. In contrast, 18:2n-6 was a poor substrate for direct elongation as evidenced by limited accumulation of 20:2n-6. However, 20:3n-6 accumulation was significant and this is consistent with the earlier observation of limited $\Delta 5$ desaturase activity towards 20:3n-6 in the [$1\text{-}^{14}\text{C}$]-labelling experiment (Table 2).

Further metabolism of 18:3n-3 to produce 20:5n-3 and 22:6n-3 or of 18:2n-6 to produce 20:4n-6 was negligible. This was surprising given the results of the [$1\text{-}^{14}\text{C}$]-labelling experiment which showed significant synthesis of 20:5n-3 from 18:3n-3 (Table 2). To determine whether this apparent anomaly was due to limited incorporation of higher concentrations of LC-PUFA into the cellular phospholipids, the FHM cells were supplied with various concentrations of either 20:5n-3, 20:4n-6 or 22:6n-3 (Figs. 5, 6). Incorporation of 20:5n-3 into the cell phospholipids increased approximately linearly with the increase in the concentration in the growth medium up to a concentration of 5 μM whereas at higher concentrations it reached a plateau. The accumulation of 20:5n-3 was paralleled by an accumulation of 22:5n-3, the direct elongation product of 20:5n-3. Further metabolism of




Fig. 3 β -oxidation activity of the FHM cell line with various fatty acids. The cells were incubated for 24 h at 25 °C with 1 μ Ci (2 μ M) of the [¹⁴C]-labelled fatty acids in L-15 medium containing no FBS. Data points with different letters represent values that are significantly different from one another at the $P = 0.05$ level

20:5n-3 to produce 22:6n-3 by chain shortening was negligible.

Incorporation of 20:4n-6 was less straightforward (Fig. 5). At lower concentrations up to 5 μ M, there was some evidence of increasing incorporation with increasing concentration in the medium. However, at higher concentrations in the medium (5–20 μ M) there was no increase in incorporation of 20:4n-6 into the cells. This was not unexpected given that the cells were cultured in the presence of 2% (v/v) FBS which is relatively rich in 20:4n-6 and therefore the capacity of the cell phospholipids to incorporate 20:4n-6 may already have reached its maximum level.

Incorporation of 22:6n-3 into the cell phospholipids reached its maximum level at a very low concentration of only 2 μ M in the growth medium (Fig. 6). This was lower than the concentration of approximately 5 μ M observed for 20:5n-3. This was consistent with the ¹⁴C-labelling data

which showed significantly lower incorporation of 22:6n-3 into total lipid (Table 3). Interestingly, there was also some evidence for retro-conversion of 22:6n-3 to 20:5n-3 (Fig. 6).

Effect of Fatty Acid Unsaturation on Cell Viability and Lipid Peroxidation

There was no significant difference in cell viability between any of the fatty acid treatments except for the treatment with 18:0 which showed significantly higher cell viability (Fig. 7). For TBARS, the concentration was significantly greater in the cells supplemented with either 20:5n-3 or 22:6n-3 as compared with the cells supplemented with either 16:0, 18:0 or 18:3n-3. Thus, increasing fatty acid unsaturation increased the susceptibility of the cells to lipid peroxidation but did not decrease cell viability.

Discussion

The aim of this study was to develop a cell culture model for the impact of fish oil replacement with vegetable oils on the oxidative stability of fish lipids. Fish oils are rich in LC-PUFA with 5 or 6 double bonds whereas vegetable oils are rich in C₁₈ PUFA with only 2 or 3 double bonds [3] and since the fatty acid composition of fish lipids reflects the fatty acid composition of their diet and fatty acid susceptibility to peroxidation increases with increasing unsaturation, we hypothesized that replacing LC-PUFA with C₁₈ PUFA would increase the oxidative stability of fish lipids. Previous studies had shown that fish cells in culture have decreased proportions of n-3 LC-PUFA in their lipids as compared with whole fish or fish fillets and this was

Table 3 Incorporation of [¹⁴C]-labelled fatty acids into total lipid and lipid classes in FHM cells

	Substrate					
	[¹⁴ C]16:0	[¹⁴ C]18:1n-9	[¹⁴ C]18:2n-6	[¹⁴ C]18:3n-3	[¹⁴ C]20:5n-3	[¹⁴ C]22:6n-3
Total incorporation (pmol mg ⁻¹ protein)	315 ± 12 ^a	382 ± 37 ^b	432 ± 36 ^b	561 ± 13 ^c	430 ± 13 ^b	239 ± 24 ^d
Incorporation into lipid classes (% of total)						
PtdCho	48.7 ± 3.5 ^{ab}	50.2 ± 3.0 ^a	43.8 ± 1.7 ^b	38.8 ± 0.4 ^c	31.8 ± 2.2 ^d	18.8 ± 0.3 ^e
PtdEtn	16.2 ± 1.6 ^a	21.5 ± 0.5 ^b	18.2 ± 0.2 ^a	40.3 ± 0.8 ^c	52.9 ± 1.9 ^d	49.7 ± 0.7 ^e
PtdSer	5.3 ± 0.1 ^a	6.1 ± 0.8 ^{ab}	6.8 ± 0.3 ^b	6.9 ± 0.3 ^b	7.0 ± 0.4 ^b	23.5 ± 0.7 ^c
PtdIns	5.7 ± 0.2 ^a	8.3 ± 0.9 ^b	12.2 ± 1.9 ^c	2.7 ± 0.1 ^d	3.7 ± 0.2 ^{ad}	3.3 ± 0.2 ^d
PtdOH/Ptd ₂ Gro	2.9 ± 1.4 ^a	2.0 ± 0.3 ^a	10.7 ± 0.3 ^b	7.1 ± 0.8 ^c	1.5 ± 0.1 ^a	1.5 ± 0.1 ^a
CerPCho	13.4 ± 3.4 ^a	5.2 ± 3.0 ^b	2.9 ± 1.0 ^{bc}	1.0 ± 0.3 ^{bc}	0.5 ± 0.2 ^c	0.5 ± 0.1 ^c
TN	7.7 ± 0.2 ^a	6.7 ± 0.2 ^b	5.4 ± 0.3 ^c	3.3 ± 0.0 ^d	2.7 ± 0.2 ^e	2.7 ± 0.1 ^e

Data are the means ± SD ($n = 4$). Within each row, values sharing the same superscript are not significantly different at the $P = 0.05$ level. CerPCho sphingomyelin, PtdCho phosphatidylcholine, PtdEtn phosphatidylethanolamine, PtdIns phosphatidylinositol, PtdOH phosphatidic acid, PtdSer phosphatidylserine, Ptd₂Gro cardiolipin, TN total neutral lipids




Fig. 4 The effect of increasing 18:3n-3 or 18:2n-6 concentration in the growth medium on the PUFA content of the FHM cell phospholipids. The growth medium contained 2% (v/v) BSA and increasing concentrations of either 18:3n-3 or 18:2n-6 and the cells were cultured for 24 h at 25 °C. Each data point represents the mean

of 3 flasks and the vertical bars represent the standard error of the mean. For each fatty acid, data points with different letters represent values that are significantly different from one another at the $P = 0.05$ level

attributed to the low concentrations of n-3 LC-PUFA in the mammalian serum which is the usual source of fatty acids for cells in culture [13, 22, 29]. The present study confirmed this and in particular we found that the proportions of the beneficial n-3 LC-PUFA, 20:5n-3 and 22:6n-3, were much lower in the FHM cells in culture than in fish tissues in general.

The effects of fish oil replacement on the fatty acid profile of fish flesh are determined by fatty acid metabolism and incorporation into cellular lipids. Previous studies had shown that cell lines derived from fresh water or anadromous fish species (Atlantic salmon, rainbow trout and common carp) were readily able to synthesize 20:5n-3 from 18:3n-3 whereas cell lines derived from carnivorous marine fish species (turbot and gilthead sea bream) were not [11–14]. In turbot this was attributed to low C_{18} to C_{20} elongase activity whereas in gilthead sea bream it was attributed to low $\Delta 5$ desaturase activity. Importantly, none of these cell lines could synthesize significant quantities of 22:6n-3. In the present study, the FHM cell line was readily able to synthesize 20:5n-3, 22:5n-3 and 24:6n-3 from 18:3n-3 and/or 20:5n-3 but 22:6n-3 synthesis was negligible. Thus, the FHM cell line expressed substantial $\Delta 5$ and $\Delta 6$ desaturase activities as well as C_{18-20} , C_{20-22} and C_{22-24}

elongase activities but negligible peroxisomal chain shortening activity. This was similar to cell lines from other freshwater fish species.

The FHM cells were also readily able to desaturate 18:2n-6 to 18:3n-6 and further elongate this to 20:3n-6 but they synthesized little 20:4n-6. This suggested limited $\Delta 5$ desaturase activity towards n-6 fatty acids, which was in contrast to the results obtained with the n-3 substrates. Thus, the $\Delta 5$ desaturase of the FHM cell line appeared to prefer n-3 over n-6 fatty acids. Previous studies with cell lines from Atlantic salmon, rainbow trout, turbot and gilt-head sea bream also showed limited $\Delta 5$ desaturase activity towards 20:3n-6 [11–13]. Thus, this appears to be a general phenomenon for fish cell lines regardless of whether they originate from freshwater, anadromous or marine fish species.

Fatty acid utilization via β -oxidation in the FHM cells was investigated by supplying them with [$1-^{14}C$]-labelled 16:0, 18:1n-9, 18:2n-6, 18:3n-3, 20:5n-3 or 22:6n-3. The β -oxidation rates were similar for 16:0, 18:2n-6 and 22:6n-3 but significantly lower for 18:1n-9 and significantly higher for 18:3n-3 and 20:5n-3. In a previous study with Atlantic salmon hepatocytes isolated from fish fed a fish oil-based diet, β -oxidation rates were similar for 16:0 and 18:2n-6,




Fig. 5 The effect of increasing 20:5n-3 or 20:4n-6 concentration in the growth medium on the PUFA content of the FHM cell phospholipids. The growth medium contained 2% (v/v) BSA and increasing concentrations of either 20:5n-3 or 20:4n-6 and the cells were cultured

for 24 h at 25 °C. Each data point represents the mean of 3 flasks and the vertical bars represent the standard error of the mean. For each fatty acid, data points with different letters represent values that are significantly different from one another at the $P = 0.05$ level




Fig. 6 The effect of increasing 22:6n-3 concentration in the growth medium on the PUFA content of the FHM cell phospholipids. The growth medium contained 2% (v/v) BSA and increasing concentrations of 22:6n-3 and the cells were cultured for 24 h at 25 °C. Each data point represents the mean of 3 flasks and the vertical bars represent the standard error of the mean. For each fatty acid, data points with different letters represent values that are significantly different from one another at the $P = 0.05$ level

slightly lower for 18:1n-9 and much lower for 18:3n-3, 20:5n-3 and 22:6n-3 [30]. Feeding the fish a predominantly vegetable oil-based diet significantly increased the β -oxidation of 18:1n-9, 18:2n-6, 18:3n-3 and 20:5n-3 but the rank order was still similar. Clearly the metabolism of primary hepatocytes is likely to be different to that of cells




Fig. 7 The effect of increasing fatty acid unsaturation on cell viability and lipid peroxidation in the FHM cells. The cells were supplied with the various fatty acids at a concentration of 20 μ M in L-15 medium containing 2% (v/v) FBS. The control contained ethanol at the concentration present in the fatty acid supplements. Three separate flasks of cells were analyzed for each data point and the columns represent the mean while the vertical bars represent the standard deviation. For the TBARS assay, * indicates a significant difference (at the $P = 0.05$ level) between the results for 20:5n-3 and 22:6n-3 and the results for the other fatty acids. For the NR assay * indicates a significant difference (at the $P = 0.05$ level) between the results for 18:0 and the results for the other fatty acids

maintained in culture for long periods of time and therefore more work needs to be done to understand β -oxidation in fish cell lines. In particular, the rates measured using

radiolabelled substrates may be affected by the pool sizes of unlabelled substrates in the cells due to dilution of the label. The FHM cell phospholipids were particularly rich in 16:0 and 18:1n-9, with moderate levels of 18:2n-6 and 22:6n-3 and low levels of 18:3n-3 and 20:5n-3. Thus, the β -oxidation rates for 16:0 and 18:1n-9, in particular, may have been underestimated. In addition, the incorporation of 22:6n-3 into total cellular lipids was significantly lower compared with the other fatty acids and this may have resulted in an underestimation of the β -oxidation rate for this fatty acid as well. Overall, therefore, the FHM cells were readily able to oxidize all six of the different fatty acids supplied to them.

The incorporation of fatty acids into cellular lipids in the FHM cells was studied in two different ways, firstly using a trace (2 μ M) concentration of [1- 14 C]-labelled fatty acids and secondly using increasing concentrations of unlabelled fatty acids in the range 1–20 μ M. Previous studies with fish cell lines and trace concentrations of [1- 14 C]-labelled 18:2n-6, 18:3n-3 and 20:5n-3 had shown that the label was incorporated predominantly (mostly >90%) into phospholipids, especially PtdCho and PtdEtn, with very little incorporation into neutral lipids [11, 13, 14]. The FHM cell line behaved the same way. In addition, we found that [1- 14 C]-labelled 16:0, 18:1n-9 and 22:6n-3 were also incorporated predominantly into phospholipids. Interestingly, there was substantial incorporation of [1- 14 C]-labelled 22:6n-3 into PtdSer and less incorporation into PtdCho than for the other fatty acids. This has also been observed for cell lines from rainbow trout, Atlantic salmon and turbot [31–33]. PtdSer is less abundant than PtdCho in fish cell lines in general [13, 14] and this may explain the lower level of incorporation of [1- 14 C]-labelled 22:6n-3 into total lipid observed in the present study.

Increasing the concentration of unlabelled 18:3n-3 in the culture medium from 1 to 20 μ M resulted in a roughly corresponding increase in the concentration of this fatty acid in the FHM cell phospholipids but there was no corresponding increase in the conventional desaturation/elongation products 20:5n-3, 22:5n-3 or 24:6n-3. Instead, there was an accumulation of 20:3n-3. This fatty acid is produced by the direct elongation of 18:3n-3. Thus, Δ 6 desaturase activity apparently limited the flux through the conventional pathway as the 18:3n-3 concentration was increased above the trace concentration used in the [1- 14 C]-labelling experiments. This was in contrast to the results of previous studies with rainbow trout and Atlantic salmon cell lines supplied increasing concentrations of unlabelled fatty acids [34, 35]. In those studies, increasing the concentration of unlabelled 18:3n-3 from 5 to 50 μ M resulted in a greater increase in 20:5n-3 than 20:3n-3. The main difference between the present and the previous studies is that the previous studies analyzed total lipid whereas in the

present study we analyzed phospholipids. Thus, there may have been incorporation into neutral lipids which was not detected in the present study. Tocher et al. [34] noted that there were no large increases in neutral lipids in either rainbow trout or turbot cell lines supplied fatty acids at concentrations up to 20 μ M however lipid droplets did appear when the concentration was increased above 50 μ M. Thus, in the future, it will be interesting to investigate the incorporation of fatty acids into neutral lipids in the FHM cells and compare this with their incorporation into phospholipids to determine the relative distributions as fatty acid concentrations are increased.

Incorporation of 18:2n-6 into the FHM cell phospholipids with increasing concentrations of this fatty acid in the growth medium showed a similar pattern to that observed with 18:3n-3 except that 20:3n-6 was the major desaturation/elongation product incorporated. This is produced by the conventional Δ 6 desaturation of 18:2n-6 to yield 18:3n-6 followed by the conventional elongation of 18:3n-6 to yield 20:3n-6. Thus, the result obtained with unlabelled 18:2n-6 was similar to that obtained with labelled 18:2n-6 and this confirmed that Δ 5 desaturase was the limiting step in n-6 LC-PUFA synthesis in the FHM cell line. This was consistent with the results of previous work with other freshwater and anadromous fish cell lines [34, 35]. Taking the results with 18:3n-3 and 18:2n-6 together, it suggests that fish cells in culture are suitable for the study of the effects of increasing concentrations of C₁₈ PUFA derived from vegetable oils. However, in order to have a control representative of a fish oil-based diet, it will be necessary to supply the cultured cells with either 20:5n-3 or 22:6n-3 to significantly raise the levels of these fatty acids in the cellular lipids. Thus, this was also investigated.

When the FHM cells were supplied with increasing concentrations of 20:5n-3, there was a corresponding increase in the concentration of this fatty acid in their phospholipids but only up to a culture medium concentration of 10 μ M. Above this concentration, the incorporation of 20:5n-3 into the cell phospholipids did not change significantly. Thus, incorporation of 20:5n-3 reached a plateau at a lower concentration than incorporation of either of the C₁₈ fatty acids. This was in contrast to the results obtained with a cell line from turbot, a marine fish species [34]. In that case 20:5n-3 incorporation continued increasing up to a culture medium concentration of at least 50 μ M. Perhaps this reflects a greater capacity for incorporation of 20:5n-3 in marine fish than in freshwater fish but it should also be noted that total lipid was analyzed in the previous study whereas only phospholipids were analyzed in the present study. The direct incorporation of 22:6n-3 into the FHM cell phospholipids was also investigated. This reached a plateau at a very low concentration

of only 2 μM in the culture medium. This has not been studied before and it clearly warrants further investigation.

In general fish flesh products (i.e. seafood) have a shorter shelf life than terrestrial animal meat products and this is due to the greater susceptibility of fish flesh to lipid peroxidation as a result of the higher proportions of LC-PUFA in fish lipids [17]. Thus, it can be hypothesized that feeding farmed fish vegetable oils with their abundance of either 18:3n-3 or 18:2n-6 as opposed to fish oils with their abundance of 20:5n-3 and 22:6n-3 will result in extended shelf life. To test this hypothesis in our in vitro system, we enriched the FHM cells with either saturated fatty acids (16:0 or 18:0), PUFA (18:3n-3) or LC-PUFA (20:5n-3 or 22:6n-3) and then subjected them to oxidative stress to stimulate lipid peroxidation. The cells enriched with 20:5n-3 or 22:6n-3 had significantly higher concentrations of TBARS than the cells enriched with 16:0, 18:0 or 18:3n-3. TBARS is a commonly used indicator of lipid peroxidation in fish [17]. Thus, increasing the unsaturation of the FHM lipids increased the susceptibility of the cells to lipid peroxidation. Therefore in this respect, the cells are a good model for the effects of dietary fish oil replacement on the flesh quality of farmed fish. Previous studies with Atlantic salmon fed diets containing either fish oil, vegetable oil or oils enriched with either 20:5n-3 or 22:6n-3 showed that increasing dietary unsaturation increased indicators of oxidative stress and apoptosis in liver and white adipose tissue [36, 37]. In particular, fish fed diets enriched with either 20:5n-3 or 22:6n-3 had greater activities of the antioxidant enzyme superoxide dismutase and the apoptosis marker caspase 3 as well as reduced integrity of their mitochondria as evidenced by almost undetectable levels of β -oxidation of ^{14}C -palmitoyl CoA. More direct evidence for the effect of dietary fatty acid unsaturation on the oxidative stability of fish lipids comes from a study with grass carp fed diets containing either lard, plant oil or fish oil [38]. In that study, TBARS concentration in the plasma of the fish increased with increasing dietary lipid content and also with increasing unsaturation of the dietary oil. It has also been shown that supplementing the growth medium with the antioxidant α -tocopherol (vitamin E) decreased the longer term inhibitory effect of 20:5n-3 supplementation on the growth of a turbot cell line [39].

In conclusion, the FHM cells had substantial capacity to synthesize 20:5n-3 from 18:3n-3 but only limited capacity to synthesize 22:6n-3. In addition they had substantial capacity to synthesize 20:3n-6 from 18:2n-6 but only limited capacity to synthesize 20:4n-6. Together the data indicated limited chain shortening activity in the peroxisomes and limited $\Delta 5$ desaturase activity especially towards 20:3n-6 but also towards 20:4n-3. At a trace (2 μM) concentration, fatty acid incorporation into total lipid was greatest for 18:3n-3 followed closely by 18:2n-6

and 20:5n-3 with incorporation of 22:6n-3 being substantially less. Fatty acid incorporation into phospholipids reached a plateau at much lower concentrations for 20:5n-3 and 22:6n-3 than for 18:3n-3 and 18:2n-6 indicating that the cell membranes had limited capacity for incorporation of 20:5n-3 and 22:6n-3 presumably because of the need to maintain appropriate levels of fluidity. Thus, dietary fish oil replacement with vegetable oils may result in substantial displacement of LC-PUFA by C_{18} PUFA in fish cell membranes. Despite the limited capacity of the cell membranes to incorporate either 20:5n-3 or 22:6n-3, the FHM cells still showed greater susceptibility to lipid peroxidation when the culture medium was supplemented with LC-PUFA as compared with either 18:3n-3 or saturated fatty acids. Thus, the FHM cell line can be a useful model for the effects of dietary fish oil replacement on lipid stability and shelf life in farmed fish.

Acknowledgments MKG was the recipient of a Flinders University Postgraduate Research Scholarship and KAS received funding from the Flinders University Outside Studies Program.

References

1. Food and Agriculture Organization of the United Nations (FAO) (2009) The state of world fisheries and aquaculture 2008, Rome
2. Miller MM, Nichols PD, Carter CG (2008) n-3 Oil sources for use in aquaculture—alternatives to the unsustainable harvest of wild fish. *Nutr Res Rev* 21:85–96
3. Turchini GM, Torstensen BE, Ng W-K (2009) Fish oil replacement in finfish nutrition. *Rev Aquac* 1:10–57
4. Calder PC, Yaqoob P (2009) Omega-3 polyunsaturated fatty acids and human health outcomes. *BioFactors* 35:266–272
5. Tocher DR (2003) Metabolism and functions of lipids and fatty acids in teleost fish. *Rev Fish Sci* 11:107–184
6. Torstensen BE, Bell JG, Rosenlund G, Henderson RJ, Graff IE, Tochers DR, Lie Ø, Sargent JR (2005) Tailoring of Atlantic salmon (*Salmo salar* L.) flesh lipid composition and sensory quality by replacing fish oil with a vegetable oil blend. *J Agric Food Chem* 53:10166–10178
7. Zheng X, Torstensen BE, Tocher DR, Dick JR, Henderson RJ, Bell JG (2005) Environmental and dietary influences on highly unsaturated fatty acid biosynthesis and expression of fatty acyl desaturase and elongase genes in liver of Atlantic salmon (*Salmo salar*). *Biochim Biophys Acta* 1734:13–24
8. Leaver MJ, Villeneuve LAN, Obach A, Jensen L, Bron JE, Tocher DR, Taggart JB (2008) Functional genomics reveals increases in cholesterol biosynthetic genes and highly unsaturated fatty acid biosynthesis after dietary substitution of fish oil with vegetable oils in Atlantic salmon (*Salmo salar*). *BMC Genomics* 9:299
9. Pratoomyot J, Bendiksen EA, Bell JG, Tocher DR (2008) Comparison of effects of vegetable oils blended with southern hemisphere fish oil and decontaminated northern hemisphere fish oil on growth performance, composition and gene expression in Atlantic salmon (*Salmo salar* L.). *Aquaculture* 280:170–178
10. Trushenski JT, Boesenberg J (2009) Influence of dietary fish oil concentration and finishing duration on beneficial fatty acid profile restoration in sunshine bass *Morone chrysops* ♀ × *M. saxatilis* ♂. *Aquaculture* 296:277–283

11. Tocher DR, Sargent JR (1990) Effect of temperature on the incorporation into phospholipid classes and metabolism via desaturation and elongation of n-3 and n-6 polyunsaturated fatty acids in fish cells in culture. *Lipids* 25:435–442
12. Ghioni C, Tocher DR, Bell MV, Dick JR, Sargent JR (1999) Low C₁₈ to C₂₀ fatty acid elongase activity and limited conversion of stearidonic acid, 18:4(n-3), to eicosapentaenoic acid, 20:5(n-3), in a cell line from the turbot, *Scophthalmus maximus*. *Biochim Biophys Acta* 1437:170–181
13. Tocher DR, Dick JR (1999) Polyunsaturated fatty acid metabolism in a cell culture model of essential fatty acid deficiency in a freshwater fish, carp (*Cyprinus carpio*). *Fish Physiol Biochem* 21:257–267
14. Tocher DR, Ghioni C (1999) Fatty acid metabolism in marine fish: low activity of fatty acyl $\Delta 5$ desaturation in gilthead sea bream (*Sparus aurata*) cells. *Lipids* 34:433–440
15. Morais S, Monroig O, Zheng X, Leaver MJ, Tocher DR (2009) Highly unsaturated fatty acid synthesis in Atlantic salmon: characterization of ELOVL5- and ELOVL2-like elongases. *Mar Biotechnol* 11:627–639
16. Gregory MK, See VHL, Gibson RA, Schuller KA (2010) Cloning and functional characterisation of a fatty acyl elongase from southern bluefin tuna (*Thunnus maccoyii*). *Comp Biochem Physiol* 155:178–185
17. Mourente G, Bell JG, Tocher DR (2007) Does dietary tocopherol level affect fatty acid metabolism in fish? *Fish Physiol Biochem* 33:269–280
18. Ward RD, Zemlak TS, Innes BH, Last PR, Hebert PDN (2005) DNA barcoding Australia's fish species. *Philos Trans Royal Soc Lond Ser B Biol Sci* 360:1847–1857
19. Ghioni C, Tocher DR, Sargent JR (1997) The effect of culture on morphology, lipid and fatty acid composition, and polyunsaturated fatty acid metabolism of rainbow trout (*Oncorhynchus mykiss*) skin cells. *Fish Physiol Biochem* 16:499–513
20. Best CA, Laposata M, Proios VG, Szczepiorkowski ZM (2006) Method to assess fatty acid ethyl ester binding to albumin. *Alcohol Alcohol* 41:240–246
21. Folch J, Lees M, Sloane Stanley GH (1957) A simple method for the isolation and purification of total lipids from animal tissues. *J Biol Chem* 226:497–509
22. Tocher DR, Sargent JR, Frerichs GN (1988) The fatty acid compositions of established fish cell lines after long-term culture in mammalian sera. *Fish Physiol Biochem* 5:219–227
23. Christie WW (1982) *Lipid analysis*, 2nd edn. Pergamon Press, Oxford
24. Wilson R, Sargent JR (1992) High resolution separation of polyunsaturated fatty acids by argentation thin-layer chromatography. *J Chromatogr* 623:403–407
25. Tocher DR, Fonseca-Madrigal J, Dick JR, Ng W-K, Bell JG, Campbell PJ (2004) Effects of water temperature and diets containing palm oil on fatty acid desaturation and oxidation in hepatocytes and intestinal enterocytes of rainbow trout (*Oncorhynchus mykiss*). *Comp Biochem Physiol Part B* 137:49–63
26. Vitiello F, Zanetta J-P (1978) Thin-layer chromatography of phospholipids. *J Chromatogr* 166:637–640
27. Repetto G, del Peso A, Zurita JL (2008) Neutral red uptake assay for the estimation of cell viability/cytotoxicity. *Nat Protocols* 3:1125–1131
28. Hoyland DV, Taylor AJ (1991) A review of the methodology of the 2-thiobarbituric acid test. *Food Chem* 40:271–291
29. Tocher DR, Dick JR, Sargent JR (1995) Development of an in vitro model of essential fatty acid deficiency in fish cells. *Prostaglandins Leukotrienes Essent Fat Acids* 53:365–375
30. Stubhaug I, Tocher DR, Bell JG, Dick JR, Torstensen BE (2005) Fatty acid metabolism in Atlantic salmon (*Salmo salar* L.) hepatocytes and influence of dietary vegetable oil. *Biochim Biophys Acta* 1734:277–288
31. Tocher DR (1990) Incorporation and metabolism of (n-3) and (n-6) polyunsaturated fatty acids in phospholipid classes in cultured rainbow trout (*Salmo gairdneri*) cells. *Fish Physiol Biochem* 8:239–249
32. Tocher DR, Dick JR (1990) Incorporation and metabolism of (n-3) and (n-6) polyunsaturated fatty acids in phospholipid classes in cultured Atlantic salmon (*Salmo salar*) cells. *Comp Biochem Physiol Part B* 96:73–79
33. Tocher DR, Mackinlay EE (1990) Incorporation and metabolism of (n-3) and (n-6) polyunsaturated fatty acids in phospholipid classes in turbot (*Scophthalmus maximus*) cells. *Fish Physiol Biochem* 8:251–260
34. Tocher DR, Carr J, Sargent JR (1989) Polyunsaturated fatty acid metabolism in fish cells: differential metabolism of (n-3) and (n-6) series acids by cultured cells originating from a freshwater teleost fish and from a marine teleost fish. *Comp Biochem Physiol Part B* 94:367–374
35. Tocher DR, Dick JR (1990) Polyunsaturated fatty acid metabolism in cultured fish cells: incorporation and metabolism of (n-3) and (n-6) series acids by Atlantic salmon (*Salmo salar*) cells. *Fish Physiol Biochem* 8:311–319
36. Kjær MA, Todorčević M, Torstensen BE, Vegusdal A, Ruyter B (2008) Dietary n-3 HUFA affects mitochondrial fatty acid β -oxidation capacity and susceptibility to oxidative stress in Atlantic salmon. *Lipids* 43:813–827
37. Todorčević M, Kjær MA, Djaković N, Vegusdal A, Torstensen BE, Ruyter B (2009) N-3 HUFAs affect fat deposition, susceptibility to oxidative stress, and apoptosis in Atlantic salmon visceral adipose tissue. *Comp Biochem Physiol Part B* 152:135–143
38. Du ZY, Clouet P, Huang LM, Degrace P, Zheng WH, He JG, Tian LX, Liu YJ (2008) Utilization of different dietary lipid sources at high level in herbivorous grass carp (*Ctenopharyngodon idella*): mechanism related to hepatic fatty acid oxidation. *Aquac Nutr* 14:77–92
39. Tocher DR, Dick JR (1991) Effect of polyunsaturated fatty acids on the growth of fish cells in culture. *Comp Biochem Physiol Part A* 100:461–466