

# Complete moment convergence for ND random variables under the sub-linear expectations

FENG Feng-xiang<sup>1,2</sup>      WANG Ding-cheng<sup>2,\*</sup>      WU Qun-ying<sup>1</sup>

**Abstract.** In this article, we establish a general result on complete moment convergence for arrays of rowwise negatively dependent( ND ) random variables under the sub-linear expectations. As applications, we can obtain a series of results on complete moment convergence for ND random variables under the sub-linear expectations.

## §1 Introduction and notation

In the classical probability theory, the additivity of the probabilities and the expectations is assumed. But in practice, such additivity assumption is not feasible in many areas of applications because the uncertainty phenomena can not be modeled using additive probabilities or additive expectations. Non-additive probabilities and non-additive expectations are useful tools for studying uncertainties in statistics, measures of risk, superhedging in finance and non-linear stochastic calculus [7,9,12,14,15]. Peng [15,16]) introduced the general framework of the sub-linear expectation in a general function space by relaxing the linear property of the classical expectation to the sub-additivity and positive homogeneity (cf. Definition 1.1 below). Under Peng's sub-linear expectation framework, many limit theorems have been established recently, including the central limit theorem and weak law of large numbers [16,17,18], strong law of large numbers [3,10,4,22], the law of the iterated logarithm [5,24,22], the moment inequalities for the maximum partial sums and the Kolomogov strong law of large numbers [25], and so on. Investigating the limit theorems in sub-linear expectation space is of great significance in the theory and application. Because sub-linear expectation and capacity are not additive, the study of the limit theorems under sub-linear expectation becomes much more complex and challenging.

---

Received: 2019-02-16.      Revised: 2022-01-25.

MR Subject Classification: 60F15.

Keywords: complete convergence, complete moment convergence, ND random variables, sub-linear expectation.

Digital Object Identifier(DOI): <https://doi.org/10.1007/s11766-023-3759-z>.

Supported by the National Natural Science Foundation of China(71871046, 11661029) and Natural Science Foundation of Guangxi(2018JJB110010).

\*Corresponding author.

©The Author(s) 2023.

Throughout this paper,  $C$  denotes a positive constant which may differ from one place to another. Denote  $a_+ = \max\{0, a\}$ .

We use the framework and notations of [16]. Let  $(\Omega, \mathcal{F})$  be a given measurable space and let  $\mathcal{H}$  be a linear space of real functions defined on  $(\Omega, \mathcal{F})$  such that if  $X_1, \dots, X_n \in \mathcal{H}$  then  $\varphi(X_1, \dots, X_n) \in \mathcal{H}$  for each  $\varphi \in C_{l.Lip}(\mathbb{R}^n)$ , where  $C_{l.Lip}(\mathbb{R}^n)$  denotes the linear space of (local Lipschitz) functions  $\varphi$  satisfying

$$|\varphi(\mathbf{x}) - \varphi(\mathbf{y})| \leq C(1 + |\mathbf{x}|^m + |\mathbf{y}|^m)|\mathbf{x} - \mathbf{y}|, \quad \forall \mathbf{x}, \mathbf{y} \in \mathbb{R}^n,$$

for some  $C > 0, m \in \mathbb{N}$  depending on  $\varphi$ .  $\mathcal{H}$  is considered as a space of "random variables". If  $X$  is an element of set  $\mathcal{H}$ , then we denote  $X \in \mathcal{H}$ .

**Definition 1.1** A sub-linear expectation  $\widehat{\mathbb{E}}$  on  $\mathcal{H}$  is a function  $\widehat{\mathbb{E}} : \mathcal{H} \rightarrow \bar{\mathbb{R}}$  satisfying the following properties: for all  $X, Y \in \mathcal{H}$ , we have

- (a) Monotonicity: If  $X \geq Y$  then  $\widehat{\mathbb{E}}[X] \geq \widehat{\mathbb{E}}[Y]$ ;
- (b) Constant preserving:  $\widehat{\mathbb{E}}[c] = c$ ;
- (c) Sub-additivity:  $\widehat{\mathbb{E}}[X + Y] \leq \widehat{\mathbb{E}}[X] + \widehat{\mathbb{E}}[Y]$  whenever  $\widehat{\mathbb{E}}[X] + \widehat{\mathbb{E}}[Y]$  is not of the form  $+\infty - \infty$  or  $-\infty + \infty$ ;
- (d) Positive homogeneity:  $\widehat{\mathbb{E}}[\lambda X] = \lambda \widehat{\mathbb{E}}[X], \lambda > 0$ .

Here  $\bar{\mathbb{R}} = [-\infty, +\infty]$ . The triple  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  is called a sub-linear expectation space. Given a sub-linear expectation  $\widehat{\mathbb{E}}$ , let us denote the conjugate expectation  $\widehat{\mathcal{E}}$  of  $\widehat{\mathbb{E}}$  by

$$\widehat{\mathcal{E}}[X] := -\widehat{\mathbb{E}}[-X], \quad \forall X \in \mathcal{H}.$$

From the definition, we can easily get that  $\widehat{\mathcal{E}}[X] \leq \widehat{\mathbb{E}}[X], \widehat{\mathbb{E}}[X + c] = \widehat{\mathbb{E}}[X] + c, \widehat{\mathbb{E}}[X - Y] \geq \widehat{\mathbb{E}}[X] - \widehat{\mathbb{E}}[Y]$  and  $|\widehat{\mathbb{E}}[X] - \widehat{\mathbb{E}}[Y]| \leq \widehat{\mathbb{E}}[|X - Y|]$ . Further, if  $\widehat{\mathbb{E}}[|X|]$  is finite, then  $\widehat{\mathcal{E}}[X]$  and  $\widehat{\mathbb{E}}[X]$  are both finite.

**Definition 1.2** [16]

(i)(Independence) In a sub-linear expectation space  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ , a random vector  $\mathbf{Y} = (Y_1, Y_2, \dots, Y_n), Y_i \in \mathcal{H}$  is said to be independent to another random vector  $\mathbf{X} = (X_1, X_2, \dots, X_n), X_i \in \mathcal{H}$  under  $\widehat{\mathbb{E}}$  if for each test function  $\varphi \in C_{l.Lip}(\mathbb{R}^m \times \mathbb{R}^n)$  we have  $\widehat{\mathbb{E}}[\varphi(\mathbf{X}, \mathbf{Y})] = \widehat{\mathbb{E}}[\widehat{\mathbb{E}}[\varphi(\mathbf{x}, \mathbf{Y})] |_{\mathbf{x}=\mathbf{X}}]$ , whenever  $\bar{\varphi}(\mathbf{x}) := \widehat{\mathbb{E}}[|\varphi(\mathbf{x}, \mathbf{Y})|] < \infty$  for all  $\mathbf{x}$  and  $\widehat{\mathbb{E}}[|\bar{\varphi}(\mathbf{X})|] < \infty$ .

**Definition 1.3** [24]

(i) (Negative dependence) In a sub-linear expectation space  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ , a random vector  $\mathbf{Y} = (Y_1, Y_2, \dots, Y_n), Y_i \in \mathcal{H}$  is said to be negatively dependent (ND) to another random vector  $\mathbf{X} = (X_1, X_2, \dots, X_n), X_i \in \mathcal{H}$  under  $\widehat{\mathbb{E}}$  if for each pair of test functions  $\varphi_1 \in C_{l.Lip}(\mathbb{R}^m)$  and  $\varphi_2 \in C_{l.Lip}(\mathbb{R}^n)$  we have  $\widehat{\mathbb{E}}[\varphi_1(\mathbf{X})\varphi_2(\mathbf{Y})] \leq \widehat{\mathbb{E}}[\varphi_1(\mathbf{X})]\widehat{\mathbb{E}}[\varphi_2(\mathbf{Y})]$ , whenever either  $\varphi_1, \varphi_2$  are coordinatewise nondecreasing or  $\varphi_1, \varphi_2$  are coordinatewise non-increasing with  $\varphi_1(\mathbf{X}) \geq 0, \widehat{\mathbb{E}}[\varphi_2(\mathbf{Y})] \geq 0, \widehat{\mathbb{E}}[|\varphi_1(\mathbf{X})\varphi_2(\mathbf{Y})|] < \infty, \widehat{\mathbb{E}}[|\varphi_1(\mathbf{X})|] < \infty, \widehat{\mathbb{E}}[|\varphi_2(\mathbf{Y})|] < \infty$ .

(ii) (ND random variables) Let  $\{X_n; n \geq 1\}$  be a sequence of random variables in the sub-linear expectation space  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ .  $X_1, X_2, \dots$  are said to be negatively dependent if  $X_{i+1}$  is negatively dependent to  $(X_1, X_2, \dots, X_i)$  for each  $i \geq 1$ .

It is obvious that, if  $\{X_n; n \geq 1\}$  is a sequence of independent random variables and  $f_1(x), f_2(x), \dots \in C_{l.Lip}(\mathbb{R})$  then  $\{f_n(X_n); n \geq 1\}$  is also a sequence of independent random variables; if  $\{X_n; n \geq 1\}$  is a sequence of negatively dependent random variables and  $f_1(x), f_2(x), \dots$

$\cdot \in C_{l.Lip}(\mathbb{R})$  are non-decreasing (resp. non-increasing) functions, then  $\{f_n(X_n); n \geq 1\}$  is also a sequence of negatively dependent random variables.

**Definition 1.4** An array of random variables  $\{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  is called rowwise negatively dependent random variables if for every given  $n \geq 1$ ,  $\{X_{ni}; 1 \leq i \leq k_n\}$  is a sequence of negatively dependent random variables in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$ , where and in the sequel  $\{k_n, n \geq 1\}$  is a sequence of positive integers and  $k_n \rightarrow \infty$ , when  $n \rightarrow \infty$ .

Next, we introduce the capacities corresponding to the sub-linear expectations. Let  $\mathcal{G} \subset \mathcal{F}$ . A function  $V : \mathcal{G} \rightarrow [0, 1]$  is called a capacity if

$$V(\phi) = 0, V(\Omega) = 1, \text{ and } V(A) \leq V(B) \quad \forall A \subset B, A, B \in \mathcal{G}.$$

It is called to be sub-additive if  $V(A \cup B) \leq V(A) + V(B)$  for all  $A, B \in \mathcal{G}$  with  $A \cup B \in \mathcal{G}$ .

Let  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  be a sub-linear space, and  $\widehat{\mathcal{E}}$  be the conjugate expectation of  $\widehat{\mathbb{E}}$ . We denote a pair  $(\mathbb{V}, \mathcal{V})$  of capacities by

$$\mathbb{V}(A) := \inf\{\widehat{\mathbb{E}}[\xi] : I_A \leq \xi, \xi \in \mathcal{H}\}, \mathcal{V}(A) := 1 - \mathbb{V}(A^c), \quad \forall A \in \mathcal{F},$$

where  $A^c$  is the complement set of  $A$ . It is obvious that  $\mathbb{V}$  is sub-additive and

$$\begin{aligned} \mathbb{V}(A) &:= \widehat{\mathbb{E}}[I_A], \mathcal{V}(A) := \widehat{\mathcal{E}}[I_A], \text{ if } I_A \in \mathcal{H}, \\ \widehat{\mathbb{E}}[f] \leq \mathbb{V}(A) \leq \widehat{\mathbb{E}}[g], \widehat{\mathcal{E}}[f] \leq \mathcal{V}(A) \leq \widehat{\mathcal{E}}[g], \text{ if } f \leq I_A \leq g, f, g \in \mathcal{H}. \end{aligned}$$

This implies Markov inequality:  $\forall X \in \mathcal{H}$ ,

$$\mathbb{V}(|X| \geq x) \leq \widehat{\mathbb{E}}[|X|^p]/x^p, \quad \forall x > 0, p > 0$$

from  $I\{|X| \geq x\} \leq |X|^p/x^p \in \mathcal{H}$ . By Lemma 4.1 in Zhang (2016a), we have Hölder inequality inequality:  $\forall X, Y \in \mathcal{H}, p, q > 1$ , satisfying  $p^{-1} + q^{-1} = 1$ ,

$$\widehat{\mathbb{E}}[|XY|] \leq (\widehat{\mathbb{E}}[|X|^p])^{\frac{1}{p}} (\widehat{\mathbb{E}}[|Y|^q])^{\frac{1}{q}},$$

particularly, Jensen inequality:

$$(\widehat{\mathbb{E}}[|X|^r])^{\frac{1}{r}} \leq (\widehat{\mathbb{E}}[|X|^s])^{\frac{1}{s}}, \quad \text{for } 0 < r \leq s.$$

We define the Choquet integrals/expectations  $(C_{\mathbb{V}}, C_{\mathcal{V}})$  by

$$C_V(X) := \int_0^\infty V(X \geq x)dx + \int_{-\infty}^0 (V(X \geq x) - 1)dx$$

with  $V$  being replaced by  $\mathbb{V}$  and  $\mathcal{V}$ , respectively. If  $\lim_{c \rightarrow \infty} \widehat{\mathbb{E}}[(|X| - c)_+] = 0$ , then  $\widehat{\mathbb{E}}[|X|] \leq C_{\mathbb{V}}(|X|)$  [24].

Complete convergence and complete moment convergence theorems are important limit theorems in probability theory. Many of related results of complete convergence have been obtained in the probability space. We refer the reader to [1,13,11,8,21], and so on. For the results of complete moment convergence in the probability space, we refer the reader to [6,2,20,19,23], and so on.

Wu et al. [23] established the following complete moment convergence theorem for END random variables in classical probability space.

**Theorem A** Let  $q \geq 1$ ,  $\{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise END random variables and  $\{c_n\}$  be a sequence of positive real numbers. Assume that the following conditions hold:

(a)  $\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} E|X_{ni}|I(|X_{ni}| > \varepsilon) < \infty$  for any  $\varepsilon > 0$ ;

(b) for some  $\delta > 0$ , there exists  $\eta > 1$  such that

$$\sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} EX_{ni}^2 I(|X_{ni}| \leq \delta) \right)^\eta < \infty;$$

(c)  $\sum_{i=1}^{k_n} E|X_{ni}|I(|X_{ni}| > \frac{\delta}{16\eta}) \rightarrow 0$ , as  $n \rightarrow \infty$ .

Then for all  $\varepsilon > 0$

$$\sum_{n=1}^{\infty} c_n E \left\{ \left| \sum_{i=1}^{k_n} (X_{ni} - EX_{ni}I(|X_{ni}| \leq \delta)) \right| - \varepsilon \right\}_+ < \infty.$$

The main purpose of this article is to investigate more general complete moment convergence for arrays of rowwise ND random variables under the sub-linear expectations.

### §2 Main results

In this paper, we define  $g(x)$  in the following: For  $0 < \mu < 1$ , let  $g(x) \in C_{l.Lip}(\mathbb{R})$  such that  $0 \leq g(x) \leq 1$  for all  $x$  and  $g(x) = 1$  if  $|x| \leq \mu$ ,  $g(x) = 0$  if  $|x| > 1$  and when  $x > 0$ ,  $g(x) \downarrow$ . Then

$$I(|x| \leq \mu) \leq g(x) \leq I(|x| \leq 1), \quad I(|x| > 1) \leq 1 - g(x) \leq I(|x| > \mu). \tag{2.1}$$

**Theorem 2.1.** Let  $\{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise ND random variables in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  and  $\{c_n\}$  be a sequence of positive real numbers,  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be an increasing function with  $f(0) = 0$  and  $\eta \geq 1$  be a constant. There exists some constant  $\delta > 0$  and  $\mu$  is the same as in (2.1). Assume that the following conditions hold:

(a)  $\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \widehat{\mathbb{E}} [f(8\eta|X_{ni}|(1 - g(\frac{X_{ni}}{\varepsilon})))]$   
 $\leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_V [f(8\eta|X_{ni}|(1 - g(\frac{X_{ni}}{\varepsilon})))] < \infty$ , for any  $\varepsilon > 0$ ;

(b)  $\sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} [X_{ni}^2 g(\frac{X_{ni}}{\delta})] \right)^\eta < \infty$ ;

(c)  $\sum_{i=1}^{k_n} \widehat{\mathbb{E}} [ |X_{ni}| \left( 1 - g\left(\frac{16\eta X_{ni}}{\mu\delta}\right) \right) ] \rightarrow 0$ , as  $n \rightarrow \infty$ ;

(d) let  $h : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  be the inverse function for  $f(t)$ , which is  $h(f(t)) = t, t \geq 0$  and  $s(t) = \max_{\mu\delta \leq x \leq h(t)/\mu} \frac{x}{f(x)}$ . Suppose that the constants  $\eta, \delta$  and the function  $f : \mathbb{R}^+ \rightarrow \mathbb{R}^+$  satisfy the condition

$$\int_{f(\delta)}^{\infty} h^{-\eta}(t)s(t) dt < \infty.$$

Then for all  $\varepsilon > 0$

$$\sum_{n=1}^{\infty} c_n C_V \left[ f \left( \left\{ \sum_{i=1}^{k_n} \left( X_{ni} - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) \right\} - \varepsilon \right) \right] < \infty. \tag{2.2}$$

Let  $f(t) = t^q, t \geq 0, q > 0$  in Theorem 2.1, we can obtain the following interesting result:

**Corollary 2.1.** Let  $q > 0, \{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise ND random variables in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  and  $\{c_n\}$  be a sequence of positive real numbers. Assume that the following conditions hold:

(a)  $\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \widehat{\mathbb{E}} [ |X_{ni}|^q (1 - g(\frac{X_{ni}}{\varepsilon})) ]$   
 $\leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_V [ |X_{ni}|^q (1 - g(\frac{X_{ni}}{\varepsilon})) ] < \infty$  for any  $\varepsilon > 0$ ;

(b) for some  $\delta > 0$ , there exists  $\eta > \max(1, q)$  such that

$$\sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ X_{ni}^2 g \left( \frac{X_{ni}}{\delta} \right) \right] \right)^\eta < \infty;$$

(c)  $\sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g \left( \frac{16\eta X_{ni}}{\mu\delta} \right) \right) \right] \rightarrow 0$ , as  $n \rightarrow \infty$ .

Then for all  $\varepsilon > 0$

$$\sum_{n=1}^{\infty} c_n C_V \left[ \left\{ \sum_{i=1}^{k_n} \left( X_{ni} - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) - \varepsilon \right\}_+^q \right] < \infty.$$

**Remark 2.1** We can obtain the corresponding result of Theorem A when  $f(t) = t$  ( $t \geq 0$ ) in our Theorem 2.1 under the sub-linear expectations. We establish a more general complete moment convergence theorem for arrays of rowwise ND random variables under the sub-linear expectations.

**Remark 2.2** Letting  $f(t)$  be some special functions, we can obtain a series of results on complete moment convergence for ND random variables.

### §3 Proofs of main results

In order to prove our results, we need the following lemmas.

**Lemma 3.1.** Let  $\{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise negatively dependent random variables in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  with  $\widehat{\mathbb{E}}[X_{ni}] \leq 0$ , for  $1 \leq i \leq k_n, n \geq 1$ . Let  $B'_n = \sum_{i=1}^{k_n} \widehat{\mathbb{E}}[X_{ni}^2]$ . Then for any given  $n$  and for all  $x > 0, y > 0$

$$\mathbb{V} \left( \sum_{i=1}^{k_n} X_{ni} > x \right) \leq \left( \max_{1 \leq i \leq k_n} |X_{ni}| > y \right) + \exp \left\{ \frac{x}{y} - \frac{x}{y} \ln \left( 1 + \frac{xy}{B'_n} \right) \right\}.$$

**Proof** Similar to the proof of Theorem 3.1 of [24], we can have

$$\mathbb{V} \left( \sum_{i=1}^{k_n} X_{ni} > x \right) \leq \left( \max_{1 \leq i \leq k_n} |X_{ni}| > y \right) + \exp \left\{ \frac{x}{y} - \frac{x}{y} \left( \frac{B'_n}{xy} + 1 \right) \ln \left( 1 + \frac{xy}{B'_n} \right) \right\}.$$

Hence we have

$$\mathbb{V} \left( \sum_{i=1}^{k_n} X_{ni} > x \right) \leq \left( \max_{1 \leq i \leq k_n} |X_{ni}| > y \right) + \exp \left\{ \frac{x}{y} - \frac{x}{y} \ln \left( 1 + \frac{xy}{B'_n} \right) \right\}.$$

**Lemma 3.2.** Let  $\{X_{ni}; 1 \leq i \leq k_n, n \geq 1\}$  be an array of rowwise ND random variables in  $(\Omega, \mathcal{H}, \widehat{\mathbb{E}})$  and  $\{c_n\}$  be a sequence of positive real numbers. Suppose that for every  $\varepsilon > 0$  and some  $\delta > 0$ :

(i)  $\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \varepsilon) < \infty$ ;

(ii) there exists  $j \geq 1$  such that

$$\sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right)^2 \right] \right)^j < \infty.$$

Then

$$\sum_{n=1}^{\infty} c_n \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) < \infty \text{ for all } \varepsilon > 0.$$

**Proof** Denote  $S'_n = \sum_{i=1}^{k_n} \left( X_{ni} - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right)$ . Note that

$$\begin{aligned} \mathbb{V}(S'_n > \varepsilon) &\leq \mathbb{V} \left( S'_n > \varepsilon, \bigcup_{i=1}^{k_n} (|X_{ni}| > \mu\delta) \right) + \mathbb{V} \left( S'_n > \varepsilon, \bigcap_{i=1}^{k_n} (|X_{ni}| \leq \mu\delta) \right) \\ &\leq \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu\delta) + \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) \end{aligned}$$

Hence, by condition (i), it is sufficient to prove that

$$\sum_{n=1}^{\infty} c_n \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) < \infty.$$

For any  $a > 0$  and set

$$d = \min \left\{ 1, \frac{a}{6\delta} \right\}, N_1 = \left\{ n : \sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \min \left\{ \mu\delta, \mu \frac{a}{6} \right\} \right) > d \right\}, \text{ and } N_2 = \mathbb{N}/N_1.$$

Note that

$$\begin{aligned} \sum_{n \in N_1} c_n \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) \\ \leq \sum_{n \in N_1} c_n \leq \frac{1}{d} \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \min \left\{ \mu\delta, \mu \frac{a}{6} \right\} \right) < \infty. \end{aligned}$$

Hence, it remains to prove that  $\sum_{n \in N_2} c_n \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) < \infty$ .

By Lemma 3.1 we get

$$\begin{aligned} \sum_{n \in N_2} c_n \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > \varepsilon \right) \\ \leq \sum_{n \in N_2} c_n \mathbb{V} \left( \max_{1 \leq i \leq k_n} \left| X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right| > a \right) \\ + \sum_{n \in N_2} c_n \exp \left\{ \frac{\varepsilon}{a} - \frac{\varepsilon}{a} \ln \left( 1 + \frac{\varepsilon a}{B_n} \right) \right\}, \end{aligned}$$

where  $B_n = \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right)^2 \right]$ .

If  $a/6 < \delta$ , for  $n \in N_2$ , by the fact  $\left| g \left( \frac{X_{ni}}{\delta} \right) - g \left( \frac{X_{ni}}{a/6} \right) \right| \leq I \left( \mu \frac{a}{6} < |X_{ni}| \leq \delta \right)$ , we have,

$$\begin{aligned} \max_{1 \leq i \leq k_n} \left| \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right| \\ \leq \max_{1 \leq i \leq k_n} \left| \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{a/6} \right) \right] \right| + \max_{1 \leq i \leq k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g \left( \frac{X_{ni}}{\delta} \right) - g \left( \frac{X_{ni}}{a/6} \right) \right| \right] \\ \leq a/6 + \delta \sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \mu \frac{a}{6} \right) \\ \leq a/6 + \delta \sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \min \left\{ \delta, \mu \frac{a}{6} \right\} \right) \leq a/6 + \delta d \leq a/3. \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{n \in N_2} c_n \mathbb{V} \left( \max_{1 \leq i \leq k_n} \left| X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right| > a \right) \\ & \leq \sum_{n=1}^{\infty} c_n \mathbb{V} \left( \max_{1 \leq i \leq k_n} \left| X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right| > \frac{2}{3} a \right) \leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \frac{2}{3} a) < \infty. \end{aligned}$$

If  $a/6 \geq \delta$ , we can also easily get  $\sum_{n \in N_2} c_n \mathbb{V} \left( \max_{1 \leq i \leq k_n} |X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} [X_{ni} g \left( \frac{X_{ni}}{\delta} \right)]| > a \right) < \infty$ .

Letting  $a = \frac{\varepsilon}{j}$ , we have

$$\begin{aligned} & \sum_{n \in N_2} c_n \exp \left\{ \frac{\varepsilon}{a} - \frac{\varepsilon}{a} \ln \left( 1 + \frac{\varepsilon a}{B_n} \right) \right\} \leq \exp \left\{ \frac{\varepsilon}{a} \right\} \sum_{n \in N_2} c_n \left( 1 + \frac{\varepsilon a}{B_n} \right)^{-\frac{\varepsilon}{a}} \\ & \leq \exp \left\{ \frac{\varepsilon}{2a} \right\} \sum_{n \in N_2} c_n \left( \frac{B_n}{\varepsilon a} \right)^j \leq C \sum_{n=1}^{\infty} c_n B_n^j \\ & = C \sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ \left( X_{ni} g \left( \frac{X_{ni}}{\delta} \right) - \widehat{\mathbb{E}} [X_{ni} g \left( \frac{X_{ni}}{\delta} \right)] \right)^2 \right] \right)^j < \infty. \end{aligned}$$

We complete the proof of Lemma 3.2.

*Proof. Proof of Theorem 2.1.* First of all, from condition (a), we have

$$\begin{aligned} (a') \quad & \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ f \left( |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\varepsilon} \right) \right) \right) \right] \\ & \leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_{\mathbb{V}} \left[ f \left( |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\varepsilon} \right) \right) \right) \right] < \infty \text{ for any } \varepsilon > 0. \end{aligned}$$

Next, by Markov's inequality and condition (a'), we have that

$$\sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \varepsilon) \leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ f \left( |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\varepsilon} \right) \right) \right) \right] < \infty,$$

which implies that condition (i) of Lemma 3.2 holds.

For  $n \geq 1$ , denote  $S_n = \sum_{i=1}^{k_n} \left( X_{ni} - \widehat{\mathbb{E}} [X_{ni} g \left( \frac{X_{ni}}{\delta} \right)] \right)$ .

We can easily get that

$$\begin{aligned} & \sum_{n=1}^{\infty} c_n C_{\mathbb{V}} [f(\{S_n - \varepsilon\}_+)] = \sum_{n=1}^{\infty} c_n \int_0^{\infty} \mathbb{V}(S_n - \varepsilon > h(t)) dt \\ & = \sum_{n=1}^{\infty} c_n \int_0^{f(\delta)} \mathbb{V}(S_n > \varepsilon + h(t)) dt + \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V}(S_n > \varepsilon + h(t)) dt \\ & \leq f(\delta) \sum_{n=1}^{\infty} c_n \mathbb{V}(S_n > \varepsilon) + \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V}(S_n > h(t)) dt \\ & \doteq I_1 + I_2. \end{aligned} \tag{3.1}$$

By Lemma 3.2, we have that  $I_1 < \infty$ . It is enough to show that  $I_2 < \infty$ .

For  $t \geq f(\delta)$ , it is easy to get that

$$\begin{aligned} \mathbb{V}(S_n > h(t)) &\leq \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu h(t)) \\ &\quad + \mathbb{V}\left(\sum_{i=1}^{k_n} \left(X_{ni}g\left(\frac{X_{ni}}{h(t)}\right) - \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{X_{ni}}{\delta}\right)\right]\right) > h(t)\right), \end{aligned}$$

which implies that

$$\begin{aligned} I_2 &\leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \int_{f(\delta)}^{\infty} \mathbb{V}(|X_{ni}| > \mu h(t)) dt \\ &\quad + \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V}\left(\sum_{i=1}^{k_n} \left(X_{ni}g\left(\frac{X_{ni}}{h(t)}\right) - \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{X_{ni}}{\delta}\right)\right]\right) > h(t)\right) dt \tag{3.2} \\ &\doteq I_3 + I_4. \end{aligned}$$

By Markov's inequality and condition (a'), we have that

$$\begin{aligned} I_3 &\leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \int_{f(\delta)}^{\infty} \mathbb{V}(f(|X_{ni}/\mu|) > t) dt \\ &\leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_{\mathbb{V}} \left[ f\left(|X_{ni}| \left(1 - g\left(\frac{X_{ni}}{\delta}\right)\right)\right) \right] < \infty. \end{aligned} \tag{3.3}$$

For fixed  $n \geq 1, 1 \leq i \leq k_n$  and  $t \geq f(\delta)$ , denote

$$\begin{aligned} Y_{ni} &= h(t)I(X_{ni} > h(t)) + X_{ni}I(|X_{ni}| \leq h(t)) - h(t)I(X_{ni} < -h(t)) \quad \text{and} \\ Y'_{ni} &= h(t)I(X_{ni} > h(t)) - h(t)I(X_{ni} < -h(t)). \end{aligned}$$

By the definition of  $Y_{ni}$ ,  $\{Y_{ni} - EY_{ni}, 1 \leq i \leq k_n, n \geq 1\}$  is an array of rowwise ND random variables under the sub-linear expectations.

By the definition of  $Y_{ni}$ ,  $C_r$  inequality and (2.1), we have

$$\widehat{\mathbb{E}}[Y_{ni}] \leq \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{\mu X_{ni}}{h(t)}\right)\right] + h(t)\mathbb{V}(|X_{ni}| > \mu h(t))$$

Hence, for  $t \geq f(\delta)$ , we have

$$\begin{aligned} &\mathbb{V}\left(\sum_{i=1}^{k_n} \left(X_{ni}g\left(\frac{X_{ni}}{h(t)}\right) - \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{X_{ni}}{\delta}\right)\right]\right) > h(t)\right) \\ &\leq \mathbb{V}\left(\sum_{i=1}^{k_n} \left(Y_{ni} - Y'_{ni} - \widehat{\mathbb{E}}[Y_{ni}] + \widehat{\mathbb{E}}[Y_{ni}] - \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{X_{ni}}{\delta}\right)\right]\right) > h(t)\right) \tag{3.4} \\ &\leq \mathbb{V}\left(\sum_{i=1}^{k_n} \left((Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}]) + |Y'_{ni}| + \left|\widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{\mu X_{ni}}{h(t)}\right)\right] - \widehat{\mathbb{E}}\left[X_{ni}g\left(\frac{X_{ni}}{\delta}\right)\right]\right) > h(t)\right) \end{aligned}$$

By  $g(x) \downarrow, x > 0$ , condition (c) and the fact  $\left|g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right)\right| \leq I(\mu\delta < |X_{ni}| \leq h(t)/\mu)$ ,



we have

$$\begin{aligned} & \max_{t \geq f(\delta)} \frac{1}{h(t)} \sum_{i=1}^{k_n} \left| \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{\mu X_{ni}}{h(t)} \right) \right] - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right| \\ & \leq \max_{t \geq f(\delta)} \frac{1}{h(t)} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g \left( \frac{\mu X_{ni}}{h(t)} \right) - g \left( \frac{X_{ni}}{\delta} \right) \right| \right] \\ & \leq \delta^{-1} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\mu \delta} \right) \right) \right] \\ & \leq \delta^{-1} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g \left( \frac{16\eta X_{ni}}{\mu \delta} \right) \right) \right] \rightarrow 0, \end{aligned}$$

as  $n \rightarrow \infty$ .

Hence, for all  $n$  large enough,

$$\sum_{i=1}^{k_n} \left| \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{\mu X_{ni}}{h(t)} \right) \right] - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right| < \frac{h(t)}{4}, \quad t \geq f(\delta). \tag{3.5}$$

By Markov's inequality and condition (c), we have

$$\begin{aligned} & \max_{t \geq f(\delta)} \frac{1}{h(t)} \sum_{i=1}^{k_n} h(t) \mathbb{V}(|X_{ni}| > \mu h(t)) \\ & = \max_{t \geq f(\delta)} \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu h(t)) \leq \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu \delta) \\ & \leq (\mu \delta)^{-1} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\mu \delta} \right) \right) \right] \rightarrow 0, \quad \text{as } n \rightarrow \infty. \end{aligned} \tag{3.6}$$

Hence, for all  $n$  large enough,

$$\sum_{i=1}^{k_n} h(t) \mathbb{V}(|X_{ni}| > \mu h(t)) < \frac{h(t)}{4}, \quad t \geq f(\delta). \tag{3.7}$$

Combining (3.4), (3.5) and (3.7), for  $n$  large enough, we have

$$\begin{aligned} & \mathbb{V} \left( \sum_{i=1}^{k_n} \left( X_{ni} g \left( \frac{X_{ni}}{h(t)} \right) - \widehat{\mathbb{E}} \left[ X_{ni} g \left( \frac{X_{ni}}{\delta} \right) \right] \right) > h(t) \right) \\ & \leq \mathbb{V} \left( \sum_{i=1}^{k_n} (Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}]) > \frac{h(t)}{4} \right) + \mathbb{V} \left( \sum_{i=1}^{k_n} |Y'_{ni}| > \frac{h(t)}{4} \right) \end{aligned}$$

Therefore

$$\begin{aligned} I_4 & \leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \sum_{i=1}^{k_n} (Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}]) > \frac{h(t)}{4} \right) dt \\ & \quad + C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \sum_{i=1}^{k_n} |Y'_{ni}| > \frac{h(t)}{4} \right) dt \triangleq I_5 + I_6. \end{aligned} \tag{3.8}$$

By the fact  $|Y'_{ni}| \leq h(t)I(|X_{ni}| > h(t)) \leq h(t) \left( 1 - g \left( \frac{X_{ni}}{h(t)} \right) \right)$ , Markov's inequality, (2.1) and

condition (a'), we can obtain

$$\begin{aligned}
 I_6 &= C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \sum_{i=1}^{k_n} |Y'_{ni}| > \frac{h(t)}{4} \right) dt \\
 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \sum_{i=1}^{k_n} h(t) \left( 1 - g \left( \frac{X_{ni}}{h(t)} \right) \right) > \frac{h(t)}{4} \right) dt \\
 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \widehat{\mathbb{E}} \left[ \sum_{i=1}^{k_n} \left( 1 - g \left( \frac{X_{ni}}{h(t)} \right) \right) \right] dt \tag{3.9} \\
 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu h(t)) dt \\
 &\leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_{\mathbb{V}} \left[ f \left( |X_{ni}| \left( 1 - g \left( \frac{X_{ni}}{\mu \delta} \right) \right) \right) \right] < \infty.
 \end{aligned}$$

Hence to prove  $I_4 < \infty$ , it is enough to show that  $I_5 < \infty$ . Denote  $B_n = \sum_{i=1}^{k_n} \widehat{\mathbb{E}}(Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}])^2$ . Applying Lemma 3.1 with  $x = h(t)/4$  and  $y = h(t)/(4\eta)$ , we have

$$\begin{aligned}
 I_5 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \max_{1 \leq i \leq k_n} |Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}]| > \frac{h(t)}{4\eta} \right) dt \\
 &\quad + C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \exp \left\{ \eta - \eta \ln \left( 1 + \frac{h^2(t)}{16\eta B_n} \right) \right\} dt \triangleq I_7 + I_8. \tag{3.10}
 \end{aligned}$$

By Markov's inequality and condition (c), we can get that for all  $n$  large enough,

$$\sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \frac{\mu \delta}{16\eta} \right) \leq C \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g \left( \frac{16\eta X_{ni}}{\mu \delta} \right) \right) \right] < \frac{\mu}{(1 + \mu)16\eta}. \tag{3.11}$$

Thus by the fact  $\left| g \left( \frac{\mu X_{ni}}{h(t)} \right) - g \left( \frac{16\eta X_{ni}}{\delta} \right) \right| \leq I \left( \frac{\mu \delta}{16\eta} < |X_{ni}| \leq h(t)/\mu \right)$  and (3.11), we have

$$\begin{aligned}
 &\max_{t \geq f(\delta)} \max_{1 \leq i \leq k_n} \frac{1}{h(t)} |\widehat{\mathbb{E}}[Y_{ni}]| \leq \max_{t \geq f(\delta)} \max_{1 \leq i \leq k_n} \frac{1}{h(t)} \widehat{\mathbb{E}}[|Y_{ni}|] \\
 &\leq \max_{t \geq f(\delta)} \max_{1 \leq i \leq k_n} \frac{1}{h(t)} \left( \widehat{\mathbb{E}} \left[ |X_{ni}| g \left( \frac{\mu X_{ni}}{h(t)} \right) \right] + h(t) \mathbb{V}(|X_{ni}| > \mu h(t)) \right) \\
 &\leq \max_{t \geq f(\delta)} \max_{1 \leq i \leq k_n} \frac{1}{h(t)} \left( \widehat{\mathbb{E}} \left[ |X_{ni}| g \left( \frac{16\eta X_{ni}}{\delta} \right) \right] + \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g \left( \frac{\mu X_{ni}}{h(t)} \right) - g \left( \frac{16\eta X_{ni}}{\delta} \right) \right| \right] \right) \\
 &\quad + \max_{t \geq f(\delta)} \max_{1 \leq i \leq k_n} \frac{1}{h(t)} (h(t) \mathbb{V}(|X_{ni}| > \mu h(t))) \\
 &\leq \frac{1}{\delta} \frac{\delta}{16\eta} + \max_{t \geq f(\delta)} \frac{1}{h(t)} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( g \left| \frac{\mu X_{ni}}{h(t)} \right| - g \left( \frac{16\eta X_{ni}}{\delta} \right) \right) \right] + \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu \delta) \\
 &\leq \frac{1}{16\eta} + \max_{t \geq f(\delta)} \frac{1}{h(t)} \cdot \frac{h(t)}{\mu} \cdot \sum_{i=1}^{k_n} \mathbb{V} \left( |X_{ni}| > \frac{\mu \delta}{16\eta} \right) + \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu \delta) \\
 &\leq \frac{1}{16\eta} + (1/\mu + 1) \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \frac{\mu \delta}{16\eta})
 \end{aligned}$$

$$< \frac{1}{16\eta} + \frac{1}{16\eta} = \frac{1}{8\eta}.$$

Therefore by the fact  $|Y_{ni}| \leq |X_{ni}|$  and condition (a), we have

$$\begin{aligned} I_7 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \mathbb{V} \left( \max_{1 \leq i \leq k_n} |Y_{ni}| > \frac{h(t)}{8\eta} \right) dt \leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \int_{f(\delta)}^{\infty} \mathbb{V} \left( |X_{ni}| > \frac{h(t)}{8\eta} \right) dt \\ &\leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_{\mathbb{V}} \left[ f \left( 8\eta |X_{ni}| \left( 1 - g \left( \frac{8\eta X_{ni}}{\delta} \right) \right) \right) \right] < \infty. \end{aligned} \tag{3.12}$$

By  $C_r$ -inequality, we have

$$\begin{aligned} I_8 &= C \sum_{n=1}^{\infty} c_n e^{\eta} \int_{f(\delta)}^{\infty} \left( 1 + \frac{h^2(t)}{16\eta B_n} \right)^{-\eta} dt \leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) B_n^{\eta} dt \\ &= C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}}(Y_{ni} - \widehat{\mathbb{E}}[Y_{ni}])^2 \right)^{\eta} dt \\ &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}}[Y_{ni}^2] \right)^{\eta} dt. \end{aligned}$$

By the definition of  $Y_{ni}$ ,  $C_r$  inequality and (2.1), we have

$$\begin{aligned} \widehat{\mathbb{E}}[Y_{ni}^2] &\leq C \widehat{\mathbb{E}} \left[ X_{ni}^2 g \left( \frac{\mu X_{ni}}{h(t)} \right) \right] + Ch^2(t) \mathbb{V}(|X_{ni}| > \mu h(t)) \\ &\leq C \widehat{\mathbb{E}} \left[ X_{ni}^2 g \left( \frac{X_{ni}}{\delta} \right) \right] + C \widehat{\mathbb{E}} \left[ X_{ni}^2 \left| g \left( \frac{\mu X_{ni}}{h(t)} \right) - g \left( \frac{X_{ni}}{\delta} \right) \right| \right] + Ch^2(t) \mathbb{V}(|X_{ni}| > \mu h(t)). \end{aligned}$$

Hence

$$\begin{aligned} I_8 &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}}[Y_{ni}^2] \right)^{\eta} dt \\ &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ X_{ni}^2 g \left( \frac{X_{ni}}{\delta} \right) \right] \right)^{\eta} dt \\ &\quad + C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-2\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ X_{ni}^2 \left| g \left( \frac{\mu X_{ni}}{h(t)} \right) - g \left( \frac{X_{ni}}{\delta} \right) \right| \right] \right)^{\eta} dt \\ &\quad + C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} \left( \sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu h(t)) \right)^{\eta} dt \triangleq I_9 + I_{10} + I_{11}. \end{aligned} \tag{3.13}$$

By the definition of  $s(t) = \max_{\mu\delta \leq x \leq h(t)/\mu} \frac{x}{f(x)}$ , we have  $s(t) \geq \frac{\mu\delta}{f(\mu\delta)}$ . Then  $\delta \leq \frac{f(\mu\delta)}{\mu} s(t)$ . For  $t > f(\delta)$ , we have

$$\begin{aligned} h^{-2\eta}(t) &\leq h^{-\eta}(t) \delta^{-\eta} \leq h^{-\eta}(t) \delta^{-\eta-1} \frac{f(\mu\delta)}{\mu} s(t) \\ &= \delta^{-\eta-1} \frac{f(\mu\delta)}{\mu} h^{-\eta}(t) s(t) = Ch^{-\eta}(t) s(t). \end{aligned}$$

Hence by conditions (b) and (d), we have

$$I_9 = C \sum_{n=1}^{\infty} c_n \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ X_{ni}^2 g \left( \frac{X_{ni}}{\delta} \right) \right] \right)^{\eta} \int_{f(\delta)}^{\infty} h^{-\eta}(t) s(t) dt < \infty. \tag{3.14}$$

Note that

$$\left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \leq I(\mu\delta < |X_{ni}| \leq h(t)/\mu), \quad t \geq f(\delta)$$

then we have

$$I_{10} \leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-\eta}(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right] \right)^{\eta} dt.$$

By condition (c), for all  $n$  large enough, we have

$$\begin{aligned} \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right] &\leq \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g\left(\frac{X_{ni}}{\mu\delta}\right) \right) \right] \\ &\leq \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left( 1 - g\left(\frac{16\eta X_{ni}}{\mu\delta}\right) \right) \right] < 1. \end{aligned}$$

Hence by the fact  $\eta \geq 1$ , conditions (a') and (d), we get

$$\begin{aligned} I_{10} &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-\eta}(t) \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ |X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right] dt \\ &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-\eta}(t) \\ &\quad \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ \frac{|X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right|}{f\left(|X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right)} f\left(|X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right) \right] \right) dt \\ &\leq C \sum_{n=1}^{\infty} c_n \int_{f(\delta)}^{\infty} h^{-\eta}(t) s(t) \left( \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ f\left(|X_{ni}| \left| g\left(\frac{\mu X_{ni}}{h(t)}\right) - g\left(\frac{X_{ni}}{\delta}\right) \right| \right) \right] \right) dt \\ &\leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \widehat{\mathbb{E}} \left[ f\left(|X_{ni}| \left( 1 - g\left(\frac{X_{ni}}{\mu\delta}\right) \right) \right) \right] \int_{f(\delta)}^{\infty} h^{-\eta}(t) s(t) dt < \infty. \end{aligned}$$

Finally, we will show that  $I_{11} < \infty$ . For  $t \geq f(\delta)$ , by the proof of (3.6), for  $n$  large enough, we have that

$$\sum_{i=1}^{k_n} \mathbb{V}(|X_{ni}| > \mu h(t)) < 1.$$

Thus by the fact  $\eta \geq 1$  and conditions (a'), we have

$$\begin{aligned} I_{11} &\leq C \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} \int_{f(\delta)}^{\infty} \mathbb{V}(|X_{ni}| > \mu h(t)) dt \\ &\leq \sum_{n=1}^{\infty} c_n \sum_{i=1}^{k_n} C_{\mathbb{V}} \left[ f\left(|X_{ni}| \left( 1 - g\left(\frac{X_{ni}}{\mu\delta}\right) \right) \right) \right] < \infty. \end{aligned}$$

□

### Declarations

**Conflict of interest** The authors declare no conflict of interest.

**Open Access** This article is licensed under a Creative Commons Attribution 4.0 In-

ternational License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the articles Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the articles Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

## References

- [1] L E Baum, M Katz. *Convergence rates in the law of large numbers*, Trans Amer Math Soc, 1965, 120(1): 108-123.
- [2] P Y Chen, D C Wang. *Complete Moment Convergence for Sequence of Identically  $\alpha$ -mixing Random Variables*, Acta Mathematica Sinica, English Series, 2010, 26(4): 679-690.
- [3] Z J Chen. *Strong laws of large numbers for sub-linear expectations*, Sci China Math, 2016, 59(5): 945-954.
- [4] Y Q Chen, A Y Chen, K W Ng. *The strong law of large numbers for extended negatively dependent random variables*, J Appl Prob, 2010, 47: 908-922.
- [5] Z J Chen, F Hu. *A law of the iterated logarithm for sublinear expectations*, J Financ Eng, 2014, 1: 1-15.
- [6] Y Chow. *On the rate of moment convergence of sample sums and extremes*, Bull Inst Math Acad Sin, 1988, 16: 177-201.
- [7] L Denis, C Martini. *A theoretical framework for the pricing of contingent claims in the presence of model uncertainty*, Ann Appl Probab, 2006, 16(2): 827-852.
- [8] F X Feng, D C Wang, Q Y Wu. *Complete and complete moment convergence for weighted sums of  $\tilde{\rho}$ -mixing random variables*, J Math Inequal, 2018, 12(1): 201-217.
- [9] I Gilboa. *Expected utility theory with purely subjective non-additive probabilities*, J Math Econom, 1987, 16: 65-68.
- [10] C Hu. *A strong law of large numbers for sub-linear expectation under a general moment condition*, Statist Probab Lett, 2016, 119: 248-258.
- [11] H W Huang, D C Wang, Q Y Wu, Q X Zhang. *A note on the complete convergence for sequences of pairwise NQD random variables*, J Inequal Appl, 2011: 92, DOI: 10.1186/1029-242X-2011-92.
- [12] M Marinacci. *Limit laws for non-additive probabilities and their frequentist interpretation*, J Econom Theory, 1999, 84: 145-195.
- [13] M Peligrad, A Gut. *Almost sure results for a class of dependent random variables*, J Theoret Probab, 1999, 12: 87-104.
- [14] S Peng. *BSDE and related  $g$ -expectation*, Pitman Res Notes Math Ser, 1997, 364: 141-159.

- [15] S Peng. *Multi-dimensional  $G$ -Brownian motion and related stochastic calculus under  $G$ -expectation*, Stochastic Process Appl, 2008, 118(12): 2223-2253.
- [16] S Peng. *A new central limit theorem under sublinear expectations*, 2008, ArXiv: 0803.2656v1 [math.PR].
- [17] S Peng. *Nonlinear Expectations and Stochastic Calculus under Uncertainty*, 2010, ArXiv: 1002.4546 [math.PR].
- [18] S Peng. *Law of large numbers and central limit theorem under nonlinear expectations*, 2007, ArXiv: math.PR/0702358v1 [math.PR].
- [19] D H Qiu, P Y Chen. *Complete moment convergence for i.i.d. random variables*, Statist Probab Lett, 2014, 91(12): 76-82.
- [20] S H Sung. *Moment inequalities and complete moment convergence*, J Ineq App, 2009, 2009, Article ID: 271265.
- [21] X Wang, X Deng, L Zheng, S Hu. *Complete convergence for arrays of rowwise negatively superadditive-dependent random variables and its applications*, Statistics: A Journal of Theoretical and Applied Statistics, 2014, 48(4): 834-850.
- [22] Q Y Wu, Y Y Jiang. *Strong law of large numbers and Chover's law of the iterated logarithm under sub-linear expectations*, J Math Anal Appl, 2018, 460: 252-270.
- [23] Y Wu, M Ordóñez Cabrera, A Volodin. *Complete convergence and complete moment convergence for arrays of rowwise end random variables*, Glasnik Matematički, 2014, 49(69): 447-466.
- [24] L X Zhang. *Exponential inequalities under sub-linear expectations with applications to laws of the iterated logarithm*, Sci China Math, 2016, 59(12): 2503-2526.
- [25] L X Zhang. *Rosenthal's inequalities for independent and negatively dependent random variables under sub-linear expectations with applications*, Sci China Math, 2016, 59(4): 751-768.

<sup>1</sup>College of Science, Guilin University of Technology, Guilin 541004, China.

<sup>2</sup>School of Mathematical Science, University of Electronic Science and Technology, Chengdu 611731, China.

Emails: fengfengxiang2013@163.com, wangdc@uestc.edu.cn, wqy666@glut.edu.cn