




Schanuel's Lemma for Exact Categories

Martin Mathieu¹  · Michael Rosbotham¹

Received: 22 January 2022 / Accepted: 20 May 2022
© Crown 2022

Abstract

We prove an injective version of Schanuel's lemma from homological algebra in the setting of exact categories.

Keywords Cohomological dimension · Injective object · Exact structures

Mathematics Subject Classification 18A20 · 18G20 · 18G50

1 Introduction

Schanuel's lemma is a useful tool in homological algebra and category theory. It appears to have come about as a response to a question by Kaplansky, see [4, p. 166], and simplifies the definition of the projective (or, injective) homological dimension in module categories, hence in abelian categories. The typical categories that arise in functional analysis are not abelian but lately, the use of exact structures on additive categories of Banach modules and related ones has been suggested and indeed been exploited successfully.

In [3], Bühler develops homological algebra for bounded cohomology in the setting of Quillen's exact categories. In [1], exact categories of sheaves of operator modules

To the memory of Jörg Eschmeier (1956–2021) who was fond of the use of homology in Functional Analysis.

Communicated by Mihai Putinar.

This article is part of the topical collection "Multivariable Operator Theory. The Jörg Eschmeier Memorial" edited by Raul Curto, Michael Hartz, Mihai Putinar and Ernst Albrecht.

✉ Martin Mathieu
m.m@qub.ac.uk

Michael Rosbotham
mrosbotham01@qub.ac.uk

¹ Mathematical Sciences Research Centre, Queen's University Belfast, Belfast BT7 1NN, Northern Ireland

over C^* -ringed spaces are studied. Relative cohomology and cohomological dimension for (not necessarily self-adjoint) operator algebras is the topic of [6], see also [8]. In view of this, it seems beneficial to establish an injective version of Schanuel’s lemma for exact categories and show how it yields the injective dimension theorem.

When we equip an additive category \mathcal{A} with an exact structure we fix a pair $(\mathcal{M}, \mathcal{P})$ consisting of a class of monomorphisms \mathcal{M} and a class of epimorphisms \mathcal{P} such that each $\mu \in \mathcal{M}$ and $\pi \in \mathcal{P}$ form a kernel-cokernel pair which we write as

$$E \rightharpoonup^{\mu} \rightarrow F \xrightarrow{\pi} \twoheadrightarrow G$$

where E, F and G are objects in \mathcal{A} . We require that \mathcal{M} and \mathcal{P} contain all identity morphisms and are closed under composition, and term their elements as *admissible monomorphisms* and *admissible epimorphisms*, respectively. Furthermore, the push-out of an admissible monomorphism along an arbitrary morphism exists and yields an admissible monomorphism, and, likewise, the pull-back of an admissible epimorphism along an arbitrary morphism exists and yields an admissible epimorphism. If these conditions are fulfilled and $(\mathcal{M}, \mathcal{P})$ is invariant under isomorphisms, $(\mathcal{M}, \mathcal{P})$ is called an *exact structure* on \mathcal{A} and will typically be denoted by $\mathcal{E}x$. The pair $(\mathcal{A}, \mathcal{E}x)$ is said to be an *exact category*.

Unlike in abelian categories not every morphism in an exact category has a canonical factorisation into an epimorphism followed by a monomorphism. One therefore has to restrict to *admissible morphisms* which are those that arise as $\mu \circ \pi$ for some $\mu \in \mathcal{M}$ and $\pi \in \mathcal{P}$. (It is easy to check that, once such factorisation exists, it is unique up to unique isomorphism.)


The kernel-cokernel pairs replace the usual short exact sequences in abelian categories while long exact sequences are built from admissible morphisms. A very readable introduction into exact categories is given in [2].

In this note, we provide the details of how Schanuel’s lemma works in general exact categories and establish the Injective Dimension Theorem (Theorem 3.5).

2 Preliminaries

We include here the necessary terminology and initial results, for a fixed exact category $(\mathcal{A}, \mathcal{E}x)$, where $\mathcal{E}x = (\mathcal{M}, \mathcal{P})$.

Definition 2.1 An object I in an exact category $(\mathcal{A}, \mathcal{E}x)$ is *\mathcal{M} -injective* if, when given $E \rightharpoonup^{\mu} \rightarrow F$ and a morphism $f \in \text{Mor}_{\mathcal{A}}(E, I)$, for objects $E, F \in \mathcal{A}$, there exists a morphism $g \in \text{Mor}_{\mathcal{A}}(F, I)$ making the following diagram commutative



The exact category has *enough \mathcal{M} -injectives* if, for every $E \in \mathcal{A}$, there exist an \mathcal{M} -injective object I and an admissible monomorphism $E \rightarrowtail I$.

We will also make use of the following characterisations of \mathcal{M} -injective objects.

Proposition 2.2 *Let E be an object in an exact category $(\mathcal{A}, \mathcal{E}x)$. The following are equivalent.*

- (i) E is \mathcal{M} -injective;
- (ii) Every admissible monomorphism $E \rightarrowtail F$, for $F \in \mathcal{A}$, has a left inverse;
- (iii) There exist an \mathcal{M} -injective object $I \in \mathcal{A}$ and a morphism $E \rightarrow I$ with a left inverse (i.e., E is a retract of an \mathcal{M} -injective object).

The arguments are standard.

As exact categories are additive, we can form the product of any two objects (and thus, of any finite number of objects).

Proposition 2.3 *Let E, F, G be objects in an additive category \mathcal{A} . The following are equivalent:*

- (i) F is a product of E and G ;
- (ii) F is a coproduct of E and G ;
- (iii) There exist a kernel-cokernel pair in \mathcal{A} ,

$$E \rightarrowtail \xrightarrow{\mu} F \xrightarrow{\pi} \twoheadrightarrow G \tag{2.1}$$

and morphisms $\tilde{\mu} \in \text{Mor}_{\mathcal{A}}(F, E)$ and $\tilde{\pi} \in \text{Mor}_{\mathcal{A}}(G, F)$ such that $\tilde{\mu} \circ \mu = \text{id}_E$ and $\pi \circ \tilde{\pi} = \text{id}_G$, and $\mu \circ \tilde{\mu} + \tilde{\pi} \circ \pi = \text{id}_F$;

- (vi) There exist a kernel-cokernel pair in \mathcal{A} ,

$$E \rightarrowtail \xrightarrow{\mu} F \xrightarrow{\pi} \twoheadrightarrow G \tag{2.2}$$

and a morphism $\tilde{\mu} \in \text{Mor}_{\mathcal{A}}(F, E)$ such that $\tilde{\mu} \circ \mu = \text{id}_E$, the identity morphism on E .

Moreover, if these equivalent conditions are met, the kernel-cokernel pair in Diagram (2.2) will belong to every exact structure that can be placed on \mathcal{A} .

Proof Finite products, coproducts and biproducts coincide in an additive category (see, e.g., [7, Proposition 7.1–Corollary 7.3.]), and condition (iii) is just the definition of F being a biproduct of E and G . That condition (iii) is equivalent to condition (iv) can be proven in the exact same way as the ‘Splitting Lemma’ in module theory (see, e.g., [5, Proposition 4.3.]). The final statement of this proposition is a direct consequence of the conditions required for monomorphisms and epimorphisms to be admissible; see [2, Lemma 2.7.] for details.

Kernel-cokernel pairs satisfying condition (iii) of Proposition 2.3 are said to be *split*. For objects E and F in \mathcal{A} we will denote their (co)product by $E \oplus F$.

Proposition 2.4 Suppose $E \xrightarrow{\mu} F \xrightarrow{\pi} G$ is a kernel-cokernel pair in $\mathcal{E}x$.

(i) For any $A \in \mathcal{A}$, there is a kernel-cokernel pair in $\mathcal{E}x$,

$$E \oplus A \xrightarrow{\quad} F \oplus A \xrightarrow{\quad} G$$

(ii) If $F \cong E \oplus G$, then F is \mathcal{M} -injective if and only if both E and G are \mathcal{M} -injective.

Proof We first prove (i). For $A \in \mathcal{A}$, there exist split kernel-cokernel pairs

$$E \xrightarrow{\iota} E \oplus A \xrightarrow{\rho} A \quad \text{and} \quad A \xrightarrow{\theta} F \oplus A \xrightarrow{\tau} F$$

and $\pi \circ \tau \in \text{Mor}_{\mathcal{A}}(F \oplus A, G)$ is an admissible epimorphism, as a composition of morphisms in \mathcal{P} . Define $\varphi \in \text{Mor}_{\mathcal{A}}(E \oplus A, F \oplus A)$ by

$$\varphi = \tilde{\tau} \circ \mu \circ \tilde{\iota} + \theta \circ \rho$$

using the same notation as in Proposition 2.3. Then $(\varphi, \pi \circ \tau)$ is the desired kernel-cokernel pair.

To show this, it is enough to demonstrate that φ is a kernel of $\pi \circ \tau$. First note the composition

$$(\pi \circ \tau) \circ \varphi = \text{id}_F \circ (\pi \circ \mu) \circ \tilde{\iota} + \pi \circ (\tau \circ \theta) \circ \rho = 0.$$

Now suppose there exist $B \in \mathcal{A}$ and a morphism $f \in \text{Mor}_{\mathcal{A}}(B, F \oplus A)$ such that $(\pi \circ \tau) \circ f = 0$. As μ is a kernel for π , there exists a unique morphism $g' \in \text{Mor}_{\mathcal{A}}(B, E)$ such that $\mu \circ g' = \tau \circ f$. Define $g \in \text{Mor}_{\mathcal{A}}(B, E \oplus A)$ by

$$g = \iota \circ g' + \tilde{\rho} \circ \tilde{\theta} \circ f.$$

Then $\varphi \circ g = (\tilde{\tau} \circ \tau) \circ f + (\theta \circ \tilde{\theta}) \circ f = \text{id}_{F \oplus A} \circ f = f$.

To finish the proof of (i), we show that there is no other morphism $h \in \text{Mor}_{\mathcal{A}}(B, E \oplus A)$ such that $\varphi \circ h = f$. Suppose we have such a morphism h . Then, $\tilde{\theta} \circ f = \tilde{\theta} \circ \varphi \circ h = \rho \circ h$, and $\mu \circ g' = \tau \circ f = \tau \circ \varphi \circ h = \mu \circ \tilde{\iota} \circ h$, and therefore $g' = \tilde{\iota} \circ h$. Combining these facts gives:

$$h = \text{id}_{E \oplus A} \circ h = (\iota \circ \tilde{\iota} + \tilde{\rho} \circ \rho) \circ h = \iota \circ g' + \tilde{\rho} \circ \tilde{\theta} \circ f = g,$$

as required.

For assertion (ii) suppose $F \cong E \oplus G$. Then there exist morphisms $\tilde{\mu} \in \text{Mor}_{\mathcal{A}}(F, E)$ and $\tilde{\pi} \in \text{Mor}_{\mathcal{A}}(G, F)$ such that $\tilde{\mu} \circ \mu = \text{id}_E$ and $\pi \circ \tilde{\pi} = \text{id}_G$, and $\mu \circ \tilde{\mu} + \tilde{\pi} \circ \pi = \text{id}_F$. In particular, E and G are retracts of F . By Proposition 2.2, if F is \mathcal{M} -injective so are E and G . Finally, suppose E and G are \mathcal{M} -injective and there is an admissible monomorphism

$$F \xrightarrow{f} B$$

where $B \in \mathcal{A}$. Because E and G are \mathcal{M} -injective, there exist $g_E \in \text{Mor}_{\mathcal{A}}(B, E)$ such that $\tilde{\mu} = g_E \circ f$ and $g_G \in \text{Mor}_{\mathcal{A}}(B, G)$ such that $\pi = g_G \circ f$. Let $g = \mu \circ g_E + \tilde{\pi} \circ g_G$, then g is a left inverse of f , indeed:

$$g \circ f = \mu \circ (g_E \circ f) + \tilde{\pi} \circ (g_G \circ f) = \mu \circ \tilde{\mu} + \tilde{\pi} \circ \pi = \text{id}_F.$$

Hence, by Proposition 2.2, F is \mathcal{M} -injective. □

3 Schanuel's Lemma

Fix an exact category $(\mathcal{A}, \mathcal{E}x)$. The following is the injective version of Schanuel's lemma for exact categories.

Proposition 3.1 *Suppose $E \rhd \xrightarrow{\mu} I \twoheadrightarrow F$ and $E \rhd \xrightarrow{\mu'} I' \twoheadrightarrow F'$ are kernel-cokernel pairs in $\mathcal{E}x$, and that I, I' are \mathcal{M} -injective objects. Then $I \oplus F' \cong I' \oplus F$ in \mathcal{A} .*

Proof First, by the axioms of an exact structure, we can form the following push-out,

$$\begin{array}{ccc}
 E & \xrightarrow{\mu} & I \\
 \mu' \downarrow & & \downarrow h \\
 I' & \xrightarrow{h'} & C
 \end{array}
 \tag{3.1}$$

where every morphism is an admissible monomorphism. Extending this diagram to include the given cokernels, and adding in some zero morphisms, we get the following commutative diagram:

$$\begin{array}{ccccc}
 & & E & \xrightarrow{\mu} & I & \xrightarrow{\pi} & F \\
 & 0 & \downarrow \mu' & & \downarrow h & & \uparrow \\
 & & I' & \xrightarrow{h'} & C & & \\
 & & \downarrow \pi' & & \downarrow & & \\
 & & F' & & & & \\
 & & & & & & 0
 \end{array}
 \tag{3.2}$$

By the universal property of push-outs, there are a unique morphism $p \in \text{Mor}(C, F)$ such that $ph' = 0$ and $ph = \pi$, and a unique morphism $p' \in \text{Mor}(C, F')$ such that $p'h = 0$ and $p'h' = \pi'$. Hence, we have the following commutative diagram:

Corollary 3.2 *Suppose there is a diagram of morphisms in an exact category $(\mathcal{A}, \mathcal{E}x)$ of the form*

$$\begin{array}{ccccc} E & \hookrightarrow & I & \twoheadrightarrow & F \\ \cong \downarrow & & & & \\ E' & \hookrightarrow & I' & \twoheadrightarrow & F' \end{array}$$

such that I and I' are \mathcal{M} -injective, the horizontal lines are in $\mathcal{E}x$ and the vertical arrow is an isomorphism. Then $I \oplus F' \cong I' \oplus F$ in \mathcal{A} .

We extend Schanuel's lemma to injective resolutions in Proposition 3.4 below. Recall that a morphism is *admissible* if it is the composition $\mu \circ \pi$ for some $\mu \in \mathcal{M}$ and $\pi \in \mathcal{P}$. Such factorisation is unique up to unique isomorphism ([2, Lemma 8.4]).

Definition 3.3 For an object $E \in \mathcal{A}$, an \mathcal{M} -injective resolution of E is a sequence of admissible morphisms of the form:

$$\begin{array}{ccccccccccc} E & \hookrightarrow & I^0 & \longrightarrow & \cdots & \longrightarrow & I^{n-1} & \longrightarrow & I^n & \longrightarrow & \cdots \\ & \searrow \cong & \swarrow & & & & \swarrow & & \swarrow & & \swarrow \\ & & G^0 & & & & G^{n-1} & & G^n & & G^{n+1} \end{array}$$

such that, for each $n \geq 0$, the object I^n is \mathcal{M} -injective, and

$$G^n \hookrightarrow I^n \twoheadrightarrow G^{n+1}$$


forms a kernel-cokernel pair in $\mathcal{E}x$ (this is the *exactness condition at I^n*).

If \mathcal{A} has enough \mathcal{M} -injectives, we can build an injective resolution for every object in \mathcal{A} .

Proposition 3.4 *Suppose we have the following \mathcal{M} -injective resolutions of E , with the factorisation of each admissible morphism included:*

$$\begin{array}{ccccccccccc} E & \hookrightarrow & I^0 & \longrightarrow & \cdots & \longrightarrow & I^{n-1} & \longrightarrow & I^n & \longrightarrow & \cdots \\ & \searrow \cong & \swarrow & & & & \swarrow & & \swarrow & & \swarrow \\ & & G^0 & & & & G^{n-1} & & G^n & & G^{n+1} \end{array}$$

and



Then, for each $n \geq 1$, we have isomorphisms

$$I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2n-1} \oplus G^{2n} \cong J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2n-1} \oplus H^{2n}$$

and

$$I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2n-1} \oplus I^{2n} \oplus H^{2n+1} \cong J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2n-1} \oplus J^{2n} \oplus G^{2n+1}.$$

Proof We prove this by induction. For $n = 1$, first note that Corollary 3.2, applied to the diagram

$$\begin{array}{ccccc}
 G^0 & \twoheadrightarrow & I^0 & \twoheadrightarrow & G^1 \\
 \cong \downarrow & & & & \\
 H^0 & \twoheadrightarrow & J^0 & \twoheadrightarrow & H^1
 \end{array}$$

gives $I^0 \oplus H^1 \cong J^0 \oplus G^1$. By Proposition 2.4, there is a diagram of the form

$$\begin{array}{ccccc}
 I^0 \oplus H^1 & \twoheadrightarrow & I^0 \oplus J^1 & \twoheadrightarrow & H^2 \\
 \cong \downarrow & & & & \\
 J^0 \oplus G^1 & \twoheadrightarrow & J^0 \oplus I^1 & \twoheadrightarrow & G^2
 \end{array}$$

and Corollary 3.2 gives $I^0 \oplus J^1 \oplus G^2 \cong J^0 \oplus I^1 \oplus H^2$. To finish the proof for $n = 1$, we again apply Proposition 2.4 followed by Corollary 3.2, to get a diagram

$$\begin{array}{ccccc}
 I^0 \oplus J^1 \oplus G^2 & \twoheadrightarrow & I^0 \oplus J^1 \oplus I^2 & \twoheadrightarrow & G^3 \\
 \cong \downarrow & & & & \\
 J^0 \oplus I^1 \oplus H^2 & \twoheadrightarrow & J^0 \oplus I^1 \oplus J^2 & \twoheadrightarrow & H^3
 \end{array}$$

and an isomorphism $I^0 \oplus J^1 \oplus I^2 \oplus H^3 \cong J^0 \oplus I^1 \oplus J^2 \oplus G^3$.

Assume the result holds some $n \geq 1$. By Proposition 2.4, there is a diagram of the form

$$\begin{array}{ccc} I^0 \oplus \dots \oplus I^{2n} \oplus H^{2n+1} & \xrightarrow{\quad} & I^0 \oplus \dots \oplus I^{2n} \oplus J^{2n+1} \longrightarrow \twoheadrightarrow H^{2(n+1)} \\ \cong \downarrow & & \\ J^0 \oplus \dots \oplus J^{2n} \oplus G^{2n+1} & \xrightarrow{\quad} & J^0 \oplus \dots \oplus J^{2n} \oplus I^{2n+1} \longrightarrow \twoheadrightarrow G^{2(n+1)} \end{array}$$

and Corollary 3.2 gives

$$\begin{aligned} I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2(n+1)-1} \oplus G^{2(n+1)} \\ \cong J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2(n+1)-1} \oplus H^{2(n+1)}. \end{aligned}$$

One final application of Proposition 2.4 yields the following diagram:

$$\begin{array}{ccc} I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2n+1} \oplus G^{2(n+1)} & \xrightarrow{\cong} & J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2n+1} \oplus H^{2(n+1)} \\ \downarrow & & \downarrow \\ I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2n+1} \oplus I^{2(n+1)} & & J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2n+1} \oplus J^{2(n+1)} \\ \downarrow & & \downarrow \\ G^{2(n+1)+1} & & H^{2(n+1)+1} \end{array}$$

By Corollary 3.2,

$$\begin{aligned} I^0 \oplus J^1 \oplus I^2 \oplus \dots \oplus J^{2(n+1)-1} \oplus I^{2(n+1)} \oplus H^{2(n+1)+1} \\ \cong J^0 \oplus I^1 \oplus J^2 \oplus \dots \oplus I^{2(n+1)-1} \oplus J^{2(n+1)} \oplus G^{2(n+1)+1} \end{aligned}$$

as required. □

We can now prove the Injective Dimension Theorem.

Theorem 3.5 *Let \mathcal{M} be the class of admissible monomorphisms in an exact category $(\mathcal{A}, \mathcal{E}x)$. Suppose \mathcal{A} has enough \mathcal{M} -injectives. The following are equivalent for $n \geq 1$ and every $E \in \mathcal{A}$.*

(i) *If there is an exact sequence of admissible morphisms*

$$E \xrightarrow{\quad} I^0 \longrightarrow \dots \longrightarrow I^{n-1} \longrightarrow \twoheadrightarrow F \tag{3.6}$$

with each I^m , $0 \leq m \leq n - 1$ injective, then F must be injective;

(ii) *There is an exact sequence of admissible morphisms*

$$E \rightarrow I^0 \rightarrow \dots \rightarrow I^{n-1} \twoheadrightarrow I^n \tag{3.7}$$

with each I^m , $0 \leq m \leq n$ injective.

Proof Let $E \in \mathcal{A}$. First we show (i) implies (ii). As \mathcal{A} has enough \mathcal{M} -injectives, we can build an \mathcal{M} -injective resolution of E :

$$\begin{array}{ccccccccccc}
 E & \longrightarrow & J^0 & \longrightarrow & \dots & \longrightarrow & J^{n-1} & \longrightarrow & J^n & \longrightarrow & \dots \\
 \cong \searrow & & \nearrow & & \searrow & & \nearrow & & \searrow & & \nearrow \\
 & & G^0 & & G^1 & & G^{n-1} & & G^n & &
 \end{array}$$

Relabel J^k as I^k for all $0 \leq k \leq n - 1$ and G^n as I^n , this gives an exact sequence as in Diagram (3.7), and I^n must be \mathcal{M} -injective, by condition (i).

Now suppose that condition (ii) holds. There must exist an injective resolution of E of the form

$$\begin{array}{ccccccccccc}
 E & \longrightarrow & J^0 & \longrightarrow & \dots & \longrightarrow & J^{n-1} & \longrightarrow & J^n & \longrightarrow & \dots \\
 \cong \searrow & & \nearrow & & \searrow & & \nearrow & & \searrow & & \nearrow \\
 & & H^0 & & H^1 & & H^{n-1} & & J^n & & \cong
 \end{array}$$

and for any exact sequence as in Diagram (3.6), with each I^n injective, there exists an injective resolution

$$\begin{array}{ccccccccccc}
 E & \longrightarrow & I^0 & \longrightarrow & \dots & \longrightarrow & I^{n-1} & \longrightarrow & I^n & \longrightarrow & \dots \\
 \cong \searrow & & \nearrow & & \searrow & & \nearrow & & \searrow & & \nearrow \\
 & & G^0 & & G^1 & & G^{n-1} & & G^n & &
 \end{array}$$

with $G^n = F$. By Proposition 3.4, there exists a kernel-cokernel pair

$$F \xrightarrow{\mu} I \xrightarrow{\pi} G$$

and a morphism $\tilde{\mu} \in \text{Mor}_{\mathcal{A}}(I, F)$ such that $\tilde{\mu} \circ \mu = \text{id}_F$, and I is a finite product of \mathcal{M} -injective objects. Then by Proposition 2.4, I is injective and $\tilde{\mu}$ is a left inverse for μ , hence, by Proposition 2.2, F is \mathcal{M} -injective.


Definition 3.6 Let \mathcal{M} be the class of admissible monomorphisms in an exact category $(\mathcal{A}, \mathcal{E}x)$. We say $E \in \mathcal{A}$ has *finite \mathcal{M} -injective dimension* if there exists an exact sequence of admissible morphisms as in Diagram (3.7) with all I^m \mathcal{M} -injective. If E

is of finite \mathcal{M} -injective dimension we write $\text{Inj}_{\mathcal{M}\text{-dim}}(E) = 0$ if E is \mathcal{M} -injective and $\text{Inj}_{\mathcal{M}\text{-dim}}(E) = n$ if E is not \mathcal{M} -injective and n is the smallest natural number such that there exists an exact sequence of admissible morphisms as in Diagram (3.7) where every I^m is \mathcal{M} -injective. If E is not of finite \mathcal{M} -injective dimension, we write $\text{Inj}_{\mathcal{M}\text{-dim}}(E) = \infty$.

The *global dimension* of the exact category $(\mathcal{A}, \mathcal{E}x)$ is

$$\sup \{ \text{Inj}_{\mathcal{M}\text{-dim}}(E) \mid E \in \mathcal{A} \} \in \mathbb{N}_0 \cup \{ \infty \}.$$

Remark 3.7 The \mathcal{M} -injective dimension of an object E in an exact category $(\mathcal{A}, \mathcal{E}x)$ can be obtained by examining any of its \mathcal{M} -injective resolutions. Indeed, suppose the following is an \mathcal{M} -injective resolution of E (with the factorisation of each admissible morphism included):



Then, by Theorem 3.5, $\text{Inj}_{\mathcal{M}\text{-dim}}(E) \leq n$ if and only if G^n is \mathcal{M} -injective.

The original version of Schanuel’s lemma is formulated for projective resolutions, see, e.g., [4, Lemma 5.1] or [9, Theorem 3.41]. An analogous version using the epimorphisms in the class \mathcal{P} can be obtained in any exact category with exact structure $(\mathcal{M}, \mathcal{P})$.

Data Availability Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Ara, P., Mathieu, M.: Sheaf cohomology for C^* -algebras. Memoir in preparation
2. Bühler, T.: Exact categories. *Expo. Math.* **28**(1), 1–69 (2010)
3. Bühler, T.: On the algebraic foundations of bounded cohomology. *Mem. Amer. Math. Soc.*, 214(1006):xxii+97, (2011)
4. Lam, T.Y.: Lectures on modules and rings. Graduate Texts in Mathematics, vol. 189. Springer-Verlag, New York (1999)
5. Mac Lane, S.: Homology. *Classics in Mathematics*. Springer-Verlag, Berlin, (1995). Reprint of the 1975 edition

6. Mathieu, M., Rosbotham, M.: Exact structures for operator modules. *Canad. J. Math.*, to appear, [arXiv:2105.05006](https://arxiv.org/abs/2105.05006)
7. Osborne, M.S.: Basic homological algebra. Graduate Texts in Mathematics, vol. 196. Springer-Verlag, New York (2000)
8. Rosbotham, M.: Cohomological dimension for C^* -algebras. PhD. Thesis, Queen's University Belfast, Belfast, (2021)
9. Rotman, J.J.: *Notes on homological algebras*. Van Nostrand Reinhold Mathematical Studies, No. 26. Van Nostrand Reinhold Co., New York-Toronto-London, (1970)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.