

Distribution of Protists in the Deep South China Sea Revealed by High-Throughput Sequencing

LI Xinran¹⁾, WARREN Alan²⁾, JIAO Nianzhi¹⁾, and XU Dapeng¹⁾,*

1) State Key Laboratory of Marine Environmental Science, Institute of Marine Microbes and Ecospheres,

College of Ocean and Earth Sciences, Xiamen University, Xiamen 361102, China

2) Department of Life Sciences, Natural History Museum, London, SW7 5BD, UK

(Received February 9, 2019; revised April 28, 2019; accepted June 12, 2019)

© Ocean University of China, Science Press and Springer-Verlag GmbH Germany 2019

Abstract Protists (microbial eukaryotes) are indispensable members of the marine microbial food web. In recent years, organisms living in the deep sea (>1000 m water depth) have increasingly become the focus of research; however, studies on protistan assemblages are relatively scarce compared with their prokaryotic counterparts. In the present study, high-throughput sequencing of the hypervariable V9 region of the 18S rRNA gene was used to explore the community composition of protists in bathypelagic waters of the South China Sea. Based on the analysis of the alpha and beta diversities of 14 samples, we discovered: 1) members belonging to Rhizaria, Alveolata, and Excavata were the dominant groups in terms of both relative sequence abundance and operational taxonomic unit (OTU) richness in all samples, although their relative contributions differed among different samples; 2) cluster analysis showed that the distribution of protistan assemblages was related neither to the sampling location nor to the water depth, and other environmental factors might have caused the differences among the communities; 3) phototrophs, including members of the Bacillariophyta, Bolidophyceae, Dictyochophyceae, Prasinophyceae, and Prymnesiophyceae, were detected in all samples, which indicated their contributions to the downward transportation *via* the biological pump and the potential presence of phagotrophy of these phototrophic cells in the deep ocean.

Key words bathypelagic water; diversity; microbial eukaryotes; SSU rRNA gene

1 Introduction

Marine protists (microbial eukaryotes) are highly diverse (both morphologically and phylogenetically) unicellular eukaryotes that occur in almost every branch of eukaryotic evolutionary tree (Bass *et al.*, 2005; Guillou *et al.*, 2008; Song *et al.*, 2009; Caron *et al.*, 2012; Sun *et al.*, 2017; Adl *et al.*, 2019; Hu *et al.*, 2019). Their size range spans more than four orders of magnitude (Caron *et al.*, 2009). The nutritional modes of protists include phototrophy, heterotrophy, and mixotrophy, and they serve as primary producers, consumers, decomposers and trophic links in aquatic food webs which are essential for the biogeochemical cycles of the ocean (Azam *et al.*, 1983; Jiao *et al.*, 2010; Caron *et al.*, 2017).

The deep dark ocean is divided into three parts: the mesopelagic zone (200–1000 m depth), the bathypelagic zone (1000–4000 m depth), and the abyssal zone (>4000 m depth). It is considered to be the largest habitat on Earth and is characterized by high inorganic nutrient concentrations, low temperature, high pressure, and is the world's largest reservoir of dissolved organic carbon

(Aristegui *et al.*, 2009; Herndl and Reinthaler, 2013). Due to the expense and technical difficulties in sampling and performing *in situ* experiments, both prokaryotic and eukaryotic microbes living in the deep waters are understudied compared with the microbes living in the sunlit ocean (Aristegui *et al.*, 2009). Fortunately, the culture-independent techniques such as denaturing gradient gel electrophoresis (DGGE), clone library, and high-throughput sequencing using marker genes (*e.g.*, the small subunit ribosomal DNA) have provided powerful methods for studying microbial diversity in the deep ocean (Aristegui *et al.*, 2009). Nevertheless, deep sea protists have been largely neglected compared with their prokaryotic counterparts (bacteria and archaea), leaving their diversity and community composition largely unresolved. Studies on deep sea protists both on local and global scales have increased significantly in recent years, uncovering their new clades, distribution patterns and potential ecological roles (Countway *et al.*, 2007; Not *et al.*, 2007; Pernice *et al.*, 2015; Xu *et al.*, 2017a, b). However, given the vast area of the deep sea environment, much work needs to be done in order to reveal the biodiversity of protists and the roles they play in biogeochemical cycles in the deep sea.

In the present study, high-throughput sequencing of the

* Corresponding author. E-mail: dapengxu@xmu.edu.cn

hypervariable V9 region of the SSU rRNA gene was performed to investigate the community composition of protists from the bathypelagic waters of the South China Sea, one of the largest marginal seas in the world.

2 Materials and Methods

2.1 Sample Collection

Six sampling stations were chosen to represent the central part of the deep basin in the South China Sea (SCS) (Fig. 1). Seawater samples were collected using Niskin bottles attached to a CTD rosette system aboard the *R/V Dongfanghong-2* in April 2012. Six samples were taken from 1000 m depth at all stations, five samples were taken from 2000 m depth except at station H7, and three samples were taken at 4000 m depth at stations H7, O3 and O14, respectively. For each sample, two liters of sea water were prefiltered through 200 μm Nitex (Sefar) mesh and then collected by polycarbonate filters with a pore-size of 0.22 μm (Millipore). The filters were immediately frozen in liquid nitrogen and stored at -80°C until further treatment.




Fig. 1 Geographic locations of the sampling sites in the central South China Sea. The map was generated using Ocean Data View 4 software (Schlitzer, 2011).

2.2 DNA Extraction, PCR Amplification, and High-Throughput Sequencing of SSU rRNA Gene

Genomic DNA was extracted from each sample using the Power-Water DNA Isolation Kit (MoBio Laboratories, Inc., Carlsbad, CA) following the protocols from the manufacturer with minor modifications (Sun *et al.*, 2017; Xu *et al.*, 2017a). Extracted DNA was stored at -20°C for later processing. Universal primers 1389F (5'-TTGTACA CACCGCCC-3') and 1510R (5'-CCTTCYGCAGGTTACCTAC-3') were used to amplify the hypervariable V9 region of the SSU rRNA gene (Amaral-Zettler *et al.*, 2009). Six individual PCR reactions for each sample were run employing Ex Taq DNA polymerase (TaKaRa, Dalian, China) and the products were pooled to collect sufficient amplicons for sequencing. The pooled PCR products were purified using Wizard[®]SV Gel and PCR Clean-Up System (Promega, Beijing, China). Bridge amplification and paired end sequencing of the amplicons were performed with an Illumina MiSeq platform by a commercial

sequencing company. Sequence data generated have been deposited in the NCBI Sequence Read Archive with accession number SRP104547.

2.3 Sequence Analysis

Quality filtering, demultiplexing and assembly of raw data were conducted with Trimmomatic (Bolger *et al.*, 2014) and Flash software (Magoč and Salzberg, 2011) and criteria employed followed Li *et al.* (2018). Potential chimeras were identified and removed using UCHIME (Edgar *et al.*, 2011) applying both *de novo* and reference-based chimera searches against the Protist Ribosomal Database 2 (PR2) (de Vargas *et al.*, 2015). Singletons (reads present as a single copy) were removed from downward analysis. Operational taxonomic units (OTUs) were clustered at a 95% similarity threshold using UPARSE (Edgar, 2013). Taxonomy assignment of each OTU were achieved using BLAST method against PR2 as implemented in QIIME (Caporaso *et al.*, 2010). Non-protistan OTUs (*e.g.*, bacteria, archaea, metazoa, fungi, plastids, or OTUs classified as 'Unassigned') were removed.

All datasets were sub-sampled at a uniform depth of 13842 sequences (the lowest sequence count for all samples) before downstream analysis. An OTU table was generated and alpha diversity indices (Chao1, Shannon, ACE, Inverse of Simpson, and Phylogenetic Diversity) were calculated using QIIME (Caporaso *et al.*, 2010). Non-metric multidimensional scaling (NMDS) was constructed using the Vegan package as implemented in R based on the Bray-Curtis coefficient. Weighted Unifrac metric, which compares samples based on the phylogenetic relatedness of OTUs in a community and considers relative OTU abundance, was used to infer the grouping of samples (Lozupone and Knight, 2005).

3 Results

To infer the diversity of protistan assemblages in the bathypelagic waters of the central South China Sea, a total of 14 samples from 6 stations were collected (Fig. 1). After quality-screening, a total of 857544 sequences with an average length of 123 bp (mean \pm std: 123.0545 \pm 16.4996) were generated. After removal of potential chimeras, singletons, and sequences that were not affiliated with protists, there were a total of 366951 sequences, ranging 13842 to 45271 reads per sample (Table 1). All datasets were rarefied at a uniform depth of 13842 sequences (the lowest number of sequences for all samples). The final OTU table containing 1911 OTUs and 193788 sequences was used to analyze alpha and beta diversities and taxonomic affiliations of bathypelagic protists.

3.1 Alpha and Beta Diversities of Bathypelagic Protists

Alpha diversity estimates are shown in Table 1. The richness observed varied among samples, ranging from 927 to 1577 OTUs per sample. The average Shannon index was 5.57 ± 0.40 , while the highest one (7.24) was found at 2000 m deep waters of station O1 and the lowest

Table 1 Diversity estimates of South China Sea samples

Sample ID	Site	Depth (m)	Coordinates	Clean reads	OTU _{0.05} *	Chao1*	ACE*	Shannon*	Simpson*	PD*
G4.1000	G4	1000	115.975°E, 17.966°N	20920	1123	1862	2027	4.41	0.003	161.7
G4.2000	G4	2000		13842	995	1703	1794	4.38	0.003	150.5
H7.1000	H7	1000	112.987°E, 17.002°N	14580	1014	1824	1961	5.14	0.005	152.4
H7.1500	H7	1500		45271	1195	1856	1913	5.77	0.006	167.5
O1.1000	O1	1000	117.004°E, 16.414°N	24601	1577	2327	2381	7.15	0.012	213.5
O1.2000	O2	2000		28665	1467	2348	2441	7.24	0.024	196.8
O14.1000	O14	1000		14722	1369	2250	2374	5.79	0.005	180.3
O14.2000	O14	2000	118.809°E, 15.101°N	43760	951	1631	1748	3.25	0.002	138.5
O14.4000	O14	4000		21128	1433	2057	2105	7.81	0.031	186.3
O3.1000	O3	1000		30349	985	1704	1928	3.50	0.002	142.1
O3.2000	O3	2000	115.242°E, 14.724°N	16970	1063	1692	1817	5.26	0.005	159.9
O3.4000	O3	4000		27149	1157	1788	1854	6.83	0.027	165.0
O7.1000	O7	1000	117°E, 13.457°N	28379	1501	2437	2435	7.07	0.014	199.7
O7.2000	O7	2000		36615	927	1742	1677	4.31	0.005	137.9

Notes: OTU_{0.05}, Operational taxonomic unit at 95% 18S rRNA V9 gene sequence identity. * Standardized numbers based on subsampling of 13842 sequences without replacement.




Fig.2 Alpha diversity of protists in bathypelagic waters of the South China Sea as inferred by clustering reads at 95% similarity. (A) Rarefaction curves for each sample. (B) Global rarefaction curve.




Fig.3 Alpha diversity measures (Observed species, Phylogenetic Diversity, Shannon and Chao1) for the pooled samples grouped by water depths. The box delimits the 25th and 75th percentiles. The whisker shows the range and the line in every box plot indicates the median.

one was at 2000m deep waters of station O14. Phylogenetic diversity (PD), which measures the total branch length connecting all OTUs in the SSU rRNA gene phylogeny, ranged from 213.5 (1000m deep waters at station O1) to 137.9 (2000m deep waters at station O7) (Table 1). Rarefaction curves showed that all samples were not fully sampled (Fig.2A). However, a rarefaction curve for the pooled dataset from all samples showed a symbol saturation that a maximal richness of bathypelagic protists was approximately 2000 OTUs (Fig.2B).

To reveal the differences of alpha diversity at different depths, samples retrieved from the same depth were pooled and the nonparametric test (Monte Carlo permutations) was conducted based on the Observed species, Phylogenetic diversity, Shannon and Chao1. As a general trend, the average alpha diversity decreased as water depth

increased although there was no significant difference among sample groups (Fig.3).

A non-metric multidimensional scaling ordination (NMDS) analysis, based on Bray-Curtis similarity, was implemented to explore the community composition of bathypelagic protists among all 14 samples. The samples were clustered into three groups. Group 1 contained samples O14.4000 and G4.2000; group 2 contained samples O14.1000, O3.1000, and H7.1000; and group 3 contained the rest of the samples (Fig.4A). This grouping pattern was also supported by the principle component analysis (PCoA) of community taxonomic relatedness quantified by the Weighted Unifrac metric (Fig.4B). Statistical analyses showed the composition of the three groups was significantly different (ANOSIM, $R=0.999$, $P=0.001$ for the NMDS and $R=0.9343$, $P=0.001$ for the PCoA analysis).




Fig.4 Plots of nonmetric multidimensional scaling (nMDS) ordination (A) and unweighted unifrac principal coordinates analysis (PCoA) (B).




Fig.5 Venn diagram shows the OTUs shared among group 1 (samples G4.2000 and O14.4000), group 2 (samples H7.1000, O3.1000, and O14.1000), and group 3 (the rest samples) of protistan communities as indicated in Fig.4. Pie charts show the unique OTUs in each group and OTUs occurred in all three groups.

OTUs found only in group 1 were dominated by members in Alveolata (about 74%), followed by Hacrobia (about 14%). OTUs occurred only in group 2 were mainly composed of members in Alveolata (about 56%), followed by Rhizaria (about 14%) and Stramenopiles (13%). Group 3 specific OTUs were dominated by members in Excavata (about 71%), followed by Alveolata (about 15%). Approximately 45% of all OTUs were discovered from all three groups and were mainly composed of members in Rhizaria (55%), followed by members in Alveolata (25%) and Excavata (11%) (Fig.5).

3.2 Community Composition of Bathypelagic Protists

Pooling all data from all samples gave the first insight of the protistan community composition in the bathypelagic waters of the South China Sea. The reads affiliated with Rhizaria represented over half of the total protists, followed by Alveolata (26%) and Excavata (13%) (Fig.6A). The other supergroups, including Stramenopiles, Opisthokonta, Archaeplastida, Amoebozoa, Hacrobia and Picozoa, collectively contributed less than 8% to the total reads (Fig.6A). The community composition varied substantially in each sample, e.g., Rhizaria dominated protistan communities in most samples except those retrieved

from waters with 2000 m depth at site G4 (G4.2000), 4000m depth at site O14 (O14.4000), and 1000m depth at site O3 (O3.1000) in which Alveolata replaced Rhizaria as the dominant taxon (Fig.6B). In at least half samples, Excavata surpassed Alveolata to be the second-most dominant group (Fig.6B). In terms of OTU richness, the dominant supergroup in the pooled dataset was Alveolata (about 47%), followed by Excavata (about 23%), Rhizaria (about 12%), and Stramenopiles (about 9%) (Fig.6C). This trend was consistent in all individual samples except G4.2000 and O14.4000 (Fig.6D).

The three dominant supergroups, including Rhizaria, Alveolata and Excavata, were further examined at lower taxonomic ranks (Fig.7). For Rhizaria, Spumellarida was the most dominant group (on average 85% of rhizarian reads) in all samples, followed by RAD-A (on average 10%) (Figs.7A, B). For Alveolata, Dinophyceae and MALV-I were the two most dominant groups, representing 35% and 27% of all reads, respectively. Members in MALV-II contributed 18% of Alveolata-affiliated reads. Apicomplexa was the fourth most dominant supergroup; however, in samples O3.1000 and O3.2000, it replaced Dinophyceae to be the most dominant alveolate group with a contribution of 40% (Figs.7C, D). For Excavata, members in Diplonemea and Neobodonid contributed equally and they constituted 91% of all Excavata-affiliated reads (Figs.7E, F). Phototrophic groups, including Bacillariophyta, Bolido-

phyceae, Dictyochophyceae, Prasinophyceae, and Prymnesiophyceae, were retrieved from all samples and their contributions to total reads ranged from 0.3% to 16% in each sample (Figs.7G, H).

The ten most abundant OTUs in the pooled dataset belonged to Spumellarida (three OTUs), Discoba (three OTUs), RAD-A (two OTUs), and Dinophyceae (two OTUs). These OTUs had high similarities with environmental sequences in the GenBank database, and seven of them showed 100% similarities (Table 2). OTU_1 was the most abundant OTU and was a polycystine radiolarian that was 94.3% similar to *Cladococcus viminalis*. It was followed by OTU_3091 which showed 92.8% similarity to the same species whereas OTU_25 was identical to *C. viminalis*. OTU_10 and OTU_37 showed rather low similarities with their nearest named neighbor in the GenBank database, namely *Sticholonche* sp. (HQ651785) (79.4% and 80.3%, respectively). OTU_31 and OTU_35 were members of Discoba and showed 77.6% and 77.8% similarities with their nearest named neighbors *Hemistasia phaeocysticola* (AB948221) and *Rhynchopus* sp. (AF38 0997), respectively (Table 2). OTU_15 was also a member of Discoba and showed 100% similarity with *Neobodo designis* (DQ207580). OTU_21 and OTU_74 showed high similarities with their nearest named neighbors, *Lepidodinium viride* (AB686256) (100%) and the toxic bloom-forming dinophyceae, *Karlodinium micrum* (JF79 1049) (99.2%).




Fig.6 Relative abundance of reads in pooled samples (A) and in each sample (B). Relative OTU richness in pooled samples (C) and in each sample (D).




Fig.7 Pie charts calculated for the percentage of total reads in all samples of different taxa from Rhizaria (A), Alveolata (C), Excavata (E), and phototrophic (G) groups. Histogram shows the relative abundances of reads from Rhizaria (B), Alveolata (D), Excavata (F), and phototrophic (H) groups in each sample. Different groups are depicted in different colors.

Table 2 List of the top ten most abundant OTUs

OTU ID	Reads	Group	NN	%-S	NNN	%-S
OTU_1	48068 (24.9%)	Spumellarida	KJ761720	100.0	<i>Cladococcus viminalis</i> (HQ651782)	94.3
OTU_3091	28221 (14.6%)	Spumellarida	KJ760941	99.3	<i>Cladococcus viminalis</i> (HQ651782)	92.8
OTU_10	5963 (3.1%)	RAD-A	KJ762813	99.3	<i>Sticholonche</i> sp. (HQ651785)	79.4
OTU_25	5082 (2.6%)	Spumellarida	KF129727	100.0	<i>Cladococcus viminalis</i> (HQ651782)	100.0
OTU_31	2700 (1.4%)	Discoba	KJ762626	100.0	<i>Hemistasia phaeocysticola</i> (AB948221)	77.6
OTU_21	2675 (1.4%)	Dinophyceae	KU743849	100.0	<i>Lepidodinium viride</i> (AB686256)	100.0
OTU_37	2473 (1.3%)	RAD-A	KF129909	100.0	<i>Sticholonche</i> sp. (AY268045)	80.3
OTU_15	1799 (0.9%)	Discoba	EF432544	100.0	<i>Neobodo designis</i> (DQ207580)	100.0
OTU_35	1766 (0.9%)	Discoba	KJ762516	99.2	<i>Rhynchopus</i> sp. (AF380997)	77.8
OTU_74	1687 (0.9%)	Dinophyceae	KF130115	100.0	<i>Karlodinium micrum</i> (JF791049)	99.2

Notes: %-S, similarities; NN, the nearest neighbor; NNN, the nearest named neighbor.

4 Discussion

High-throughput sequencing, which can detect species at rather low abundance and bypass the difficulties of conventional morphology-based methods, was used to reveal the composition of protist communities in bathypelagic waters of the central South China Sea. The main aim was to characterize the community structure and diversity patterns of protists, which is helpful for understanding their roles in deep oceanic waters.

4.1 Protistan Community Composition in the Bathyal Zone of the South China Sea

In the present study, the average alpha diversity was found to decrease as water depth increased, although no significant difference was found among sample groups. Previous studies have shown that bacterial richness declines by about 25% from the epipelagic layer to the bathypelagic waters (Moeseneder *et al.*, 2001; Hewson *et al.*, 2006) and temperature was proposed to be a major stratifying factor of bacterial communities overriding hydrostatic pressure (Martin-Cuadrado *et al.*, 2007). Studies have also shown that protistan diversity declines with depth along the water columns (Countway *et al.*, 2007; Xu *et al.*, 2017b; Giner *et al.*, 2019). The decline of protistan diversity with depth has been interpreted as an indication of a finite number of ecological niches present in the deep ocean for protists (Countway *et al.*, 2007).

In terms of relative sequence abundances, members affiliated with Rhizaria constituted the most dominant group followed by Alveolata, though there were variabilities among individual samples. Excavata surpassed Stramenopiles to be the third most dominant group (Figs. 5A, B). In terms of OTU richness, Alveolata was the most dominant group followed by Excavata and Rhizaria (Figs. 5C, D). Rhizarians have been repeatedly identified to dominate deep sea protistan communities and, in some cases, they were also reported to possess the highest richness (Pernice *et al.*, 2015; Xu *et al.*, 2017a). In the present study, the dominant groups of rhizarians were RAD_A, B, C and Polycystinea, which is consistent with previous analyses of bathypelagic radiolarian communities (Pernice *et al.*, 2015). A survey using clone library construction performed on the samples collected from the same cruise also showed the dominance of Rhizaria in deep waters with Spumellarida being the greatest contributor in this supergroup (Xu *et al.*, 2017a). It is noteworthy that the cell-size of rhizarians is usually larger than the majority of planktonic protists, the standing stock of which in the top 200 m of the world's oceans is estimated to be equivalent to 5.2% of the total oceanic biota carbon reservoir (Biard *et al.*, 2016). Furthermore, rhizarians also contribute to the downward transportation of carbon via the so-called biological pump. For example, large colonial rhizarians of the order Collodaria have been shown to correlate significantly with downward fluxes of carbon (Guidi *et al.*, 2016). Additionally, the large-celled

members of Phaeodaria are reported to play a major role in carbon flux attenuation in the twilight zone in the California Current Ecosystem (Stukel *et al.*, 2018). Another rhizarian group, Acantharia, has been shown to contribute to the particulate organic carbon flux *via* cyst formation down to meso-/bathy-pelagic waters in the Southern Ocean and the Atlantic Ocean (Decelle *et al.*, 2013; Belcher *et al.*, 2018). Thus, the rhizarian-affiliated sequences retrieved from the deep sea were thought to derive from extracellular material of larger cells sinking from the upper waters (Not *et al.*, 2009). Other data, however, do not support this conclusion. For example, novel rhizarian OTUs/clades have been reported, both in the deep oxygenated meso-/bathy-pelagic waters and in anoxic basins (Not *et al.*, 2007a; Gilg *et al.*, 2010; Orsi *et al.*, 2011; Xu *et al.*, 2017a). Furthermore, RNA-based sequence data, which is not subject to the same distortions as DNA-based sequence data (*e.g.*, sequences generated from the inactive/dead cells and extracellular nucleic acids), have confirmed the presence of active rhizarians in deep oceanic waters (Hu *et al.*, 2016; Xu *et al.*, 2017b). Radiolarians, which have characteristic cell structures and photosynthetic symbionts, are classified as photosynthetic organisms in surface waters. However, little knowledge has been known for their lifestyle in the bathypelagic zone where photosynthetic symbionts are not expected (Stoecker *et al.*, 2009; Ishitani *et al.*, 2012). Thus the unknown rhizarian species dwelling in the deep waters needs to be further studied. When combined with paired, *i.e.*, RNA- and DNA-based, molecular sequencing approaches, these data will serve as a basis to gain a better understanding of the ecological role of rhizarians and other protists in the deep sea.

It has been suggested that the universal primers targeting the SSU rRNA gene favor some eukaryotic clades over others, leading to bias in the resulting molecular datasets (Amaral-Zettler *et al.*, 2009). Compared with the V4 region of the SSU rRNA gene, it has been suggested that the shorter V9 region might mitigate such biases (Pawlowski *et al.*, 2011). This finding appears to be especially true for excavates and foraminiferans (Pawlowski *et al.*, 2011). Recent studies have shown that diplomonads (Excavata, Diplonemea), a major component of Excavata, is the most diverse and the sixth most abundant eukaryotic taxon in the global plankton, with a huge undiscovered diversity in the deep ocean (de Vargas *et al.*, 2015; Flegontova *et al.*, 2016). In the present study, Excavata was found to be the third most abundant group in terms of relative sequence abundance, and the second most diverse group in terms of OTU richness in the deep waters of the South China Sea. Members in Diplonemea were the major contributors of Excavata which is consistent with previous reports (Figs. 6E, F). Members of the family Neobodonidae (Excavata, Kinetoplastida) were also retrieved from the deep waters of the South China Sea and constituted about 8% of all sequences affiliated with Excavata (Figs. 6E, F). In the deep dark ocean, protistan grazing is considered to be one of the major causes

of prokaryote mortality, arguably surpassing viral lysis (Nagata *et al.*, 2010). The deep sea protistan grazers are mostly small, nano-sized, flagellated cells. They are commonly referred to as heterotrophic nanoflagellates (HNFs), which include a wide range of phylogenetically diverse microbial eukaryotes. Members in Neobododid are considered to be important protistan bacterivores in the ocean (Chavez-Dozal *et al.*, 2013; Mukherjee *et al.*, 2015). Using catalyzed reporter deposition fluorescence *in situ* hybridization (CARD-FISH), it has been reported that kinetoplastids represented a significant fraction (average 21.8%) of total eukaryotic microbes in the deep subtropical North Atlantic (Morgan-Smith *et al.*, 2011, 2013). It has also been reported that neobodonids are dominant kinetoplastids in the global ocean (Flegontova *et al.*, 2018). Thus neobodonids are probably a major component of deep sea HNF communities and may play key roles in deep sea biogeochemical cycling through active grazing on prokaryotes.

4.2 Phototrophic Protists in the Bathy Zone of the South China Sea

In recent years, signals from deep sea phototrophic cells (both prokaryotes and eukaryotes) have been repeatedly reported in studies using various approaches including microscopy, flow cytometry and DNA/RNA-based sequencing (Scharek *et al.*, 1999; Jiao *et al.*, 2014; Pernice *et al.*, 2015; Xu *et al.*, 2017b). For instance, viable diatoms were retrieved from the global deep ocean collected by customized bottle-net sampler followed by vital stain and DNA-based sequencing (Agusti *et al.*, 2015). Using both SSU rRNA and p23S rRNA gene analysis, phototrophic signals were found in the Challenger Deep of the Mariana Trench (Guo *et al.*, 2018). By integrating epifluorescence microscopy observation and sequencing of SSU rRNA and psbA gene transcripts, live/active pigmented nano-sized eukaryotes (primarily haptophytes) were detected in the deep Western Pacific Ocean (Xu *et al.*, 2018). In the present study, the contribution of phototrophic signals (mainly from Bacillariophyta, Bolidophyceae, Dictyochophyceae, Prasinophyceae, and Prymnesiophyceae) to the total microbial eukaryote biota ranged from 0.3% to 16%, with an average value of 3% (Figs.6G, H). Previous studies have shown that phototrophic microbes make a significant contribution to the downward transportation of carbon via various fast-sinking mechanisms (Jiao *et al.*, 2014; Cai *et al.*, 2015; Pernice *et al.*, 2015). Based on DNA rather than RNA sequencing, the present study further confirmed the presence of phototrophic microbial eukaryotes in the deep sea. It has also been proposed that deep sea phototrophic signals are possibly from mixotrophs as many planktonic protists have a mixotrophic mode of nutrition (Stoecker *et al.*, 2017; Xu *et al.*, 2018). In the present study, members of Prymnesiophyceae, some of which are mixotrophic organisms, were recovered. It indicates the possible existence of heterotrophy of these cells that can make themselves adapt to and survive in the bathypelagic envi-

ronment.

Acknowledgements

We thank the captain and crew of the *R/V Dongfanghong-2* for successful sampling. This work was supported by the National Natural Science Foundation of China (Nos. 41876142, 91751207 and 41861144018) and the Senior User Project of *RV KEXUE* (No. KEXUE2019 G08) funded by the Center for Ocean Mega-Science, Chinese Academy of Sciences.

References

- Adl, S. M., Bass, D., Lane, C. E., Lukeš, J., Schoch, C. L., Smirnov, A., Agatha, S., Berney, C., Brown, M. W., Burki, F., Cárdenas, P., Čepička, I., Chistyakova, L., Del Campo, J., Dunthorn, M., Edvardsen, B., Eglit, Y., Guillou, L., Hampl, V., Heiss, A. A., Hoppenrath, M., James, T. Y., Karnkowska, A., Karpov, S., Kim, E., Kolisko, M., Kudryavtsev, A., Lahr, D. J. G., Lara, E., Le Gall, L., Lynn, D. H., Mann, D. G., Massana, R., Mitchell, E. A. D., Morrow, C., Park, J. S., Pawlowski, J. W., Powell, M. J., Richter, D. J., Rueckert, S., Shadwick, L., Shimano, S., Spiegel, F. W., Torruella, G., Youssef, N., Zlatogursky, V., and Zhang, Q., 2019. Revisions to the classification, nomenclature, and diversity of eukaryotes. *Journal of Eukaryotic Microbiology*, **66**: 4-119.
- Agusti, S., Gonzalez-Gordillo, J. I., Vaque, D., Estrada, M., Cerezo, M. I., Salazar, G., Gasol, J. M., and Duarte, C. M., 2015. Ubiquitous healthy diatoms in the deep sea confirm deep carbon injection by the biological pump. *Nature Communications*, **6**: 7608.
- Amaral-Zettler, L. A., McCliment, E. A., Ducklow, H. W., and Huse, S. M., 2009. A method for studying protistan diversity using massively parallel sequencing of V9 hypervariable regions of small-subunit ribosomal RNA genes. *PLoS One*, **4**: e6372.
- Aristegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J., 2009. Microbial oceanography of the dark ocean's pelagic realm. *Limnology and Oceanography*, **54**: 1501-1529.
- Azam, F., Fenchel, T., Field, J. G., Gray, J. S., Meyer-Rei, L. A., and Thingstad, F., 1983. The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series*, **10**: 257-263.
- Bass, D., Moreira, D., López-García, P., Polet, S., Chao, E. E., von der Heyden, S., Pawlowski, J., and Cavalier-Smith, T., 2005. Polyubiquitin insertions and the phylogeny of Cercozoa and Rhizaria. *Protist*, **156**: 149-161.
- Belcher, A., Manno, C., Thorpe, S., and Tarling, G., 2018. Acantharian cysts: High flux occurrence in the bathypelagic zone of the Scotia Sea, Southern Ocean. *Marine Biology*, **165**: 117.
- Biard, T., Stemann, L., Picheral, M., Mayot, N., Vandromme, P., Hauss, H., Gorsky, G., Guidi, L., Kiko, R., and Not, F., 2016. *In situ* imaging reveals the biomass of giant protists in the global ocean. *Nature*, **532**: 504-507.
- Bolger, A. M., Lohse, M., and Usadel, B., 2014. Trimmomatic: A flexible trimmer for Illumina Sequence Data. *Bioinformatics*, **30**: 2114-2120.
- Cai, P., Zhao, D., Wang, L., Huang, B., and Dai, M., 2015. Role of particle stock and phytoplankton community structure in regulating particulate organic carbon export in a large marginal sea. *Journal of Geophysical Research: Oceans*, **120**:

- 2063-2095.
- Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K., Fierer, N., Peña, A. G., Goodrich, J. K., Gordon, J. I., Huttley, G. A., Kelley, S. T., Knights, D., Koenig, J. E., Ley, R. E., Lozupone, C. A., McDonald, D., Muegge, B. D., Pirrung, M., Reeder, J., Sevinsky, J. R., Turnbaugh, P. J., Walters, W. A., Widmann, J., Yatsunenko, T., Zaneveld, J., and Knight, R., 2010. QIIME allows analysis of high-throughput community sequencing data. *Nature Methods*, **7**: 335-336.
- Caron, D. A., Alexander, H., Allen, A. E., Archibald, J. M., Armbrust, E. V., Bachy, C., Bharti, A., Bell, C. J., Dyhrman, S. T., Guida, S., Heidelberg, K. B., Kaye, J. Z., Metzner, J., Smith, S. R., and Worden, A. Z., 2017. Probing the evolution, ecology and physiology of marine protists using transcriptomics. *Nature Reviews Microbiology*, **15**: 6-20.
- Caron, D. A., Countway, P. D., Jones, A. C., Kim, D. Y., and Schnetzer, A., 2012. Marine protistan diversity. *Annual Review of Marine Science*, **4**: 467-493.
- Caron, D. A., Worden, A. Z., Countway, P. D., Demir, E., and Heidelberg, K. B., 2009. Protists are microbes too: A perspective. *The ISME Journal*, **3**: 4-12.
- Chavez-Dozal, A., Gorman, C., Erken, M., Steinberg, P. D., McDougald, D., and Nishiguchi, M. K., 2013. From the laboratory into the field: Testing defense mechanisms of bacterial biofilms against protozoan grazing. *Applied and Environmental Microbiology*, **79**: 553-558.
- Countway, P. D., Gast, R. J., Dennett, M. R., Savai, P., Rose, J. M., and Caron, D. A., 2007. Distinct protistan assemblages characterize the euphotic zone and deep sea (2500 m) of the western North Atlantic (Sargasso Sea and Gulf Stream). *Environmental Microbiology*, **9**: 1219-1232.
- Decelle, J., Martin, P., Paborstava, K., Pond, D. W., Tarling, G., Mahé, F., de Vargas, C., Lampitt, R., and Not, F., 2013. Diversity, ecology and biogeochemistry of cyst-forming Acantharia (Radiolaria) in the oceans. *PLoS One*, **8**: e53598.
- de Vargas, C., Audic, S., Henry, N., Decelle, J., Mahé, F., Logares, R., Lara, E., Berney, C., Le Bescot, N., Probert, I., Carmichael, M., Poulain, J., Romac, S., Colin, S., Aury, J. M., Bittner, L., Chaffron, S., Dunthorn, M., Engelen, S., Flegontova, O., Guidi, L., Horák, A., Jaillon, O., Lima-Mendez, G., Lukeš, J., Malviya, S., Morard, R., Mulot, M., Scalco, E., Siano, R., Vincent, F., Zingone, A., Dimier, C., Picheral, M., Searson, S., Kandels-Lewis, S., and Tara Oceans Coordinators, Acinas, S. G., Bork, P., Bowler, C., Gorsky, G., Grimsley, N., Hingamp, P., Iudicone, D., Not, F., Ogata, H., Pesant, S., Raes, J., Sieracki, M. E., Speich, S., Stemmann, L., Sunagawa, S., Weissenbach, J., Wincker, P., and Karsenti, E., 2015. Eukaryotic plankton diversity in the sunlit ocean. *Science*, **348**: 1261605.
- Edgar, R. C., 2013. UPARSE: Highly accurate OTU sequences from microbial amplicon reads. *Nature Methods*, **10**: 996-998.
- Edgar, R. C., Haas, B. J., Clemente, J. C., Quince, C., and Knight, R., 2011. UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics*, **27**: 2194-2200.
- Flegontova, O., Flegontov, P., Malviya, S., Audic, S., Wincker, P., de Vargas, C., Bowler, C., Lukeš, J., and Horák, A., 2016. Extreme diversity of diplomonid eukaryotes in the ocean. *Current Biology*, **26**: 3060-3065.
- Flegontova, O., Flegontov, P., Malviya, S., Poulain, J., de Vargas, C., Bowler, C., Lukeš, J., and Horák, A., 2018. Neobodonids are dominant kinetoplastids in the global ocean. *Environmental Microbiology*, **20**: 878-889.
- Gilg, I. C., Amaral-Zettler, L. A., Countway, P. D., Moorthi, S., Schnetzer, A., and Caron, D. A., 2010. Phylogenetic affiliations of mesopelagic Acantharia and acantharian-like environmental 18S rRNA genes off the southern California coast. *Protist*, **161**: 197-211.
- Giner, C. R., Balagué, V., Pernice, M. C., Duarte, C. M., Gasol, J. M., Logares, R., and Massana, R., 2019. Marked changes in diversity and relative activity of picoeukaryotes with depth in the global ocean. *BioRxiv*, DOI: <https://doi.org/10.1101/552604>.
- Guidi, L., Chaffron, S., Bittner, L., Eveillard, D., Larhlimi, A., Roux, S., Darzi, Y., Audic, S., Berline, L., Brum, J., Coelho, L. P., Espinoza, J. C. I., Malviya, S., Sunagawa, S., Dimier, C., Kandels-Lewis, S., Picheral, M., Poulain, J., Searson, S., Tara Oceans coordinators, Stemmann, L., Not, F., Hingamp, P., Speich, S., Follows, M., Karp-Boss, L., Boss, E., Ogata, H., Pesant, S., Weissenbach, J., Wincker, P., Acinas, S. G., Bork, P., de Vargas, C., Iudicone, D., Sullivan, M. B., Raes, J., Karsenti, E., Bowler, C., and Gorsky, G., 2016. Plankton networks driving carbon export in the oligotrophic ocean. *Nature*, **532**: 465-470.
- Guillou, L., Viprey, M., Chambouvet, A., Welsh, R. M., Kirkham, A. R., Massana, R., Scanlan, D. J., and Worden, A. Z., 2008. Widespread occurrence and genetic diversity of marine parasitoids belonging to Syndiniales (Alveolata). *Environmental Microbiology*, **10**: 3349-3365.
- Guo, R., Liang, Y., Xin, Y., Wang, L., Mou, S., Cao, C., Xie, R., Zhang, C., Tian, J., and Zhang, Y., 2018. Insight into the pico- and nano-phytoplankton communities in the deepest biosphere, the Mariana Trench. *Frontiers in Microbiology*, **9**: 2289.
- Herndl, G. J., and Reinthaler, T., 2013. Microbial control of the dark end of the biological pump. *Nature Geoscience*, **6**: 718-724.
- Hewson, I., Steele, J. A., Capone, D. G., and Fuhrman, J. A., 2006. Remarkable heterogeneity in meso- and bathypelagic bacterioplankton assemblage composition. *Limnology and Oceanography*, **51**: 1274-1283.
- Hu, S. K., Campbell, V., Connell, P., Gellene, A. G., Liu, Z., Terrado, R., and Caron, D. A., 2016. Protistan diversity and activity inferred from RNA and DNA at a coastal ocean site in the eastern North Pacific. *FEMS Microbiology Ecology*, **92**: ffw050.
- Hu, X., Lin, X., and Song, W., 2019. *Ciliates Atlas: Species Found in the South China Sea*. Science Press, Beijing, 932pp.
- Ishitani, Y., Ujiie, Y., de Vargas, C., Not, F., and Takahashi, K., 2012. Phylogenetic relationships and evolutionary patterns of the order Collodaria (Radiolaria). *PLoS One*, **7**: e35775.
- Jiao, N., Herndl, G. J., Hansell, D. A., Benner, R., Kattner, G., Wilhelm, S. W., Kirchman, D. L., Weinbauer, M. G., Luo, T. W., Chen, F., and Azam, F., 2010. Microbial production of recalcitrant dissolved organic matter: Long-term carbon storage in the global ocean. *Nature Reviews Microbiology*, **8**: 593-599.
- Jiao, N., Luo, T., Zhang, R., Yan, W., Lin, Y., Johnson, Z. I., Tian, J., Yuan, D., Yang, Q., Sun, J., Hu, D., and Wang, P., 2014. Presence of *Prochlorococcus* in the aphotic waters of the Western Pacific Ocean. *Biogeosciences*, **11**: 2391-2400.
- Li, R., Jiao, N., Warren, A., and Xu, D., 2018. Changes in community structure of active protistan assemblages from the lower Pearl River to coastal waters of the South China Sea. *European Journal of Protistology*, **63**: 72-82.
- Lozupone, C., and Knight, R., 2005. UniFrac: A new phylogenetic method for comparing microbial communities. *Applied and Environmental Microbiology*, **71**: 8228-8235.
- Magoč, T., and Salzberg, S. L., 2011. FLASH: Fast length ad-

- justment of short reads to improve genome assemblies. *Bioinformatics*, **27**: 2957-2963.
- Martín-Cuadrado, A. B., López-García, P., Alba, J. C., Moreira, D., Monticelli, L., Strittmatter, A., Gottschalk, G., and Rodríguez-Valera, F., 2007. Metagenomics of the deep Mediterranean, a warm bathypelagic habitat. *PLoS One*, **2**: e914.
- Moeseneder, M. M., Winter, C., and Herndl, G. J., 2001. Horizontal and vertical complexity of attached and free-living bacteria of the eastern Mediterranean Sea, determined by 16S rDNA and 16S rRNA fingerprints. *Limnology and Oceanography*, **46**: 95-107.
- Morgan-Smith, D., Herndl, G., van Aken, H., and Bochkansky, A., 2011. Abundance of eukaryotic microbes in the deep subtropical North Atlantic. *Aquatic Microbial Ecology*, **65**: 103-115.
- Morgan-Smith, D., Clouse, M. A., Herndl, G. J., and Bochkansky, A., 2013. Diversity and distribution of microbial eukaryotes in the deep tropical and subtropical North Atlantic Ocean. *Deep Sea Research Part I*, **78**: 58-69.
- Mukherjee, I., Hodoki, Y., and Nakano, S., 2015. Kinetoplastid flagellates overlooked by universal primers dominate in the oxygenated hypolimnion of Lake Biwa, Japan. *FEMS Microbiology Ecology*, **91**: fiv083.
- Nagata, T., Tamburini, C., Aristegui, J., Baltar, F., Bochkansky, A. B., Fonda-Umani, S., Fukuda, H., Gogou, A., Hansell, D. A., Hansman, R. L., Herndl, G. J., Panagiotopoulos, C., Reinthaler, T., Sohrin, R., Verdugo, P., Yamada, N., Yamashita, Y., Yokokawa, T., and Bartlett, D. H., 2010. Emerging concepts on microbial processes in the bathypelagic ocean – Ecology, biogeochemistry, and genomics. *Deep Sea Research Part II: Topical Studies in Oceanography*, **57**: 1519-1536.
- Not, F., del Campo, J., Balagué, V., de Vargas, C., and Massana, R., 2009. New insights into the diversity of marine picoeukaryotes. *PLoS One*, **4**: e7143.
- Not, F., Gausling, R., Azam, F., Heidelberg, J. F., and Worden, A. Z., 2007. Vertical distribution of picoeukaryotic diversity in the Sargasso Sea. *Environmental Microbiology*, **9**: 1233-1252.
- Orsi, W., Edgcomb, V., Jeon, S., Leslin, C., Bunge, J., Taylor, G. T., Varela, R., and Epstein, S., 2011. Protistan microbial observatory in the Cariaco Basin, Caribbean. II. Habitat specialization. *The ISME Journal*, **5**: 1357-1373.
- Pawlowski, J., Christen, R., Lecroq, B., Bachar, D., Shahbazkia, H. R., Amaral-Zettler, L., and Guillou, L., 2011. Eukaryotic richness in the abyss: Insights from pyrotag sequencing. *PLoS One*, **6**: e18169.
- Pernice, M. C., Giner, C. R., Logares, R., Perera-Bel, J., Acinas, S. G., Duarte, C. M., Gasol, J. M., and Massana, R., 2015. Large variability of bathypelagic microbial eukaryotic communities across the world's oceans. *The ISME Journal*, **10**: 945-958.
- Scharek, R., Tupas, L. M., and Karl, D. M., 1999. Diatom fluxes to the deep sea in the oligotrophic North Pacific gyre at station ALOHA. *Marine Ecology Progress Series*, **182**: 55-67.
- Schlitzer, R., 2011. Ocean data view. <http://odv.awi.de>.
- Song, W., Warren, A., and Hu, X., 2009. *Free-Living Ciliates in the Bohai and Yellow Seas, China*. Science Press, Beijing, 518pp.
- Stoecker, D. K., Johnson, M. D., deVargas, C., and Not, F., 2009. Acquired phototrophy in aquatic protists. *Aquatic Microbial Ecology*, **57**: 279-310.
- Stoecker, D. K., Hansen, P. J., Caron, D. A., and Mitra, A., 2017. Mixotrophy in the marine plankton. *Annual Review of Marine Science*, **9**: 311-335.
- Stukel, M. R., Biard, T., Krause, J., and Ohman, M. D., 2018. Large Phaeodaria in the twilight zone: Their role in the carbon cycle. *Limnology and Oceanography*, **63**: 2579-2594.
- Sun, P., Huang, L., Xu, D., Chen, N., and Warren, A., 2017. Marked seasonality and high spatial variation in estuarine ciliates are driven by exchanges between the 'abundant' and 'intermediate' biospheres. *Scientific Reports*, **7**: 9494.
- Xu, D., Jiao, N., Ren, R., and Warren, A., 2017a. Distribution and diversity of microbial eukaryotes in bathypelagic waters of the South China Sea. *Journal of Eukaryotic Microbiology*, **64**: 370-382.
- Xu, D., Li, R., Hu, C., Sun, P., Jiao, N., and Warren, A., 2017b. Microbial eukaryote diversity and activity in the water column of the South China Sea based on DNA and RNA high throughput sequencing. *Frontiers in Microbiology*, **8**: 1121.
- Xu, D., Sun, P., Zhang, Y., Li, R., Huang, B., Jiao, N., Warren, A., and Wang, L., 2018. Pigmented microbial eukaryotes fuel the deep sea carbon pool in the tropical Western Pacific Ocean. *Environmental Microbiology*, **20**: 3811-3824.

(Edited by Qiu Yantao)