

TWISTED PARTIAL ACTIONS OF HOPF ALGEBRAS *

BY

MARCELO MUNIZ S. ALVES

*Departamento de Matemática, Universidade Federal do Paraná
81531-980, Curitiba, PR, Brazil
e-mail: marcelomsa@ufpr.br*

AND

ELIEZER BATISTA

*Departamento de Matemática, Universidade Federal de Santa Catarina
88040-900, Florianópolis, SC, Brazil
e-mail: ebatista@mtm.ufsc.br*

AND

MICHAEL DOKUCHAEV

*Instituto de Matemática e Estatística, Universidade de São Paulo
05508-090, São Paulo, SP, Brazil
e-mail: dokucha@ime.usp.br*

AND

ANTONIO PAQUES

*Instituto de Matemática, Universidade Federal do Rio Grande do Sul
91509-900, Porto Alegre, RS, Brazil
e-mail: paques@mat.ufrgs.br*

* The first and second authors were partially supported by Fundação Araucária of Brazil, 490/16032. The third and fourth authors were partially supported by Fapesp of Brazil. The first and third authors were also partially supported by CNPq of Brazil.

Received December 19, 2011

ABSTRACT

In this work, the notion of a twisted partial Hopf action is introduced as a unified approach for twisted partial group actions, partial Hopf actions and twisted actions of Hopf algebras. The conditions on partial cocycles are established in order to construct partial crossed products, which are also related to partially cleft extensions of algebras. Examples are elaborated using algebraic groups.

1. Introduction

The desire to endow important classes of C^* -algebras generated by partial isometries with a structure of a more general crossed product led to the concept of a partial group action, introduced in [19], [27], [22], [23]. The new structure permitted to obtain relevant results on K -theory, ideal structure and representations of the algebras under consideration, as well as to treat amenability questions, especially amenability of C^* -algebraic bundles (also called Fell bundles), using both partial actions and the related concept of a partial representation. Amongst prominent classes of C^* -algebras endowed with the structure of non-trivial crossed products by partial actions one may list the Bunce–Deddens and the Bunce–Deddens–Toeplitz algebras [20], the approximately finite dimensional algebras [21], the Toeplitz algebras of quasi-ordered groups, as well as the Cuntz–Krieger algebras [26], [31].

The algebraic study of partial actions and partial representations was initiated in [23], [13] and [12], motivating investigations in diverse directions. In particular, the Galois theory of partial group actions developed in [16] inspired further Galois theoretic results in [9], as well as the introduction and study of partial Hopf actions and coactions in [10]. The latter paper became in turn the starting point for further investigation of partial Hopf (co)actions in [2], [3] and [4]. The Galois theoretic treatment in [9] was based on a coring \mathcal{C} constructed for an idempotent partial action of a finite group. The coring \mathcal{C} was shown to fit the general theory of cleft bicomodules in [7], and, in addition, in [8] descent theory for corings was applied, using \mathcal{C} , to define non-Abelian Galois cohomology ($i = 0, 1$) for idempotent partial Galois actions of finite groups.

The general notion of a (continuous) twisted partial action of a locally compact group on a C^* -algebra (a twisted partial C^* -dynamical system) and the

corresponding crossed products were given by R. Exel in [22]. The new construction permitted to show that any second countable C^* -algebraic bundle, which satisfies a certain regularity condition (automatically verified if the unit fiber algebra is stable), is a C^* -crossed product of the unit fiber algebra by a continuous partial action of the base group. The algebraic version of the latter fact was established in [14]. The importance of partial actions and partial representations was reinforced by R. Exel in [24] where, among other results, it was proved that given a field K of characteristic 0, a group G and subgroups $H, N \subseteq G$ with N normal in G and H normal in N , there is a twisted partial action θ of G/N on the group algebra $K(N/H)$ such that the Hecke algebra $\mathcal{H}(G, H)$ is isomorphic to the crossed product $K(N/H) *_{\theta} G/N$. More recent algebraic results on twisted partial actions and corresponding crossed products were obtained in [5], [15] and [30]. The algebraic concept of twisted partial actions also motivated the study of projective partial group representations, the corresponding partial Schur Multiplier and the relation to partial group actions with K -valued twistings in [17] and [18], contributing towards the elaboration of a background for a general cohomological theory based on partial actions. Further information around partial actions may be consulted in the survey [11].

The aim of this article is to introduce and study twisted partial Hopf actions on rings. The general definitions are given in Section 2, including that of a partial crossed product. The cocycle and normalization conditions are needed in order to make the partial crossed product be both associative and unital. As expected, restrictions of usual (global) twisted Hopf actions naturally result in twisted partial Hopf actions. Idempotent twisted partial actions of groups give natural examples of twisted partial actions of Hopf group algebras. Less evident examples may be obtained using algebraic groups, as shown in Section 3. Actions of an affine algebraic group on affine varieties give rise to coactions of the corresponding commutative Hopf algebra H on the coordinate algebras of the varieties, restrictions of which produce concrete examples of partial Hopf coactions. Then one may dualize in order to obtain partial Hopf actions. This works theoretically, but the elaboration of a concrete example needs some work. One possibility is to try to identify the finite dual H^0 for a specific H obtained

this way. A more flexible possibility is to find a concrete Hopf algebra H_1 such that H and H_1 form a dual pairing. Then Proposition 8 from [2] produces a partial action of H_1 . One still wishes to transform it into a twisted one, which in the setting specified in Section 3 is not difficult. A concrete example is elaborated in Proposition 5.

In order to treat the convolution invertibility of the partial cocycle in a manageable way, we introduce symmetric twisted partial Hopf actions in Section 4 and establish some useful technical formulas. Our definition is inspired by the case of twisted partial group actions. We also show that a restriction of a global twisted Hopf action with convolution invertible cocycle gives a symmetric twisted partial Hopf action. Theorem 1 relates isomorphisms of crossed products by symmetric twisted partial actions with a kind of “partial coboundaries,” establishing an analogue of a corresponding result known in the global case.

The last Section 5 is dedicated to the notion of partial cleft extensions and its relation with partial crossed products, in a quite similar fashion as is done in classical Hopf algebra theory. The definition of a partial cleft extension reflects the “partiality” in more than one way, incorporating, in particular, some equalities already proved to be significant in the study of partial group actions and partial representations (see Remark 2). Then the main result, Theorem 2, states that the partial cleft extensions over the coinvariants A are exactly the crossed products by symmetric twisted partial Hopf actions on A .

2. Twisted partial actions and partial crossed products

In this paper, except Section 3, κ will denote an arbitrary (associative) unital commutative ring and unadorned \otimes will stand for \otimes_κ , and also $\text{Hom}(V, W)$ will mean $\text{Hom}_\kappa(V, W)$ for any κ -modules V and W .

Definition 1: Let H be a Hopf κ -algebra, A a unital κ -algebra with unity element 1_A . Let furthermore $\alpha : H \otimes A \rightarrow A$ and $\omega : H \otimes H \rightarrow A$ be two κ -linear maps. We will write $\alpha(h \otimes a) := h \cdot a$, and $\omega(h \otimes l) := \omega(h, l)$, where $a \in A$ and $h, l \in H$.

The pair (α, ω) is called a **twisted partial action** of H on A if the following conditions hold:

- (1) $1_H \cdot a = a,$
- (2) $h \cdot (ab) = \sum (h_{(1)} \cdot a)(h_{(2)} \cdot b),$

$$(3) \quad \sum (h_{(1)} \cdot (l_{(1)} \cdot a))\omega(h_{(2)}, l_{(2)}) = \sum \omega(h_{(1)}, l_{(1)})(h_{(2)}l_{(2)} \cdot a),$$

$$(4) \quad \omega(h, l) = \sum \omega(h_{(1)}, l_{(1)})(h_{(2)}l_{(2)} \cdot \mathbf{1}_A),$$

for all $a, b \in A$ and $h, l \in H$.

If H, A and (α, ω) satisfy Definition 1, then we shall also say that (A, \cdot, ω) is a **twisted partial H -module algebra**.

PROPOSITION 1: *If (α, ω) is a twisted partial action, then the following identities hold:*

$$(5) \quad \omega(h, l) = \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))\omega(h_{(2)}, l_{(2)}) = \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, l).$$

Proof. The first identity is obtained from (3) by taking $a = 1$:

$$\sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))\omega(h_{(2)}, l_{(2)}) = \sum \omega(h_{(1)}, l_{(1)})(h_{(2)}l_{(2)} \cdot \mathbf{1}_A) \stackrel{(4)}{=} \omega(h, l).$$

For the second identity notice that

$$\begin{aligned} \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, l) &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot (l_{(1)} \cdot \mathbf{1}_A))\omega(h_{(3)}, l_{(2)}) \\ &= \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))\omega(h_{(2)}, l_{(2)}) = \omega(h, l) \end{aligned}$$

is obtained by using the first identity and (2). ■

We say that the map ω is **trivial**, if the following condition holds:

$$(6) \quad h \cdot (l \cdot \mathbf{1}_A) = \omega(h, l) = \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}l \cdot \mathbf{1}_A)$$

for all $h, l \in H$. In this case, the twisted partial action (α, ω) turns out to be a partial action of H on A , as introduced in [10]. Indeed, if (6) holds then the condition (4) is superfluous, and for all $h, l \in H$ and $a \in A$ we have

$$\begin{aligned} h \cdot (l \cdot a) &\stackrel{(2)}{=} \sum (h_{(1)} \cdot (l_{(1)} \cdot a))(h_{(2)} \cdot (l_{(2)} \cdot \mathbf{1}_A)) \\ &\stackrel{(6)}{=} \sum (h_{(1)} \cdot (l_{(1)} \cdot a))\omega(h_{(2)}, l_{(2)}) \stackrel{(3)}{=} \sum \omega(h_{(1)}, l_{(1)})(h_{(2)}l_{(2)} \cdot a) \\ &\stackrel{(6)}{=} \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}l_{(1)} \cdot \mathbf{1}_A)(h_{(3)}l_{(2)} \cdot a) \stackrel{(2)}{=} (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}l \cdot a). \end{aligned}$$

Observe also that if $h \cdot \mathbf{1}_A = \varepsilon(h)\mathbf{1}_A$ for all $h \in H$, then the condition (4) is a trivial consequence of the counit's properties and the κ -linearity of ω , and so we recover the classical notion of a twisted (global) action of H on A (see, for instance, [29]).

Example 1: This example is inspired by [10, Proposition 4.9] and suits¹ Definition 2.1 of [14]. An idempotent twisted partial action of a group G on a κ -algebra A is a triple

$$(\{D_g\}_{g \in G}, \{\alpha_g\}_{g \in G}, \{w_{g,h}\}_{(g,h) \in G \times G}),$$

where for each $g \in G$, D_g is an ideal of A generated by a central idempotent 1_g of A , $\alpha_g : D_{g^{-1}} \rightarrow D_g$ is an isomorphism of unital κ -algebras, and for each $(g, h) \in G \times G$, $w_{g,h}$ is an element of $D_g D_{gh}$, and the following statements are satisfied:

$$(7) \quad 1_e = \mathbf{1}_A \quad \text{and} \quad \alpha_e = I_A,$$

$$(8) \quad \alpha_g(\alpha_h(a1_{h^{-1}})1_{g^{-1}})w_{g,h} = w_{g,h}\alpha_{gh}(a1_{(gh)^{-1}}),$$

for all $a \in A$ and $g, h \in G$, where e denotes the identity element of G and I_A the identity map of A .

Let $\alpha : \kappa G \otimes A \rightarrow A$ and $\omega : \kappa G \otimes \kappa G \rightarrow A$ be the κ -linear maps given respectively by $\alpha(g \otimes a) = \alpha_g(a1_{g^{-1}})$ and $\omega(g, h) = w_{g,h}$, for all $a \in A$ and $g, h \in G$. This pair (α, ω) is a twisted partial action of κG on A . Indeed, conditions (2) and (4) are obvious since each α_g is multiplicative and each $w_{g,h}$ lives in $D_g D_{gh}$. Conditions (1) and (3) follow easily from (7) and (8), respectively. Notice in addition that $g \cdot \mathbf{1}_A = 1_g$ is central, for all $g \in G$.

Conversely, consider a twisted partial action (α, ω) of κG on A and set $\alpha(g \otimes a) = g \cdot a$ and $w_{g,h} = \omega(g, h)$, for all $g, h \in G$ and $a \in A$. By (2) we have that $1_g := g \cdot \mathbf{1}_A$ is an idempotent of A and by (4) and Proposition 1, $w_{g,h} \in (1_g A) \cap (A1_{gh})$, for all $g, h \in G$. Since by (1), $1_e = \mathbf{1}_A$, we have $w_{g,g^{-1}} \in 1_g A$ for all $g \in G$.

Now assume, in addition, that 1_g is central in A and $w_{g,g^{-1}}$ is invertible in $1_g A$, for all $g \in G$. Thus, $1_g A$ is a unital κ -algebra, $w_{g,h} \in (1_g A)(1_{gh} A)$,

$$g \cdot 1_{g^{-1}} = g \cdot (g^{-1} \cdot \mathbf{1}_A) \stackrel{(3)}{=} w_{g,g^{-1}} \mathbf{1}_A w_{g,g^{-1}}^{-1} = 1_g$$

and

$$g \cdot (1_{g^{-1}} a) \stackrel{(2)}{=} (g \cdot 1_{g^{-1}})(g \cdot a) = 1_g(g \cdot a) = (g \cdot \mathbf{1}_A)(g \cdot a) \stackrel{(2)}{=} g \cdot a,$$

for all $g, h \in G$ and $a \in A$.

¹ If one assumes in Definition 2.1 of [14] that each D_g is generated by a central idempotent, then the definition below is more general, as neither the invertibility in $D_g D_{gh}$ of each $w_{g,h}$ is required, nor the 2-cocycle equality.

Hence, α induces by restriction a map of κ -algebras $\alpha_g : 1_{g^{-1}}A \rightarrow 1_gA$, given by $\alpha_g(1_{g^{-1}}a) = g \cdot (1_{g^{-1}}a) = g \cdot a$, for all $g \in G$ and $a \in A$. Furthermore, it follows from (3) that $\alpha_g \circ \alpha_{g^{-1}}(1_ga) = \omega_{g,g^{-1}}(1_ga)\omega_{g,g^{-1}}^{-1}$, that is, $\alpha_g \circ \alpha_{g^{-1}}$ is an inner automorphism of 1_gA . In particular, $\alpha_{g^{-1}}$ is injective and α_g is surjective, for all $g \in G$. Consequently, α_g is an isomorphism and $1_gA = g \cdot A$, for all $g \in G$. Finally, (1) and (3), imply the above conditions (7) and (8), respectively. Therefore,

$$\{g \cdot A\}_{g \in G}, \{g \cdot \cdot\}_{g \in G}, \{\omega(g, h)\}_{(g, h) \in G \times G}$$

is a twisted partial action of G on A , as defined above. ■

Example 2 (Induced twisted partial action): Let B be a unital κ -algebra measured by an action $\beta : H \otimes B \rightarrow B$, denoted by $\beta(h, b) = h \triangleright b$, which is twisted by a map $u : H \otimes H \rightarrow B$, i.e.,

$$(9) \quad h \triangleright (ab) = \sum (h_{(1)} \triangleright a)(h_{(2)} \triangleright b),$$

$$(10) \quad h \triangleright 1_B = \varepsilon(h)1_B,$$

$$(11) \quad \sum (h_{(1)} \triangleright (k_{(1)} \triangleright a))u(h_{(2)}, k_{(2)}) = \sum u(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \triangleright a),$$

for all $h, k \in H$ and $a, b \in B$. Assume furthermore that

$$(12) \quad 1_H \triangleright a = a$$

for all $a \in A$. Here u is neither supposed to be convolution invertible, nor to satisfy the 2-cocycle equality.

Suppose that $\mathbf{1}_A$ is a non-trivial central idempotent of B , and let A be the ideal generated by $\mathbf{1}_A$. Given $a \in A, h \in H$, define a map $\cdot : H \otimes A \rightarrow A$ by

$$(13) \quad h \cdot a = \mathbf{1}_A(h \triangleright a).$$

It is clear that (12) implies (1), and (2) follows from the fact that $\mathbf{1}_A b = b \mathbf{1}_A$ for all $b \in B$. We still have to define a map $\omega : H \otimes H \rightarrow A$.

From equation (11) we obtain

$$\begin{aligned} \sum (h_{(1)} \cdot (k_{(1)} \cdot a))u(h_{(2)}, k_{(2)}) &= \sum \mathbf{1}_A(h_{(1)} \triangleright \mathbf{1}_A(k_{(1)} \triangleright a))u(h_{(2)}, k_{(2)}) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \triangleright (k_{(1)} \triangleright a))u(h_{(3)}, k_{(2)}) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \triangleright a) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot a), \end{aligned}$$

where the last equality follows from the fact that $\sum(h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k)$ lies in A . In particular, for $a = \mathbf{1}_A$ we obtain

$$(14) \quad \sum(h_{(1)} \cdot (k_{(1)} \cdot \mathbf{1}_A))u(h_{(2)}, k_{(2)}) = \sum(h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot \mathbf{1}_A),$$

which, in view of conditions (3) and (4), suggests we define ω by

$$(15) \quad \omega(h, k) = \sum(h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot \mathbf{1}_A).$$

With ω thus defined, (3) and (4) are clearly satisfied, and (A, \cdot, ω) is a twisted partial H -module algebra.

In particular, when B is an H -module algebra, i.e., when u is the trivial cocycle $u(h, k) = \varepsilon(h)\varepsilon(k)1_B$, it follows from (14) that ω is also trivial, i.e., ω satisfies (6). Therefore, in this case A becomes a partial H -module algebra as defined in [10]. ■

Given any two κ -linear maps $\alpha : H \otimes A \rightarrow A$, $h \otimes a \mapsto h \cdot a$, and $\omega : H \otimes H \rightarrow A$, we can define on the κ -module $A \otimes H$ a product, given by the multiplication

$$(a \otimes h)(b \otimes l) = \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, l_{(1)}) \otimes h_{(3)}l_{(2)},$$

for all $a, b \in A$ and $h, l \in H$. Write $A\#_{(\alpha, \omega)}H = (A \otimes H)(\mathbf{1}_A \otimes 1_H)$. It is readily seen that this corresponds to the κ -submodule of $A \otimes H$ generated by the elements of the form $a\#h := \sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)}$, for all $a \in A$ and $h \in H$.

In general $A \otimes H$, with this above-defined product, is neither associative nor unital. The following proposition gives necessary and sufficient conditions under which $A \otimes H$ (so also $A\#_{(\alpha, \omega)}H$) is associative and $A\#_{(\alpha, \omega)}H$ is unital with $\mathbf{1}_A\#1_H = \mathbf{1}_A \otimes 1_H$ as the identity element. This proposition is a generalization of [6, Lemmas 4.4 and 4.5] to the setting of twisted partial Hopf algebra actions.

PROPOSITION 2: *Let A be a unital κ -algebra, H a Hopf κ -algebra, $\omega : H \otimes H \rightarrow A$ and $\alpha : H \otimes A \rightarrow A$, $h \otimes a \mapsto h \cdot a$, two κ -linear maps satisfying the conditions (1), (2) and (4).*

(i) $\mathbf{1}_A\#1_H$ is the unity of $A\#_{(\alpha, \omega)}H$ if and only if, for all $h \in H$,

$$(16) \quad \omega(h, 1_H) = \omega(1_H, h) = h \cdot \mathbf{1}_A.$$

(ii) Suppose that $\omega(h, 1_H) = h \cdot \mathbf{1}_A$, for all $h \in H$. Then $A \otimes H$ is associative if and only if the condition (3) holds and, for all $h, l, m \in H$,

$$(17) \quad \sum(h_{(1)} \cdot \omega(l_{(1)}, m_{(1)}))\omega(h_{(2)}, l_{(2)}m_{(2)}) = \sum\omega(h_{(1)}, l_{(1)})\omega(h_{(2)}l_{(2)}, m).$$

Proof. The proof is similar to that of [6, Lemmas 4.4 and 4.5].

(i) Assume that $\omega(h, 1_H) = \omega(1_H, h) = h \cdot 1_A$. Then

$$\begin{aligned} (\mathbf{1}_A \# 1_H)(a \# h) &= (\mathbf{1}_A \otimes 1_H) \left(\sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} \right) \\ &= \sum \mathbf{1}_A(1_H \cdot (a(h_{(1)} \cdot \mathbf{1}_A)))\omega(1_H, h_{(2)}) \otimes 1_H h_{(3)} \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A) \otimes h_{(3)} \\ &\stackrel{(2)}{=} \sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} = a \# h \end{aligned}$$

and

$$\begin{aligned} (a \# h)(\mathbf{1}_A \# 1_H) &= \left(\sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} \right) (\mathbf{1}_A \otimes 1_H) \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A)\omega(h_{(3)}, 1_H) \otimes h_{(4)} 1_H \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A)(h_{(3)} \cdot \mathbf{1}_A) \otimes h_{(4)} \\ &\stackrel{(2)}{=} \sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} = a \# h \end{aligned}$$

for every $a \in A$ and $h \in H$.

Conversely, if $\mathbf{1}_A \# 1_H$ is the unity of $A \#_{(\alpha, \omega)} H$, then applying $I_A \otimes \varepsilon$ to the equalities

$$\sum (h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} = \mathbf{1}_A \# h = (\mathbf{1}_A \# 1_H)(\mathbf{1}_A \# h) = \sum \omega(1_H, h_{(1)}) \otimes h_{(2)}$$

and

$$\begin{aligned} \sum (h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} &= \mathbf{1}_A \# h = (\mathbf{1}_A \# h)(\mathbf{1}_A \# 1_H) \\ &= \left(\sum (h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} \right) (\mathbf{1}_A \otimes 1_H) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A)\omega(h_{(3)}, 1_H) \otimes h_{(4)} \\ &\stackrel{(2)}{=} \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, 1_H) \otimes h_{(3)} \\ &\stackrel{(4)}{=} \sum \omega(h_{(1)}, 1_H) \otimes h_{(2)}, \end{aligned}$$

we obtain

$$h \cdot \mathbf{1}_A = \sum (h_{(1)} \cdot \mathbf{1}_A)\varepsilon(h_{(2)}) = \sum \omega(1_H, h_{(1)})\varepsilon(h_{(2)}) = \omega(1_H, h),$$

$$h \cdot \mathbf{1}_A = \sum (h_{(1)} \cdot \mathbf{1}_A)\varepsilon(h_{(2)}) = \sum \omega(h_{(1)}, 1_H)\varepsilon(h_{(2)}) = \omega(h, 1_H).$$

(ii) Assume that (3) and (17) hold. Then, for all $a, b, c \in A$ and $h, l, m \in H$ we have

$$\begin{aligned}
 & (a \otimes h)[(b \otimes l)(c \otimes m)] \\
 &= \sum a[h_{(1)} \cdot (b(l_{(1)} \cdot c)\omega(l_{(2)}, m_{(1)}))] \omega(h_{(2)}, l_{(3)}m_{(2)}) \otimes h_{(3)}l_{(4)}m_{(3)} \\
 &\stackrel{(2)}{=} \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot (l_{(1)} \cdot c))(h_{(3)} \cdot \omega(l_{(2)}, m_{(1)})) \\
 &\quad \times \omega(h_{(4)}, l_{(3)}m_{(2)}) \otimes h_{(5)}l_{(4)}m_{(3)} \\
 &\stackrel{(17)}{=} \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot (l_{(1)} \cdot c))\omega(h_{(3)}, l_{(2)})\omega(h_{(4)}l_{(3)}, m_{(1)}) \otimes h_{(5)}l_{(4)}m_{(2)} \\
 &\stackrel{(3)}{=} \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, l_{(1)})(h_{(3)}l_{(2)} \cdot c)\omega(h_{(4)}l_{(3)}, m_{(1)}) \otimes h_{(5)}l_{(4)}m_{(2)} \\
 &= [(a \otimes h)(b \otimes l)](c \otimes m).
 \end{aligned}$$

Conversely, we have by assumption that

$$(\mathbf{1}_A \otimes h)[(\mathbf{1}_A \otimes l)(a \otimes \mathbf{1}_H)] = [(\mathbf{1}_A \otimes h)(\mathbf{1}_A \otimes l)](a \otimes \mathbf{1}_H)$$

and

$$(\mathbf{1}_A \otimes h)[(\mathbf{1}_A \otimes l)(\mathbf{1}_A \otimes m)] = [(\mathbf{1}_A \otimes h)(\mathbf{1}_A \otimes l)](\mathbf{1}_A \otimes m),$$

for all $a \in A$ and $h, l, m \in H$.

Using mainly condition (4) (and the hypothesis on ω in the first case only) one easily obtains, by a straightforward calculation, from the first equality

$$\sum (h_{(1)} \cdot (l_{(1)} \cdot a))\omega(h_{(2)}, l_{(2)}) \otimes h_{(3)}l_{(3)} = \sum \omega(h_{(1)}, l_{(1)})(h_{(2)}l_{(2)} \cdot a) \otimes h_{(3)}l_{(3)},$$

and from the second

$$\begin{aligned}
 & \sum (h_{(1)} \cdot \omega(l_{(1)}, m_{(1)}))\omega(h_{(2)}, l_{(2)}m_{(2)}) \otimes h_{(3)}l_{(3)}m_{(3)} \\
 &= \sum \omega(h_{(1)}, l_{(1)})\omega(h_{(2)}l_{(2)}, m_{(1)}) \otimes h_{(3)}l_{(3)}m_{(2)}.
 \end{aligned}$$

Now, applying $I_A \otimes \varepsilon$ to both sides of these equalities the conditions (3) and (17) follow, respectively. ■

Given a twisted partial action (α, ω) of a Hopf κ -algebra H on a κ -algebra A , the κ -algebra $A \#_{(\alpha, \omega)} H$ is called a **crossed product by a twisted partial action** (in brief, a **partial crossed product**) if the additional conditions (16) and (17) hold.

In order to establish some notation, we give the following lemma.

LEMMA 1: *In $A \#_{(\alpha, \omega)} H$ we have the following identities:*

- (i) $a\#h = \sum a(h_{(1)} \cdot \mathbf{1}_A)\#h_{(2)}$.
- (ii) $(a\#h)(b\#k) = \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)})\#h_{(3)}k_{(2)}$.

Proof. Item (i) is straightforward:

$$\begin{aligned} \sum a(h_{(1)} \cdot \mathbf{1}_A)\#h_{(2)} &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A) \otimes h_{(3)} \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} = a\#h. \end{aligned}$$

For item (ii), we have

$$\begin{aligned} (a\#h)(b\#k) &= \left(\sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} \right) \left(\sum b(k_{(1)} \cdot \mathbf{1}_A) \otimes k_{(2)} \right) \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot (b(k_{(1)} \cdot \mathbf{1}_A)))\omega(h_{(3)}, k_{(2)}) \otimes h_{(4)}k_{(3)} \\ &= \sum a(h_{(1)} \cdot (b(k_{(1)} \cdot \mathbf{1}_A)))\omega(h_{(2)}, k_{(2)}) \otimes h_{(3)}k_{(3)} \\ &= \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot (k_{(1)} \cdot \mathbf{1}_A))\omega(h_{(3)}, k_{(2)}) \otimes h_{(4)}k_{(3)} \\ &\stackrel{(5)}{=} \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)}) \otimes h_{(3)}k_{(2)} \\ &\stackrel{(4)}{=} \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot \mathbf{1}_A) \otimes h_{(4)}k_{(3)} \\ &= \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)})\#h_{(3)}k_{(2)}. \quad \blacksquare \end{aligned}$$

If, in particular, ω is trivial, then the multiplication in $A\#_{(\alpha, \omega)}H$ becomes

$$\begin{aligned} (a\#h)(b\#l) &= \left(\sum a(h_{(1)} \cdot \mathbf{1}_A) \otimes h_{(2)} \right) \left(\sum b(l_{(1)} \cdot \mathbf{1}_A) \otimes l_{(2)} \right) \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot (b(l_{(1)} \cdot \mathbf{1}_A)))\omega(h_{(3)}, l_{(2)}) \otimes h_{(4)}l_{(3)} \\ &\stackrel{(6)}{=} \sum a(h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot (b(l_{(1)} \cdot \mathbf{1}_A)))(h_{(3)} \cdot (l_{(2)} \cdot \mathbf{1}_A)) \otimes h_{(4)}l_{(3)} \\ &\stackrel{(2)}{=} \sum a(h_{(1)} \cdot (b(l_{(1)} \cdot \mathbf{1}_A))) \otimes h_{(2)}l_{(2)} \end{aligned}$$

for all $a, b \in A$ and $h, l \in H$. Consequently, in this case we recover the partial smash product introduced in [10].

Remark 1: If (α, ω) is a twisted partial action of κG on a κ -algebra A arisen from a twisted partial action of a group G on A , as defined in Example 1, and the conditions (16) and (17) also hold in this case, then $A\#_{(\alpha, \omega)}\kappa G$ is a slight generalization of the partial crossed product introduced in [14].

Example 3: Consider an induced partial twisted H -module structure as given in Example 2, and suppose that the map $u : H \rightarrow A$ is a **normalized cocycle**,

i.e., assume that

$$(18) \quad \sum (h_{(1)} \triangleright u(k_{(1)}, l_{(1)}))u(h_{(2)}, k_{(2)}l_{(2)}) = u(h_{(1)}, k_{(1)})u(h_{(2)}k_{(2)}, l),$$

$$(19) \quad u(h, 1_H) = u(1_H, h) = \varepsilon(h)1_B,$$

for all $h, k, l \in H$. It is clear that the induced map ω (see equality (15)) satisfies condition (16), and we will show that (17) is also satisfied. In what follows, note that $\sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot x) = \sum (h_{(1)} \cdot x)(h_{(2)} \cdot \mathbf{1}_A)$, and that if $a, b \in A$, then $a(h \cdot b) = a\mathbf{1}_A(h \triangleright b) = a(h \triangleright b)$. Using equality (18),

$$\begin{aligned} & \sum (h_{(1)} \cdot \omega(l_{(1)}, m_{(1)}))\omega(h_{(2)}, l_{(2)}m_{(2)}) = \\ & = \sum (h_{(1)} \cdot [(l_{(1)} \cdot \mathbf{1}_A)u(l_{(2)}, m_{(1)})(l_{(3)}m_{(2)} \cdot \mathbf{1}_A)](h_{(2)} \cdot \mathbf{1}_A) \\ & \quad \times u(h_{(3)}, l_{(4)}m_{(3)})(h_{(4)}l_{(5)}m_{(4)} \cdot \mathbf{1}_A)) \\ & = \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))((h_{(2)} \cdot u(l_{(2)}, m_{(1)}))(h_{(3)} \cdot (l_{(3)}m_{(2)} \cdot \mathbf{1}_A)) \\ & \quad \times (h_{(4)} \cdot \mathbf{1}_A)u(h_{(5)}, l_{(4)}m_{(3)})(h_{(6)}l_{(5)}m_{(4)} \cdot \mathbf{1}_A)) \\ & = \sum \underbrace{(h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))(h_{(2)} \cdot \mathbf{1}_A)}_{(14)}((h_{(3)} \cdot u(l_{(2)}, m_{(1)})) \\ & \quad \times (h_{(4)} \cdot (l_{(3)}m_{(2)} \cdot \mathbf{1}_A))u(h_{(5)}, l_{(4)}m_{(3)})(h_{(6)}l_{(5)}m_{(4)} \cdot \mathbf{1}_A)) \\ & = \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))(h_{(2)} \cdot u(l_{(2)}, m_{(1)})) \times \\ & \quad \times \underbrace{(h_{(3)} \cdot (l_{(3)}m_{(2)} \cdot \mathbf{1}_A))u(h_{(4)}, l_{(4)}m_{(3)})(h_{(5)}l_{(5)}m_{(4)} \cdot \mathbf{1}_A)}_{(14)} \\ & \stackrel{(14)}{=} \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))(h_{(2)} \cdot u(l_{(2)}, m_{(1)}))(h_{(3)} \cdot \mathbf{1}_A)u(h_{(4)}, l_{(3)}m_{(2)}) \\ & \quad \times (h_{(5)}l_{(4)}m_{(3)} \cdot \mathbf{1}_A)) \\ & = \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))(h_{(2)} \cdot u(l_{(2)}, m_{(1)}))u(h_{(3)}, l_{(3)}m_{(2)})(h_{(4)}l_{(4)}m_{(3)} \cdot \mathbf{1}_A) \\ & = \sum (h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A)) \underbrace{(h_{(2)} \triangleright u(l_{(3)}, m_{(1)}))u(h_{(3)}, l_{(4)}m_{(2)})(h_{(4)}l_{(5)}m_{(4)} \cdot \mathbf{1}_A)}_{(18)} \\ & \stackrel{(18)}{=} \sum \underbrace{(h_{(1)} \cdot (l_{(1)} \cdot \mathbf{1}_A))u(h_{(2)}, l_{(2)})}_{(14)}u(h_{(3)}l_{(3)}, m_{(1)})(h_{(4)}l_{(4)}m_{(2)} \cdot \mathbf{1}_A) \\ & \stackrel{(14)}{=} \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, l_{(1)})(h_{(3)}l_{(2)} \cdot \mathbf{1}_A)u(h_{(4)}l_{(3)}, m_{(1)})(h_{(5)}l_{(4)}m_{(2)} \cdot \mathbf{1}_A) \\ & = \sum \omega(h_{(1)}, l_{(1)})\omega(h_{(2)}, l_{(2)}m). \end{aligned}$$

If one forms the usual crossed product $B\#_uH$, then it is easy to see that $\mathbf{1}_A\#\mathbf{1}_H$ is an idempotent of this algebra, that

$$(\mathbf{1}_A\#\mathbf{1}_H)(B\#_uH) = A \otimes H$$

and that

$$(\mathbf{1}_A \# \mathbf{1}_H)(B \#_u H)(\mathbf{1}_A \# \mathbf{1}_H) = A \#_{(\alpha, \omega)} H.$$

3. Examples of (twisted) partial actions via algebraic groups

In this section we use the relation between algebraic groups and commutative Hopf algebras (see [1], [32]) to explain a way of producing examples of partial Hopf (co)actions. A concrete example is elaborated to which we attach a twisting resulting in a twisted partial Hopf action.

We shall extract our example from central notions of the theory of algebraic groups such as maximal tori, Cartan subgroups and the Weyl group. Let κ be an algebraically closed field and let \mathbf{G} be a linear algebraic group over κ , by which we mean a subgroup of $\text{GL}_n(\kappa)$ (for some positive integer n), which is closed in the Zariski topology of $\text{GL}_n(\kappa)$. Let T be a maximal torus in G . We recall that a torus is a connected diagonalizable linear algebraic group. Let C be the centralizer of T in \mathbf{G} and N be the normalizer of T in \mathbf{G} . Then C is the Cartan subgroup of \mathbf{G} and $W = N/C$ is the corresponding Weyl group (which is finite). Then evidently N acts on itself by left multiplication and this permutes the left cosets of N by C . Then taking the Hopf algebra H which corresponds to N , its comultiplication $\Delta : H \rightarrow H \otimes H$ is the right coaction of H on itself, which corresponds to the left action of N on itself. Let $Y \subsetneq N$ be a union of some left cosets of N by C . Then N acts (globally) on N and only partially on Y . Then one may take a two-sided ideal A in H determined by Y (see the concrete example below) so that one comes to a partial coaction $\rho : A \rightarrow A \otimes H$ which is obtained by the restriction $\rho = (1_A \otimes 1_H)\Delta$. Then taking a Hopf algebra H_1 such that there exists a pairing $\langle \cdot, \cdot \rangle_{H_1 \otimes H} : H_1 \otimes H \rightarrow \kappa$, one can dualize to obtain a partial action of H_1 on A , as given in [2, Prop. 8]. In particular, H_1 can be the finite dual of H .

For a concrete example we take one of the most classical cases, in which $\mathbf{G} = \text{GL}_n(\kappa)$ and $T \subseteq \text{GL}_n(\kappa)$ is the group of all diagonal matrices of $\text{GL}_n(k)$. Then $T \cong (\kappa^*)^n$, where κ^* is the multiplicative group of the field κ . It is directly verified that in this case $C = T$ and N is formed by the monomial matrices, that is, the matrices whose rows and columns have only one nonvanishing entry. The Weyl group W can be identified with the group of $n \times n$ permutation matrices, which is isomorphic to the symmetric group S_n .

The group N is an algebraic group and is isomorphic to the semidirect product of T by the action of the Weyl group

$$N \cong T \rtimes W = T \rtimes S_n.$$

Here, the left action of S_n on T is given by conjugation, whose net effect is the permutation of the diagonal matrix entries. By the fact that all these groups are algebraic groups, one can associate to the action

$$\begin{aligned} \alpha : S_n \times T &\rightarrow T, \\ (g, x) &\mapsto g \cdot x = gxg^{-1}, \end{aligned}$$

a left coaction of the corresponding Hopf algebras. It is a basic fact that the Hopf algebra which corresponds to a finite group is the dual of the group algebra, i.e. in our case it is $(\kappa S_n)^*$. It is also basic that the algebra corresponding to κ^* is the Hopf algebra of the Laurent polynomials $\kappa[t, t^{-1}]$. Since tensor products of Hopf algebras correspond to direct product of algebraic groups, it follows that the Hopf algebra corresponding to T is $\kappa[t, t^{-1}]^{\otimes n}$. Consequently there is a left coaction of $(\kappa S_n)^*$ on $\kappa[t, t^{-1}]^{\otimes n}$, which corresponds to the above action of S_n on T , i.e. $\kappa[t, t^{-1}]^{\otimes n}$ turns out to be a left $(\kappa S_n)^*$ -comodule coalgebra. Since $N \cong T \rtimes S_n$, it follows by [1, p. 143, p. 208] that the Hopf algebra associated to the group N is the co-semidirect product

$$\kappa[t, t^{-1}]^{\otimes n} \blacktriangleleft (\kappa S_n)^*.$$

A typical element of $\kappa[t, t^{-1}]^{\otimes n}$ is a tensor polynomial of the form

$$\sum_{N \in \mathbb{Z}} \sum_{k_1 + \dots + k_n = N} \lambda_N t^{k_1} \otimes \dots \otimes t^{k_n}.$$

In order to simplify the notation, write

$$(20) \quad t_i = 1 \otimes \dots \otimes 1 \otimes t \otimes 1 \otimes \dots \otimes 1,$$

where t belongs to the i -copy of $\kappa[t, t^{-1}]$. Then we have

$$t_1^{k_1} \dots t_n^{k_n} = t^{k_1} \otimes \dots \otimes t^{k_n}.$$

Since S_n operates on T by permuting the entries, it follows by a direct verification that the left $(\kappa S_n)^*$ -coaction on $\kappa[t, t^{-1}]^{\otimes n}$ is given by

$$\delta(t_1^{k_1} \dots t_n^{k_n}) = \sum_{g \in S_n} p_g \otimes t_{g^{-1}(1)}^{k_1} \dots t_{g^{-1}(n)}^{k_n}.$$

With this coaction, $\kappa[t, t^{-1}]^{\otimes n}$ is a left $(\kappa S_n)^*$ -comodule coalgebra, and the comultiplication of the cosemidirect product

$$H = \kappa[t, t^{-1}]^{\otimes n} \rtimes (\kappa S_n)^*$$

is given explicitly by

$$\begin{aligned} \Delta(t_1^{k_1} \cdots t_n^{k_n} \otimes p_g) &= \sum_{s, f \in S_n} t_1^{k_1} \cdots t_n^{k_n} \otimes p_s p_f \otimes t_{s^{-1}(1)}^{k_1} \cdots t_{s^{-1}(n)}^{k_n} \otimes p_{f^{-1}g} \\ &= \sum_{s \in S_n} t_1^{k_1} \cdots t_n^{k_n} \otimes p_s \otimes t_{s^{-1}(1)}^{k_1} \cdots t_{s^{-1}(n)}^{k_n} \otimes p_{s^{-1}g}. \end{aligned}$$

The cosemidirect product coacts on the right on itself by the comultiplication. In order to construct a partial coaction one can simply project over a two-sided ideal.

Let X be a subset of S_n which is not a subgroup. Write $L = \kappa[t, t^{-1}]^{\otimes n}$. Then evidently

$$(21) \quad e_X = 1_L \otimes \left(\sum_{g \in X} p_g \right)$$

is a central idempotent in $H = L \rtimes (\kappa S_n)^*$, and the algebra $A = e_X H$ is a two-sided ideal. Write $e_X \cdot$ for the map $H \rightarrow A$ given by multiplication by e_X . Then the restriction $\rho : A \rightarrow A \otimes H$, $\rho = (e_X \cdot \otimes I) \circ \Delta$, of $\Delta_H : H \rightarrow H \otimes H$ is a right partial coaction of H given by

$$(22) \quad \rho(t_1^{k_1} \cdots t_n^{k_n} \otimes p_g) = \sum_{s \in X} t_1^{k_1} \cdots t_n^{k_n} \otimes p_s \otimes t_{s^{-1}(1)}^{k_1} \cdots t_{s^{-1}(n)}^{k_n} \otimes p_{s^{-1}g},$$

where $g \in X$. Since $X \subseteq S_n$ is not a subgroup, it is readily seen that ρ is not a (global) coaction (if X was a subgroup then one would have $\rho : A \rightarrow A \otimes A$).

We shall obtain a partial action from a partial coaction using Proposition 8 from [2], which we recall for the reader's convenience:

PROPOSITION 3: *Let H_1 and H_2 be two Hopf algebras with a pairing between them:*

$$\begin{aligned} \langle \cdot, \cdot \rangle : H_1 \otimes H_2 &\rightarrow \kappa \\ h \otimes p &\mapsto \langle h, p \rangle. \end{aligned}$$

Then a partial right H_2 -comodule algebra A , acquires a structure of partial left H_1 -module algebra by the partial action

$$h \cdot a = \sum a^{[0]} \langle h, a^{[1]} \rangle,$$

where $\rho(a) = \sum a^{[0]} \otimes a^{[1]}$ is the partial right coaction of H_2 on A .

Assume now that κ is an isomorphic copy of the complex numbers \mathbb{C} , and let \mathbb{S}^1 be the unit circle group. The elements of \mathbb{S}^1 can be viewed as the complex roots of 1; however, we assume that κ and $\mathbb{S}^1 \subseteq \mathbb{C}$ are disjoint and consider the group algebra $\kappa\mathbb{S}^1$ of \mathbb{S}^1 over κ , so that the roots of unity $\chi \in \mathbb{S}^1$ are linearly independent over κ . Then S_n acts on $(\kappa\mathbb{S}^1)^{\otimes n}$ by permutation of roots, which gives an action of the group Hopf algebra κS_n on $(\kappa\mathbb{S}^1)^{\otimes n}$. Note that κS_n and $(\kappa\mathbb{S}^1)^{\otimes n}$ are both cocommutative. Then we may consider the smash product Hopf algebra

$$H_1 = (\kappa\mathbb{S}^1)^{\otimes n} \rtimes \kappa S_n.$$

Write

$$(23) \quad \chi_{\theta_1, \dots, \theta_n} = \chi_{\theta_1} \otimes \dots \otimes \chi_{\theta_n} \in (\kappa\mathbb{S}^1)^{\otimes n},$$

where $\chi_{\theta_i} \in \mathbb{S}^1$ is the root of 1 whose angular coordinate is θ_i and which belongs to the i -factor of $(\kappa\mathbb{S}^1)^{\otimes n}$. Then evidently the elements $\chi_{\theta_1, \dots, \theta_n} \otimes u_g$ ($g \in S_n$) form a κ -basis of H_1 . With this notation define the map $\langle, \rangle : H_1 \otimes H \rightarrow \kappa$, by setting

$$\langle \chi_{\theta_1, \dots, \theta_n} \otimes u_g, t_1^{k_1} \dots t_n^{k_n} \otimes p_s \rangle = \delta_{g,s} \exp\{ik_1\theta_1\} \dots \exp\{ik_n\theta_n\},$$

where $\delta_{g,s}$ is the Kronecker delta and $i^2 = -1$. It is an easy straightforward verification that this defines a pairing of Hopf algebras. Observe that it is non-degenerate, however we do not need to use this property.

Now using the coaction $\rho : A \rightarrow A \times H$ we obtain by Proposition 3 a partial action $H_1 \times A \rightarrow A$. To specify it, take $h = \chi_{\theta_1, \dots, \theta_n} \otimes u_g \in H_1$ and $a = t_1^{k_1} \dots t_n^{k_n} \otimes p_s \in A$ and check by the formula in Proposition 3 that the partial action is explicitly given by

$$\begin{aligned} h \cdot a &= \sum_{f \in X} t_1^{k_1} \dots t_n^{k_n} \otimes p_f \langle \chi_{\theta_1, \dots, \theta_n} \otimes u_g, t_{f^{-1}(1)}^{k_1} \dots t_{f^{-1}(n)}^{k_n} \otimes p_{f^{-1}s} \rangle \\ &= \sum_{f \in X} t_1^{k_1} \dots t_n^{k_n} \otimes p_f \{ \delta_{g, f^{-1}s} \exp\{ik_1\theta_{f^{-1}(1)}\} \dots \exp\{ik_n\theta_{f^{-1}(n)}\} \} \\ &= \exp\{ik_1\theta_{g^{-1}s(1)}\} \dots \exp\{ik_n\theta_{g^{-1}s(n)}\} t_1^{k_1} \dots t_n^{k_n} \otimes p_{s^{-1}g}, \end{aligned}$$

where $g \in S_n$ and $x \in X$.

Since any finite group G can be seen as a subgroup of S_n for some n , we may replace in the above considerations S_n by an arbitrary finite group G as follows. Fix a monomorphism $G \rightarrow S_n$ so that G will be considered as a subgroup of

S_n . Then the formula

$$\delta(t_1^{k_1} \cdots t_n^{k_n}) = \sum_{g \in G} p_g \otimes t_{g^{-1}(1)}^{k_1} \cdots t_{g^{-1}(n)}^{k_n}$$

gives a structure of a left $(\kappa G)^*$ -comodule coalgebra on $L = \kappa[t, t^{-1}]^{\otimes n}$, and one can take the cosemidirect product

$$H_2 = L \rtimes (\kappa G)^*$$

with comultiplication given by

$$\Delta(t_1^{k_1} \cdots t_n^{k_n} \otimes p_g) = \sum_{s \in G} t_1^{k_1} \cdots t_n^{k_n} \otimes p_s \otimes t_{s^{-1}(1)}^{k_1} \cdots t_{s^{-1}(n)}^{k_n} \otimes p_{s^{-1}g} \quad (g \in G).$$

Clearly, $H_2 = eH$, where $e = 1_L \otimes (\sum_{g \in G} p_g)$.

Let now X be an arbitrary subset of G which is not a subgroup. The element e_X defined by the formula (21) is obviously a central idempotent in H_2 and the algebra $A' = e_X H_2$ is a two-sided ideal. The restriction $\rho' : A' \rightarrow A' \otimes H$, $\rho' = e_X \Delta_{H_2}$ of $\Delta_{H_2} : H_2 \rightarrow H_2 \otimes H_2$ is a right partial coaction of H_2 given by exactly the same formula (22) which was used for ρ . The elements of $G \subseteq S_n$ act on $(\kappa \mathbb{S}^1)^{\otimes n}$, as above, by permutation of roots, and we have the smash product

$$H'_1 = (\kappa \mathbb{S}^1)^{\otimes n} \rtimes \kappa G.$$

Then the formula above which defined the left partial action of H_1 on A gives a left partial action $H'_1 \times A' \rightarrow A'$:

$$(24) \quad (\chi_{\theta_1, \dots, \theta_n} \otimes u_g) \cdot (t_1^{k_1} \cdots t_n^{k_n} \otimes p_s) = \exp\{i(k_1 \theta_{g^{-1}s(1)} + \cdots + k_n \theta_{g^{-1}s(n)})\} t_1^{k_1} \cdots t_n^{k_n} \otimes p_{s^{-1}g},$$

where $g \in G$ and $s \in X \subseteq G$.

In order to turn the partial action (24) into a twisted one, take a finite group G whose Schur Multiplier over $\kappa = \mathbb{C}$ is not trivial. Then there exists a 2-cocycle $\gamma : G \times G \rightarrow \mathbb{C}^*$ which is not a coboundary. The 2-cocycle equality means that

$$(25) \quad \gamma(x, y)\gamma(xy, z) = \gamma(x, yz)\gamma(y, z) \quad \forall x, y, z \in G.$$

Assume also that γ is normalized, i.e.

$$(26) \quad \gamma(g, 1) = \gamma(1, g) = 1.$$

For arbitrary $h = \chi_{\theta_1, \dots, \theta_n} \otimes u_g$ and $l = \chi_{\theta'_1, \dots, \theta'_n} \otimes u_s$ in H'_1 set

$$(27) \quad \omega(h, l) = \gamma(g, s) (h \cdot l \cdot \mathbf{1}_{A'}).$$

The fact that (24) and (27) define a twisted partial action of $H'_1 = (\kappa\mathbb{S}^1)^{\otimes n} \rtimes \kappa G$ on A' , which satisfies (16) and (17), will follow from the next easy:

PROPOSITION 4: *Let G be a finite group and L be a cocommutative Hopf algebra over a field κ , such that L is a left κG -module algebra. Suppose that there is a left partial action of the smash product $H = L \rtimes \kappa G$ on a κ -algebra A :*

$$H \otimes A \ni h \otimes a \mapsto h \cdot a \in A.$$

If $\gamma : G \times G \rightarrow \kappa^*$ is a normalized 2-cocycle, then the map

$$(28) \quad \omega(h, m) = \sum \gamma(g, s) (h \cdot m \cdot 1_A),$$

where $h = \sum l \otimes g, m = \sum l' \otimes s \in L \rtimes \kappa G$, turns the partial action $H \otimes A \rightarrow A$ into a twisted one such that (16) and (17) are satisfied.

Proof. One needs to check (3), (4), (16) and (17). It is obviously enough to verify these properties for the elements $h, m, k \in H$ of the form $h = l \otimes g, m = l' \otimes s$, and $k = l'' \otimes f$, with $l, l', l'' \in L, g, s, f \in G$. Recall from [1, p. 142] that

$$\Delta_H(l \otimes g) = \sum (l_{(1)} \otimes g) \otimes (l_{(2)} \otimes g).$$

Then using (26) and (25) it is readily seen that the properties (3), (4), (16) and (17) correspond respectively to the following equalities:

$$\begin{aligned} \sum (h_{(1)} \cdot m_{(1)} \cdot a)(h_{(2)} \cdot m_{(2)} \cdot \mathbf{1}_A) &= \sum (h_{(1)} \cdot m_{(1)} \cdot \mathbf{1}_A)(h_{(2)} m_{(2)} \cdot a), \\ h \cdot m \cdot \mathbf{1}_A &= \sum (h_{(1)} \cdot m_{(1)} \cdot \mathbf{1}_A)(h_{(2)} m_{(2)} \cdot \mathbf{1}_A), \\ h \cdot 1_H \cdot \mathbf{1}_A &= 1_H \cdot h \cdot \mathbf{1}_A = h \cdot \mathbf{1}_A \end{aligned}$$

and

$$\begin{aligned} \sum (h_{(1)} \cdot m_{(1)} \cdot k_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot m_{(2)} k_{(2)} \cdot \mathbf{1}_A) \\ = \sum (h_{(1)} \cdot m_{(1)} \cdot \mathbf{1}_A)(h_{(2)} m_{(2)} \cdot k_{(2)} \cdot \mathbf{1}_A), \end{aligned}$$

for all $h, k, m \in H$ and $a \in A$.

The first three equalities are immediate consequences of the definition of a (non-twisted) partial action. With respect to the last one, we have

$$\begin{aligned}
 & \sum (h_{(1)} \cdot m_{(1)} \cdot k_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot m_{(2)}k_{(2)} \cdot \mathbf{1}_A) \\
 &= \sum h \cdot [(m_{(1)} \cdot k_{(1)} \cdot \mathbf{1}_A)(m_{(2)}k_{(2)} \cdot \mathbf{1}_A)] \\
 &= \sum h \cdot [(m_{(1)} \cdot \mathbf{1}_A)(m_{(2)}k_{(1)} \cdot \mathbf{1}_A)(m_{(3)}k_{(2)} \cdot \mathbf{1}_A)] \\
 &= \sum h \cdot [(m_{(1)} \cdot \mathbf{1}_A)(m_{(2)}k_{(1)} \cdot \mathbf{1}_A)] \\
 &= h \cdot (m \cdot (k \cdot \mathbf{1}_A)) \\
 &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}m \cdot k \cdot \mathbf{1}_A) \\
 &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}m_{(1)} \cdot \mathbf{1}_A)(h_{(3)}m_{(2)} \cdot k \cdot \mathbf{1}_A) \\
 &= \sum (h_{(1)} \cdot m_{(1)} \cdot \mathbf{1}_A)(h_{(2)}m_{(2)} \cdot k_{(2)} \cdot \mathbf{1}_A),
 \end{aligned}$$

which completes the proof. ■

Note that if in the proposition above we do not assume (25) and (26), i.e., we take an arbitrary map $\gamma : G \times G \rightarrow \kappa^*$, then we obtain a twisted partial action which in general does not satisfy (16) and (17).

The above example can be made more specific by taking a concrete group G . The smallest finite group with non-trivial Schur Multiplier is the Klein-four group $G = \langle a \rangle \times \langle b \rangle$, $a^2 = b^2 = 1$. In this case the Schur Multiplier $M(G)$ has order 2, and a 2-cocycle γ , which is not a coboundary, can be easily obtained by considering the covering group G^* of G , which is the quaternion group of order 8. In order to obtain γ one takes a function $\phi : G \rightarrow G^*$, which is a choice of representatives of cosets of G^* by G , and defines $\tilde{\gamma}(g, s) = \phi(g)\phi(s)(\phi(gs))^{-1}$, $g, s \in G$. Denote by $\varphi : \mathcal{Z}(G^*) \rightarrow \langle -1 \rangle$ the isomorphism between the center of G^* (which has order 2) and $\langle -1 \rangle \subseteq \mathbb{C}$. Then $\gamma = \varphi \circ \tilde{\gamma}$ is a 2-cocycle which is not a coboundary. One readily checks that this gives the cocycle $\gamma : G \times G \rightarrow \kappa^*$ with $\gamma(g, 1) = \gamma(1, g) = 1$, for all $g \in G$ and $\gamma(a, a) = \gamma(a, ab) = \gamma(b, a) = \gamma(b, b) = \gamma(ab, b) = \gamma(ab, ab) = -1$, $\gamma(a, b) = \gamma(b, ab) = \gamma(ab, a) = 1$.

We resume the example of this section in the next:

PROPOSITION 5: *Let κ be an isomorphic copy of the complex numbers \mathbb{C} and let $\mathbb{S}^1 \subseteq \mathbb{C}$ be the circle group, i.e., the group of all complex roots of 1. Let, furthermore, G be an arbitrary finite group seen as a subgroup of S_n for some n . Taking the action of $G \subseteq S_n$ on $(\kappa\mathbb{S}^1)^{\otimes n}$ by permutation of roots, consider*

the smash product Hopf algebra

$$H'_1 = (\kappa\mathbb{S}^1)^{\otimes n} \rtimes \kappa G.$$

Let $X \subseteq G$ be an arbitrary subset which is not a subgroup, and consider the subalgebra $\tilde{A} = (\sum_{g \in X} p_g)(\kappa G)^* \subseteq (\kappa G)^*$, and write $A' = \kappa[t, t^{-1}]^{\otimes n} \otimes \tilde{A}$. Then with the notation established in (20) and (23), the formula

$$(\chi_{\theta_1, \dots, \theta_n} \otimes u_g) \cdot (t_1^{k_1} \cdots t_n^{k_n} \otimes p_s) \\ = \exp\{i(k_1\theta_{g^{-1}s(1)} + \cdots + k_n\theta_{g^{-1}s(n)})\} t_1^{k_1} \cdots t_n^{k_n} \otimes p_{s^{-1}g},$$

where $g \in G$ and $s \in X \subseteq G$, gives a left partial action $\alpha : H'_1 \times A' \rightarrow A'$. Assume now that the Schur Multiplier of G is non-trivial and take a normalized (see (26)) 2-cocycle $\gamma : G \times G \rightarrow \kappa^*$ which is not a coboundary. For arbitrary $h = \chi_{\theta_1, \dots, \theta_n} \otimes u_g$ and $l = \chi_{\theta'_1, \dots, \theta'_n} \otimes u_s$ in H'_1 set

$$\omega(h, l) = \gamma(g, s) (h \cdot l \cdot \mathbf{1}_{A'}).$$

Then the pair (α, ω) forms a twisted partial action of $H'_1 = (\kappa\mathbb{S}^1)^{\otimes n} \rtimes \kappa G$ on A' , which satisfies (16) and (17).

4. Symmetric twisted partial actions

In [14] a twisted partial action of a group G over a unital κ -algebra A was defined as a triple

$$(\{D_g\}_{g \in G}, \{\alpha_g\}_{g \in G}, \{w_{g,h}\}_{(g,h) \in G \times G}),$$

where for each $g, h \in G$, D_g is an ideal of A and $w_{g,h}$ is a multiplier of $D_g D_{gh}$ with some properties. If each D_g is generated by a central idempotent 1_g , then, as we have seen in Example 1, this matches our concept of a partial action of the group Hopf algebra κG over A , and in this case, $1_g = g \cdot 1_A$. Then, from now on, unless explicitly stated, we are going to consider only partial actions of a Hopf algebra H over some unital algebra A such that the map $\mathbf{e} \in \text{Hom}(H, A)$, given by $\mathbf{e}(h) = (h \cdot 1_A)$, is central with respect to the convolution product. These partial actions are, in some sense, more akin to partial group actions.

The second point of interest in twisted partial group actions is the case where the cocycles $\omega_{g,h}$ are invertible in $D_g D_{gh}$, for all $g, h \in G$. If the group action is global, then every element $\omega_{g,h}$ is an invertible element in A ; this is automatically translated into the Hopf algebra setting by saying that the cocycle

$\omega \in \text{Hom}(H \otimes H, A)$ is convolution invertible. In the partial case, we have to search for more suitable conditions to replace the convolution invertibility for the cocycle.

Let $A = (A, \cdot, \omega)$ be a twisted partial H -module algebra. From the definition it follows that $f_1(h, k) = (h \cdot \mathbf{1}_A)\varepsilon(k)$ and $f_2(h, k) = (hk \cdot \mathbf{1}_A)$ are both (convolution) idempotents in $\text{Hom}(H \otimes H, A)$. We also have that \mathbf{e} is an idempotent in $\text{Hom}(H, A)$ (and $f_1(h, k) = \mathbf{e}(h)\varepsilon(k)$).

Let us assume that both f_1 and f_2 are central in $\text{Hom}(H \otimes H, A)$. In this case condition (4) of the definition of a twisted partial action reads as

$$\sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k_{(2)}) = \omega(h, k),$$

and by Proposition 1 one also has

$$\sum \omega(h_{(1)}, k)(h_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k) = \omega(h, k).$$

Notice that this actually says that ω is an element of the ideal $\langle f_1 * f_2 \rangle \subset \text{Hom}(H \otimes H, A)$ generated by $f_1 * f_2$. Clearly $f_1 * f_2$ is the unity element of $\langle f_1 * f_2 \rangle$. Observe also that the centrality of f_1 evidently implies that of $\mathbf{e} \in \text{Hom}(H, A)$.

Definition 2: Let $A = (A, \cdot, \omega)$ be a twisted partial H -module algebra. We will say that the partial action is **symmetric** if

- (i) f_1 and f_2 are central in $\text{Hom}(H \otimes H, A)$;
- (ii) ω is a normalized cocycle which is an *invertible* element of the ideal $\langle f_1 * f_2 \rangle \subset \text{Hom}(H \otimes H, A)$, i.e., ω satisfies conditions (16) and (17) and has a convolution inverse ω' in $\langle f_1 * f_2 \rangle$;
- (iii) $h \cdot (k \cdot \mathbf{1}_A) = \sum (h_1 \cdot \mathbf{1}_A)(h_2k \cdot \mathbf{1}_A)$, for every $h, k \in H$.

We remark once more that to say that $\omega' : H \otimes H \rightarrow A$ lies in $\langle f_1 * f_2 \rangle$ is equivalent to requiring the equalities

$$(29) \quad \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot \mathbf{1}_A) = \omega'(h, k) = \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A),$$

and that ω' is the inverse of ω in $\langle f_1 * f_2 \rangle$ if and only if

$$(30) \quad (\omega * \omega')(h, k) = (\omega' * \omega)(h, k) = \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k \cdot \mathbf{1}_A).$$

It readily follows from (29) and (30) that ω' is also normalized, i.e. $\omega'(1_H, h) = \omega'(h, 1_H) = h \cdot \mathbf{1}_A$ for all $h \in H$.

Multiplying equality (3) on the right by ω' and using (iii) of Definition 2, we obtain

$$(31) \quad h \cdot (k \cdot a) = \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)})$$

for all $h, k \in H$ and $a \in A$, which is an expression analogous to that of global twisted actions of Hopf algebras and also of partial twisted actions of groups. It is easy to prove that if one assumes that the first two items of Definition 2 and equality (31) hold, then item (iii) of Definition 2 follows.

Formula (31) also provides another equality for ω' which is similar to (3). Multiplying (31) by ω' on the left and using the centrality of e , we obtain

$$\begin{aligned} & \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot (k_{(2)} \cdot a)) \\ &= \sum (\omega' * \omega)(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)}) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k_{(1)} \cdot \mathbf{1}_A)(h_{(3)}k_{(2)} \cdot a)\omega'(h_{(4)}, k_{(3)}) \\ &= \sum (h_{(1)}k_{(1)} \cdot a)(h_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(3)}, k_{(2)}) \stackrel{(29)}{=} \sum (h_{(1)}k_{(1)} \cdot a)\omega'(h_{(2)}, k_{(2)}). \end{aligned}$$

Therefore, ω' satisfies

$$(32) \quad \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot (k_{(2)} \cdot a)) = \sum (h_{(1)}k_{(1)} \cdot a)\omega'(h_{(2)}, k_{(2)})$$

for all $h, k \in H$ and $a \in A$.

We shall need expressions for $h \cdot \omega(h, k)$ and $h \cdot \omega'(h, k)$, and for this we prove first an intermediate result, which is interesting on its own.

LEMMA 2: *Let \mathcal{S} be a semigroup and let v, e, e' be elements of \mathcal{S} . If there is an element $v' \in \mathcal{S}$ such that*

$$(33) \quad vv' = e, \quad v'v = e' \quad \text{and} \quad v'e = v',$$

then $v' \in \mathcal{S}$ satisfying (33) is unique.

Proof. In fact, assume that v' is a solution of (33). Then v' also satisfies

$$(34) \quad e'v' = v',$$

because

$$e'v' = (v'v)v' = v'(vv') = v'e = v'.$$

Suppose that v'' is another solution. It follows from (33) and (34) that

$$v'' = v''e = v''(vv') = (v''v)v' = e'v' = v'. \quad \blacksquare$$

PROPOSITION 6: *Let $(A, \cdot, (\omega, \omega'))$ be a symmetric twisted partial H -module algebra. Then*

$$(35) \quad h \cdot \omega(k, m) = \sum \omega(h_{(1)}, k_{(1)})\omega(h_{(2)}k_{(2)}, m_{(1)})\omega'(h_{(3)}, k_{(3)}m_{(2)}),$$

$$(36) \quad h \cdot \omega'(k, m) = \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega'(h_{(3)}, k_{(3)}).$$

Proof. To prove (35), multiply (17) by ω' on the right, obtaining

$$\begin{aligned} \sum (h_{(1)} \cdot \omega(k_{(1)}, m_{(1)}))(\omega * \omega')(h_{(2)}, k_{(2)}m_{(2)}) \\ = \sum \omega(h_{(1)}, k_{(1)})\omega(h_{(2)}k_{(2)}, m_{(1)})\omega'(h_{(2)}, k_{(2)}m_{(2)}). \end{aligned}$$

Since the left-hand side equals

$$\begin{aligned} \sum (h_{(1)} \cdot (\omega(k_{(1)}, m_{(1)}))(h_{(2)} \cdot (k_{(2)}m_{(2)} \cdot \mathbf{1}_A)) \\ = h \cdot \left(\sum \omega(k_{(1)}, m_{(1)})(k_{(2)}m_{(2)} \cdot \mathbf{1}_A) \right) = h \cdot \omega(k, m), \end{aligned}$$

equation (35) follows. The proof of (36) is a bit more involved and uses Lemma 2. Consider $\text{Hom}(H^{\otimes 3}, A)$ as a multiplicative semigroup, and take the elements

$$v(h, k, m) = h \cdot \omega(k, m),$$

$$e(h, k, m) = e'(h \otimes k \otimes m) = (h \cdot (k \cdot (m \cdot \mathbf{1}_A))) = \sum h \cdot [(k_{(1)} \cdot \mathbf{1}_A)(k_{(2)}m \cdot \mathbf{1}_A)].$$

We will show that

$$v'(h, k, m) = h \cdot \omega'(k, m),$$

$$v''(h, k, m) = \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega'(h_{(3)}, k_{(3)})$$

satisfy

$$v * v' = e, \quad v' * v = e, \quad v' * e = v'$$

and

$$v * v'' = e, \quad v'' * v = e, \quad v'' * e = v'',$$

thus proving, via Lemma 2, that $v' = v''$.

Keep in mind that e can be written also as

$$\begin{aligned} e(h, k, m) &= \sum h \cdot [(k_{(1)} \cdot \mathbf{1}_A)(k_{(2)}m \cdot \mathbf{1}_A)] \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k_{(1)} \cdot \mathbf{1}_A)(h_{(3)}k_{(2)}m \cdot \mathbf{1}_A). \end{aligned}$$

The equalities involving v' are straightforward. For instance,

$$(v' * e)(h, k, m) = h[(\omega'(k, m))(k \cdot (m \cdot \mathbf{1}_A))] = h \cdot (\omega'(k, m)) = v'(h \otimes k \otimes m).$$

As for v'' , we first compute $v * v''$, using (35), the centrality of e and (3) of Definition 1:

$$\begin{aligned} (v * v'')(h, k, m) &= \sum \underbrace{(h_{(1)} \cdot \omega(k_{(1)}, m_{(1)}))\omega(h_{(2)}, k_{(2)}m_{(2)})}_{\omega(h_{(1)}, k_{(1)})\omega(h_{(2)}k_{(2)}, m_{(1)})} \omega'(h_{(3)}k_{(3)}, m_{(3)})\omega'(h_{(4)}, k_{(4)}) \\ &\stackrel{(17)}{=} \sum \omega(h_{(1)}, k_{(1)})\omega(h_{(2)}k_{(2)}, m_{(1)})\omega'(h_{(3)}, k_{(3)}m_{(2)})\omega'(h_{(4)}, k_{(4)}) \\ &\stackrel{2.\text{iii},(30)}{=} \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot (m \cdot \mathbf{1}_A))\omega'(h_{(3)}, k_{(3)}) \\ &= (h \cdot (k \cdot (m \cdot \mathbf{1}_A))) = e(h, k, m). \end{aligned}$$

Observe next that given $m \in H$, the linear function $\nu_m : H \otimes H \rightarrow A$ given by $h \otimes k \mapsto \omega(h, km)$ lies in $\text{Hom}(H \otimes H, A)$ and therefore commutes with f_2 , i.e.,

$$(37) \quad \sum \omega(h_{(1)}, k_{(1)}m)(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k_{(2)}m),$$

for all $h, k, m \in H$. Then we calculate $v'' * v$:

$$\begin{aligned} (v'' * v)(h \otimes k \otimes m) &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega'(h_{(3)}, k_{(3)}) (h_{(4)} \cdot \omega(k_{(4)}, m_{(3)})) \\ &\stackrel{(35)}{=} \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)}) \underbrace{\omega'(h_{(3)}, k_{(3)})\omega(h_{(4)}, k_{(4)})}_{\omega(h_{(5)}k_{(5)}, m_{(3)})} \omega'(h_{(6)}, k_{(6)}m_{(4)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})(h_{(3)} \cdot \mathbf{1}_A)(h_{(4)}k_{(3)} \cdot \mathbf{1}_A) \\ &\quad \times \omega(h_{(5)}k_{(4)}, m_{(3)})\omega'(h_{(6)}, k_{(5)}m_{(4)}) \\ &= \sum \underbrace{\omega(h_{(1)}, k_{(1)}m_{(1)})(h_{(2)} \cdot \mathbf{1}_A)}_{\omega(h_{(1)}, k_{(1)}m_{(1)})} \underbrace{\omega'(h_{(3)}k_{(2)}, m_{(2)})(h_{(4)}k_{(3)} \cdot \mathbf{1}_A)}_{\omega'(h_{(3)}k_{(2)}, m_{(2)})} \\ &\quad \times \omega(h_{(5)}k_{(4)}, m_{(3)})\omega'(h_{(6)}, k_{(5)}m_{(4)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)}) \underbrace{\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega(h_{(3)}k_{(3)}, m_{(3)})}_{\omega'(h_{(2)}k_{(2)}, m_{(2)})} \omega'(h_{(4)}, k_{(4)}m_{(4)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) \underbrace{(h_{(3)}k_{(3)}m_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(4)}, k_{(4)}m_{(3)})}_{(h_{(3)}k_{(3)}m_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(4)}, k_{(4)}m_{(3)})} \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(3)}, k_{(3)}m_{(2)}) \end{aligned}$$

$$\begin{aligned} &\stackrel{(37)}{=} \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k_{(2)}m_{(1)})\omega'(h_{(3)}, k_{(3)}m_{(2)}) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k_{(1)} \cdot \mathbf{1}_A)(h_{(3)}k_{(2)}m \cdot \mathbf{1}_A) = e(h, k, m). \end{aligned}$$

And finally, the expression for $v'' * e$ can be obtained as follows

$$\begin{aligned} &v'' * e(h, k, m) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\underbrace{\omega'(h_{(3)}, k_{(3)})(h_{(4)} \cdot \mathbf{1}_A)} \\ &\quad \times \underbrace{(h_{(5)}k_{(4)} \cdot \mathbf{1}_A)(h_{(6)}k_{(5)}m_{(3)} \cdot \mathbf{1}_A)} \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega'(h_{(3)}, k_{(3)})(h_{(4)}k_{(4)} \cdot (m_{(3)} \cdot \mathbf{1}_A)) \\ &\stackrel{(31)}{=} \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\underbrace{\omega'(h_{(3)}, k_{(3)})\omega(h_{(4)}, k_{(4)})} \\ &\quad \times (h_{(5)}k_{(5)}m_{(3)} \cdot \mathbf{1}_A)\omega'(h_{(6)}, k_{(6)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})(h_{(3)} \cdot \mathbf{1}_A)(h_{(4)}k_{(3)} \cdot \mathbf{1}_A) \\ &\quad \times (h_{(5)}k_{(4)}m_{(3)} \cdot \mathbf{1}_A)\omega'(h_{(6)}, k_{(5)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})(h_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(3)}k_{(2)}, m_{(2)})(h_{(4)}k_{(3)} \cdot \mathbf{1}_A) \\ &\quad \times (h_{(5)}k_{(4)}m_{(3)} \cdot \mathbf{1}_A) \times \omega'(h_{(6)}, k_{(5)}) \\ &= \sum \omega(h_{(1)}, k_{(1)}m_{(1)})\omega'(h_{(2)}k_{(2)}, m_{(2)})\omega'(h_{(3)}, k_{(3)}) = v''(h, k, m). \end{aligned}$$

Therefore, Lemma 2 implies (36). ■

Example 4: Consider a twisted H -module algebra B as in Example 2, and assume that the map $u : H \otimes H \rightarrow B$, which twists the action, is a normalized *invertible* cocycle with convolution inverse u^{-1} . Suppose, furthermore, that B has a nontrivial central idempotent $\mathbf{1}_A$, and consider the twisted partial H -module structure on the ideal $A = \mathbf{1}_A B$ as was done in Example 2: the partial action and the cocycle ω are defined by

$$\begin{aligned} &h \cdot a = \mathbf{1}_A(h \triangleright a), \\ &\omega(h, k) = \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot \mathbf{1}_A). \end{aligned}$$

Suppose also that $f_1(h \otimes k) = (h \cdot \mathbf{1}_A)\varepsilon(k)$ and $f_2(h \otimes k) = (hk \cdot \mathbf{1}_A)$ are central in $\text{Hom}(H \otimes H, A)$. Under this hypothesis, the functions $h \otimes k \mapsto \mathbf{1}_A u(h, k)$ and $h \otimes k \mapsto \mathbf{1}_A u^{-1}(h, k)$ commute with \mathbf{e} and f_2 , and it is obvious that

$$\omega'(h, k) = \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)u^{-1}(h_{(2)}, k_{(2)})(h_{(3)} \cdot \mathbf{1}_A)$$

is the inverse of ω in $\langle f_1 * f_2 \rangle$. Note also that

$$\begin{aligned}
 h \cdot (k \cdot a) &= \mathbf{1}_A(h \triangleright (\mathbf{1}_A(k \triangleright a))) = \mathbf{1}_A(h \triangleright (\mathbf{1}_A(k \triangleright a)\mathbf{1}_A)) \\
 &= \mathbf{1}_A \sum (h_{(1)} \triangleright \mathbf{1}_A)(h_{(2)} \triangleright (k \triangleright a))(h_{(3)} \triangleright \mathbf{1}_A) \\
 &= \mathbf{1}_A \sum (h_{(1)} \triangleright \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \triangleright a)u^{-1}(h_{(4)}, k_{(3)})(h_{(5)} \triangleright \mathbf{1}_A) \\
 &= \mathbf{1}_A \sum (h_{(1)} \triangleright \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \triangleright \mathbf{1}_A)(h_{(4)}k_{(3)} \triangleright a)(h_{(5)}k_{(3)} \triangleright \mathbf{1}_A) \\
 &\quad \times u^{-1}(h_{(6)}, k_{(4)})(h_{(7)} \triangleright \mathbf{1}_A) \\
 &= \sum \underbrace{(h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}, k_{(1)})(h_{(3)}k_{(2)} \cdot \mathbf{1}_A)}_{\omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)}(h_{(4)}k_{(3)} \cdot a) \\
 &\quad \times \underbrace{(h_{(5)}k_{(3)} \cdot \mathbf{1}_A)u^{-1}(h_{(6)}, k_{(4)})(h_{(7)} \cdot \mathbf{1}_A)}_{\omega'(h_{(3)}, k_{(3)})} \\
 &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)}),
 \end{aligned}$$

and it follows that

$$\begin{aligned}
 h \cdot (k \cdot \mathbf{1}_A) &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A)\omega'(h_{(3)}, k_{(3)}) \\
 &= \sum \omega(h_{(1)}, k_{(1)})\omega'(h_{(2)}, k_{(2)}) = \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k \cdot \mathbf{1}_A),
 \end{aligned}$$

proving that we have a symmetric twisted partial H -module algebra.

Another important point arising in the context of symmetric twisted partial Hopf actions is to give criteria in order to decide whether two twisted partial actions give rise to the same crossed product. In the classical case, two crossed products are isomorphic if, and only if, the associated twisted (global) actions can be transformed one into another by some kind of coboundary (see, for instance, [29] for the main results of the classical case). In the case of abelian groups, there is, indeed, a cohomology theory involved, and the cocycles performing the twisted actions are related by coboundaries. What we shall see now is an analogue of Theorem 7.3.4 of [29] for twisted partial Hopf actions; this result opens a window for a cohomological point of view of the twisted cocycles presented above.

THEOREM 1: *Let A be a unital algebra and H a Hopf algebra with two symmetric twisted partial actions on A , $h \otimes a \mapsto h \cdot a$, and $h \otimes a \mapsto h \bullet a$, with cocycles ω and σ , respectively. Suppose that there is an algebra isomorphism*

$$\Phi : A\#_{\omega}H \rightarrow A\#_{\sigma}H$$

which is also a left A -module and right H -comodule map. Then there exists linear maps $u, v \in \text{Hom}(H, A)$ such that, for all $h, k \in H, a \in A$,

- (i) $u * v(h) = h \cdot \mathbf{1}_A$,
- (ii) $u(h) = \sum u(h_{(1)})(h_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)})$,
- (iii) $h \bullet a = \sum v(h_{(1)})(h_{(2)} \cdot a)u(h_{(3)})$,
- (iv) $\sigma(h, k) = \sum v(h_{(1)})(h_{(2)} \cdot v(k_{(1)}))\omega(h_{(3)}, k_{(2)})u(h_{(4)}k_{(3)})$,
- (v) $\Phi(a\#_\omega h) = \sum au(h_{(1)})\#_\sigma h_{(2)}$.

Conversely, given maps $u, v \in \text{Hom}(H, A)$ satisfying (i),(ii),(iii) and (iv), and in addition $u(1_H) = v(1_H) = \mathbf{1}_A$, then the map Φ , as presented in (v), is an isomorphism of algebras.

Proof. (\Rightarrow) The left A -module structure on the crossed products is given by the left multiplication:

$$a \blacktriangleright (b\#h) = (a\#1_H)(b\#h) = ab\#h,$$

and the right H -comodule structure is given by $\rho = I_A \otimes \Delta$. Let

$$\Phi : A\#_\omega H \rightarrow A\#_\sigma H$$

be the algebra isomorphism which also is a left A -module and right H -comodule map. Define $u, v \in \text{Hom}(H, A)$ as

$$u(h) = (I_A \otimes \varepsilon)\Phi(1_A\#_\omega h), \quad \text{and} \quad v(h) = (I_A \otimes \varepsilon)\Phi^{-1}(1_A\#_\sigma h).$$

Let us verify that the maps u, v , as defined above, satisfy the items (i) to (v). For the item (v) we have, for all $a \in A$ and $h \in H$,

$$\begin{aligned} \Phi(a\#_\omega h) &= a \blacktriangleright ((\Phi(1_A\#_\omega h))) = a \blacktriangleright \{(I_A \otimes \varepsilon \otimes I_H)(I_A \otimes \Delta)\Phi(1_A\#_\omega h)\} \\ &= a \blacktriangleright \left\{ (I_A \otimes \varepsilon \otimes I_H)\Phi \otimes I_H \left(\sum 1_A\#_\omega h_{(1)} \right) \otimes h_{(2)} \right\} \\ &= a \blacktriangleright \left\{ (I_A \otimes \varepsilon)\Phi \left(\sum 1_A\#_\omega h_{(1)} \right) \otimes h_{(2)} \right\} \\ &= a \blacktriangleright \left(\sum u(h_{(1)})\#_\sigma h_{(2)} \right) = \sum au(h_{(1)})\#_\sigma h_{(2)}. \end{aligned}$$

With a totally similar reasoning, we can conclude that

$$\Phi^{-1}(a\#_\sigma h) = \sum av(h_{(1)})\#_\omega h_{(2)}.$$

Notice that we readily obtain from the above that $u(1_H) = v(1_H) = \mathbf{1}_A$.

For item (i) consider the expression

$$\begin{aligned} \sum (h_{(1)} \cdot \mathbf{1}_A) \#_{\omega} h_{(2)} &= \mathbf{1}_A \#_{\omega} h = \Phi^{-1}(\Phi(\mathbf{1}_A \#_{\omega} h)) \\ &= \Phi^{-1}\left(\sum u(h_{(1)}) \#_{\sigma} h_{(2)}\right) \\ &= \sum u(h_{(1)})v(h_{(2)}) \#_{\omega} h_{(3)}. \end{aligned}$$

Applying $(\text{Id} \otimes \varepsilon)$ on both sides, we obtain

$$\sum u(h_{(1)})v(h_{(2)}) = h \cdot \mathbf{1}_A.$$

Analogously, we can conclude that

$$\sum v(h_{(1)})u(h_{(2)}) = h \bullet \mathbf{1}_A.$$

Item (ii) is easily obtained by applying $I_A \otimes \varepsilon$ on both sides of the equality

$$\begin{aligned} \sum u(h_{(1)}) \#_{\sigma} h_{(2)} &= \Phi(\mathbf{1}_A \#_{\omega} h) = \Phi\left(\sum (h_{(1)} \cdot \mathbf{1}_A) \#_{\omega} h_{(2)}\right) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)u(h_{(2)}) \#_{\sigma} h_{(3)}. \end{aligned}$$

The absorption of $h \cdot \mathbf{1}_A$ on the other side in (ii) comes from the fact that the twisted partial action is symmetric.

In order to prove items (iii) and (iv), we use the fact that Φ^{-1} is an algebra morphism, as so is Φ . Therefore

$$\Phi^{-1}((a \#_{\sigma} h)(b \#_{\sigma} k)) = \Phi^{-1}(a \#_{\sigma} h)\Phi^{-1}(b \#_{\sigma} k),$$

which gives

$$\begin{aligned} \sum a(h_{(1)} \bullet b)\sigma(h_{(2)}, k_{(1)})v(h_{(3)}k_{(2)}) \#_{\omega} h_{(4)}k_{(3)} \\ = \sum av(h_{(1)})(h_{(2)} \cdot (bv(k_{(1)})))\omega(h_{(3)}, k_{(2)}) \#_{\omega} h_{(4)}k_{(3)}. \end{aligned}$$

Applying $I_A \otimes \varepsilon$ on both sides, we get

(38)

$$\sum a(h_{(1)} \bullet b)\sigma(h_{(2)}, k_{(1)})v(h_{(3)}k_{(2)}) = \sum av(h_{(1)})(h_{(2)} \cdot (bv(k_{(1)})))\omega(h_{(3)}, k_{(2)}).$$

Using this formula for $a = \mathbf{1}_A$ and $k = 1_H$ we obtain

$$\sum (h_{(1)} \bullet b)v(h_{(2)}) = \sum v(h_{(1)})(h_{(2)} \cdot b).$$

The expression (iii) is finally obtained multiplying convolutively on the right by u :

$$\sum (h_{(1)} \bullet b)v(h_{(2)})u(h_{(3)}) = \sum v(h_{(1)})(h_{(2)} \cdot b)u(h_{(3)}),$$

and using the fact that $v * u(h) = h \bullet \mathbf{1}_A$. This gives

$$h \bullet b = \sum v(h_{(1)})(h_{(2)} \cdot b)u(h_{(3)}).$$

On the other hand, putting $a = b = \mathbf{1}_A$ in (38) we get

$$\sum \sigma(h_{(1)}, k_{(1)})v(h_{(2)}k_{(2)}) = \sum v(h_{(1)})(h_{(2)} \cdot v(k_{(1)}))\omega(h_{(3)}, k_{(2)}).$$

Therefore

$$\begin{aligned} \sum \sigma(h_{(1)}, k_{(1)})v(h_{(2)}k_{(2)})u(h_{(3)}k_{(3)}) \\ = \sum v(h_{(1)})(h_{(2)} \cdot v(k_{(1)}))\omega(h_{(3)}, k_{(2)})u(h_{(4)}k_{(3)}). \end{aligned}$$

Remembering that the cocycle σ has the absorption property

$$\sigma(h, k) = \sum \sigma(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \bullet \mathbf{1}_A),$$

we obtain

$$\sigma(h, k) = \sum v(h_{(1)})(h_{(2)} \cdot v(k_{(1)}))\omega(h_{(3)}, k_{(2)})u(h_{(4)}k_{(3)}).$$

(\Leftarrow) Conversely, let us consider unit preserving maps $u, v \in \text{Hom}(H, A)$, satisfying the items (i) to (iv) in the statement. We shall verify that $\Phi : A \#_{\omega} H \rightarrow A \#_{\sigma} H$ given by

$$\Phi(a \#_{\omega} h) = \sum au(h_{(1)}) \#_{\sigma} h_{(2)}$$

is indeed an algebra morphism.

We see immediately that $\Phi(\mathbf{1}_A \#_{\omega} \mathbf{1}_H) = \mathbf{1}_A \#_{\sigma} \mathbf{1}_H$. For the multiplicativity, we have

$$\begin{aligned} &\Phi(a \#_{\omega} h) \Phi(b \#_{\omega} k) \\ &= \sum (au(h_{(1)}) \#_{\sigma} h_{(2)})(bu(k_{(1)}) \#_{\sigma} k_{(2)}) \\ &= \sum au(h_{(1)})(h_{(2)} \bullet (bu(k_{(1)}))) \sigma(h_{(3)}, k_{(2)}) \#_{\sigma} h_{(4)}k_{(3)} \\ &= \sum au(h_{(1)})v(h_{(2)})(h_{(3)} \cdot (bu(k_{(1)})))u(h_{(4)})v(h_{(5)})(h_{(6)} \cdot v(k_{(2)})) \\ &\quad \times \omega(h_{(7)}, k_{(3)})u(h_{(8)}k_{(4)}) \#_{\sigma} h_{(9)}k_{(5)} \\ &= \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot u(k_{(1)}))(h_{(3)} \cdot v(k_{(2)}))\omega(h_{(4)}, k_{(3)})u(h_{(5)}k_{(4)}) \#_{\sigma} h_{(6)}k_{(5)} \\ &= \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot (u(k_{(1)})v(k_{(2)})))\omega(h_{(3)}, k_{(3)})u(h_{(4)}k_{(4)}) \#_{\sigma} h_{(5)}k_{(5)} \\ &= \sum a(h_{(1)} \cdot b)(h_{(2)} \cdot (k_{(1)} \cdot \mathbf{1}_A))\omega(h_{(3)}, k_{(2)})u(h_{(4)}k_{(3)}) \#_{\sigma} h_{(5)}k_{(4)} \end{aligned}$$

$$\begin{aligned}
 &= \sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)})u(h_{(3)}k_{(2)})\#_{\sigma}h_{(4)}k_{(3)} \\
 &= \Phi\left(\sum a(h_{(1)} \cdot b)\omega(h_{(2)}, k_{(1)})\#_{\omega}h_{(3)}k_{(2)}\right) \\
 &= \Phi((a\#_{\omega}h)(b\#_{\omega}k)).
 \end{aligned}$$

Now, it remains to show that Φ is invertible. Consider the map

$$\Psi : A\#_{\sigma}H \rightarrow A\#_{\omega}H$$

given by

$$\Psi(a\#_{\sigma}h) = \sum av(h_{(1)})\#_{\omega}h_{(2)}.$$

Then, we have

$$\Psi(\Phi(a\#_{\omega}h)) = \sum au(h_{(1)})v(h_{(2)})\#_{\omega}h_{(3)} = \sum a(h_{(1)} \cdot \mathbf{1}_A)\#_{\omega}h_{(2)} = a\#_{\omega}h.$$

From (ii) and (iii), we easily conclude that $v * u(h) = h \bullet \mathbf{1}_A$, and then

$$\Phi(\Psi(a\#_{\sigma}h)) = \sum av(h_{(1)})u(h_{(2)})\#_{\sigma}h_{(3)} = \sum a(h_{(1)} \bullet \mathbf{1}_A)\#_{\sigma}h_{(2)} = a\#_{\sigma}h.$$

Therefore, $\Psi = \Phi^{-1}$ as we wanted to prove. \blacksquare

5. Partial cleft extensions

It is a well-known simple fact that a group graded algebra $\mathcal{B} = \bigoplus_{g \in G} \mathcal{B}_g$ is isomorphic to a crossed product $\mathcal{A} * G$, where $\mathcal{A} = \mathcal{B}_e$ and $e \in G$ is the neutral element of the group G , exactly when each \mathcal{B}_g contains an element u_g which is invertible in \mathcal{B} . Evidently, the inverse v_g of u_g belongs to $\mathcal{B}_{g^{-1}}$. Thus we have the maps $\gamma : G \rightarrow \mathcal{B}$, $g \mapsto u_g \in \mathcal{B}_g$ and $\gamma' : G \rightarrow \mathcal{B}$, $g \mapsto v_g \in \mathcal{B}_{g^{-1}}$, and γ' is in some sense inverse to γ . This becomes precise if we recall that \mathcal{B} is a κG -module algebra, and a more general result for a Hopf algebra H says that an H -comodule algebra B is isomorphic to a smash product $A\#H$, $A = B^{coH}$, if and only if $A \subseteq B$ is a cleft extension, which means that there exists a κ -linear map $\gamma : H \rightarrow B$ which fits into an appropriate commutative diagram and possesses a convolution inverse $\gamma' : H \rightarrow B$.

The partial case is essentially more complicated. One of the results in [14] gives a criteria for a non-degenerate G -graded algebra $\mathcal{B} = \bigoplus_{g \in G} \mathcal{B}_g$ to have the structure of a crossed product $\mathcal{A} * G$ by a twisted partial action of G on $\mathcal{A} = \mathcal{B}_e$. More specifically, if \mathcal{B} satisfies

$$(39) \quad \mathcal{B}_g\mathcal{B}_{g^{-1}}\mathcal{B}_g = \mathcal{B}_g, \quad (\forall g \in G)$$

then using the multiplication in \mathcal{B} it is possible to define, for each $g \in G$, idempotent ideals $\mathcal{D}_g = \mathcal{B}_g \mathcal{B}_{g^{-1}}$, $\mathcal{D}_{g^{-1}} = \mathcal{B}_{g^{-1}} \mathcal{B}_g$ of \mathcal{B}_e , a unital $\mathcal{D}_g - \mathcal{D}_{g^{-1}}$ bimodule \mathcal{B}_g and a unital $\mathcal{D}_{g^{-1}} - \mathcal{D}_g$ bimodule $\mathcal{B}_{g^{-1}}$, such that they constitute a Morita context. The main ingredients used to the construction of the crossed product are operators u_g and v_g in the multiplier algebra of the context algebra

$$\mathcal{C}_g = \begin{pmatrix} \mathcal{D}_g & \mathcal{B}_g \\ \mathcal{B}_{g^{-1}} & \mathcal{D}_{g^{-1}} \end{pmatrix}$$

such that

$$(40) \quad u_g v_g = e_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad \text{and} \quad v_g u_g = e_{22} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then it turns out that a non-degenerate G -graded algebra \mathcal{B} is isomorphic as a graded algebra to the crossed product $\mathcal{A} * G$ by a twisted partial action exactly when (39) is satisfied and for each $g \in G$ there exist multipliers u_g and v_g of \mathcal{C}_g such that (40) holds.

Now it becomes natural to treat this topic in the context of twisted partial Hopf actions, which is the purpose of the present section. The ‘‘partiality’’ is reflected now on the properties of γ . Instead of assuming that γ is convolution invertible, one declares the existence of a map γ' which is related to γ by conditions which are weaker than that of the convolution invertibility. Some of them match equalities which already played a crucial role in the study of partial actions and partial representations (see Remark 2).

Definition 3: Let B be a right H -comodule unital algebra with coaction given by $\rho : B \rightarrow B \otimes H$ and let A be a subalgebra of B . We will say that $A \subset B$ is an H -extension if $A = B^{coH}$. An H -extension $A \subset B$ is **partially cleft** if there is a pair of k -linear maps $\gamma, \gamma' : H \rightarrow B$ such that

- (i) $\gamma(1_H) = 1_B$,
- (ii) the diagrams below are commutative:

$$(41) \quad \begin{array}{ccc} H & \xrightarrow{\gamma} & B \\ \downarrow \Delta & & \downarrow \rho \\ H \otimes H & \xrightarrow{\gamma \otimes I_H} & B \otimes H \end{array} \qquad \begin{array}{ccc} H & \xrightarrow{\gamma'} & B \\ \downarrow \Delta^{cop} & & \downarrow \rho \\ H \otimes H & \xrightarrow{\gamma' \otimes S} & B \otimes H \end{array}$$

- (iii) $(\gamma * \gamma') \circ M$ is a central element in the convolution algebra $\text{Hom}(H \otimes H, A)$, where $M : H \otimes H \rightarrow H$ is the multiplication in H , and $(\gamma' * \gamma)(h)$ commutes with every element of A for each $h \in H$,

and, for all $b \in B$ and $h, k \in H$, if we write $e_h = (\gamma * \gamma')(h)$ and $\tilde{e}_h = (\gamma' * \gamma)(h)$, then

- (iv) $\sum b_{(0)} \gamma'(b_{(1)}) \gamma(b_{(2)}) = b$,
- (v) $\gamma(h) e_k = \sum e_{h_{(1)} k} \gamma(h_{(2)})$,
- (vi) $\gamma'(k) \tilde{e}_h = \sum \tilde{e}_{h k_{(1)}} \gamma'(k_{(2)})$,
- (vii) $\sum \gamma(h k_{(1)}) \tilde{e}_{k_{(2)}} = \sum e_{h_{(1)}} \gamma(h_{(2)} k)$.

Note that item (iii) makes sense because item (ii) implies that $(\gamma * \gamma')(h) \in A$, for all $h \in H$, and therefore $\gamma * \gamma' \in \text{Hom}(H, A)$.

With respect to items (v), (vi) and (vii) we make the following:

Remark 2: Let $\gamma : G \rightarrow \mathcal{B}$ be a partial representation of a group G into a κ -algebra \mathcal{B} , i.e. a κ -linear map such that $\gamma(1_G) = 1_B$, $\gamma(g)\gamma(s)\gamma(s^{-1}) = \gamma(gs)\gamma(s^{-1})$ and $\gamma(g^{-1})\gamma(g)\gamma(s) = \gamma(g^{-1})\gamma(gs)$, for all $g, s \in G$. Then by (2) of [13] the following equality holds:

$$\gamma(g) e_r = e_{gr} \gamma(g),$$

where $e_g = \gamma(g)\gamma(g^{-1})$. This corresponds to item (v) if we take $H = \kappa G$. The above equality plays a crucial role for the interaction between partial actions and partial representations (see [12]), as well as for an analogous interaction in the context of partial projective representations (see [17], [18]). Now writing $\gamma'(g) = \gamma(g^{-1})$ and $\tilde{e}_g = e_{g^{-1}}$, we readily obtain from the above equality that

$$\gamma'(g) \tilde{e}_s = \tilde{e}_{sg} \gamma'(g) \quad \text{and} \quad \gamma(gs) \tilde{e}_s = e_g \gamma(gs),$$

for all $g, s \in G$, which are exactly items (vi) and (vii) above with $H = \kappa G$.

Note that in the case of a cleft extension, with a convolution invertible map γ , the axioms for partial cleft extensions are automatically satisfied if we take γ' to be the convolution inverse of γ .

Observe furthermore that given a partial cleft extension, we also have

$$(42) \quad \gamma'(1_H) = 1_B,$$

since by (iv) of Definition 3 we see that

$$1_B = \sum (1_B)_{(0)} \gamma'((1_B)_{(1)}) \gamma((1_B)_{(2)}) = (1_B) \gamma'(1_H) \gamma(1_H) = \gamma'(1_H).$$

Moreover, since by (41) γ is a morphism of comodules, we have that $\rho^2(\gamma(h)) = (\mathbb{I}_A \otimes \Delta)\rho(\gamma(h)) = \sum \gamma(h_{(1)}) \otimes h_{(2)} \otimes h_{(3)}$. Then applying (iv) of Definition 3 to $b = \gamma(h)$, we conclude that

$$(43) \quad \gamma * \gamma' * \gamma = \gamma.$$

The latter will be quite important in what follows. In particular, multiplying this equality by γ' on the right we obtain that $\gamma * \gamma'$ is an idempotent, and, moreover, multiplying (43) by γ' on the left, we see that $\gamma' * \gamma$ is also idempotent. Furthermore, since any linear function $\tau \in \text{Hom}(H, A)$ can be seen as a function $h \otimes k \mapsto \tau(h)$ in $\text{Hom}(H \otimes H, A)$, item (iii) of Definition 3 implies that $(\gamma * \gamma') * \tau = \tau * (\gamma * \gamma')$, so that we have:

Remark 3: Given a partially cleft extension, $\gamma * \gamma'$ is a central idempotent in the convolution algebra $\text{Hom}(H, A)$.

The map γ' may not satisfy an equality similar to (43), but it always can be replaced by another map $\bar{\gamma}$ that does, and the pair $(\gamma, \bar{\gamma})$ still satisfies properties (i)–(vii), as seen in the following:

LEMMA 3: We may assume that γ' in Definition 3 satisfies the equality

$$(44) \quad \gamma' * \gamma * \gamma' = \gamma'.$$

Proof. Consider the map $\bar{\gamma} = \gamma' * \gamma * \gamma'$. Since $\gamma' * \gamma$ is an idempotent,

$$\bar{\gamma} * \gamma * \bar{\gamma} = (\gamma' * \gamma * \gamma') * \gamma * (\gamma' * \gamma * \gamma') = (\gamma' * \gamma) * (\gamma' * \gamma) * (\gamma' * \gamma) * \gamma' = \gamma' * \gamma * \gamma' = \bar{\gamma}.$$

We will show that the pair $(\gamma, \bar{\gamma})$ satisfies the properties (i)–(vii). Item (i) is immediate in view of (42). Item (ii) holds, since

$$\begin{aligned} \rho(\bar{\gamma}(h)) &= \sum \rho(\gamma'(h_{(1)}))\gamma(h_{(2)})\gamma'(h_{(3)}) = \sum \rho(\gamma'(h_{(1)}))\rho(\gamma(h_{(2)}))\rho(\gamma'(h_{(3)})) \\ &= \sum (\gamma'(h_{(2)}) \otimes S(h_{(1)}))(\gamma(h_{(3)}) \otimes h_{(4)})(\gamma'(h_{(6)}) \otimes S(h_{(5)})) \\ &= \sum \gamma'(h_{(2)})\gamma(h_{(3)})\gamma'(h_{(6)}) \otimes S(h_{(1)})h_{(4)}S(h_{(5)}) \\ &= \sum \gamma'(h_{(2)})\gamma(h_{(3)})\gamma'(h_{(4)}) \otimes S(h_{(1)}) \\ &= \sum \bar{\gamma}(h_{(2)}) \otimes S(h_{(1)}) = (\bar{\gamma} \otimes S)\Delta^{cop}(h). \end{aligned}$$

Item (iii) immediately follows from

$$(45) \quad \gamma * \bar{\gamma} = \gamma * \gamma', \quad \bar{\gamma} * \gamma = \gamma' * \gamma.$$

Item (iv) holds because, given $b \in B$,

$$\begin{aligned} \sum b_{(0)}\bar{\gamma}(b_{(1)})\gamma(b_{(2)}) &= \sum b_{(0)}\gamma'(b_{(1)})(\gamma(b_{(2)})\gamma'(b_{(3)})\gamma(b_{(4)})) \\ &= \sum b_{(0)}\gamma'(b_{(1)})\gamma(b_{(2)}) = b. \end{aligned}$$

It remains to check (v)–(vii). Notice that $e_h = (\gamma * \bar{\gamma})(h)$ and $\tilde{e}_h = (\bar{\gamma} * \gamma)(h)$, thanks to (45). Thus for $\bar{\gamma}$ we need to verify only (vi). We compute

$$\begin{aligned} \bar{\gamma}(k)\tilde{e}_h &= \gamma'(k_{(1)})e_{k_{(2)}}\tilde{e}_h \stackrel{\text{(iii)}}{=} \gamma'(k_{(1)})\tilde{e}_h e_{k_{(2)}} \\ &\stackrel{\text{(vi)}}{=} \tilde{e}_{hk_{(1)}}\gamma'(k_{(2)})e_{k_{(3)}} = \tilde{e}_{hk_{(1)}}\bar{\gamma}(k_{(2)}), \end{aligned}$$

taking into account that $e_k \in A$. ■

Since ρ is an algebra morphism, applying (iv) of Definition 3 to $b = \gamma(h)\gamma(k)$ and also to $b = \gamma(h)\gamma(k)a$, we obtain for any $a \in A = B^{coH}$ and $h, k \in H$ the following equalities:

$$(46) \quad \gamma(h)\gamma(k) = \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)})\gamma(h_{(3)}k_{(3)}),$$

$$(47) \quad \gamma(h)\gamma(k)a = \sum \gamma(h_{(1)})\gamma(k_{(1)})a\gamma'(h_{(2)}k_{(2)})\gamma(h_{(3)}k_{(3)}).$$

Then taking $k = 1_H$ in (47) we have

$$(48) \quad \gamma(h)a = \sum \gamma(h_{(1)})a\gamma'(h_{(2)})\gamma(h_{(3)}).$$

PROPOSITION 7: *If $(A, \cdot, (\omega, \omega'))$ is a symmetric partial twisted H -module algebra, then $A \subset A\#_{(\alpha, \omega)}H$ is a partially cleft H -extension.*

Proof. We see that $A\#_{(\alpha, \omega)}H$ is a right comodule algebra via the mapping $\rho = (I \otimes \Delta) : A\#_{(\alpha, \omega)}H \rightarrow (A\#_{(\alpha, \omega)}H) \otimes H$. It is easy to see that $(A\#_{(\alpha, \omega)}H)^{coH} = A \otimes 1_H$, which we will identify with A via the canonical monomorphism $A \rightarrow A \otimes 1_H$.

Consider the maps $\gamma, \gamma' : H \rightarrow A\#_{(\alpha, \omega)}H$ given by

$$(49) \quad \gamma(h) = \mathbf{1}_A\#h = (\mathbf{1}_A \otimes h)(\mathbf{1}_A \otimes 1_H),$$

$$(50) \quad \gamma'(h) = \sum \omega'(S(h_{(2)}), h_{(3)})\#S(h_{(1)}).$$

From the definition of γ we have $\gamma(1_H) = \mathbf{1}_A\#1_H = \mathbf{1}_{A\#_{(\alpha, \omega)}H}$, which gives (i) of Definition 3. With respect to item (ii), the equality $\rho\gamma = (\gamma \otimes I)\Delta$ follows

directly by the definition of ρ . As for the second diagram in (ii), we have:

$$\begin{aligned} \rho\gamma'(h) &= \sum \rho(\omega'(S(h_{(2)}), h_{(3)})\#S(h_{(1)})) \\ &= \sum \omega'(S(h_{(3)}), h_{(4)})\#S(h_{(2)}) \otimes S(h_{(1)}) \\ &= \sum \gamma'(h_{(2)}) \otimes S(h_{(1)}) = (\gamma' \otimes S)\Delta^{cop}(h), \end{aligned}$$

which completes the proof of (ii) of the definition of partial cleft extension.

Now,

$$\begin{aligned} (\gamma * \gamma')(h) &= \sum (\mathbf{1}_A \# h_{(1)}) (\omega'(S(h_{(3)}), h_{(4)})\#S(h_{(2)})) \\ &= \sum (h_{(1)} \cdot \omega'(S(h_{(6)}), h_{(7)})) \omega(h_{(2)}, S(h_{(5)})) \# \underbrace{h_{(3)} S(h_{(4)})} \\ &= \sum (h_{(1)} \cdot \omega'(S(h_{(4)}), h_{(5)})) \omega(h_{(2)}, S(h_{(3)})) \# \mathbf{1}_H \\ &\stackrel{(36)}{=} \sum \omega(h_{(1)}, \underbrace{S(h_{(8)})h_{(9)}}) \omega'(h_{(2)}S(h_{(7)}), h_{(10)}) \\ &\quad \times \underbrace{\omega'(h_{(3)}, S(h_{(6)})) \omega(h_{(4)}, S(h_{(5)}))} \# \mathbf{1}_H \\ &\stackrel{(30),(16)}{=} \sum (h_{(1)} \cdot \mathbf{1}_A) \underbrace{\omega'(h_{(2)}S(h_{(6)}), h_{(7)})(h_{(3)} \cdot \mathbf{1}_A)} \underbrace{(h_{(4)}S(h_{(5)})) \cdot \mathbf{1}_A} \# \mathbf{1}_H \\ &\stackrel{(29)}{=} \sum (h_{(1)} \cdot \mathbf{1}_A) \omega'(h_{(2)}S(h_{(3)}), h_{(4)}) \# \mathbf{1}_H = (h \cdot \mathbf{1}_A) \# \mathbf{1}_H. \end{aligned}$$

Hence $(\gamma * \gamma')(hk) = f_2(h, k) \# \mathbf{1}_H$, and this implies that $(\gamma * \gamma') \circ M$ is central in $\text{Hom}(H \otimes H, A)$ thanks to the convolution centrality of f_2 . Observe also that $(\gamma * \gamma')(h)$ commutes with every element of A , since each $a \in A$ gives rise to a linear map $\tau_a : H \rightarrow A$ defined by $\tau_a(h) = \varepsilon(h)a$, and $\mathbf{e}(h) = (h \cdot \mathbf{1}_A)$ is central in $\text{Hom}(H, A)$ by assumption. Hence

$$(h \cdot \mathbf{1}_A)a = \sum (h_{(1)} \cdot \mathbf{1}_A)\varepsilon(h_{(2)})a = (\mathbf{e} * \tau_a)(h) = (\tau_a * \mathbf{e})(h) = a(h \cdot \mathbf{1}_A).$$

With respect to $\gamma' * \gamma$,

$$\begin{aligned} \gamma' * \gamma(h) &= (\omega'(S(h_{(2)}), h_{(3)})\#S(h_{(1)}))(\mathbf{1}_A \# h_{(4)}) \\ &= \omega'(S(h_{(4)}), h_{(5)})(S(h_{(3)}) \cdot \mathbf{1}_A) \omega(S(h_{(2)}), h_{(6)}) \# S(h_{(1)})h_{(7)} \\ &= \omega'(S(h_{(3)}), h_{(4)}) \omega(S(h_{(2)}), h_{(5)}) \# S(h_{(1)})h_{(6)} \\ &= (S(h_{(3)})h_{(4)} \cdot \mathbf{1}_A)(S(h_{(2)}) \cdot \mathbf{1}_A) \# S(h_{(1)})h_{(5)} \\ &= (S(h_{(2)}) \cdot \mathbf{1}_A) \# S(h_{(1)})h_{(3)}, \end{aligned}$$

and this expression implies

$$\begin{aligned}
 & (\gamma' * \gamma)(h)(a\#1_H) \\
 &= \sum ((S(h_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})h_{(3)})(a\#1_H) \\
 &= \sum (S(h_{(4)}) \cdot \mathbf{1}_A)(S(h_{(3)})h_{(5)} \cdot a)\omega(S(h_{(2)})h_{(6)}, 1_H)\#S(h_{(1)})h_{(7)} \\
 &= \sum \underbrace{(S(h_{(4)})h_{(5)} \cdot a)(S(h_{(3)}) \cdot \mathbf{1}_A)}\omega(S(h_{(2)})h_{(6)}, 1_H)\#S(h_{(1)})h_{(7)} \\
 &= \sum a(S(h_{(3)}) \cdot \mathbf{1}_A)\omega(S(h_{(2)})h_{(4)}, 1_H)\#S(h_{(1)})h_{(5)} \\
 &= \sum a\omega(S(h_{(3)})h_{(4)}, 1_H)(S(h_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})h_{(5)} \\
 &= \sum a(S(h_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})h_{(3)} \\
 &= \sum (a\#1_H)((S(h_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})h_{(3)}) = (a\#1_H)(\gamma' * \gamma)(h),
 \end{aligned}$$

proving item (iii).

For item (iv), consider $b = a\#h$ in $A\#_{(\alpha, \omega)}H$. Applying $\rho^2 = (I_A \otimes \Delta)\rho$ to b we obtain

$$\sum b_{(0)} \otimes b_{(1)} \otimes b_{(2)} = \sum (a\#h_{(1)}) \otimes h_{(2)} \otimes h_{(3)},$$

and therefore

$$\begin{aligned}
 & \sum b_{(0)}\gamma'(b_{(1)})\gamma(b_{(2)}) \\
 &= \sum (a\#h_{(1)})\gamma'(h_{(2)})\gamma(h_{(3)}) \\
 &= \sum (a\#h_{(1)})(\omega'(S(h_{(3)}) \otimes h_{(4)})\#S(h_{(2)}))(\mathbf{1}_A\#h_{(5)}) \\
 &= \sum (a\#1_H) \underbrace{(\mathbf{1}_A\#h_{(1)})(\omega'(S(h_{(3)}) \otimes h_{(4)})\#S(h_{(2)}))}_{(47), (48)}(\mathbf{1}_A\#h_{(5)}) \\
 &\stackrel{(47), (48)}{=} \sum (a\#1_H)(\gamma * \gamma')(h_{(1)})(\mathbf{1}_A\#h_{(2)}) \\
 &= \sum (a\#1_H)((h_{(1)} \cdot \mathbf{1}_A)\#1_H)(\mathbf{1}_A\#h_{(2)}) = a\#h = b.
 \end{aligned}$$

Next we check (v), using (iii) of Definition 2, as follows:

$$\begin{aligned}
 \sum e_{h_{(1)}k}\gamma(h_{(2)}) &= \sum (h_{(1)}k \cdot \mathbf{1}_A\#1_H)(\mathbf{1}_A\#h_{(2)}) \stackrel{(47)}{=} \sum (h_{(1)}k \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A)\#h_{(3)} \\
 &= \sum (h_{(1)} \cdot (k \cdot \mathbf{1}_A))\#h_{(2)} \\
 &= \sum (h_{(1)} \cdot (k \cdot \mathbf{1}_A))(h_{(2)} \cdot \mathbf{1}_A)\#h_{(3)} \\
 &= (\mathbf{1}_A\#h)(k \cdot \mathbf{1}_A\#1_H) = \gamma(h)(\gamma * \gamma')(k) = \gamma(h)e_k.
 \end{aligned}$$

In order to establish (vi) we compute, using again (iii) of Definition 2, that

$$\begin{aligned}
 \gamma'(h)\tilde{e}_k &= \left(\sum \omega'(S(h_{(2)}), h_{(3)})\#S(h_{(1)}) \left(\sum S(k_{(2)}) \cdot \mathbf{1}_A\#S(k_{(1)})k_{(3)} \right) \right) \\
 &= \sum \omega'(S(h_{(4)}), h_{(5)}) \underbrace{(S(h_{(3)}) \cdot (S(k_{(3)}) \cdot \mathbf{1}_A))}_{\omega(S(h_{(2)}), S(k_{(2)})k_{(4)})\#S(h_{(1)})S(k_{(1)})k_{(5)}} \\
 &\quad \times \omega(S(h_{(2)}), S(k_{(2)})k_{(4)})\#S(h_{(1)})S(k_{(1)})k_{(5)} \\
 &= \sum \omega'(S(h_{(5)}), h_{(6)}) \underbrace{(S(h_{(4)}) \cdot \mathbf{1}_A)(S(h_{(3)})S(k_{(3)}) \cdot \mathbf{1}_A)}_{\omega(S(h_{(2)}), S(k_{(2)})k_{(4)})\#S(h_{(1)})S(k_{(1)})k_{(5)}} \\
 &\quad \times \omega(S(h_{(2)}), S(k_{(2)})k_{(4)})\#S(h_{(1)})S(k_{(1)})k_{(5)} \\
 &\stackrel{(29)}{=} \sum \omega'(S(h_{(4)}), h_{(5)}) \\
 &\quad \times \underbrace{(S(h_{(3)})S(k_{(3)}) \cdot \mathbf{1}_A)\omega(S(h_{(2)}), S(k_{(2)})k_{(4)})\#S(h_{(1)})S(k_{(1)})k_{(5)}}.
 \end{aligned}$$

With respect to the underbraced product, for a fixed $m \in H$ consider the function $\tau_m : H \otimes H \rightarrow A$ given by $h \otimes k \mapsto \omega(h, km)$. Since f_2 is central, we have

$$\begin{aligned}
 &\sum (S(h_{(2)})S(k_{(2)}) \cdot \mathbf{1}_A)\omega(S(h_{(1)}), S(k_{(1)})m) \\
 &= (f_2 * \tau_m)(S(h) \otimes S(k)) = (\tau_m * f_2)(S(h) \otimes S(k)) \\
 &= \sum \omega(S(h_{(2)}), S(k_{(2)})m)(S(h_{(1)})S(k_{(1)}) \cdot \mathbf{1}_A),
 \end{aligned}$$

and consequently we obtain

$$\begin{aligned}
 \gamma'(h)\tilde{e}_k &= \sum \omega'(S(h_{(4)}), h_{(5)})\omega(S(h_{(3)}), \underbrace{S(k_{(3)})k_{(4)}}_{\omega(S(h_{(2)}), S(k_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})S(k_{(1)})k_{(5)}}) \\
 &\quad \times (S(h_{(2)})S(k_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})S(k_{(1)})k_{(5)} \\
 &= \sum \omega'(S(h_{(4)}), h_{(5)}) \underbrace{(S(h_{(3)}) \cdot \mathbf{1}_A)}_{\omega(S(h_{(2)}), S(k_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})S(k_{(1)})k_{(3)}} \\
 &\quad \times (S(h_{(2)})S(k_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})S(k_{(1)})k_{(3)} \\
 &\stackrel{(29)}{=} \sum \omega'(S(h_{(3)}), h_{(4)}) (S(h_{(2)})S(k_{(2)}) \cdot \mathbf{1}_A)\#S(h_{(1)})S(k_{(1)})k_{(3)}.
 \end{aligned}$$

To compute $\sum \tilde{e}_{kh_{(1)}}\gamma'(h_{(2)})$ consider first the function $\mu_{l,m,n} : H \rightarrow A$ defined by $h \mapsto hl \cdot \omega'(m, n)$, where $l, m, n \in H$ are fixed. Then $\mathbf{e} * \mu_{l,m,n} = \mu_{l,m,n} * \mathbf{e}$, and applying both sides of this equality to $S(kh)$, we obtain

$$\begin{aligned}
 (51) \quad &\sum (S(k_{(2)}h_{(2)}) \cdot \mathbf{1}_A) [(S(k_{(1)}h_{(1)})l) \cdot \omega'(m, n)] \\
 &= \sum [(S(k_{(2)}h_{(2)})l) \cdot \omega'(m, n)] (S(k_{(1)}h_{(1)}) \cdot \mathbf{1}_A).
 \end{aligned}$$

Similarly, taking the function $\nu_{m,n} : H \rightarrow A$, given by $h \mapsto \omega(hm, n)$, and using $\mathbf{e} * \nu_{m,n} = \nu_{m,n} * \mathbf{e}$ applied also to $S(hk)$, we also obtain

$$(52) \quad \begin{aligned} \sum (S(k_{(2)}h_{(2)}) \cdot \mathbf{1}_A) \omega(S(k_{(1)}h_{(1)})m, n) \\ = \sum \omega(S(k_{(2)}h_{(2)})m, n) (S(k_{(1)}h_{(1)}) \cdot \mathbf{1}_A). \end{aligned}$$

Then we have

$$\begin{aligned} & \sum \tilde{e}_{kh_{(1)}} \gamma'(h_{(2)}) \\ &= \left[\sum (S(k_{(2)}h_{(2)}) \cdot \mathbf{1}_A) \# S(k_{(1)}h_{(2)})k_{(3)}h_{(3)} \right] \\ & \quad \times \left[\sum \omega'(S(h_{(5)}), h_{(6)}) \# S(h_{(4)}) \right] \\ &= \sum (S(k_{(4)}h_{(4)}) \cdot \mathbf{1}_A) [S(k_{(3)}h_{(3)})k_{(5)}h_{(5)} \cdot \omega'(S(h_{(10)}), h_{(11)})] \\ & \quad \times \omega(S(k_{(2)}h_{(2)})k_{(6)}h_{(6)}, S(h_{(9)})) \# S(k_{(1)}h_{(1)})k_{(7)} \underbrace{h_{(7)}S(h_{(8)})} \\ &= \sum (S(k_{(4)}h_{(4)}) \cdot \mathbf{1}_A) [S(k_{(3)}h_{(3)})k_{(5)}h_{(5)} \cdot \omega'(S(h_{(8)}), h_{(9)})] \\ & \quad \times \omega(S(k_{(2)}h_{(2)})k_{(6)}h_{(6)}, S(h_{(7)})) \# S(k_{(1)}h_{(1)})k_{(7)} \\ &\stackrel{(51)}{=} \sum \left[\underbrace{S(k_{(4)}h_{(4)})k_{(5)}h_{(5)}} \cdot \omega'(S(h_{(8)}), h_{(9)}) \right] (S(k_{(3)}h_{(3)}) \cdot \mathbf{1}_A) \\ & \quad \omega(S(k_{(2)}h_{(2)})k_{(6)}h_{(6)}, S(h_{(7)})) \# S(k_{(1)}h_{(1)})k_{(7)} \\ &= \sum \omega'(S(h_{(6)}), h_{(7)}) \\ & \quad \times \underbrace{(S(k_{(3)}h_{(3)}) \cdot \mathbf{1}_A) \omega(S(k_{(2)}h_{(2)})k_{(4)}h_{(4)}, S(h_{(5)}))}_{\# S(k_{(1)}h_{(1)})k_{(5)}} \\ &= \sum \omega'(S(h_{(6)}), h_{(7)}) \omega \underbrace{(S(k_{(3)}h_{(3)})k_{(4)}h_{(4)}, S(h_{(5)}))}_{\# S(k_{(1)}h_{(1)})k_{(5)}} \\ & \quad \times (S(k_{(2)}h_{(2)}) \cdot \mathbf{1}_A) \# S(k_{(1)}h_{(1)})k_{(5)} \\ &= \sum \underbrace{\omega'(S(h_{(4)}), h_{(5)})(S(h_{(3)}) \cdot \mathbf{1}_A)}_{\# S(k_{(1)}h_{(1)})k_{(3)}} (S(k_{(2)}h_{(2)}) \cdot \mathbf{1}_A) \# S(k_{(1)}h_{(1)})k_{(3)} \\ &= \sum \omega'(S(h_{(3)}), h_{(4)})(S(h_{(2)})S(k_{(2)}) \cdot \mathbf{1}_A) \# S(h_{(1)})S(k_{(1)})k_{(3)}, \end{aligned}$$

which coincides with the expression obtained above for $\gamma'(h)\tilde{e}_k$, thus proving (vi).

Finally, item (vii) follows from the next calculation, in which we use again the convolution centrality of f_2 and \mathbf{e} :

$$\begin{aligned}
 & \sum \gamma(hk_{(1)})\tilde{e}_{k_{(2)}} \\
 &= \sum (\mathbf{1}_A \# hk_{(1)}) [(S(k_3) \cdot \mathbf{1}_A) \# S(k_{(2)})k_{(4)}] \\
 &= \sum (h_{(1)}k_{(1)} \cdot (S(k_{(6)}) \cdot \mathbf{1}_A)) \\
 &\quad \times \omega(h_{(2)}k_{(2)}, S(k_{(5)}) \# k_{(7)}) \# h_{(3)} \underbrace{k_{(3)}S(k_{(4)})}_{k_{(8)}} \\
 &= \sum (h_{(1)}k_{(1)} \cdot (S(k_{(4)}) \cdot \mathbf{1}_A)) \omega(h_{(2)}k_{(2)}, S(k_{(3)})k_{(5)}) \# h_{(3)}k_{(6)} \\
 &= \sum \underbrace{(h_{(1)}k_{(1)} \cdot \mathbf{1}_A)((h_{(2)}k_{(2)}S(k_{(5)}) \cdot \mathbf{1}_A)}_{\omega(h_{(3)}k_{(3)}, S(k_{(4)})k_{(6)})} \# h_{(4)}k_{(7)} \\
 &= \sum (h_{(1)}k_{(1)}S(k_{(5)}) \cdot \mathbf{1}_A)(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) \omega(h_{(3)}k_{(3)}, S(k_{(4)})k_{(6)}) \# h_{(4)}k_{(7)} \\
 &= \sum (h_{(1)}k_{(1)}S(k_{(5)}) \cdot \mathbf{1}_A) \underbrace{(h_{(2)}k_{(2)} \cdot \mathbf{1}_A)\omega(h_{(3)}k_{(3)}, S(k_{(4)})k_{(6)})}_{\# h_{(4)}k_{(7)}} \\
 &= \sum \underbrace{(h_{(1)}k_{(1)}S(k_{(4)}) \cdot \mathbf{1}_A)\omega(h_{(2)}k_{(2)}, S(k_{(3)})k_{(5)})}_{\# h_{(3)}k_{(6)}} \\
 &= \sum \omega(h_{(1)}k_{(1)}, \underbrace{S(k_{(4)})k_{(5)}}_{(h_{(2)}k_{(2)}S(k_{(3)}) \cdot \mathbf{1}_A)}) \# h_{(3)}k_{(6)} \\
 &= \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A) \# h_{(3)}k_{(2)} \\
 &\quad \times \sum (h_{(1)} \cdot \mathbf{1}_A) (h_{(2)}k_{(1)} \cdot \mathbf{1}_A) \# h_{(3)}k_{(2)} \\
 &= \sum (h_{(1)} \cdot \mathbf{1}_A) \omega(1_H, h_{(2)}k_{(1)}) \# h_{(3)}k_{(2)} \\
 &\stackrel{(5)}{=} \sum ((h_{(1)} \cdot \mathbf{1}_A) \# 1_H) (\mathbf{1}_A \# h_{(2)}k) \\
 &= \sum e_{h_{(1)}} \gamma(h_{(2)}k). \quad \blacksquare
 \end{aligned}$$

THEOREM 2: *Let B be an H -comodule algebra and let $A = B^{coH}$. Then the H -extension $A \subset B$ is partially cleft if and only if B is isomorphic to a partial crossed product $A \#_{(\alpha, \omega)} H$ with respect to a symmetric twisted partial H -module structure on A .*

Proof. We have already proved half of this statement in Proposition 7. So, assume that B is partially cleft by the pair of maps $\gamma, \gamma' : H \rightarrow B$. The pair (γ, γ') allows us to define a twisted partial action of H on $A = B^{coH}$ as follows. Given $h, k \in H$ and $a \in A$, set

$$h \cdot a = \sum \gamma(h_{(1)})a\gamma'(h_{(2)}),$$

$$\begin{aligned} \omega(h, k) &= \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)}), \\ \omega'(h, k) &= \sum \gamma(h_{(1)}k_{(1)})\gamma'(k_{(2)})\gamma'(h_{(2)}). \end{aligned}$$

Before anything else, we must check that these elements lie in A , but this is quite simple:

$$\begin{aligned} \rho(h \cdot a) &= \sum \rho(\gamma(h_{(1)}))\rho(a)\rho(\gamma'(h_{(2)})) \\ &= \sum (\gamma(h_{(1)}) \otimes h_{(2)})(a \otimes 1_H)(\gamma'(h_{(4)}) \otimes S(h_{(3)})) \\ &= \sum (\gamma(h_{(1)})a\gamma'(h_{(4)}) \otimes h_{(2)}S(h_{(3)})) \\ &= \sum (\gamma(h_{(1)})a\gamma'(h_{(2)}) \otimes 1_H) = (h \cdot a) \otimes 1_H, \end{aligned}$$

and thus $a \in B^{coH} = A$. In an analogous fashion, one may check that both $\omega(h, k)$ and $\omega'(h, k)$ lie in A for every h, k in H . For instance,

$$\begin{aligned} \rho(\omega(h, k)) &= \sum (\gamma(h_{(1)}) \otimes h_{(2)})(\gamma(k_{(1)}) \otimes k_{(2)})(\gamma'(h_{(4)}k_{(4)}) \otimes S(h_{(3)}k_{(3)})) \\ &= \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)}) \otimes 1_H \\ &= \omega(h, k) \otimes 1_H, \end{aligned}$$

and similarly for $\omega'(h, k)$.

Note that, since $A = B^{coH}$, then $1_B = 1_A$, and since $\gamma(1_H) = 1_B = 1_A = \gamma'(1_H)$, we have $1_H \cdot a = a$ for all $a \in A$.

Next, given $h \in H$ and $a, b \in A$, we see that

$$\begin{aligned} h \cdot ab &= \sum \underbrace{\gamma(h_{(1)})a}_{\text{in } A} b\gamma'(h_{(2)}) \stackrel{(48)}{=} \sum \gamma(h_{(1)})a\gamma'(h_{(2)})\gamma(h_{(3)})b\gamma'(h_{(4)}) \\ &= \sum (h_{(1)} \cdot a)(h_{(2)} \cdot b). \end{aligned}$$

The partial action is twisted by (ω, ω') , since

$$\begin{aligned} h \cdot (k \cdot a) &= \sum \underbrace{\gamma(h_{(1)})\gamma(k_{(1)})a}_{\text{in } A} \gamma'(k_{(2)})\gamma'(h_{(2)}) \\ &\stackrel{(47)}{=} \sum \underbrace{\gamma(h_{(1)})\gamma(k_{(1)})}_{\text{in } A} a\gamma'(h_{(2)}k_{(2)})\gamma(h_{(3)}k_{(3)})\gamma'(k_{(4)})\gamma'(h_{(4)}) \\ &\stackrel{(46)}{=} \sum [\gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)})] \gamma(h_{(3)}k_{(3)})a\gamma'(h_{(4)}k_{(4)}) \\ &\quad \times [\gamma(h_{(5)}k_{(5)})\gamma'(k_{(6)})\gamma'(h_{(6)})] \\ &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)}) \end{aligned}$$

for every $a \in A$ and $h, k \in H$.

With respect to ω and ω' , first we observe that

$$\omega(h, 1_H) = \sum \gamma(h_{(1)})\gamma(1_H)\gamma'(h_{(2)}) = \gamma(h_{(1)})\gamma'(h_{(2)}) = h \cdot \mathbf{1}_A,$$

and also $\omega(1_H, h) = h \cdot \mathbf{1}_A$, showing that ω is normalized. Note, furthermore, that

$$(53) \quad \sum \omega(h_{(1)}, k)(h_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k) = \omega(h, k),$$

because $h \cdot \mathbf{1}_A = (\gamma * \gamma')(h)$ and $\mathbf{e} = \gamma * \gamma'$ is central in $\text{Hom}(H, A)$ by Remark 3. Therefore

$$\begin{aligned} \sum \omega(h_{(1)}, k)(h_{(2)} \cdot \mathbf{1}_A) &= \sum (h_{(1)} \cdot \mathbf{1}_A)\omega(h_{(2)}, k) \\ &= \sum \underbrace{\gamma(h_{(1)})\gamma'(h_{(2)})\gamma(h_{(3)})}_{\gamma(h_{(1)})\gamma'(h_{(2)})}\gamma(k_{(1)})\gamma'(h_{(4)}k_{(2)}) \\ &\stackrel{(43)}{=} \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)}) = \omega(h, k). \end{aligned}$$

Analogously, using (44), one shows that

$$(54) \quad \sum \omega'(h_{(1)}, k)(h_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)} \cdot \mathbf{1}_A)\omega'(h_{(2)}, k) = \omega'(h, k).$$

For $\omega * \omega'$ we have

$$\begin{aligned} (\omega * \omega')(h \otimes k) &= \sum \underbrace{\gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)})\gamma(h_{(3)}k_{(3)})}_{\gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)})}\gamma'(k_{(4)})\gamma'(h_{(4)}) \\ &\stackrel{(46)}{=} \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(k_{(2)})\gamma'(h_{(2)}) = h \cdot (k \cdot \mathbf{1}_A). \end{aligned}$$

For the evaluation of $\omega' * \omega$ we use (vi) and (vii) of Definition 3 to compute

$$\begin{aligned} (\omega' * \omega)(h \otimes k) &= \sum \gamma(h_{(1)}k_{(1)})\gamma'(k_{(2)})\underbrace{\gamma'(h_{(2)})\gamma(h_{(3)})}_{\gamma'(h_{(2)})\gamma(h_{(3)})}\gamma(k_{(3)})\gamma'(h_{(4)}k_{(4)}) \\ &= \sum \gamma(h_{(1)}k_{(1)})\underbrace{\gamma'(k_{(2)})\tilde{e}_{h_{(2)}}}_{\gamma'(k_{(2)})\tilde{e}_{h_{(2)}}}\gamma(k_{(3)})\gamma'(h_{(3)}k_{(4)}) \\ &\stackrel{(vi)}{=} \sum \underbrace{\gamma(h_{(1)}k_{(1)})\tilde{e}_{h_{(2)}k_{(2)}}}_{\gamma(h_{(1)}k_{(1)})\tilde{e}_{h_{(2)}k_{(2)}}}\gamma'(k_{(3)})\gamma(k_{(4)})\gamma'(h_{(3)}k_{(5)}) \\ &\stackrel{(44)}{=} \sum \gamma(h_{(1)}k_{(1)})\gamma'(k_{(2)})\gamma(k_{(3)})\gamma'(h_{(2)}k_{(4)}) \\ &= \sum \underbrace{\gamma(h_{(1)}k_{(1)})\tilde{e}_{k_{(2)}}}_{\gamma(h_{(1)}k_{(1)})\tilde{e}_{k_{(2)}}}\gamma'(h_{(2)}k_{(3)}) \\ &\stackrel{(vi)}{=} \sum e_{h_{(1)}}\gamma(h_{(2)}k_{(1)})\gamma'(h_{(3)}k_{(2)}) \\ &= \sum \gamma(h_{(1)})\gamma'(h_{(2)})\gamma(h_{(3)}k_{(1)})\gamma'(h_{(4)}k_{(2)}) \\ &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k \cdot \mathbf{1}_A). \end{aligned}$$

We use this to obtain the initial form of the twisting condition given in (3) of Definition 1:

$$\begin{aligned}
 \sum (h_{(1)} \cdot (k_{(1)} \cdot a))\omega(h_{(2)}, k_{(2)}) &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)})\omega(h_{(4)}, k_{(4)}) \\
 &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)(h_{(3)} \cdot \mathbf{1}_A)(h_{(4)}k_{(3)} \cdot \mathbf{1}_A) \\
 &= \sum \underbrace{\omega(h_{(1)}, k_{(1)})(h_{(2)} \cdot \mathbf{1}_A)(h_{(3)}k_{(2)} \cdot a)(h_{(4)}k_{(3)} \cdot \mathbf{1}_A)} \\
 &= \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a).
 \end{aligned}$$

Using again $\omega' * \omega$, we obtain the similar twisting equality for ω' :

$$(55) \quad \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot (k_{(2)} \cdot a)) = \sum (h_{(1)}k_{(1)} \cdot a)\omega'(h_{(2)}, k_{(2)}).$$

Indeed, multiplying the above obtained equality

$$h \cdot (k \cdot a) = \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)})$$

by ω' on the left, and using the convolution centrality of \mathbf{e} and (54), we have

$$\begin{aligned}
 \sum \omega'(h_{(1)}, k_{(1)})(h \cdot (k \cdot a)) &= \sum (\omega' * \omega)(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot a)\omega'(h_{(3)}, k_{(3)}) \\
 &= \sum (h_{(1)} \cdot \mathbf{1}_A) \underbrace{(h_{(2)}k_{(1)} \cdot \mathbf{1}_A)(h_{(3)}k_{(2)} \cdot a)} \omega'(h_{(4)}, k_{(3)}) \\
 &= \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k_{(1)} \cdot a)\omega'(h_{(3)}, k_{(2)}) \\
 &= \sum (h_{(1)}k_{(1)} \cdot a) \underbrace{(h_{(2)} \cdot \mathbf{1}_A)} \omega'(h_{(3)}, k_{(2)}) \\
 &= \sum (h_{(1)}k_{(1)} \cdot a)\omega'(h_{(2)}, k_{(2)}),
 \end{aligned}$$

as desired. Note now that by (v) and (iii) of Definition 3 we have

$$\begin{aligned}
 h \cdot (k \cdot \mathbf{1}_A) &= \sum \gamma(h_{(1)})e_k\gamma'(h_{(2)}) = \sum e_{h_{(1)}k}\gamma(h_{(2)})\gamma'(h_{(3)}) \\
 &= \sum (h_{(1)}k \cdot \mathbf{1}_A)(h_{(2)} \cdot \mathbf{1}_A) = \sum (h_{(1)} \cdot \mathbf{1}_A)(h_{(2)}k \cdot \mathbf{1}_A),
 \end{aligned}$$

which gives (iii) of Definition 2. Next, we see that ω absorbs $hk \cdot \mathbf{1}_A$ on the

right:

$$\begin{aligned} &\sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) \\ &= \sum \gamma(h_{(1)})\gamma(k_{(1)}) \underbrace{\gamma'(h_{(2)}k_{(2)})\gamma(h_{(3)}k_{(3)})\gamma'(h_{(4)}k_{(4)})}_{(44)} \\ &\stackrel{(44)}{=} \sum \gamma(h_{(1)})\gamma(k_{(1)})\gamma'(h_{(4)}k_{(4)}) = \omega(h, k), \end{aligned}$$

showing that (4) holds. Then using the twisting condition (3) we see that

$$\sum (h_{(1)} \cdot (k_{(1)} \cdot \mathbf{1}_A))\omega(h_{(2)}, k_{(2)}) = \sum \omega(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) = \omega(h, k).$$

Thus we have that $\omega(h, k)$ absorbs the elements $h \cdot k \cdot \mathbf{1}_A$, $h \cdot \mathbf{1}_A$ and $hk \cdot \mathbf{1}_A$ from both sides for any $h, k \in H$. In particular, ω is contained in the ideal $\langle f_1 * f_2 \rangle$.

Now, ω' absorbs $(h \cdot (k \cdot \mathbf{1}_A))$ on the right, which we see by using (vi) of Definition 3:

$$\begin{aligned} &\sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot (k_{(2)} \cdot \mathbf{1}_A)) \\ &= \sum \gamma(h_{(1)}k_{(1)})\gamma'(k_{(2)})\gamma'(h_{(2)})\gamma(h_{(3)})\gamma(k_{(3)})\gamma'(k_{(4)})\gamma'(h_{(4)}) \\ &= \sum \gamma(h_{(1)}k_{(1)}) \underbrace{\gamma'(k_{(2)})\tilde{e}_{h_{(2)}}}_{(43),(44)} \gamma(k_{(3)})\gamma'(k_{(4)})\gamma'(h_{(3)}) \\ &\stackrel{\text{Def. 3. (vi)}}{=} \sum \underbrace{\gamma(h_{(1)}k_{(1)})\tilde{e}_{h_{(2)}k_{(2)}}}_{(43),(44)} \underbrace{\gamma'(k_{(3)})\gamma(k_{(4)})\gamma'(k_{(5)})\gamma'(h_{(3)})}_{(43),(44)} \\ &\stackrel{(43),(44)}{=} \sum \gamma(h_{(1)}k_{(1)})\gamma'(k_{(2)})\gamma'(h_{(2)}) = \omega'(h, k). \end{aligned}$$

By (54) and the convolution centrality of \mathbf{e} , this implies

$$\sum \omega'(h_{(1)}, k_{(1)})(h_{(2)}k_{(2)} \cdot \mathbf{1}_A) = \omega'(h, k),$$

and, moreover, it follows using (55) that

$$\omega'(h, k) = \sum \omega'(h_{(1)}, k_{(1)})(h_{(2)} \cdot (k_{(2)} \cdot \mathbf{1}_A)) = \sum (h_{(1)}k_{(1)} \cdot \mathbf{1}_A)\omega'(h_{(2)}, k_{(2)}).$$

Consequently, ω' also absorbs the elements $h \cdot k \cdot \mathbf{1}_A$, $h \cdot \mathbf{1}_A$ and $hk \cdot \mathbf{1}_A$ from both sides. In particular, (29) is satisfied, i.e. ω' belongs to $\langle f_1 * f_2 \rangle$.

We check the cocycle equality (17) for ω , taking into account that $\gamma' * \gamma$ commutes with each element of A , as follows:

$$\begin{aligned} &\sum [h_{(1)} \cdot \omega(k_{(1)}, l_{(1)})] \omega(h_{(2)}, k_{(2)}l_{(2)}) \\ &= \sum \gamma(h_{(1)})\omega(k_{(1)}, l_{(1)})[(\gamma' * \gamma)(h_{(2)})]\gamma(k_{(3)}l_{(3)})\gamma'(h_{(3)}k_{(2)}l_{(2)}) \\ &= \sum \underbrace{\gamma(h_{(1)})[(\gamma' * \gamma)(h_{(2)})]}_{(17)} \omega(k_{(1)}, l_{(1)})\gamma(k_{(3)}l_{(3)})\gamma'(h_{(3)}k_{(2)}l_{(2)}) \end{aligned}$$

$$\begin{aligned}
 &\stackrel{(43)}{=} \sum \gamma(h_{(1)}) \underbrace{\gamma(k_{(1)})\gamma(l_{(1)})\gamma'(k_{(2)}l_{(2)})\gamma(k_{(3)}l_{(3)})}_{\gamma'(h_{(2)}k_{(4)}l_{(4)})} \\
 &\stackrel{(46)}{=} \sum \underbrace{\gamma(h_{(1)}) \gamma(k_{(1)}) \gamma(l_{(1)})}_{\gamma'(h_{(2)}k_{(2)}l_{(2)})} \\
 &\stackrel{(46)}{=} \sum [\gamma(h_{(1)}) \gamma(k_{(1)})\gamma'(h_{(2)}k_{(2)})][\gamma(h_{(3)}k_{(3)})\gamma(l_{(1)})\gamma'(h_{(4)}k_{(4)}l_{(2)})] \\
 &= \sum \omega(h_{(1)}, k_{(1)}) \omega(h_{(2)}k_{(2)}, l).
 \end{aligned}$$

This completes the proof of the fact that $A = (A, \cdot, \omega, \omega')$ is a symmetric twisted partial H -module algebra.

Finally, we claim that

$$\begin{aligned}
 \Phi : A\#_{(\alpha,\omega)}H &\rightarrow B \\
 a\#h &\mapsto a\gamma(h)
 \end{aligned}$$

is an algebra isomorphism, with inverse given by

$$\begin{aligned}
 \Psi : B &\rightarrow A\#_{(\alpha,\omega)}H \\
 b &\mapsto \sum b_{(0)}\gamma'(b_{(1)})\#b_{(2)}.
 \end{aligned}$$

In fact, Φ is an algebra map since it obviously takes unity to unity and

$$\begin{aligned}
 \Phi(a\#h)\Phi(b\#k) &= a\gamma(h)b\gamma(k) \\
 &\stackrel{(48)}{=} \sum a\gamma(h_{(1)})b\gamma'(h_{(2)})\gamma(h_{(3)})\gamma(k) \\
 &\stackrel{(46)}{=} \sum a[\gamma(h_{(1)})b\gamma'(h_{(2)})][\gamma(h_{(3)})\gamma(k_{(1)})\gamma'(h_{(4)}k_{(2)})] \\
 &\quad \times \gamma(h_{(5)}k_{(3)}) \\
 &= \sum a(h_{(1)} \cdot b)\omega(h_{(2)} \otimes k_{(1)})\gamma(h_{(3)}k_{(2)}) \\
 &= \Phi\left(\sum a(h_{(1)} \cdot b)\omega(h_{(2)} \otimes k_{(1)})\right)\#h_{(3)}k_{(2)} \\
 &= \Phi((a\#h)\Phi(b\#k))
 \end{aligned}$$

In order to prove that $\Psi = \Phi^{-1}$, first note that $\sum b_{(0)}\gamma'(b_{(1)})$ lies in A , because item (ii) of Definition 3 implies that

$$\begin{aligned}
 \rho\left(\sum b_{(0)}\gamma'(b_{(1)})\right) &= \sum \rho(b_{(0)})(\rho \circ \gamma')(b_{(1)}) \\
 &= \sum b_{(0)}(\gamma'(b_{(3)}) \otimes b_{(1)})S(b_{(2)}) \\
 &= \sum b_{(0)}\gamma(b_{(1)}) \otimes 1_H.
 \end{aligned}$$

Now, $\Phi\Psi = \text{Id}_B$ is just condition (iv) of Definition 3. For the other composition, given $a\#h \in A\#_{(\alpha,\omega)}H$, since γ is a comodule morphism and $A = B^{\text{co}H}$ it follows that

$$\begin{aligned}\Psi(\Phi(a\#h)) &= \Psi(a\gamma(h)) = \sum a\gamma(h_{(1)})\gamma'(h_{(2)})\#h_{(3)} \\ &= \sum a(h_{(1)} \cdot \mathbf{1}_A)\#h_{(2)} = a\#h. \quad \blacksquare\end{aligned}$$

References

- [1] E. Abe, *Hopf Algebras*, Cambridge University Press, Cambridge–New York, 1980.
- [2] M. M. S. Alves and E. Batista, *Enveloping actions for partial Hopf actions*, *Communications in Algebra* **38** (2010), 2872–2902.
- [3] M. M. S. Alves and E. Batista, *Partial Hopf actions, partial invariants and a Morita context*, *Algebra and Discrete Mathematics* **3** (2009), 1–19.
- [4] M. M. S. Alves and E. Batista, *Globalization theorems for partial Hopf (co)actions and some of their applications*, *Contemporary Mathematics* **537** (2011), 13–30.
- [5] D. Bagio, J. Lazzarin and A. Paques, *Crossed products by twisted partial actions: separability, semisimplicity and Frobenius properties*, *Communications in Algebra* **38** (2010), 496–508.
- [6] R. J. Blattner, M. Cohen and S. Montgomery, *Crossed products and inner actions of Hopf algebras*, *Transactions of the American Mathematical Society* **208** (1986), 671–711.
- [7] G. Böhm and J. Vercruyssen, *Morita theory for coring extensions and cleft bicomodules*, *Advances in Mathematics* **209** (2007), 611–648.
- [8] T. Brzeziński, *Descent cohomology and corings*, *Communications in Algebra* **36** (2008), 1894–1900.
- [9] S. Caenepeel and E. D. Groot, *Galois corings applied to partial Galois theory*, *Proc. ICMA-2004*, Kuwait University 2005, pp. 117–134.
- [10] S. Caenepeel and K. Janssen, *Partial (co)actions of Hopf algebras and partial Hopf-Galois theory*, *Communications in Algebra* **36** (2008), 2923–2946.
- [11] M. Dokuchaev, *Partial actions: a survey*, *Contemporary Mathematics* **537** (2011), 173–184.
- [12] M. Dokuchaev and R. Exel, *Associativity of crossed products by partial actions, enveloping actions and partial representations*, *Transactions of the American Mathematical Society* **357** (2005), 1931–1952.
- [13] M. Dokuchaev, R. Exel and P. Piccione, *Partial representations and partial group algebras*, *Journal of Algebra*, **226** (2000), 251–268.
- [14] M. Dokuchaev, R. Exel and J. J. Simón, *Crossed products by twisted partial actions and graded algebras*, *Journal of Algebra* **320** (2008), 3278–3310.
- [15] M. Dokuchaev, R. Exel and J. J. Simón, *Globalization of twisted partial actions*, *Transactions of the American Mathematical Society* **362** (2010), 4137–4160.
- [16] M. Dokuchaev, M. Ferrero and A. Paques, *Partial Actions and Galois Theory*, *Journal of Pure and Applied Algebra* **208** (2007), 77–87.

- [17] M. Dokuchaev and B. Novikov, *Partial projective representations and partial actions*, Journal of Pure and Applied Algebra **214** (2010), 251–268.
- [18] M. Dokuchaev and B. Novikov, *Partial projective representations and partial actions II*, Journal of Pure and Applied Algebra **216** (2012), 438–455.
- [19] R. Exel, *Circle actions on C^* -algebras, partial automorphisms and generalized Pimsner-Voiculescu exact sequences*, Journal of Functional Analysis **122** (1994), 361–401.
- [20] R. Exel, *The Bunce–Deddens algebras as crossed products by partial automorphisms*, Bol. Soc. Brasil. Mat. (N.S.), **25** (1994), 173–179.
- [21] R. Exel, *Approximately finite C^* -algebras and partial automorphisms*, Mathematica Scandinavica **77** (1995), 281–288.
- [22] R. Exel, *Twisted partial actions: a classification of regular C^* -algebraic bundles*, Proceedings London Mathematical Society **74** (1997), 417–443.
- [23] R. Exel, *Partial actions of groups and actions of inverse semigroups*, Proceedings of the American Mathematical Society **126** (1998), 3481–3494.
- [24] R. Exel, *Hecke algebras for protonormal subgroups*, Journal of Algebra **320** (2008), 1771–1813.
- [25] R. Exel and M. Laca, *Cuntz–Krieger Algebras for Infinite Matrices*, Journal für die Reine und Angewandte Mathematik **512** (1999), 119–172.
- [26] R. Exel, M. Laca and J. Quigg, *Partial dynamical systems and C^* -algebras generated by partial isometries*, The Journal of Operator Theory **47** (2002), 169–186.
- [27] K. McClanahan, *K -theory for partial crossed products by discrete groups*, Journal of Functional Analysis **130** (1995), 77–117.
- [28] S. Majid, *Foundations of Quantum Group Theory*, Cambridge University Press, Cambridge, 1995.
- [29] S. Montgomery, *Hopf Algebras and Their Actions on Rings*, CBMS Regional Conference Series in Mathematics, Vol. 82, American Mathematical Society, Providence, RI, 1993.
- [30] A. Paques and A. Sant’Ana, *When is a crossed product by a twisted partial action is Azumaya?*, Communications in Algebra **38** (2010), 1093–1103.
- [31] J. C. Quigg and I. Raeburn, *Characterisations of crossed products by partial actions*, Journal of Operator Theory **37** (1997), 311–340.
- [32] W. C. Waterhouse, *Introduction to Affine Group Schemes*, Graduate Texts in Mathematics, Vol. 66, Springer, New York–Berlin, 1979.