

THE INDEPENDENT NEIGHBORHOODS PROCESS

BY

TOM BOHMAN*

*Department of Mathematical Sciences, Carnegie Mellon University
Pittsburgh, PA 15213, USA
e-mail: tbohman@math.cmu.edu*

AND

DHRUV MUBAYI**

*Department of Mathematics, Statistics, and Computer Science
University of Illinois at Chicago, Chicago, IL 60607, USA
e-mail: mubayi@uic.edu*

AND

MICHAEL PICOLLELLI

*Department of Mathematics, California State University San Marcos
San Marcos, CA 92096, USA
e-mail: mpicollelli@csusm.edu*

ABSTRACT

A triangle $T^{(r)}$ in an r -uniform hypergraph is a set of $r+1$ edges such that r of them share a common $(r-1)$ -set of vertices and the last edge contains the remaining vertex from each of the first r edges. Our main result is that the random greedy triangle-free process on n points terminates in an r -uniform hypergraph with independence number $O((n \log n)^{1/r})$. As a consequence, using recent results on independent sets in hypergraphs, the Ramsey number $r(T^{(r)}, K_s^{(r)})$ has order of magnitude $s^r / \log s$. This answers questions posed in [4, 10] and generalizes the celebrated results of Ajtai–Komlós–Szemerédi [1] and Kim [9] to hypergraphs.

* Research supported by NSF grant DMS-1100215.

** Research supported by NSF grant DMS-1300138.

Received August 7, 2014 and in revised form March 17, 2015

1. Introduction

An r -uniform hypergraph H (r -graph for short) is a collection of r -element subsets of a vertex set $V(H)$. Given r -graphs G and H , the Ramsey number $r(G, H)$ is the minimum n such that every red/blue-edge coloring of the complete r -graph $K_n^{(r)} := \binom{[n]}{r}$ contains a red copy of G or a blue copy of H (often we will write K_n for $K_n^{(r)}$). Determining these numbers for graphs ($r = 2$) is known to be notoriously difficult, indeed the order of magnitude (for fixed t) of $r(K_t, K_s)$ is wide open when $t \geq 4$. The case $t = 3$ is one of the celebrated results in graph Ramsey theory:

$$(1) \quad r(K_3, K_s) = \Theta(s^2 / \log s).$$

The upper bound was proved by Ajtai–Komlós–Szemerédi [1] as one of the first applications of the semi-random method in combinatorics (simpler proofs now exist due to Shearer [12, 13]). The lower bound, due to Kim [9], was also achieved by using the semi-random or nibble method. More recently, the first author [3] showed that a lower bound for $r(K_3, K_s)$ could also be obtained by the triangle-free process, which is a random greedy algorithm. This settled a question of Spencer on the independence number of the triangle-free process. Still more recently, Bohman–Keevash [6] and Fiz Pontiveros–Griffiths–Morris [8] have analyzed the triangle-free process more carefully and improved the constants obtained so that the gap between the upper and lower bounds for $r(K_3, K_s)$ is now asymptotically a multiplicative factor of 4.

Given the difficulty of these basic questions in graph Ramsey theory, one would expect that the corresponding questions for hypergraphs are hopeless. This is not always the case. Hypergraphs behave quite differently for asymmetric Ramsey problems; for example, there exist $K_4^{(3)}$ -free 3-graphs on n points with independence number of order $\log n$, so $r(K_4^{(3)}, K_s^{(3)})$ is exponential in s unlike the graph case. Consequently, to obtain r -graph results parallel to (1), one must consider problems $r(G, K_s)$ where G is much sparser than a complete graph. A recent result in this vein due to Kostochka–Mubayi–Verstraëte [10] is that there are positive constants c_1, c_2 with

$$\frac{c_1 s^{3/2}}{(\log s)^{3/4}} < r(C_3^{(3)}, K_s^{(3)}) < c_2 s^{3/2}$$

where $C_3^{(3)}$ is the loose triangle, comprising 3 edges that have pairwise intersections of size one and have no point in common. The authors in [10] conjectured

that $r(C_3^{(3)}, K_s^{(3)}) = o(s^{3/2})$ and the order of magnitude remains open. Another result of this type for hypergraphs due to Phelps and Rödl [11] is that $r(P_t^{(3)}, K_s^{(3)}) = \Theta(s^2/\log s)$, where $P_t^{(3)}$ is the tight path with t edges. Recently, the second author and Cooper [7] prove that for fixed $t \geq 4$, the behavior of this Ramsey number changes and we have $r(P_t^{(3)}, K_s^{(3)}) = \Theta(s^2)$; the growth rate for $t = 3$ remains open. These are the only nontrivial hypergraph results of polynomial Ramsey numbers, and in this paper we add to this list with an extension of (1).

Definition 1: An r -uniform triangle $T^{(r)}$ is a set of $r + 1$ edges b_1, \dots, b_r, a with $b_i \cap b_j = R$ for all $i < j$ where $|R| = r - 1$ and $a = \bigcup_i (b_i - R)$. In other words, r of the edges share a common $(r - 1)$ -set of vertices, and the last edge contains the remaining point in all these previous edges.

When $r = 2$, then $T^{(2)} = K_3$, so in this sense $T^{(r)}$ is a generalization of a graph triangle. We may view a $T^{(r)}$ -free r -graph as one in which all neighborhoods are independent sets, where the neighborhood of an $R \in \binom{V(H)}{r-1}$ is $\{x : R \cup \{x\} \in H\}$. Frieze and the first two authors [4] proved that for fixed $r \geq 2$, there are positive constants c_1 and c_2 with

$$c_1 \frac{s^r}{(\log s)^{r/(r-1)}} < r(T^{(r)}, K_s^{(r)}) < c_2 s^r.$$

They conjectured that the upper bound could be improved to $o(s^r)$ and believed that the log factor in the lower bound could also be improved. Kostochka–Mubayi–Verstraëte [10] partially achieved this by improving the upper bound to

$$r(T^{(r)}, K_s^{(r)}) = O(s^r / \log r)$$

and believed that the log factor was optimal.

In this paper we verify this assertion by analyzing the $T^{(r)}$ -free (hyper)graph process. This process begins with an empty hypergraph $G(0)$ on n vertices. Given $G(i - 1)$, the hypergraph $G(i)$ is then formed by adding an edge e_i selected uniformly at random from the r -sets of vertices which neither form edges of $G(i - 1)$ nor create a copy of $T^{(r)}$ in the hypergraph $G(i - 1) + e_i$. The process terminates with a maximal $T^{(r)}$ -free graph $G(M)$ with a random number M of edges. Our main result is the following:

THEOREM 1: *For $r \geq 3$ fixed, the $T^{(r)}$ -free process on n points produces an r -graph with independence number $O((n \log n)^{1/r})$ with high probability.*

This result together with the aforementioned result of Kostochka–Mubayi–Verstraëte give the following generalization of (1) to hypergraphs.

COROLLARY 2: *For fixed $r \geq 3$ there are positive constants c_1 and c_2 with*

$$c_1 \frac{s^r}{\log s} < r(T^{(r)}, K_s^{(r)}) < c_2 \frac{s^r}{\log s}.$$

Graph processes that iteratively add edges chosen uniformly at random subject to the condition that some graph property is maintained have been used to generate interesting combinatorial objects in a number of contexts. In addition to the lower bound on the Ramsey number $r(K_3, K_s)$ given by the triangle-free graph process (discussed above), the H -free graph process gives the best known lower bound on the Ramsey number $r(K_t, K_s)$ for $t \geq 4$ fixed and the best known lower bound on the Turán numbers for some bipartite graphs [5]. The process that forms a subset of \mathbb{Z}_n by iteratively choosing elements to be members of the set uniformly at random subject to the condition that the set does not contain a k -term arithmetic progression produces a set that has interesting properties with respect to the Gowers norm [2].

The $T^{(r)}$ -free (hyper)graph process can be viewed as an instance of the random greedy hypergraph independent set process. Let H be a hypergraph. An independent set in H is a set of vertices that contains no edge of H . The random greedy independent set process forms such a set by starting with an empty set of vertices and iteratively choosing vertices uniformly at random subject to the condition that the set of chosen vertices continues to be an independent set. We study the random greedy independent set process for the hypergraph $\mathcal{H}_{T^{(r)}}$ which has vertex set $\binom{[n]}{r}$ and edge set consisting of all copies of $T^{(r)}$ on vertex set $[n]$. Note that, since an independent set in $\mathcal{H}_{T^{(r)}}$ gives a $T^{(r)}$ -free r -graph on point set $[n]$, the random greedy independent set process on $\mathcal{H}_{T^{(r)}}$ is equivalent to the $T^{(r)}$ -free process. Our analysis of the $T^{(r)}$ -free process is based on recent work on the random greedy hypergraph independent set process due to Bennett and Bohman [2].

The remainder of the paper is organized as follows. In the following Section we establish some notation and recall the necessary facts from [2]. The proof of Theorem 1 is given in the Section that follows, modulo the proofs of some technical lemmas. These lemmas are proved in the final Section by application of the differential equations method for proving dynamic concentration.

2. Preliminaries

Let \mathcal{H} be a hypergraph on vertex set $V = V(\mathcal{H})$. For each set of vertices $A \subseteq V$, let $N_{\mathcal{H}}(A)$ denote the **neighborhood** of A in \mathcal{H} , the family of all sets $Y \subseteq V \setminus A$ for which $A \cup Y \in \mathcal{H}$. We then define the **degree** of A in \mathcal{H} to be $d_{\mathcal{H}}(A) = |N_{\mathcal{H}}(A)|$. For a nonnegative integer a , we define $\Delta_a(\mathcal{H})$ to be the maximum of $d_{\mathcal{H}}(A)$ over all $A \in \binom{V}{a}$. Next, for a pair of (not necessarily disjoint) sets $A, B \subseteq V$, we define the **codegree** of A and B to be the number of sets $X \subseteq V \setminus (A \cup B)$ for which $A \cup X, B \cup X$ both lie in \mathcal{H} .

Recall that we define $G(i)$ to be the r -graph produced through i steps of the $T^{(r)}$ -free process. We let \mathcal{F}_i denote the natural filtration determined by the process (see [3], for example). We also simplify our notation somewhat and write $N_i(A)$ in place of $N_{G(i)}(A)$, $d_i(A)$ in place of $d_{G(i)}(A)$, etc., when appropriate.

The r -graph $G(i)$ partitions $\binom{[n]}{r}$ into three sets $E(i), O(i), C(i)$. The set $E(i)$ is simply the set of i edges chosen in the first i steps of the process. The set $O(i)$ consists of the **open** r -sets: all $e \in \binom{[n]}{r} \setminus E(i)$ for which $G(i) + e$ is $T^{(r)}$ -free. The r -sets in $C(i) := \binom{[n]}{r} \setminus (E(i) \cup O(i))$ are **closed**. Finally, for each open r -set $e \in O(i)$, we define the set $C_e(i)$ to consist of all open r -sets $f \in O(i)$ such that the graph $G(i) + e + f$ contains a copy of $T^{(r)}$ using both e and f as edges. (That is, $C_e(i)$ consists of the open r -sets whose selection as the next edge e_{i+1} would result in $e \in C(i+1)$.)

We now introduce some notation in preparation for our application of the results in [2]. Set

$$N := \binom{n}{r}, \quad D := (r + 1) \cdot \binom{n - r}{r - 1}, \quad s := \frac{N}{D^{1/r}}.$$

Note that N is the size of the vertex set of the hypergraph $\mathcal{H}_{T^{(r)}}$ and D is the vertex degree of $\mathcal{H}_{T^{(r)}}$ (in other words, every r -set in $[n]$ is in D copies of $T^{(r)}$). The parameter s is the ‘scaling’ for the length of the process. This choice is motivated by the heuristic that $E(i)$ should be pseudorandom; that is, $E(i)$ should resemble in some ways a collection of r -sets chosen uniformly at random (without any further condition). If this is indeed the case, then the probability that a given r -set is open would be roughly

$$\left(1 - \left(\frac{i}{N}\right)^r\right)^D \approx \exp \left\{ - \left(\frac{i}{N}\right)^r D \right\}$$

and a substantial number of r -sets are closed when roughly s edges have been added. In order to discuss the evolution in more detail, we pass to a limit by introducing a continuous time variable t where $t = t(i) = i/s$.

The evolution of key parameters of the process closely follows trajectories given by the functions

$$q(t) := \exp\{-t^r\} \quad \text{and} \quad c(t) := -q'(t) = rt^{r-1}q(t).$$

We introduce small constants ζ, γ such that $\zeta \ll \gamma \ll 1/r$. (The notation $\alpha \ll \beta$ here means that α is chosen to be sufficiently small relative to β .) The point where we stop tracking the process is given by

$$i_{\max} := \zeta \cdot ND^{-1/r}(\log^{1/r} N) \quad \text{and} \quad t_{\max} := i_{\max}/s = \zeta \log^{1/r} N.$$

For $i^* \geq 0$, let \mathcal{T}_{i^*} denote the event that the following estimates hold for all steps $0 \leq i \leq i^*$:

$$(2) \quad |O(i)| = (q(t) \pm N^{-\gamma})N,$$

and for every open r -set $e \in O(i)$

$$(3) \quad |C_e(i)| = (c(t) \pm N^{-\gamma})D^{1/r}.$$

It follows from the results of Bennett and Bohman that $\mathcal{T}_{i_{\max}}$ holds with high probability. We now recall the results of [2] in order to verify that this is indeed the case.

Bennett and Bohman studied the random greedy independent set process applied to an \hat{r} -uniform, D -regular hypergraph \mathcal{H} . As we discuss above, the $T^{(r)}$ -free process is identical to the random greedy independent set process on the hypergraph $\mathcal{H}_{T^{(r)}}$. Note that $\mathcal{H}_{T^{(r)}}$ is $(r + 1)$ -uniform, and so in our application of Bennett–Bohman we have $\hat{r} = r + 1$. Define the $(\hat{r} - 1)$ -codegree of a pair of distinct vertices v, v' in the hypergraph \mathcal{H} to be the number of edges $e, e' \in \mathcal{H}$ such that $v \in e \setminus e', v' \in e' \setminus e$ and $|e \cap e'| = \hat{r} - 1$. We let $\Gamma(\mathcal{H})$ be the maximum $(\hat{r} - 1)$ -codegree of \mathcal{H} .

THEOREM 3 (Theorem 1.1 of [2]): *Let \hat{r} and $\epsilon > 0$ be fixed. Let \mathcal{H} be an \hat{r} -uniform, D -regular hypergraph on N vertices such that $D > N^\epsilon$. If*

$$(4) \quad \Delta_\ell(\mathcal{H}) < D^{\frac{\hat{r}-\ell}{\hat{r}-1}-\epsilon} \quad \text{for } \ell = 2, \dots, \hat{r} - 1$$

and $\Gamma(\mathcal{H}) < D^{1-\epsilon}$; then the random greedy independent set algorithm produces an independent set I in \mathcal{H} with

$$(5) \quad |I| = \Omega\left(N \cdot \left(\frac{\log N}{D}\right)^{\frac{1}{r-1}}\right)$$

with probability $1 - \exp\{-N^{\Omega(1)}\}$.

Note that $\Delta_\ell(\mathcal{H}_{T^{(r)}}) = \Theta(n^{r-\ell}) = \Theta(n^{\hat{r}-1-\ell})$ and $\Gamma(\mathcal{H}_{T^{(r)}}) = 0$, and therefore the work of Bennett–Bohman applies to the $T^{(r)}$ -free process. We require more detailed information from [2]. Theorem 3 is proved by tracking key parameters of the process. These include the following, where we let $I(i)$ be the independent set that has been formed through i steps of the random greedy independent set algorithm:

- The size of the set $V(i)$ of vertices of \mathcal{H} that remain available for inclusion in the independent set after i vertices have been added to the independent set. Note that for the $T^{(r)}$ -free process, the set of vertices that remain available for inclusion in the independent set in $\mathcal{H}_{T^{(r)}}$ is precisely the collection of open edges $O(i)$.
- For every vertex v available at step i , the number $d_2(v, i)$ of available vertices $u \neq v$ with the property that there is some edge $e \in \mathcal{H}$ such that $u, v \in e$ and $|e \cap I(i)| = \hat{r} - 2$. Note that such an available vertex v in the vertex set of $\mathcal{H}_{T^{(r)}}$ is an open edge in the $T^{(r)}$ -free process, and the collection of vertices u that satisfy these conditions in $\mathcal{H}_{T^{(r)}}$ is the set of open edges in the $T^{(r)}$ -free process in the set $C_e(i)$.

Bounds on $|V(i)|$ and $d_2(v, i)$ are given in equations (8) and (9), respectively, of [2]. These bounds immediately give the estimates (2) and (3) quoted above. Note that $s_2 = s_2^+ - s_2^-$, $d_2 = d_2^+ - d_2^-$, and the error functions f_v, f_2 can be bounded above by D to an arbitrarily small constant, uniformly in t . Moreover, s_2 (defined on page 11 of [2] and translated to our notation) is equal to $rD^{1/r}t^{r-1}q$, which matches the main term in (3) as $c(t) = rt^{r-1}q(t)$.

We will also make use of the following fact regarding r -graphs that appear as subgraphs of the $T^{(r)}$ -free process.

LEMMA 4 (Lemma 5.1 of [2]): *Fix a constant L and suppose $e_1, \dots, e_L \in \binom{[n]}{r}$ form a $T^{(r)}$ -free hypergraph. Then for all steps $j \leq i_{\max}$,*

$$\mathbb{P}[\{e_1, \dots, e_L\} \subseteq E(j)] = (j/N)^L \cdot (1 + o(1)).$$

Note that the fact that $\mathcal{T}_{i_{\max}}$ holds with high probability does not prove that the independence number of $G(M)$ is $O((n \log n)^{1/r})$ with high probability. This is the main result in this work; it is proved below. Before commencing with the details of the proof, we briefly observe that the desired bound on the independence number of $G(M)$ can be viewed as a pseudorandom property of the r -graph $G(i)$. Indeed, if $G(i)$ resembles a collection of r -sets chosen uniformly at random, then the expected number of independent sets of size k would be

$$\binom{n}{k} \left(1 - \frac{\binom{k}{r}}{\binom{n}{r}}\right)^i = \exp \left\{ \Theta(k \log n) - \Theta\left(i \frac{k^r}{n^r}\right) \right\}.$$

If the process lasts through $i = \Theta(ND^{-1/r}(\log^{1/r} N)) = \Theta(n^{r-1+1/r} \log^{1/r} n)$ steps, then we would anticipate an independence number of $O((n \log n)^{1/r})$. In the remainder of the paper we make this heuristic calculation rigorous.

3. Independence number: Proof of Theorem 1

We expand the list of constants given in the previous section by introducing large constants κ and W , and small constant ϵ such that

$$(6) \quad \frac{1}{\kappa} \ll \zeta \ll \frac{1}{W} \ll \epsilon \ll \gamma.$$

In the course of the argument we introduce dynamic concentration phenomena that will be stated in terms of the error function

$$f(t) := \exp\{W(t^r + t)\}.$$

Define the constant $\lambda := \frac{\kappa - \gamma}{2}$, and then let

$$k := \kappa(n \log n)^{1/r} \quad \text{and} \quad \ell := \lambda(n \log n)^{1/r},$$

noting that as γ is small, $k \approx 2\ell$. Our aim is to show that the independence number of $G(i_{\max})$ is at most k with high probability. To do so, we will show that, provided κ is suitably large, w.h.p. for every step $0 \leq i \leq i_{\max}$, every k -element set of vertices has at least $\Omega(q(t)\binom{k}{r})$ open r -sets. As equation (2) establishes $(1+o(1))q(t)N$ open r -sets in total w.h.p., the probability that $\mathcal{T}_{i_{\max}}$ holds and a given k -set remains independent over all i_{\max} steps is then at most

$$\prod_{i=1}^{i_{\max}} \left(1 - \Omega\left(\frac{q(t)k^r}{q(t)N}\right)\right) = \left(1 - \Omega\left(\frac{\kappa^r \log n}{n^{r-1}}\right)\right)^{i_{\max}} = \exp\{-\zeta \kappa^r \cdot \Omega(n^{1/r} \log^{1+1/r} n)\},$$

where our $O(\cdot), \Omega(\cdot), \Theta(\cdot)$ notation does not suppress any constant that appears in (6). Since

$$n^k = \exp\{\kappa \cdot O(n^{1/r} \log^{1+1/r} n)\},$$

this suffices by the union bound, provided κ is suitably large with respect to r and ζ .

There is a significant obstacle to proving that every set of k vertices contains the ‘right’ number of open r -sets. Since the forbidden r -graph $T^{(r)}$ consists of an $(r - 1)$ -set (R) along with an edge (a) contained in its neighborhood, it follows that all r -sets within the neighborhood of an $(r - 1)$ -set in $G(i)$ must be closed. (That is, if $A \in \binom{[n]}{r-1}$ then $\binom{N_i(A)}{r} \subseteq C(i)$.) So a set of k vertices that has a large intersection with the neighborhood of an $(r - 1)$ -set does not have the ‘right’ number of open r -sets. To overcome this obstacle, we extend the argument in [3] for bounding the independence number of the triangle-free process. Our argument has two steps:

- (1) We apply the differential equations method for establishing dynamic concentration to show that unless a certain ‘bad’ condition occurs, a pair of disjoint ℓ -sets will have the ‘right’ number of open r -sets that are contained in the union of the pair of ℓ -sets and intersect both ℓ -sets, that is about $q(t) \cdot [\binom{2\ell}{r} - 2\binom{\ell}{r}]$ open r -sets. Note that $\binom{2\ell}{r} - 2\binom{\ell}{r} > \frac{1}{3} \binom{k}{r}$, say, as γ is small.
- (2) We then argue that w.h.p., every k -set contains a (disjoint) pair of ℓ -sets which is ‘good’, i.e., for which the bad condition does not occur.

We formalize this with the notion of r -sets which are open ‘with respect to’ a pair of disjoint ℓ -sets.

Definition 2: Fix a disjoint pair $A, B \in \binom{[n]}{\ell}$. The stopping time $\tau_{A,B}$ is the minimum of i_{\max} and the first step i for which there exists a $(r - 1)$ -set X such that

$$N_i(X) \cap A \neq \emptyset, \quad N_i(X) \cap B \neq \emptyset, \quad \text{and} \quad |N_i(X) \cap (A \cup B)| \geq k/n^{2\epsilon}.$$

Definition 3: For each step $i \geq 0$, we say that an r -set $e \subseteq A \cup B$ is **open with respect to the pair A, B in $G(i)$** if $e \cap A \neq \emptyset, e \cap B \neq \emptyset$, and either

- $e \in O(i)$ or
- $e \in O(i - 1) \cap C(i)$ and $i = \tau_{A,B}$.

Let $Q_{A,B}(i)$ count the number of r -sets which are open with respect to the pair A, B in $G(i)$.

LEMMA 5: *With high probability, for every disjoint pair $A, B \in \binom{[n]}{\ell}$ and all steps $0 \leq i \leq \tau_{A,B}$,*

$$(7) \quad Q_{A,B}(i) = \left(q(t) \pm \frac{f(t)}{n^\varepsilon} \right) \cdot \left[\binom{2\ell}{r} - 2 \binom{\ell}{r} \right].$$

LEMMA 6: *With high probability, for every step $0 \leq i < i_{\max}$ and every set $K \in \binom{[n]}{k}$, there exists a pair of disjoint ℓ -sets A, B contained in K for which $\tau_{A,B} > i$.*

Lemmas 5 and 6, respectively, complete steps 1 and 2 of the proof outlined above. The ‘bad’ condition for a pair A, B of disjoint ℓ -sets is the event that we have reached the stopping time $\tau_{A,B}$; that is, the bad condition is that there is some $(r - 1)$ -set whose neighborhood intersects both A and B and has large intersection with $A \cup B$. Note that if $i < \tau_{A,B}$, then $Q_{A,B}$ is equal to the number of open r -sets that are contained in $A \cup B$ and intersect both A and B . Thus, Lemma 5 says that if we do not have the ‘bad’ condition then we have the ‘right’ number of such sets. Lemma 6 then says that every k -set contains a pair disjoint pair A, B of ℓ -sets for which the ‘bad’ condition does not hold. Taken together, Lemmas 5 and 6 yield that w.h.p., for every step $0 \leq i < i_{\max}$, every k -set contains at least $q(t)(1 + o(1)) \left[\binom{2\ell}{r} - 2 \binom{\ell}{r} \right] = \Omega(q(t) \binom{k}{r})$ open r -sets, as required. We now prove Lemma 6 modulo the proof of Lemma 7 which bounds the maximum degree of an $(r - 1)$ -set. Lemmas 5 and 7 are proved in the next Section.

Proof of Lemma 6. We require a bound on the maximum degree of $(r - 1)$ -sets of vertices. For each step $i \geq 0$ let \mathcal{D}_i denote the event that

$$\Delta_{r-1}(G(i)) \leq \varepsilon(n \log n)^{1/(r-1)}.$$

LEMMA 7: $\mathcal{T}_{i_{\max}} \wedge \mathcal{D}_{i_{\max}}$ holds with high probability.

The proof of Lemma 7 is given in the next Section.

Fix a step $0 \leq i < i_{\max}$, and a set $K \in \binom{[n]}{k}$. Note that, by Lemma 7, we may assume that \mathcal{D}_i holds. We also note that the maximum co-degree of a pair of sets $A, B \in \binom{[n]}{r-1}$ is at most $5r$ with high probability. This follows from

Lemma 4 and the union bound:

$$(8) \quad \Pr \left(\exists A, B \in \binom{[n]}{r-1} \text{ with co-degree } 5r \right) \leq \binom{n}{r-1} \binom{n}{r-1} n^{5r} \left(\frac{i}{N} \right)^{10r} \\ = n^{8-3r+o(1)} = o(1).$$

Given these two facts (i.e., these degree and co-degree bounds for $(r-1)$ -sets), the remainder of the proof is deterministic.

To begin, define the set

$$\mathcal{X} := \left\{ X \in \binom{[n]}{r-1} : |N_i(X) \cap K| \geq k/n^{2\varepsilon} \right\}.$$

CLAIM 1: $|\mathcal{X}| < 2n^{2\varepsilon}$.

Proof. Suppose $\exists \mathcal{Y} \subseteq \mathcal{X}$ with $|\mathcal{Y}| = 2n^{2\varepsilon}$. Let $N = \bigcup_{Y \in \mathcal{Y}} (N_i(Y) \cap K)$. By inclusion-exclusion,

$$k \geq |N| \geq |\mathcal{Y}| \cdot (k/n^{2\varepsilon}) - |\mathcal{Y}|^2 5r \geq 2k - 20rn^{4\varepsilon},$$

a contradiction as ε is small and $k = n^{1/r+o(1)}$. ■

Next, we ‘discard’ from K the vertices which are common neighbors of $(r-1)$ -sets in \mathcal{X} : let

$$K_{\text{bad}} := \{v \in K : \exists X, Y \in \mathcal{X} \text{ with } X \neq Y \text{ and } v \in N_i(X) \cap N_i(Y)\}$$

and $K_{\text{good}} := K \setminus K_{\text{bad}}$. Then

$$|K_{\text{bad}}| \leq |\mathcal{X}|^2 5r \leq 20rn^{4\varepsilon} < \frac{\gamma}{2} \cdot (n \log n)^{1/r},$$

say, for large n .

We find disjoint ℓ -subsets A, B of K_{good} as follows, noting

$$|K_{\text{good}}| \geq 2\ell + (\gamma/2)(n \log n)^{1/r}.$$

For each subset $\mathcal{Y} \subseteq \mathcal{X}$, let

$$N(\mathcal{Y}) = \bigcup_{Y \in \mathcal{Y}} N_i(Y) \cap K_{\text{good}}.$$

Now, choose a maximal subset $\mathcal{X}^* \subseteq \mathcal{X}$ subject to $|N(\mathcal{X}^*)| \leq \ell$. If $\mathcal{X}^* = \mathcal{X}$, then let A, B be ℓ -sets satisfying $N(\mathcal{X}^*) \subseteq A \subseteq K_{\text{good}}$ and $B \subseteq K_{\text{good}} \setminus A$.

Otherwise, pick any set $X^* \in \mathcal{X} \setminus \mathcal{X}^*$, so

$$\ell < |N(\mathcal{X}^* \cup \{X^*\})| < \ell + \varepsilon(n \log n)^{1/r};$$

let $A \subseteq N(\mathcal{X}^* \cup \{X^*\})$ and $B \subseteq K_{\text{good}} \setminus N(\mathcal{X}^* \cup \{X^*\})$ be ℓ -sets.

Observe now that if $\mathcal{X}^* = \mathcal{X}$, then $N_i(X) \cap B = \emptyset$ for all $X \in \mathcal{X}$. Otherwise, if $X \in \mathcal{X}^* \cup \{X^*\}$, $N_i(X) \cap B = \emptyset$, but if $X \in \mathcal{X} \setminus (\mathcal{X}^* \cup \{X^*\})$, then $N_i(X) \cap A = \emptyset$ as we are working within K_{good} . In either case, for every $(r - 1)$ -set X for which $|N_i(X) \cap (A \cup B)| \geq k/n^{2^\varepsilon}$ holds, either $N_i(X) \cap A = \emptyset$ or $N_i(X) \cap B = \emptyset$, and $\tau_{A,B} > i$ follows. ■

4. Dynamic Concentration

In this section we prove Lemmas 5 and 7. Both of these statements assert dynamic concentration of key parameters of the $T^{(r)}$ -free process. We apply the differential equations method for proving dynamic concentration, which we now briefly sketch.

Suppose we have a combinatorial stochastic process based on a ground set of size n that generates a natural filtration $\mathcal{F}_0, \mathcal{F}_1, \dots$. Suppose further that we have a sequence of random variables A_0, A_1, \dots and that we would like to prove a dynamic concentration statement of the form

$$(9) \quad A_i \leq T_i + E_i \quad \text{for all } 0 \leq i \leq m(n) \quad \text{with high probability,}$$

where T_0, T_1, \dots is the expected trajectory of the sequence of random variables A_i and E_0, E_1, \dots is a sequence of error functions. (One is often interested in proving a lower bound on A_i in conjunction with (9). The argument for proving this is essentially the same as the upper bound argument that we discuss here.) We often make this statement in the context of a limit that we define in terms of a continuous time t given by $t = i/s$ where s is the **time scaling** of the process. The limit of the expected trajectory is determined by setting

$$T_i = f(t)S(n)$$

where $S = S(n)$ is the **order scaling** of the random variable A_i . Given these assumptions we should have

$$\mathbb{E}[A_{i+1} - A_i \mid \mathcal{F}_i] = T_{i+1} - T_i = [f(t + 1/s) - f(t)]S \approx f'(t) \cdot \frac{S}{s}.$$

Thus the trajectory is determined by the expected one-step change in A_i .

We prove (9) by applying facts regarding the probability of large deviations in martingales with bounded differences. In particular, we consider the sequence

$$D_i = A_i - T_i - E_i.$$

Note that if we set $T_0 = A_0$ (which is often the natural initial condition), then $D_0 = -E_0$. If we can establish that the sequence D_i is a supermartingale and E_0 is sufficiently large, then it should be unlikely that D_i is ever positive, and (9) follows. In order to complete such a proof we show that the sequence D_i is a supermartingale, a fact that is sometimes called the **trend hypothesis** (see Wormald [14]). The trend hypothesis will often impose a condition that the sequence of error functions E_i is growing sufficiently quickly (i.e., the derivative of the limit of the error function is sufficiently large). We then show that the one-step changes in D_i are bounded in some way (this is sometimes called the **boundedness hypothesis**). This puts us in the position to apply a martingale inequality. In order to get good bounds from the martingale inequality one generally needs to make E_0 large.

In this section we appeal to the following pair of martingale inequalities (see [3]). For positive reals b, B , the sequence A_0, A_1, \dots is said to be (b, B) -**bounded** if $A_i - b \leq A_{i+1} \leq A_i + B$ for all $i \geq 0$.

LEMMA 8: *Suppose $b \leq B/10$ and $0 < a < bm$. If A_0, A_1, \dots is a (b, B) -bounded submartingale, then $\mathbb{P}[A_m \leq A_0 - a] \leq \exp\{-a^2/3bmB\}$.*

LEMMA 9: *Suppose $b \leq B/10$ and $0 < a < bm$. If A_0, A_1, \dots is a (b, B) -bounded supermartingale, then $\mathbb{P}[A_m \geq A_0 + a] \leq \exp\{-a^2/3bmB\}$.*

Our applications of these Lemmas make use of stopping times. Formally speaking, a stopping time is simply a positive integer-valued random variable τ for which $\{\tau \leq n\} \in \mathcal{F}_n$. In other words, τ is a stopping time if the event $\tau \leq n$ is determined by the first n steps of the process. We consider the stopped process $(D_{i \wedge \tau})$, where $x \wedge y := \min\{x, y\}$, in place of the sequence D_0, D_1, \dots . Our stopping time τ is the first step in the process when any condition on some short list of conditions fails to hold, where the condition $D_i \leq 0$ is one of the conditions in the list. Note that, since the variable $(D_{i \wedge \tau})$ does not change once we reach the stopping time τ , we can assume that all conditions in the list hold when we are proving the trend and boundedness hypotheses. Also note that if the stopping time τ' is simply the minimum of i_{\max} and the first step for which $D_i > 0$, then $\{D_{i_{\max} \wedge \tau'} > 0\}$ contains the event $\{\exists i \leq i_{\max} : D_i > 0\}$.

4.1. PROOF OF LEMMA 7. For each set $A \in \binom{[n]}{r-1}$ and step $i \geq 0$, let

$$O_A(i) := \{e \in O(i) : A \subseteq e\}, \quad \text{and} \quad Q_A(i) = |O_A(i)|.$$

We define sequences of random variables

$$\begin{aligned}
 Y_A^+(i) &:= q(t) \cdot n - Q_A(i) + f(t) \cdot n^{1-\varepsilon}, \\
 Y_A^-(i) &:= q(t) \cdot n - Q_A(i) - f(t) \cdot n^{1-\varepsilon}, \\
 Z_A(i) &:= d_i(A) - t \cdot D^{-1/r} n - f(t)q(t)^{-1} \cdot n^{1/r-\varepsilon}.
 \end{aligned}$$

Finally, we define the stopping time τ to be the minimum of $\binom{n}{r}$, the first step i where \mathcal{T}_i fails, or where any of $Y_A^+(i) < 0$, $Y_A^-(i) > 0$, or $Z_A(i) > 0$ holds for some $A \in \binom{[n]}{r-1}$.

To prove Lemma 7, we show that for each $A \in \binom{[n]}{r-1}$,

(10) $\mathbb{P}[Y_A^+(i_{\max} \wedge \tau) < 0] = o(n^{-(r-1)}),$

(11) $\mathbb{P}[Y_A^-(i_{\max} \wedge \tau) > 0] = o(n^{-(r-1)}),$ and

(12) $\mathbb{P}[Z_A(i_{\max} \wedge \tau) > 0] = o(n^{-(r-1)}).$

Consider the event $\tau \leq i_{\max}$. This event is the union of the event that $\mathcal{T}_{i_{\max}}$ fails and the event that there exists $A \in \binom{[n]}{r-1}$ such that $Y_A^+(i_{\max} \wedge \tau) < 0$ or $Y_A^-(i_{\max} \wedge \tau) > 0$ or $Z_A(i_{\max} \wedge \tau) > 0$. Since $\mathcal{T}_{i_{\max}}$ holds with high probability, it follows from (10)–(12) and the union bound that w.h.p. $\tau > i_{\max}$. In particular, $Z_A(i) \leq 0$ for all $(r - 1)$ -sets A and steps $0 \leq i \leq i_{\max}$. It then follows—since $\zeta \ll \min\{1/W, \varepsilon\}$ implies that we may bound $f(t_{\max}) < n^{\varepsilon/2}$, say—that we have

$$\begin{aligned}
 \Delta_{r-1}(G(i_{\max})) &\leq t_{\max} D^{-1/r} n + f(t_{\max}) n^{1/r-\varepsilon/2} \\
 &= \zeta \cdot O((n \log n)^{1/r}) \leq \varepsilon (n \log n)^{1/r},
 \end{aligned}$$

for n sufficiently large. (We remark in passing that the bounds on $Y_A^\pm(i)$ given when $i < \tau$ are necessary for our proof of the bounds on $Z_A(i)$.)

For the remainder of this argument, fix a set $A \in \binom{[n]}{r-1}$. We first prove (10) and (11).

CLAIM 2: For n sufficiently large, the variables $Y_A^+(0), \dots, Y_A^+(i_{\max} \wedge \tau)$ form an $(O(n/s), O(n^{1-\frac{1}{2r}}))$ -bounded submartingale, and the variables $Y_A^-(0), \dots, Y_A^-(i_{\max} \wedge \tau)$ form an $(O(n/s), O(n^{1-\frac{1}{2r}}))$ -bounded supermartingale.

Proof. We begin by fixing a step $0 \leq i \leq i_{\max}$, and we assume that $i < \tau$. Throughout we write $t = t(i)$, and note that $t(i + 1) = t + s^{-1}$ and that $s^{-1} = D^{1/r}/N = \Theta(n^{1-1/r-r})$.

To aid the calculations to follow, we begin by estimating the quantity

$$\Xi := f(t + s^{-1}) - f(t).$$

Since $f(t) = \exp(Wt^r + Wt)$, $f'(t)$ and $f''(t)$ are products of $f(t)$ with polynomials in t . As $\zeta \ll \max\{1/W, \varepsilon\}$, t_{\max} is polylogarithmic in n , and n is large, we have the crude bounds $f(t) \leq n^{\varepsilon/2}$ and $f''(t) \leq n^{o(1)}f'(t)$. Thus, by Taylor's Theorem,

$$(13) \quad \left| \Xi - \frac{f'(t)}{s} \right| = O\left(\frac{\max_{t^* \leq t_{\max}} f''(t^*)}{s^2}\right) = o\left(\frac{f'(t)}{s}\right).$$

Observe now that we may write

$$Y_A^\pm(i + 1) - Y_A^\pm(i) = (q(t + s^{-1}) - q(t)) \cdot n - (Q_A(i + 1) - Q_A(i)) \pm \Xi \cdot n^{1-\varepsilon}.$$

(Note that this stands for the pair of equations in which each \pm is replaced with $+$ or with $-$, respectively.) We begin by establishing the boundedness claims: it is routine to verify that $c(t)$ and $c'(t)$

are bounded over the reals, implying

$$(14) \quad |q(t + s^{-1}) - q(t) - c(t) \cdot s^{-1}| = O(s^{-2}),$$

and so

$$0 \geq (q(t + s^{-1}) - q(t)) \cdot n \geq -O\left(\frac{n}{s}\right).$$

As we have the bound $|f'(t)| = n^{\varepsilon/2+o(1)}$ and (13), we have $|\Xi| \cdot n^{1-\varepsilon} = o(n/s)$, and the lower bound in the boundedness claims follows. To establish the upper bounds, it remains to bound $Q_A(i) - Q_A(i + 1)$. Consider the 'next' edge $e_{i+1} \in O(i)$ and observe that

$$Q_A(i) - Q_A(i + 1) = |(\{e_{i+1}\} \cup C_{e_{i+1}}(i)) \cap O_A(i)|.$$

We bound $|C_{e_{i+1}}(i) \cap O_A(i)|$ by considering five cases depending on $|e_{i+1} \cap A|$:

Case 1: $|e_{i+1} \cap A| = 0$. Let $f \in O_A(i) \cap C_{e_{i+1}}(i)$: then $f = A \cup \{v\}$ for some vertex v , and since $G(i) + e_{i+1} + f$ contains a copy of $T^{(r)}$, $v \in e_{i+1}$ must hold. (Recall that every pair of edges in $T^{(r)}$ either shares exactly one or $r - 1$ vertices.) In this case, $|C_{e_{i+1}}(i) \cap O_A(i)| \leq |e_{i+1}| = r$.

Case 2: $|e_{i+1} \cap A| = r - 1$. In this case, we may write $e_{i+1} = A \cup \{u_1\}$. Now, let $f = A \cup \{v\} \in O_A(i) \cap C_{e_{i+1}}(i)$: since $f \cap e_{i+1} = A$ and $f \in C_{e_{i+1}}(i)$, there must exist vertices $u_2, \dots, u_{r-1} \in N_i(A)$ so that $\{u_1, \dots, u_{r-1}, v\} \in E(i)$. As then $v \in N_i(\{u_1, \dots, u_{r-1}\})$, we may bound the number of such choices of v

(and hence of f) in this case above by $\Delta_{r-1}(G(i))^{r-1} \leq \zeta^{r-1}(n \log n)^{(r-1)/r}$. (Note that the bound on the maximum degree follows as $Z_A(i) \leq 0$ since $i < \tau$.)

Case 3: $|e_{i+1} \cap A| = 1$. Write $A = \{x_1, \dots, x_{r-1}\}$, where we take $e_{i+1} \cap A = \{x_1\}$. Let $f = A \cup \{v\} \in C_{e_{i+1}}(i) \cap O_A(i)$, and suppose $v \notin e_{i+1}$ (as there are at most $r - 1$ such v), so $f \cap e_{i+1} = \{x_1\}$. Consider a copy of $T^{(r)}$ in $G(i) + e_{i+1} + f$ using both e_{i+1} and f as edges: without loss of generality, we may assume that one of e_{i+1}, f maps to the edge b_1 of $T^{(r)}$, the other to the edge a .

If e_{i+1} maps to b_1 , then the $(r - 1)$ -set $e_{i+1} \setminus \{x_1\}$ maps to the common intersection B of b_1, \dots, b_r . Consequently $v \in N_i(e_{i+1} \setminus \{x_1\})$ must hold, and so there are at most $\Delta_{r-1}(G(i))$ such r -sets $f \in C_{e_{i+1}}(i) \cap O_A(i)$.

Otherwise, if e_{i+1} maps to the edge a and f maps to b_1 , then $\{x_2, \dots, x_{r-1}, v\}$ maps to the common intersection B . Thus, for each $u \in e_{i+1} \setminus \{x_1\}$ we have $\{u, x_2, \dots, x_{r-1}, v\} \in E(i)$, implying $v \in N_i(\{u, x_2, \dots, x_{r-1}\})$ and (as e_{i+1} is fixed) there are again at most $\Delta_{r-1}(G(i))$ such choices of f . Thus, in this case we have $|C_{e_{i+1}}(i) \cap O_A(i)| \leq 2 + 2\Delta_{r-1}(G(i)) = n^{1/r+o(1)}$.

Case 4: $1 < |e_{i+1} \cap A| = r - 2$. Let $f = A \cup \{v\} \in O_A(i) \cap C_{e_{i+1}}(i)$. Since $|f \cap e_{i+1}| \geq |A \cap e_{i+1}| > 1$, $|f \cap e_{i+1}| = r - 1$ must hold, implying $v \in e_{i+1}$ and so $|O_A(i) \cap C_{e_{i+1}}(i)| \leq r$ as in Case 1.

Case 5: $2 \leq |e_{i+1} \cap A| \leq r - 3$. In this case, $|C_{e_{i+1}}(i) \cap O_A(i)| = 0$, as every $f \in O_A(i)$ satisfies $1 \leq |f \cap e_{i+1}| \leq r - 2$.

From the cases above it follows that

$$Q_A(i) - Q_A(i + 1) \leq n^{(r-1)/r+o(1)},$$

and combining the above bounds, it follows that the sequences

$$Y_A^\pm(0), \dots, Y_A^\pm(i_{\max} \wedge \tau)$$

are $(O(n/s), O(n^{1-\frac{1}{2r}}))$ -bounded.

We turn now to the sub- and supermartingale claims: all expectation calculations to follow are implicitly conditioned on the history of the process up to step i , and we recall that we assume $i < \tau$. For each open r -set $f \in O_A(i)$, we have $f \notin O_A(i + 1)$ if and only if $e_{i+1} \in C_f(i) \cup \{f\}$. Thus,

$$\mathbb{E} [Y_A^\pm((i + 1)) - Y_A^\pm(i)] = (q(t + s^{-1}) - q(t)) \cdot n + \sum_{f \in O_A(i)} \frac{|C_f(i)| + 1}{|O(i)|} \pm \Xi \cdot n^{1-\varepsilon}.$$

To establish the submartingale claim, consider the following chain of inequalities:

$$\begin{aligned} \sum_{f \in O_A(i)} \frac{|C_f(i)| + 1}{|O(i)|} &\geq (q(t) - f(t)n^{-\varepsilon}) \cdot n \cdot \frac{(c(t) - N^{-\gamma}) \cdot D^{1/r}}{(q(t) + N^{-\gamma}) \cdot N} \\ &= \left(1 - \frac{N^{-\gamma} + f(t)n^{-\varepsilon}}{q(t) + N^{-\gamma}}\right) (c(t) - N^{-\gamma}) \cdot \frac{n}{s} \\ &\geq (1 - 2q(t)^{-1}f(t)n^{-\varepsilon})(c(t) - N^{-\gamma}) \cdot \frac{n}{s} \\ &\geq (c(t) - 2c(t)q(t)^{-1}f(t)n^{-\varepsilon} - N^{-\gamma}) \cdot \frac{n}{s} \\ &\geq (c(t) - (2c(t)q(t)^{-1} + 1) \cdot f(t)n^{-\varepsilon}) \cdot \frac{n}{s} \end{aligned}$$

The first inequality follows from the bounds given by (2) and (3) on the event \mathcal{T}_i and as $Y_A^-(i) \leq 0$, since $i < \tau$. In the second and fourth inequalities we bounded $N^{-\gamma} < f(t)n^{-\varepsilon}$, valid as $f(t) \geq 1$ and $\varepsilon \ll \gamma$. Thus, applying this bound and (14) gives

$$\begin{aligned} \mathbb{E} [Y_A^+(i + 1) - Y_A^+(i)] &\geq \Xi \cdot n^{1-\varepsilon} - (2c(t)q(t)^{-1} + 1)f(t) \frac{n^{1-\varepsilon}}{s} - O\left(\frac{1}{s^2}\right) \\ &\geq \Xi \cdot n^{1-\varepsilon} - (2c(t)q(t)^{-1} + 2)f(t) \frac{n^{1-\varepsilon}}{s} \\ &= ((1 + o(1))f'(t) - (2c(t)q(t)^{-1} + 2)f(t)) \cdot \frac{n^{1-\varepsilon}}{s} \end{aligned}$$

by (13). Since $f'(t) = (Wrt^{r-1} + W)f(t)$ and $2c(t)q(t)^{-1} = 2rt^{r-1}$, this final bound is nonnegative for large n as W is large, and so $Y_A^+(0), \dots, Y_A^+(i_{\max} \wedge \tau)$ forms a submartingale.

We similarly bound $\mathbb{E} [Q_A(i) - Q_A(i + 1)]$ above to establish the supermartingale claim: as $1 < N^{-\gamma}D^{1/r}$ for large n , and as \mathcal{T}_i holds and $Y_A^+(i) \geq 0$,

$$\begin{aligned} \sum_{f \in O_A(i)} \frac{|C_f(i)| + 1}{|O(i)|} &\leq (q(t) + f(t)n^{-\varepsilon}) \cdot n \cdot \frac{(c(t) + 2N^{-\gamma}) \cdot D^{1/r}}{(q(t) - N^{-\gamma}) \cdot N} \\ &= \left(1 + \frac{N^{-\gamma} + f(t)n^{-\varepsilon}}{q(t) - N^{-\gamma}}\right) (c(t) + 2N^{-\gamma}) \cdot \frac{n}{s} \\ &\leq (1 + 4q(t)^{-1}f(t)n^{-\varepsilon})(c(t) + 2N^{-\gamma}) \cdot \frac{n}{s} \\ &\leq (c(t) + (4c(t)q(t)^{-1} + 4)f(t)n^{-\varepsilon}) \cdot \frac{n}{s} \end{aligned}$$

In addition to the bound $N^{-\gamma} \leq f(t)n^{-\varepsilon}$ used above, in the second inequality, we bounded $q(t) - N^{-\gamma} \geq q(t)/2$, and in the final we bounded

$$2N^{-\gamma}(1 + 4q(t)^{-1}f(t)n^{-\varepsilon}) \leq 4f(t)n^{-\varepsilon}$$

as

$$q(t)^{-1}f(t)n^{-\varepsilon} \leq 1$$

which holds as $2W\zeta^r < \varepsilon$ and n is large.

Thus,

$$\begin{aligned} \mathbb{E} [Y_A^-(i + 1) - Y_A^-(i)] &\leq -\Xi \cdot n^{1-\varepsilon} + (4c(t)q(t)^{-1} + 4)f(t)\frac{n^{1-\varepsilon}}{s} + O\left(\frac{1}{s^2}\right) \\ &\leq -\Xi \cdot n^{1-\varepsilon} + (4c(t)q(t)^{-1} + 5)f(t)\frac{n^{1-\varepsilon}}{s} \\ &= -(1 + o(1))f'(t) + (4c(t)q(t)^{-1} + 5)f(t) \cdot \frac{n^{1-\varepsilon}}{s}, \end{aligned}$$

and again, as W is large, this is strictly negative for n sufficiently large. Thus, the sequence $Y_A^-(0), \dots, Y_A^-(i_{\max} \wedge \tau)$ forms a supermartingale, completing the proof. ■

Since $Q_A(0) = n - r + 1$, $Y_A^+(0) = r - 1 + n^{1-\varepsilon}$ and $Y_A^-(0) = r - 1 - n^{1-\varepsilon}$. Applying Lemmas 8 and 9, respectively, we have

$$\begin{aligned} \mathbb{P}[Y_A^+(i_{\max} \wedge \tau) < 0] &\leq \exp\left\{-\Omega\left(\frac{n^{2-2\varepsilon}}{\frac{n}{s} \cdot \zeta s \log^{1/r} N \cdot n^{1-\frac{1}{2r}}}\right)\right\} \\ &= \exp\{-n^{\frac{1}{2r}-2\varepsilon+o(1)}\} \\ &< \exp\{-n^{\frac{1}{4r}}\} \end{aligned}$$

(valid for large n as ε is small), and an identical calculation yields

$$\mathbb{P}[Y_A^-(i_{\max} \wedge \tau) > 0] \leq \exp\{-n^{\frac{1}{4r}}\}.$$

We have established (10) and (11).

It remains to prove (12).

CLAIM 3: *The variables $Z_A(0), \dots, Z_A(i_{\max} \wedge \tau)$ form a $(2n/N, 2)$ -bounded supermartingale.*

Proof. We begin by fixing a step $0 \leq i \leq i_{\max}$, and we assume that $i < \tau$. Throughout we write $t = t(i)$. Let $f_1(t) = f(t)q(t)^{-1} = \exp((W + 1)t^r + Wt)$, and let $\Xi_1 := f_1(t + s^{-1}) - f_1(t)$.

By the same reasoning given in Claim 2, we may bound $|f_1(t)| < n^{\varepsilon/2}$, say, for large n , and $f_1''(t) \leq n^{o(1)} f_1'(t)$, and so

$$(15) \quad \left| \Xi_1 - \frac{f_1'(t)}{s} \right| = O\left(\frac{\max_{t^* < t_{\max}} f_1''(t^*)}{s^2}\right) = o\left(\frac{f_1'(t)}{s}\right).$$

Next, we observe that

$$Z_A(i + 1) - Z_A(i) = d_{i+1}(A) - d_i(A) - \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon}.$$

The boundedness claim then follows for n sufficiently large as

$$0 \leq d_A(i + 1) - d_A(i) \leq 1$$

and as

$$|\Xi_1| \cdot n^{1/r-\varepsilon} \leq n^{\varepsilon/2+o(1)} \cdot n^{1/r-\varepsilon} \cdot s^{-1} < n/N$$

as

$$s^{-1} = D^{1/r}/N = \Theta(n^{1-1/r}/N).$$

Turning to the supermartingale condition, observe that $d_{i+1}(A) = d_i(A) + 1$ if and only if e_{i+1} lies in the set of open r -sets counted by $Q_A(i)$. Conditioned on the history of the process up to step i , it follows that

$$(16) \quad \begin{aligned} \mathbb{E}[Z_A(i + 1) - Z_A(i)] &= \frac{Q_A(i)}{|O(i)|} - \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \\ &\leq \frac{(q(t) + f(t)n^{-\varepsilon}) \cdot n}{(q(t) - N^{-\gamma}) \cdot N} - \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \\ &= \frac{N^{-\gamma} + f(t)n^{-\varepsilon}}{(q(t) - N^{-\gamma})} \cdot \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \\ &\leq (N^{-\gamma} + f(t)n^{-\varepsilon}) \cdot 2q(t)^{-1} \cdot \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \\ &= (2q(t)^{-1}N^{-\gamma} + 2f_1(t)n^{-\varepsilon}) \cdot \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \\ &\leq 4f_1(t) \cdot n^{-\varepsilon} \cdot \frac{n}{N} - \Xi_1 \cdot n^{1/r-\varepsilon} \end{aligned}$$

Note that the first inequality holds as \mathcal{T}_i and $Y_A^+(i) \geq 0$ since $i < \tau$, the second as $q(t) - N^{-\gamma} \geq q(t)/2$ since $\zeta \ll \gamma$, and the final as $N^{-\gamma} \leq f(t) \cdot n^{-\varepsilon}$, since $f(t) \geq 1$ and $\varepsilon \ll \gamma$. Noting that for large n , $D \geq n^{r-1}/r^r$ and so

$s^{-1} \geq n^{1-1/r}/(rN)$, by (15) we have

$$\begin{aligned} \Xi_1 \cdot n^{1/r-\varepsilon} &= (1 + o(1)) \cdot \frac{f'_1(t)}{s} \cdot n^{1/r-\varepsilon} \\ &\geq (1 + o(1)) \cdot \frac{W f_1(t) \cdot n^{1-1/r}}{rN} n^{1/r-\varepsilon} \\ &> \frac{W}{2r} \cdot f_1(t) \cdot n^{-\varepsilon} \cdot \frac{n}{N}. \end{aligned}$$

Thus, since we assume W is large, the supermartingale condition follows now from (16). ■

Finally, to show (12), we apply Lemma 9 to yield

$$\begin{aligned} \mathbb{P}[Z_A(i_{\max} \wedge \tau) > 0] &\leq \exp \left\{ -\Omega \left(\frac{n^{2/r-2\varepsilon}}{\frac{n}{N} \cdot \zeta s \log^{1/r} N} \right) \right\} \\ &= \exp \left\{ -\frac{n^{2/r-2\varepsilon}}{n^{1-(r-1)/r+o(1)}} \right\} \\ &= \exp \{ -n^{1/r-2\varepsilon-o(1)} \} \end{aligned}$$

which suffices as ε is small. This completes the proof of Lemma 7.

4.2. PROOF OF LEMMA 5. We begin by letting

$$S = S(n) = \binom{2\ell}{r} - 2 \binom{\ell}{r},$$

and we note that $S = \Theta(k^r)$.

We fix a pair A, B of disjoint ℓ -element subsets of $[n]$, and define the following sequences of random variables: for each step $i \geq 0$, let

$$\begin{aligned} X^+(i) &= q(t) \cdot S - Q_{A,B}(i) + f(t) \cdot S n^{-\varepsilon}, \text{ and} \\ X^-(i) &= q(t) \cdot S - Q_{A,B}(i) - f(t) \cdot S n^{-\varepsilon}. \end{aligned}$$

We next define the stopping time τ^* to be the minimum of $\tau_{A,B}$ and the first step i for which $X^+(i) \leq 0$, $X^-(i) \geq 0$, or the event \mathcal{T}_i fails to hold.

CLAIM 4: *The sequence*

$$X^+(0), \dots, X^+(i_{\max} \wedge \tau^*)$$

forms a $(O(k^r/s), O(k^{r-1}/n^{4\varepsilon}))$ -bounded submartingale, and the sequence

$$X^-(0), \dots, X^-(i_{\max} \wedge \tau^*)$$

forms a $(O(k^r/s), O(k^{r-1}/n^{4\varepsilon}))$ -bounded supermartingale.

Proof. We fix a step $0 \leq i \leq i_{\max}$, and we suppose that $i < \tau^*$. Throughout we write $t = t(i)$, and note that $t(i + 1) = t + s^{-1}$ and that

$$s^{-1} = D^{1/r}/N = \Theta(n^{1-1/r-r}).$$

To aid the calculations to follow, we begin by estimating the quantity $\Xi := f(t + s^{-1}) - f(t)$. Recall equation (13):

$$\left| \Xi - \frac{f'(t)}{s} \right| = O\left(\frac{\max_{t^* \leq t \leq t_{\max}} f''(t^*)}{s^2}\right) = o\left(\frac{f'(t)}{s}\right).$$

Observe that we may write

$$X^\pm(i + 1) - X^\pm(i) = (q(t + s^{-1}) - q(t)) \cdot S - (Q_{A,B}(i + 1) - Q_{A,B}(i)) \pm \Xi \cdot Sn^{-\varepsilon}.$$

(As above, this stands for the pair of equations in which each \pm is replaced with $+$ or with $-$, respectively.) We begin by establishing the boundedness claims: by (14) and as $S = \Theta(k^r)$, we have

$$0 \geq (q(t + s^{-1}) - q(t)) \cdot S \geq -O\left(\frac{k^r}{s}\right).$$

Next, bounding $|f'(t)| \leq n^{\varepsilon/2+o(1)}$,

$$|\Xi| \cdot Sn^{-\varepsilon} \leq n^{-\varepsilon/2+o(1)} \cdot \frac{k^r}{s}.$$

In order to establish the boundedness part of the claim, it remains to bound the quantity $Q_{A,B}(i + 1) - Q_{A,B}(i)$. Let $O_{A,B}(i)$ denote the set of r -sets that are open with respect to the pair A, B in $G(i)$, and let O_τ denote the set of all open r -sets whose selection as e_{i+1} would result in $\tau_{A,B} = i + 1$.

Now, if $e_{i+1} \in O_\tau$, then $Q_{A,B}(i + 1) - Q_{A,B}(i) = 0$ by definition, and, otherwise, we have

$$Q_{A,B}(i + 1) - Q_{A,B}(i) = -|O_{A,B}(i) \cap (C_{e_{i+1}}(i) \cup \{e_{i+1}\})|.$$

It suffices, then, to bound the quantity $|C_e(i) \cap O_{A,B}(i)|$ for all $e \in O(i) \setminus O_\tau$: fix such an open r -set e . Now, for any $f \in C_e(i) \cap O_{A,B}(i)$, there is a copy $T_{r,f}$ of $T^{(r)}$ in the graph $G(i) + e + f$ using both e and f as edges. Up to isomorphism, there are only three possibilities for the pair (e, f) in that copy: (e, f) maps to (b_1, b_2) , or to (b_1, a) , or to (a, b_1) . We treat these three cases separately.

Case 1: (e, f) maps to (b_1, b_2) . In this case, the $r - 1$ vertices that map to the set R lie entirely in e , and f is the union of those $r - 1$ vertices along with another vertex lying in $A \cup B$. Thus, we may bound the total number of such f above by rk .

Case 2: (e, f) maps to (b_1, a) . Let $R' = e - f$, the set of $r - 1$ vertices shared by all edges b_j in this copy of $T^{(r)}$. Since f maps to a , it follows that $f \subseteq N_i(R')$, and as $f \in O_{A,B}(i)$, we know that $f \subseteq A \cup B$, $f \cap A \neq \emptyset$, and $f \cap B \neq \emptyset$. Consequently, $N_i(R')$ intersects both A and B : since $e \notin O_\tau$, it follows that $|N_i(R') \cap (A \cup B)| \leq (k/n^{2\varepsilon})$ must hold.

Thus, by first selecting $R' \subseteq e$, which then identifies the sole vertex in $e \cap f$, and then selecting the $r - 1$ vertices comprising $f - e$ from $N_i(R') \cap (A \cup B)$, we can therefore bound the total number of such open r -sets f above by $r(k/n^{2\varepsilon})^{r-1}$.

Case 3: (e, f) maps to (a, b_1) . There exists an $(r - 1)$ -set $R' \subseteq A \cup B$ and a vertex $v \in e$ so that $f = R' \cup \{v\}$ and so that $e \setminus \{v\} \subseteq N_i(R')$. To bound the number of such f , it suffices to bound the number of $(r - 1)$ -sets $R' \subseteq A \cup B$ for which $N_i(R')$ contains $(r - 1)$ vertices from e .

To that end, fix a vertex $v \in e$ and let \mathcal{H}_v denote the $(r - 1)$ -uniform hypergraph on $(A \cup B) \setminus e$ whose edges are the $(r - 1)$ -subsets X for which $N_i(X) \supseteq e \setminus \{v\}$. We claim that

$$\Delta_{r-2}(\mathcal{H}_v) < 4r.$$

Suppose to the contrary that this does not hold: then there exist an $(r - 2)$ -set $Y \subseteq (A \cup B) \setminus e$ and vertices $x_1, x_2, \dots, x_{4r} \in (A \cup B) \setminus (Y \cup e)$ so that for each vertex $u \in e \setminus \{v\}$, $\{u\} \cup Y \cup \{x_j\} \in E(i)$ for $1 \leq j \leq 4r$. It follows from Lemma 4 that such a configuration does not appear in $G(i)$. Indeed, as this configuration spans $6r - 3$ vertices and has $4r(r - 1)$ edges, the probability that such a configuration appears is at most

$$n^{6r-3} \left(\frac{i}{N} \right)^{4r(r-1)} = n^{6r-3-4(r-1)^2+o(1)} = o(1).$$

It follows that $|\mathcal{H}_v| < 4r \binom{k}{r-2}$, and thus the total number of such open r -sets f as above is less than $4r^2 k^{r-2}$.

As ε is small and as $k = n^{1/r+o(1)}$, it follows that for large n we have

$$|C_e(i) \cap O_{A,B}(i)| \leq rk + r \cdot (k/n^{2\varepsilon})^{r-1} + 4r^2 k^{r-2} = O(k^{r-1}/n^{2\varepsilon(r-1)}),$$

and as $r \geq 3$ we conclude that

$$0 \geq Q_{A,B}(i + 1) - Q_{A,B}(i) = -O(k^{r-1}/n^{4\varepsilon}).$$

Thus, it follows that the sequences $X^\pm(0), \dots, X^\pm(i_{\max} \wedge \tau^*)$ are $(O(k^r/s), O(k^{r-1}/n^{4\varepsilon}))$ -bounded as claimed.

We now turn to the sub- and supermartingale claims, and we remark that all expectation and probability calculations to follow are implicitly conditioned on the history of the process up to step i . We begin by bounding the expected value of $Q_{A,B}(i + 1) - Q_{A,B}(i)$. Recall that we assume $i < \tau_{A,B}$ and that $O_\tau \subseteq O(i)$ consists of the open r -sets whose selection as e_{i+1} would yield $\tau_{A,B} = i + 1$. We claim that

$$(17) \quad |O_\tau| \leq 4n^{2\varepsilon} \cdot k$$

To see this, let

$$\mathcal{R} := \left\{ X \in \binom{[n]}{r-1} : |N_i(X) \cap (A \cup B)| \geq k/(2n^{2\varepsilon}) \right\}.$$

Then $|\mathcal{R}| < 4n^{2\varepsilon}$, which can be argued as follows. Suppose by way of contradiction that $\exists \mathcal{S} \subseteq \mathcal{R}$ with $|\mathcal{S}| = 4n^{2\varepsilon}$. Let $N = \bigcup_{Y \in \mathcal{S}} (N_i(Y) \cap (A \cup B))$. By inclusion-exclusion and the fact that Lemma 4 implies that the co-degree of any pair of $(r - 1)$ -sets is at most $5r$ (see (8)), we have

$$k \geq 2\ell = |A \cup B| \geq |N| \geq |\mathcal{S}| \cdot k/(2n^{2\varepsilon}) - |\mathcal{S}|^2 5r \geq 2k - 80rn^{4\varepsilon},$$

a contradiction as ε is small and $k = n^{1/r+o(1)}$. To deduce (17) it suffices to observe that each open r -set $e \in O_\tau$ can be written $e = \{v\} \cup X$ for some vertex $v \in A \cup B$ and $(r - 1)$ -set X satisfying $|N_i(X) \cap (A \cup B)| \geq k/n^{2\varepsilon} - 1$ (and thus $X \in \mathcal{R}$).

Conditioning on the event $e_{i+1} \notin O_\tau$ then yields

$$\mathbb{E} [Q_{A,B}(i + 1) - Q_{A,B}(i)] = - \sum_{e \in O_{A,B}(i)} \frac{|C_e(i) \setminus O_\tau|}{|O(i)|}$$

by linearity of expectation. Consequently,

$$\mathbb{E} [X^\pm(i + 1) - X^\pm(i)] = (q(t+s^{-1}) - q(t)) \cdot S + \sum_{e \in O_{A,B}(i)} \frac{|C_e(i) \setminus O_\tau|}{|O(i)|} \pm \Xi \cdot Sn^{-\varepsilon}.$$

To establish the submartingale claim, we note first that as $r \geq 3$ and $\varepsilon \ll \gamma \ll 1/r$, from (17) we have

$$|O_\tau| = n^{1/r+2\varepsilon+o(1)} < N^{-\gamma} \cdot D^{1/r}.$$

Now, as $i < \tau^*$, \mathcal{T}_i and $X^-(i) \leq 0$ hold, we have

$$\begin{aligned} \sum_{e \in O_{A,B}(i)} \frac{|C_e(i) \setminus O_\tau|}{|O(i)|} &\geq \left(q(t) - \frac{f(t)}{n^\varepsilon} \right) \cdot S \cdot \frac{(c(t) - 2N^{-\gamma})D^{1/r}}{(q(t) + N^{-\gamma})N} \\ &= \left(1 - \frac{N^{-\gamma} + f(t)n^{-\varepsilon}}{q(t) + N^{-\gamma}} \right) (c(t) - 2N^{-\gamma}) \cdot \frac{S}{s} \\ &\geq (1 - 2q(t)^{-1}f(t)n^{-\varepsilon})(c(t) - 2N^{-\gamma}) \cdot \frac{S}{s} \\ &\geq (c(t) - 2c(t)q(t)^{-1}f(t)n^{-\varepsilon} - 2N^{-\gamma}) \cdot \frac{S}{s} \\ &\geq (c(t) - (2c(t)q(t)^{-1} + 1)f(t)n^{-\varepsilon}) \cdot \frac{S}{s}. \end{aligned}$$

Note that these bounds follow for large n since $f(t) \geq 1$ and $\varepsilon \ll \gamma$ imply $N^{-\gamma} \leq f(t)n^{-\varepsilon}/2$. Applying this and (14) gives

$$\begin{aligned} \mathbb{E} [X^+(i+1) - X^+(i)] &\geq \Xi \cdot Sn^{-\varepsilon} - (2c(t)q(t)^{-1} + 1)f(t) \frac{Sn^{-\varepsilon}}{s} - O\left(\frac{1}{s^2}\right) \\ &\geq \Xi \cdot Sn^{-\varepsilon} - (2c(t)q(t)^{-1} + 2)f(t) \frac{Sn^{-\varepsilon}}{s} \\ &= \left((1 + o(1))f'(t) - (2c(t)q(t)^{-1} + 2)f(t) \right) \cdot \frac{Sn^{-\varepsilon}}{s} \end{aligned}$$

by (13). Since $f'(t) = (Wrt^{r-1} + W)f(t)$ and $2c(t)q(t)^{-1} = 2rt^{r-1}$, this final bound is nonnegative for large n as W is large, and so $X^+(0), \dots, X^+(i_{\max} \wedge \tau)$ forms a submartingale.

Turning to the supermartingale claim, we take a similar approach and begin by noting that as \mathcal{T}_i holds and $X^+(i) \geq 0$,

$$\begin{aligned} \sum_{e \in O_{A,B}(i)} \frac{|C_e(i) \setminus O_\tau|}{|O(i)|} &\leq \left(q(t) + \frac{f(t)}{n^\varepsilon} \right) \cdot S \cdot \frac{(c(t) + N^{-\gamma})D^{1/r}}{(q(t) - N^{-\gamma})N} \\ &= \left(1 + \frac{N^{-\gamma} + f(t)n^{-\varepsilon}}{q(t) - N^{-\gamma}} \right) (c(t) + N^{-\gamma}) \cdot \frac{S}{s} \\ &\leq (1 + 2q(t)^{-1}f(t)n^{-\varepsilon})(c(t) + N^{-\gamma}) \cdot \frac{S}{s} \\ &\leq (c(t) + (2c(t)q(t)^{-1} + 1)f(t)n^{-\varepsilon}) \cdot \frac{S}{s}. \end{aligned}$$

The supermartingale condition then follows in essentially the same way as the submartingale condition above. ■

Now, as $X^+(0) = Sn^{-\varepsilon}$, $X^-(0) = -Sn^{-\varepsilon}$, $S = \Theta(k^r)$ and $i_{\max} = s \cdot n^{o(1)}$, it follows from Claim 4 and Lemmas 8 and 9 that

$$\mathbb{P}[X^+(i_{\max} \wedge \tau^*) \leq 0] \leq \exp \left\{ -\Omega \left(\frac{S^2 n^{-2\varepsilon}}{\frac{k^r}{s} \cdot \frac{k^r-1}{n^{4\varepsilon}} \cdot sn^{o(1)}} \right) \right\} = \exp \{ -k \cdot n^{2\varepsilon - o(1)} \}.$$

Similarly, we have

$$\mathbb{P}[X^-(i_{\max} \wedge \tau^*) \geq 0] \leq \exp \{ -k \cdot n^{2\varepsilon - o(1)} \}.$$

Since there are fewer than $n^{2k} = \exp\{2k \log n\}$ choices of the pair of sets A and B , Lemma 5 follows from the union bound.

References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, *A note on Ramsey numbers*, Journal of Combinatorial Theory. Series A **29** (1980), 354–360.
- [2] P. Bennett and T. Bohman, *A note on the random greedy independent set algorithm*, submitted, <http://arxiv.org/abs/1308.3732>.
- [3] T. Bohman, *The triangle-free process*, Advances in Mathematics **221** (2009), 1653–1677.
- [4] T. Bohman, A. Frieze and D. Mubayi, *Coloring H -free hypergraphs*, Random Structures & Algorithms **36** (2010), 11–25.
- [5] T. Bohman and P. Keevash, *The early evolution of the H -free process*, Inventiones Mathematicae **181** (2010), 291–336.
- [6] T. Bohman and P. Keevash, *Dynamic concentration of the triangle-free process*, submitted, <http://arxiv.org/abs/1302.5963>.
- [7] J. Cooper and D. Mubayi, *Coloring sparse hypergraphs*, SIAM Journal on Discrete Mathematics, accepted, <http://arxiv.org/abs/1404.2895>.
- [8] G. Fiz Pontiveros, S. Griffiths and R. Morris, *The triangle-free process and $R(3, k)$* , submitted, <http://arxiv.org/abs/1302.6279>.
- [9] J. H. Kim, *The Ramsey number $R(3, t)$ has order of magnitude $t^2 / \log t$* , Random Structures & Algorithms **7** (1995), 173–207.
- [10] A. Kostochka, D. Mubayi and J. Verstraëte, *On independent sets in hypergraphs*, Random Structures & Algorithms **44** (2014), 224–239.
- [11] K. T. Phelps and V. Rödl, *Steiner triple systems with minimum independence number*, Ars Combinatoria **21** (1986), 167–172.
- [12] J. B. Shearer, *A note on the independence number of triangle-free graphs*, Discrete Mathematics **46** (1983), 83–87.
- [13] J. B. Shearer, *A note on the independence number of triangle-free graphs II*, Journal of Combinatorial Theory. Series B **2** (1991), 300–307.
- [14] N. Wormald, *The differential equation method for random graph processes and greedy algorithms*, in *Lectures on Approximation and Randomized Algorithms, Proceedings of the Berlin–Poznań Summer School*, Polish Scientific Publishers PWN, Warsaw, 1999, pp. 73–155.