

MODEL-THEORETIC APPLICATIONS OF COFINALITY SPECTRUM PROBLEMS

BY

MARYANTHE MALLIARIS*

*Department of Mathematics, University of Chicago
5734 S. University Avenue, Chicago, IL 60637, USA*

and

*Mathematical Sciences Research Institute
17 Gauss Way, Berkeley, CA 94720, USA
e-mail: mem@math.uchicago.edu*

AND

SAHARON SHELAH**

*Einstein Institute of Mathematics, Edmond J. Safra Campus, Givat Ram
The Hebrew University of Jerusalem, Jerusalem, 91904, Israel*

and

*Department of Mathematics, Hill Center - Busch Campus
Rutgers, The State University of New Jersey
110 Frelinghuysen Road, Piscataway, NJ 08854-8019 USA
e-mail: shelah@math.huji.ac.il
URL: <http://shelah.logic.at>*

* Malliaris was partially supported by NSF grants 1300634 and 1001666 and by a Sloan fellowship, and by a research membership at MSRI funded by NSF grant 0932078 000 (Spring 2014).

** Shelah was partially supported by the Israel Science Foundation grants 710/07 and 1053/11 and European Research Council grant 338821. This is manuscript 1051 in Shelah's list of publications.

Received February 28, 2015 and in revised form September 1, 2016

ABSTRACT

We apply the recently developed technology of cofinality spectrum problems to prove a range of theorems in model theory. First, we prove that any model of Peano arithmetic is λ -saturated iff it has cofinality $\geq \lambda$ and the underlying order has no (κ, κ) -gaps for regular $\kappa < \lambda$. We also answer a question about balanced pairs of models of PA. Second, assuming instances of GCH, we prove that SOP_2 characterizes maximality in the interpretability order \triangleleft^* , settling a prior conjecture and proving that SOP_2 is a real dividing line. Third, we establish the beginnings of a structure theory for $NSOP_2$, proving that $NSOP_2$ can be characterized by the existence of few so-called higher formulas. In the course of the paper, we show that $\mathfrak{p}_s = \mathfrak{t}_s$ in any weak cofinality spectrum problem closed under exponentiation (naturally defined). We also prove that the local versions of these cardinals need not coincide, even in cofinality spectrum problems arising from Peano arithmetic.

CONTENTS

Introduction 948

1. Cofinality spectrum problems: history and context 950

2. Definition of cofinality spectrum problems 952

 2.1. New definitions: weak and hereditary CSPs. 957

3. CSPs with exponentiation 960

4. On bounded arithmetic 963

5. Saturated models of Peano arithmetic 968

6. On the local cardinals $\mathfrak{p}_{s,a}$ and $\mathfrak{t}_{s,a}$ 973

7. Characterizing the \triangleleft^* -maximal class 979

8. Useful tools and additional definitions 987

9. Towards a structure theory for $NSOP_2$ 992

10. Balanced pairs of models of PA 1009

References 1013

Introduction

In a recent paper [9] we connected and solved two a priori unrelated open questions: the question from model theory of whether SOP_2 is maximal in Keisler’s order, and the question from set theory/general topology of whether $\mathfrak{p} = \mathfrak{t}$. This work was described in the research announcement [8] and the commentary [13]. In order to prove these theorems, we introduced a general

framework called cofinality spectrum problems. The big picture behind that framework and also the present theorems will be sketched in §2 below. First, however, we list the main theorems of the present paper.

In the present paper, we develop and apply cofinality spectrum problems to a range of problems in model theory, primarily on Peano arithmetic and around the strong tree property SOP_2 , also called the 2-strong order property. We prove the following theorems:

THEOREM (Theorem 5.7): *Let N be a model of Peano arithmetic, or just bounded PA, and λ an uncountable cardinal. If the reduct of N to the language of order has cofinality $> \kappa$ and no (κ, κ) -cuts for all $\kappa < \lambda$, then N is λ -saturated.*

On the earlier history, which involves [15] and [17] VI.2-3, see §5. We also address a question about balanced pairs of models of PA. The proof of Theorem 5.7 relies on the answer to a question arising from [9], of intrinsic interest:

THEOREM (Theorem 3.11): *Let \mathfrak{s} be a cofinality spectrum problem which is closed under exponentiation. Then $\mathfrak{p}_{\mathfrak{s}} = \mathfrak{t}_{\mathfrak{s}}$.*

As explained in §2, Theorem 3.11 complements a main theorem of [9], which showed that $\mathfrak{t}_{\mathfrak{s}} \leq \mathfrak{p}_{\mathfrak{s}}$ for any cofinality spectrum problem \mathfrak{s} . As a consequence, we are able to characterize $\mathfrak{t}_{\mathfrak{s}}$ in terms of the first symmetric cut. However, as we show in §6, the local versions of these cardinals, $\mathfrak{p}_{\mathfrak{s}, \mathfrak{a}}$, $\mathfrak{t}_{\mathfrak{s}, \mathfrak{a}}$ need not coincide unless the underlying model is saturated.

We then turn to the strong tree property SOP_2 . A major result of [9] was that SOP_2 suffices for being maximal in Keisler’s order \triangleleft . It was not proved to be a necessary condition, but we conjectured there that SOP_2 characterizes maximality in Keisler’s order. The difficulty in addressing this question may be in building ultrafilters. However, in the present paper, for a related open problem, we give a complete answer:

THEOREM (Theorem 7.13, under relevant instances of GCH): *T is \triangleleft^* -maximal if and only if it has SOP_2 .*

The ordering \triangleleft^* refines Keisler’s order, but is defined not in terms of ultra-powers but rather in terms of interpretability. Theorem 7.13 answers a very interesting question going back to Džamonja and Shelah [1] and Shelah and Usvyatsov [26]. We inherit the assumption of a case of GCH from [26], where

one direction of the theorem was proved, building on work of [1]. The direction proved here is in ZFC.

Theorem 7.13 gives decisive evidence for SOP_2 being a dividing line, by giving the equivalence of a natural inside/syntactic property and an outside property. However, this was done without developing a structure theory. So in §9, we develop the beginnings of a structure theory for $NSOP_2$. We define a notion of ‘higher formulas’ using ultrafilters and prove, for example, that:

THEOREM (Theorem 9.21): *T is $NSOP_2$ iff for all infinite A , the number of pairwise 1-contradictory higher φ -formulas over A is $\leq |A|$.*

Section 9 contains several other results, notably the “symmetry lemma” states that $NSOP_3$ can be characterized in terms of symmetry of inconsistency for these higher formulas. We also prove that SOP_2 is sufficient for a certain exact-saturation spectrum to be empty, connecting to work of Shelah [23] and Kaplan and Shelah [4]. The results seem persuasive and we believe they begin a structure theory for $NSOP_2$ theories. The introductions to each section contain further motivation for these results and discussions of prior work.

Sections 5 and 8 may be read essentially independently of prior sections. Section 7 uses one earlier theorem; the reader interested in the \triangleleft^* result may wish to begin there and refer back as needed.

We thank the referee for a very constructive reading of the paper.

1. Cofinality spectrum problems: history and context

A major interest, and contribution, of model theory has been the study of the complexity of families of definable sets, e.g. the sets definable in models of a given complete first-order theory. The **comparative** or **relative** complexity of such families often takes the following dichotomous form: research may eventually identify a so-called **dividing line** in the sense of classification theory [17], meaning a property, often a combinatorial property, whose presence signals complexity and whose absence gives a strong structure theory. With respect to a given dividing line, any given theory is either “tame” or “wild.” Few dividing lines are known, but those which are well developed such as stability/instability in [17] have been foundational for the field.

In the new millenium, in the context of our joint work on Keisler's order, a productive complementary perspective is emerging: from studying dichotomies around given dividing lines one by one, one may move to a framework in which one can directly compare the complexity of any two theories (a move which, ideally, then allows for the identification of new dividing lines in a systematic way).

Keisler's order was defined in 1967 as a pre-order on complete countable theories which sets $T_1 \triangleleft T_2$ if the regular ultrapowers of T_1 are "more likely to be saturated" than those of T_2 , see [5]. It is generally thought of as a partial order on the equivalence classes. For a recent account and current motivation; see, e.g., [10, §2]. The longstanding question is to determine the structure of Keisler's order.

Cofinality spectrum problems arose in the context of our proof that the maximum class in Keisler's order is much wider than previously thought. Formally, we proved that any theory with the model-theoretic tree property SOP_2 is maximal in Keisler's order. At the heart of this proof was an analysis of the comparative complexity (in the saturation of ultrapowers sense) of orders and trees. As explained in the introduction to [9], in the course of that work it became apparent that we were really studying the interaction of two invariants of a regular ultrafilter \mathcal{D} : $\mathfrak{p}_{\mathcal{D}}$, essentially the size of the first cut in a \mathcal{D} -ultrapower of a model of linear order, and $\mathfrak{t}_{\mathcal{D}}$, essentially the length of the smallest unbounded path in a \mathcal{D} -ultrapower of a tree. The names we gave to these invariants suggest a connection to the famous cardinal invariants of the continuum, \mathfrak{p} and \mathfrak{t} , and so to the long open question of whether it is possible that $\mathfrak{p} < \mathfrak{t}$. Here is the connection: As the analysis proceeded towards our eventual proof that $\mathfrak{p}_{\mathcal{D}}$ could not be strictly smaller than $\mathfrak{t}_{\mathcal{D}}$, we noticed that our proofs could be carried out in a context more general than ultrapowers. The proofs used only a few key facts: expansion properties available because ultrapowers commute with reducts, and pseudofiniteness assumptions available because ultraproducts of unbounded finite linear orders behave in a way that is 'pseudofinite'. The right general context was to abstract $(M, M^I/\mathcal{D})$ to an elementary pair of models of linear order admitting certain expansions (in which certain trees are definable) and capturing the relevant pseudofiniteness. Formally, such a pair is called a cofinality spectrum problem; before we begin the proofs of the present paper, the definition and some of its main consequences will be recalled below.

Now to any cofinality spectrum problem \mathbf{s} , one can naturally associate invariants $\mathfrak{p}_{\mathbf{s}}$ and $\mathfrak{t}_{\mathbf{s}}$, measuring certain cuts and certain paths through trees respectively, which specialize to $\mathfrak{p}_{\mathcal{D}}$ and $\mathfrak{t}_{\mathcal{D}}$ in the case where the csp arises as an ultrapower. The first main theorem of [9], already mentioned above, is that for any csp \mathbf{s} , $\mathfrak{t}_{\mathbf{s}} \leq \mathfrak{p}_{\mathbf{s}}$. In [9], for a certain well chosen csp consisting of a model of set theory and its generic ultrapower, we had naturally that $\mathfrak{p}_{\mathbf{s}} \leq \mathfrak{p}$ and $\mathfrak{t} \leq \mathfrak{t}_{\mathbf{s}}$, for the cardinal invariants of the continuum \mathfrak{p} and \mathfrak{t} . From this it followed that *if $\mathfrak{p} < \mathfrak{t}$ in some model of set theory, then $\mathfrak{p}_{\mathbf{s}} < \mathfrak{t}_{\mathbf{s}}$ there*, violating the ZFC theorem just mentioned. This proved that $\mathfrak{p} = \mathfrak{t}$ in ZFC, settling the question in a surprising way [8]. (A parallel reduction to regular ultrapowers proved the other main theorem of that paper already mentioned, concerning the maximality of SOP_2 in Keisler’s order.)

In the course of writing that paper and afterwards in discussing and presenting the proofs, we understood that cofinality spectrum problems were likely to be useful in other contexts besides the two, i.e., regular and generic ultrapowers, applications of [9] just sketched. As stated in the introduction, the present paper develops their theory with an eye towards model theoretic applications, along the way solving some relevant technical questions left outstanding from [9]. These are explained further in the sections below.

We now proceed to define cofinality spectrum problems.

2. Definition of cofinality spectrum problems

Informally, a cofinality spectrum problem

$$\mathbf{s} = (M, M_1, M^+, M_1^+, \Delta)$$

involves:

- an elementary pair of models (M, M_1) which can be expanded to the elementary pair (M^+, M_1^+) , and
- a set Δ of formulas in $\tau(M)$ defining discrete linear orders, closed under finite Cartesian products,

such that in the expanded model M_1^+ ,

- each instance of a formula in Δ defines a “pseudofinite” linear order, meaning that each of its M_1^+ -definable subsets has a first and last element,

- on at least one of the Cartesian products, the ordering is well behaved, e.g., like the Gödel pairing function,¹ and
- (M^+, M_1^+, Δ) has “enough set theory for trees,” meaning essentially that for each $\varphi \in \Delta$ there is $\psi \in \tau(M_1^+)$ so that for each linear order defined by an instance of φ , there is a tree defined by an instance of ψ :
 - whose nodes are functions from that order to itself, of length bounded by a distinguished element d of the order, and
 - the basic operations on this tree (the partial order on nodes given by initial segment, the length of a node, i.e., size of its domain, the value of a node at an element of its domain, and concatenation of an additional value) are likewise uniformly definable.

Natural and motivating examples of cofinality spectrum problems may be constructed beginning with pairs of models of Peano arithmetic or pairs of models of set theory for (M^+, M_1^+) with M being a reduct to a language containing linear order; or beginning with a pair (M, M_1) where M_1 is an ultrapower of M , using the theorem that ultrapowers commute with reducts. See [9] §2.3 for more details on these examples. The fruitful idea was simply that, in some sufficiently rich model, one may study the amount of saturation in an underlying linear order in relation to the fullness of the derived trees.

Here are the formal definitions, originally given in [9].

Definition 2.1 (Enough set theory for trees, [9] Definition 2.3): Let M_1 be a model and Δ a nonempty set of formulas in the language of M_1 . We say that (M_1, Δ) has **enough set theory for trees** when the following conditions are true:

- (1) Δ consists of first-order formulas $\varphi(\bar{x}, \bar{y}; \bar{z})$, with $\ell(\bar{x}) = \ell(\bar{y})$.
- (2) For each $\varphi \in \Delta$ and each parameter $\bar{c} \in {}^{\ell(\bar{z})}M_1$, $\varphi(\bar{x}, \bar{y}, \bar{c})$ defines a discrete linear order on $\{\bar{a} : M_1 \models \varphi(\bar{a}, \bar{a}, \bar{c})\}$ with a first and last element.

¹ Gödel’s pairing function orders pairs of ordinals first by maximum, then by first coordinate, then by second coordinate. If, e.g., there is a total linear order on the model, it makes sense to compare (“maximum”) elements in any two such ordered sets; if not, this requirement may be satisfied by some $\mathbf{a} \times \mathbf{a}$, that is the Cartesian product of a given ordered set with itself.

- (3) The family of all linear orders defined in this way will be denoted $\text{Or}(\Delta, M_1)$. Specifically, each $\mathbf{a} \in \text{Or}(\Delta, M_1)$ is a tuple $(X_{\mathbf{a}}, \leq_{\mathbf{a}}, \varphi_{\mathbf{a}}, \bar{c}_{\mathbf{a}}, d_{\mathbf{a}})$, where:
- (a) $X_{\mathbf{a}}$ denotes the underlying set $\{\bar{a} : M_1 \models \varphi_{\mathbf{a}}(\bar{a}, \bar{a}, \bar{c}_{\mathbf{a}})\}$,
 - (b) $\bar{x} \leq_{\mathbf{a}} \bar{y}$ abbreviates the formula $\varphi_{\mathbf{a}}(\bar{x}, \bar{y}, \bar{c}_{\mathbf{a}})$,
 - (c) $d_{\mathbf{a}} \in X_{\mathbf{a}}$ is a bound for the length of elements in the associated tree; it is often, but not always, $\max X_{\mathbf{a}}$. If $d_{\mathbf{a}}$ is finite, we call \mathbf{a} trivial.
- (4) For each $\mathbf{a} \in \text{Or}(\mathbf{a})$, $(X_{\mathbf{a}}, \leq_{\mathbf{a}})$ is pseudofinite, meaning that any bounded, nonempty, M_1 -definable subset has a $\leq_{\mathbf{a}}$ -greatest and $\leq_{\mathbf{a}}$ -least element.
- (5) For each pair \mathbf{a} and \mathbf{b} in $\text{Or}(\Delta, M_1)$, there is $\mathbf{c} \in \text{Or}(\Delta, M_1)$ such that:
- (a) there exists an M_1 -definable bijection $\text{Pr} : X_{\mathbf{a}} \times X_{\mathbf{b}} \rightarrow X_{\mathbf{c}}$ such that the coordinate projections are M_1 -definable,
 - (b) if $d_{\mathbf{a}}$ is not finite in $X_{\mathbf{a}}$ and $d_{\mathbf{b}}$ is not finite in $X_{\mathbf{b}}$, then also $d_{\mathbf{c}}$ is not finite in $X_{\mathbf{c}}$.
- (6) For some nontrivial $\mathbf{a} \in \text{Or}(\Delta, M_1)$, there is $\mathbf{c} \in \text{Or}(\Delta, M_1)$ such that $X_{\mathbf{c}} = \text{Pr}(X_{\mathbf{a}} \times X_{\mathbf{a}})$ and the ordering $\leq_{\mathbf{c}}$ satisfies:

$$M_1 \models (\forall x \in X_{\mathbf{a}})(\exists y \in X_{\mathbf{c}})(\forall x_1, x_2 \in X_{\mathbf{a}})(\max\{x_1, x_2\} \leq_{\mathbf{a}} x \iff \text{Pr}(x_1, x_2) \leq_{\mathbf{c}} y).$$

- (7) To the family of distinguished orders, we associate a family of trees, as follows. For each formula $\varphi(\bar{x}, \bar{y}; \bar{z})$ in Δ there are formulas ψ_0, ψ_1, ψ_2 of the language of M_1 such that for any $\mathbf{a} \in \text{Or}(\mathbf{s})$ with $\varphi_{\mathbf{a}} = \varphi$:
- (a) $\psi_0(\bar{x}, \bar{c}_{\mathbf{a}})$ defines a set, denoted $\mathcal{T}_{\mathbf{a}}$, of functions from $X_{\mathbf{a}}$ to $X_{\mathbf{a}}$.
 - (b) $\psi_1(\bar{x}, \bar{y}, \bar{c})$ defines a function $\text{lg}_{\mathbf{a}} : \mathcal{T}_{\mathbf{a}} \rightarrow X_{\mathbf{a}}$ satisfying:
 - (i) for all $b \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b) \leq_{\mathbf{a}} d_{\mathbf{a}}$,
 - (ii) for all $b \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b) = \max(\text{dom}(b))$.
 - (c) $\psi_2(\bar{x}, \bar{y}, \bar{c})$ defines a function from $\{(b, a) : b \in \mathcal{T}_{\mathbf{a}}, a \in X_{\mathbf{a}}, a <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b)\}$ into $X_{\mathbf{a}}$ whose value is called $\text{val}_{\mathbf{a}}(b, c)$, and abbreviated $b(a)$.
 - (i) If $c \in \mathcal{T}_{\mathbf{a}}$ and $\text{lg}_{\mathbf{a}}(c) < d_{\mathbf{a}}$ and $a \in X_{\mathbf{a}}$, then $c \hat{\ } \langle a \rangle$ exists, i.e., there is $c' \in \mathcal{T}_{\mathbf{a}}$ such that $\text{lg}_{\mathbf{a}}(c') = \text{lg}_{\mathbf{a}}(c) + 1$, $c'(\text{lg}_{\mathbf{a}}(c)) = a$, and

$$(\forall a <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(c))(c(a) = c'(a)).$$

- (ii) $\psi_0(\bar{x}, \bar{c})$ implies that if $b_1 \neq b_2 \in \mathcal{T}_{\mathbf{a}}$, $\text{lg}_{\mathbf{a}}(b_1) = \text{lg}_{\mathbf{a}}(b_2)$, then for some $n <_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_1)$,

$$b_1(n) \neq b_2(n).$$

- (d) $\psi_3(\bar{x}, \bar{y}, \bar{c})$ defines the partial order $\triangleleft_{\mathbf{a}}$ on $\mathcal{T}_{\mathbf{a}}$ given by the initial segment, that is, such that $b_1 \triangleleft_{\mathbf{a}} b_2$ implies:
 - (i) for all $b, c \in \mathcal{T}_{\mathbf{a}}$, $b \triangleleft c$ implies $\text{lg}_{\mathbf{a}}(b) \leq_{\mathbf{a}} \text{lg}_{\mathbf{a}}(c)$,
 - (ii) $\text{lg}_{\mathbf{a}}(b_1) \leq_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_2)$,
 - (iii) $(\forall a \triangleleft_{\mathbf{a}} \text{lg}_{\mathbf{a}}(b_1))(b_2(a) = b_1(a))$.

The family of all $\mathcal{T}_{\mathbf{a}}$ defined this way will be denoted $\text{Tr}(\Delta, M_1)$. We refer to elements of this family as trees.

Notice the pairing requirement in condition (6). While we need Cartesian products to exist, it is largely unimportant what exactly the order on these products is (as long as it satisfies the other requirements). It's sufficient for the results of [9] that one such order behave well, like the usual Gödel pairing function. In the course of the present paper, we will consider weakenings of this requirement.

Definition 2.2 (Cofinality spectrum problems, [9] Definition 2.5): We call the six-tuple

$$\mathbf{s} = (M, M_1, M^+, M_1^+, T, \Delta)$$

a **cofinality spectrum problem** when:

- (1) $M \preceq M_1$.
- (2) $T \supseteq \text{Th}(M)$ is a theory in a possibly larger vocabulary.
- (3) Δ is a set of formulas in the language of M , i.e., we are interested in studying the orders of M, M_1 in the presence of the additional structure of the expansion.
- (4) M^+, M_1^+ expand M, M_1 respectively so that

$$M^+ \preceq M_1^+ \models T$$

and (M_1^+, Δ) has enough set theory for trees.

- (5) We may refer to the components of \mathbf{s} as $M^{\mathbf{s}}, \Delta^{\mathbf{s}}$, etc. for definiteness. When $T = \text{Th}(M)$, $M = M^+$, $M_1 = M_1^+$, or Δ is the set of all formulas $\varphi(x, y, \bar{z})$ in the language of T which satisfy 2.1(2)–(4), these may be omitted.

Remark 2.3: The identities of M^+ and M_1 are not essential to many arguments.

Definition 2.4 (The cardinals $\mathfrak{p}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}}$ and the cut spectrum, [9] Definition 2.9): For a cofinality spectrum problem \mathbf{s} we define the following:

- (1) $\text{Or}(\mathfrak{s}) = \text{Or}(\Delta^{\mathfrak{s}}, M_1^{\mathfrak{s}})$.
- (2) $\mathcal{C}^{\text{ct}}(\mathfrak{s}) = \{(\kappa_1, \kappa_2) : \text{for some } \mathfrak{a} \in \text{Or}(\mathfrak{s}), (X_{\mathfrak{a}}, \leq_{\mathfrak{a}}) \text{ has a } (\kappa_1, \kappa_2)\text{-cut}\}$.

Note that the κ_{ℓ} are infinite.

- (3) $\text{Tr}(\mathfrak{s}) = \{\mathcal{T}_{\mathfrak{a}} : \mathfrak{a} \in \text{Or}(\mathfrak{s})\} = \text{Tr}(\Delta^{\mathfrak{s}}, M_1^{\mathfrak{s}})$.
- (4) $\mathcal{C}^{\text{ttp}}(\mathfrak{s}) = \{\kappa : \kappa \geq \aleph_0, \mathfrak{a} \in \text{Or}(\mathfrak{s}),$

and there is in the tree $\mathcal{T}_{\mathfrak{a}}$ a strictly increasing sequence
of cofinality κ with no upper bound}

- (5) Let $\mathfrak{t}_{\mathfrak{s}}$ be $\min \mathcal{C}^{\text{ttp}}(\mathfrak{s})$ and let $\mathfrak{p}_{\mathfrak{s}}$ be

$$\min\{\kappa : (\kappa_1, \kappa_2) \in \mathcal{C}^{\text{ct}}(\mathfrak{s}) \text{ and } \kappa = \kappa_1 + \kappa_2\}.$$

A key role will be played by $\mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}})$, where this means:

- (6) For λ an infinite cardinal, write

$$\mathcal{C}(\mathfrak{s}, \lambda) = \{(\kappa_1, \kappa_2) : \kappa_1 + \kappa_2 < \lambda, (\kappa_1, \kappa_2) \in \mathcal{C}^{\text{ct}}(\mathfrak{s})\}.$$

By definition, both $\mathfrak{t}_{\mathfrak{s}}$ and $\mathfrak{p}_{\mathfrak{s}}$ are regular. The main engine of the paper [9] was the following theorem, proved by model-theoretic means. Note that it entails $\mathfrak{t}_{\mathfrak{s}} \leq \mathfrak{p}_{\mathfrak{s}}$.

THEOREM A ([9] Theorem 9.1): *For any cofinality spectrum problem \mathfrak{s} ,*

$$\mathcal{C}(\mathfrak{s}, \mathfrak{t}_{\mathfrak{s}}) = \emptyset.$$

Discussion 2.5: We now resume the discussion following Definition 2.1. In [9], Cartesian products were used in two ways. First, we needed the simple existence of Cartesian products of pairs of elements of $\text{Or}(\mathfrak{s})$, with no restrictions on the ordering of the pairs other than: (i) pseudofiniteness and (ii) the property that if $d_{\mathfrak{a}}, d_{\mathfrak{b}}$ are nonstandard then so is $d_{\mathfrak{a} \times \mathfrak{b}}$ (call such a product **nontrivial**). This was needed for the basic arguments connecting behavior across all orders: notably, establishing existence of the lower cofinality function. To rule out symmetric cuts, we needed only to be able to take the Cartesian product of an order with itself. For the main lemma ruling out asymmetric cuts, we needed (a) the existence of \mathfrak{a} and \mathfrak{a}' such that \mathfrak{a} is coverable as a pair by \mathfrak{a}' (see below), and then (b) the existence of the nontrivial product $\mathfrak{a} \times \mathfrak{a} \times \mathfrak{a} \times \mathfrak{a}' \times \mathfrak{a}' \times \mathfrak{a}'$. One may take as a definition of “coverable as a pair” its key property, Corollary 5.7 of [9] [quoted here with the assumptions of nontriviality made there explicitly added in]:

- (**) there is $a \in X_{\mathbf{a}}$, with $a < d_{\mathbf{a}}$ and a not a finite successor of $0_{\mathbf{a}}$, such that the Gödel codes for functions from $[0, a]_{\mathbf{a}}$ to $[0, a]_{\mathbf{a}} \times [0, a]_{\mathbf{a}}$ may be definably identified with a definable subset of $X_{\mathbf{a}'}$ whose greatest element is $< d_{\mathbf{a}'}$.

For property (**), it suffices to have $\mathbf{a} \in \text{Or}(\mathbf{s})$ such that the order on $\mathbf{a} \times \mathbf{a}$ is given by the Gödel pairing function and $d_{\mathbf{a} \times \mathbf{a}}$ is nonstandard, by the arguments of [9]. We will give some alternate sufficient conditions in Section 8.

The following summarizes our conventions.

Convention 2.6 (Key conventions on CSPs):

- (a) Recall that definable means in the sense of $M_1^+ = M_1^+[\mathbf{s}]$, unless otherwise stated.²
- (b) Recall that for $\mathcal{T}_{\mathbf{a}} \in \text{Tr}(\mathbf{s})$, $\eta \in \mathcal{T}_{\mathbf{a}}$ implies that $\text{lg}(\eta) \leq d_{\mathbf{a}}$, and we have closure under concatenation, i.e., if $\text{lg}(\eta) < d_{\mathbf{a}}$ and $a \in X_{\mathbf{a}}$ then $\eta \frown \langle a \rangle \in \mathcal{T}_{\mathbf{a}}$.
- (c) When \mathcal{T} is a definable subtree of $\mathcal{T}_{\mathbf{a}}$, we will write $(\mathcal{T}, \triangleleft_{\mathbf{a}})$ or just $(\mathcal{T}, \triangleleft)$ to mean $(\mathcal{T}, \triangleleft_{\mathbf{a}} \upharpoonright_{\mathcal{T}})$.
- (d) When η is a sequence of n -tuples, $t < \text{lg}(\eta)$, and $k < n$, write $\eta(t, k)$ for the k th element of $\eta(t)$.
- (e) Generalizing [9], we use “internal map” in the present paper for any map definable in M_1^+ , not necessarily an element of some $\mathcal{T} \in \text{Tr}(\mathbf{s})$.
- (f) We call a definable, discrete order **pseudofinite** when every nonempty, bounded, definable subset has a first and last element.
- (g) We call $\mathbf{a} \in \text{Or}(\mathbf{s})$ **nontrivial** when $d_{\mathbf{a}}$ is not a finite successor of $0_{\mathbf{a}}$, and we call \mathbf{s} nontrivial when at least one $\mathbf{a} \in \text{Or}(\mathbf{s})$ is nontrivial.

2.1. NEW DEFINITIONS: WEAK AND HEREDITARY CSPs. We now give some new definitions weakening and extending CSPs looking towards applications.

First, as regards applying CSPs it will be useful to keep track of whether or not the assumption on closure under Cartesian products is needed. For example, in dealing with a CSP arising from a model $M = M^+ = M_1^+$ whose domain is linearly ordered, it may be that we care primarily about cuts occurring in the linear order $\text{dom}(M)$. Thus, we introduce ‘weak’ CSPs in Definition 2.7 and in the present paper, use them where possible.

² The main exception being that the elements of $\text{Or}(\mathbf{s})$ are a priori required to be Δ -definable; this is relaxed if \mathbf{s} is hereditarily closed, defined below.

Definition 2.7 (Weak CSP):

- (1) Call \mathbf{s} a **weak CSP** if it satisfies 2.2 omitting the requirement that $\text{Or}(\mathbf{s})$ be closed under Cartesian product, 2.1(5)+(6).
- (2) For \mathbf{s} a weak CSP, $\mathbf{a} \in \text{Or}(\mathbf{s})$, let $Y_{\mathbf{a}} = Y_{\mathbf{a},\mathbf{s}} = \{a \in X_{\mathbf{a}} : a <_{\mathbf{a}} d_{\mathbf{a}}\}$.

Second, in [9] Convention 5.1 we had observed that CSPs have available an internal notion of cardinality.³ In the present paper, we use the more general definition:

Definition 2.8: Let \mathbf{s} be a cofinality spectrum problem. Whenever A, B are definable subsets of M_1 (with parameters), we write

$$“|A| \leq |B|”$$

to mean “there exists an M_1^+ -definable partial 1-to-1 function h with $A \subseteq \text{dom}(h)$ and $\text{range}(h) \subseteq B$ ”. Likewise, we write

$$“|A| < |B|”$$

to mean “ $(|A| \leq |B|) \wedge \neg(|B| \leq |A|)$,” i.e., $|A| \leq |B|$ and there does not exist an M_1^+ -definable injection from B into A .

The referee has raised the interesting question of whether some kind of Schröder–Bernstein property holds for this weak notion of cardinality. In this vein, in §8 below we will show that from suitable internal injections or partial 1-to-1 functions between sets (in the sense of Definition 2.8) it is possible to recover an order-isomorphism at least on an initial segment: see, for example, Claim 8.3. The order is essential in our current setup, so this is a useful recovery and, as explained there, allows for some technical simplifications. Otherwise, the results so far have used very few ‘cardinality’-type properties of this definition: in particular, in our earlier paper, the related notion of internal cardinality ([9] 5.1) was used primarily to preserve distance inequalities for the proof by contradiction in [9, Theorem 8.1].

³ Which a priori need not satisfy all properties of cardinality. There it was required that the function be an element of the tree $\mathcal{T}_{\mathbf{a} \times \mathbf{b}}$. Depending on the amount of set theory available in M_1^+ , the present weakening is an a priori loss in definability: when functions are required to belong to a definable tree, we are always able to quantify over them, whereas in the definition below the quantification over all such functions may be strictly informal (not first order).

Let $(W, <_W)$ be a definable, pseudofinite linear order in the model. If there is an internal injection f of sets from W into some initial segment of N , then for some $\mathbf{b} \in \text{Or}(\mathbf{s})$ with $X_{\mathbf{b}} \subseteq N$ there exists an internal order-isomorphism g from $(W, <_W)$ onto an initial segment of $Y_{\mathbf{b}}$.

Third, while the definition of CSP carefully lays out the ‘canonical’ orders and trees, it is natural to add some internal closure conditions, stating essentially that objects which are internally isomorphic to canonical ones also count.

Definition 2.9: Let \mathbf{s} be a CSP or weak CSP.

- (1) We call \mathbf{b} a **pseudo-order** of \mathbf{s} and write $\mathbf{b} \in \text{PsOr}(\mathbf{s})$ when it satisfies all of the requirements on elements of $\text{Or}(\mathbf{s})$ from Definition 2.1 in the case where Δ is the set of all formulas in the language.⁴ (The point is that possibly the formulas defining it are not from $\Delta_{\mathbf{s}}$.)
- (2) Given $\mathbf{b}_2 \in \text{PsOr}(\mathbf{s})$, we say \mathbf{b}_1 is a **suborder** of \mathbf{b}_2 and write $\mathbf{b}_1 \in \text{Sub}(\mathbf{b}_2)$ when: $\mathbf{b}_1 \in \text{PsOr}(\mathbf{s})$, $X_{\mathbf{b}_1} \subseteq X_{\mathbf{b}_2}$ as linear orders, and there is $d \leq_{\mathbf{b}_2} d_{\mathbf{b}_2}$ and $\eta \in \mathcal{T}_{\mathbf{b}_2}$ with $\max \text{dom}(\eta) = d = d_{\mathbf{b}_1}$ which lists $\langle c \in X_{\mathbf{b}_1} : c <_{\mathbf{b}_1} d_{\mathbf{b}_1} \rangle$ in increasing order.⁵

Definition 2.10 (Hereditary closure): Let \mathbf{s} be a CSP or weak CSP.

- (1) Let $\mathbf{b}_1, \mathbf{b}_2 \in \text{PsOr}(\mathbf{s})$. We say (f, g, h) is an accurate isomorphism from \mathbf{b}_1 onto \mathbf{b}_2 when it respects both the order structure below $d_{\mathbf{b}_i}$ and the tree structure, that is:
 - (a) f is a one-to-one mapping, definable in M_1^+ , from $X_{\mathbf{b}_1}$ to $X_{\mathbf{b}_2}$,
 - (b) g is a one-to-one mapping, definable in M_1^+ , from $\{a : a <_{\mathbf{b}_1} d_{\mathbf{b}_1}\}$ onto $\{a : a <_{\mathbf{b}_2} d_{\mathbf{b}_2}\}$, such that

$$a <_{\mathbf{b}_1} c <_{\mathbf{b}_1} d_{\mathbf{b}_1} \iff g(a) <_{\mathbf{b}_2} g(c) <_{\mathbf{b}_2} d_{\mathbf{b}_2},$$

- (c) h is a one-to-one function definable in M_1^+ from $\mathcal{T}_{\mathbf{b}_1}$ onto $\mathcal{T}_{\mathbf{b}_2}$ such that

$$\eta \triangleleft_{\mathcal{T}_{\mathbf{b}_1}} \nu \iff h(\eta) <_{\mathcal{T}_{\mathbf{b}_2}} h(\nu)$$

and $\eta = \nu \frown \langle a \rangle$ in \mathbf{b}_1 iff $h(\eta) = h(\nu) \frown \langle f(a) \rangle$ in \mathbf{b}_2 .

- (2) Call \mathbf{s} **hereditary** if $\text{Or}(\mathbf{s})$ is closed under Sub and accurate isomorphism.

⁴ In particular, it must have an associated, definable tree, with definable length functions and concatenation, and so forth.

⁵ So, e.g., $\mathbf{b}_1 \notin \text{Sub}(\mathbf{b}_2)$ when $X_{\mathbf{b}_1}$ is an initial segment of $X_{\mathbf{b}_2}$ and $d_{\mathbf{b}_2} < \max(X_{\mathbf{b}_1})$, since there is not enough room for the list.

- (3) Say “ s_2 is the hereditary closure of s_1 ” when s_1, s_2 are the same except that $Or(s_2)$ is the closure of $Or(s_1)$ under members of Sub and under accurate isomorphism.

Observation 2.11: If s is a weak CSP then so is its hereditary closure.

Observation 2.12: As a consequence of Definition 2.9(3), if s is a weak CSP which is hereditarily closed, then for any $a \in Or(s)$, any $X \subseteq Y_a$ and any $d \in X \cap [0_a, d_a]$, there is $b \in Or(s)$ with $X_b = X, <_b = <_a \upharpoonright X$, and $d = d_b$.

3. CSPs with exponentiation

A feature of CSPs is that one can always develop a certain amount of Peano arithmetic internally, as in [9] §5. That observation motivates the main definition of this section, ‘closure under exponentiation.’ In this section we work out that CSPs of this kind have the very nice property that $p_s = t_s$. For context, recall that the main theorem of [9] that $C(s, t_s) = \emptyset$ only implies that $t_s \leq p_s$. For the other direction, it’s natural to try to show that a κ -indexed path through some \mathcal{T} with no upper bound translates to a (κ, κ) -cut in some $a \in Or(s)$. The problem is that the natural translation produces a definable, discrete linear order which contains a (κ, κ) -cut but is not necessarily a member of $Or(s)$. After explaining the problem, we will propose the solution.

Hypothesis 3.1: In this section, unless otherwise stated, s is a weak CSP. For transparency, we assume s is hereditary, i.e., closed under Sub and full internal isomorphism. We retain the notation Y_a from 2.7.

Definition 3.2 (Flattening the tree, cf. [9] Lemma 6.2): Let $a \in Or(s)$ be given. We define the following linear order. Fix in advance two distinct elements of X_a ; without loss of generality⁶ we use $0_a, 1_a$, called $0, 1$, so $0 <_a 1$. Let \mathcal{S}_a be the set $\mathcal{T}_a \times \{0, 1\}$. Let $<_{\mathcal{S}_a}$ be the linear order on \mathcal{S}_a defined as follows:

- If $c = d$, then $(c, i) <_{\mathcal{S}_a} (d, j)$ iff $i <_a j$.
- If $c <_a d$ and $c \neq d$, then $(c, 0) <_{\mathcal{S}_a} (d, 0) <_{\mathcal{S}_a} (d, 1) <_{\mathcal{S}_a} (c, 1)$.
- If c, d are \triangleleft -incomparable, then let $e \in \mathcal{T}_a, n_c, n_d \in \mathcal{S}_a$ be such that $e = \text{cis}(c, d)$ is the common initial segment and $e \hat{\ } n_c \triangleleft_a c$ and $e \hat{\ } n_d \triangleleft_a d$. Necessarily $n_c \neq n_d$ by definition of e , so for $s, t \in \{0, 1\}$ we define

⁶ Recall that X_a is a discrete linear order with first element, so $0_a, 1_a$ have the natural meanings.

$$(c, s) <_{\mathcal{S}_a} (d, t) \iff n_c <_a n_d.$$

For each given \mathbf{a} , we refer to this ordered set $(\mathcal{S}_a, <_{\mathcal{S}_a})$ just constructed as “the order given by flattening the tree \mathcal{T}_a ”.

Convention 3.3: For the remainder of this article, given \mathbf{s} and $\mathbf{a} \in \text{Or}(\mathbf{s})$, let $(\mathcal{S}_a, <_{\mathcal{S}_a})$ denote the order constructed in 3.2.

Definition 3.4: Say that a tree $\mathcal{T}_a \in \text{Tr}(\mathbf{s})$ **witnesses treetops** if there is in \mathcal{T}_a a strictly increasing sequence of cofinality \mathfrak{t}_s with no upper bound in \mathcal{T}_a . (In other words, this tree exemplifies why \mathfrak{t}_s takes the value it does.)

Fact 3.5: Let \mathbf{s} be a weak CSP. If $\mathbf{a} \in \text{Or}(\mathbf{s})$, \mathcal{T}_a witnesses treetops and $(\mathcal{S}_a, <_{\mathcal{S}_a})$ is the order given by flattening the tree \mathcal{T}_a , then $(\mathcal{S}_a, <_{\mathcal{S}_a})$ has a $(\mathfrak{t}_s, \mathfrak{t}_s)$ -cut.

Proof. The proof when \mathbf{s} is a CSP is [9] Lemma 6.2. No assumptions are made about closure under Cartesian products in that proof, so the identical result holds for weak CSPs as well. ■

We cannot a priori conclude from Fact 3.5 that $(\mathfrak{t}_s, \mathfrak{t}_s) \in \mathcal{C}(\mathbf{s}, \mathfrak{t}_s)$, because we have not shown that \mathcal{S}_a belongs to $\text{Or}(\mathbf{s})$. However, note that $(\mathcal{S}_a, <_{\mathcal{S}_a})$ is a definable discrete linear order with a first and last element, since:

Observation 3.6: For any weak cofinality spectrum problem \mathbf{s} and $\mathbf{a} \in \text{Or}(\mathbf{s})$, there is an internal order-isomorphism between an initial segment of $(\mathcal{S}_a, <_{\mathcal{S}_a})$ and Y_a .

Proof. Let $\langle c_\alpha : \alpha < d_a \rangle$ be the sequence of elements of \mathcal{T}_a corresponding to functions which are constantly 0_a , listed in increasing order: this sequence is linearly ordered by $<_a$. By the definition of \mathcal{S}_a , we have that $\{(c_\alpha, 0) : \alpha < d_a\}, <_a$ is an initial segment of $(\mathcal{S}_a, <_{\mathcal{S}_a})$. Moreover, it is isomorphic to $(Y_a, <_a)$ via the internal map lg_a from Definition 2.1(7). ■

Definition 3.7 (Closed under exponentiation): Let \mathbf{s} be a weak CSP. We say \mathbf{s} is closed under (simple) exponentiation when for every nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ there is a nontrivial $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $(X_b, <_b)$ and $(\mathcal{S}_a, <_{\mathcal{S}_a})$ are internally isomorphic.

Definition 3.8: For \mathbf{s} a weak CSP, it will also be useful to define:

- (a) **s** is **strongly closed under exponentiation**, or has strong exponentiation, when: for every $\mathbf{a} \in \text{Or}(\mathbf{s})$ there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ is accurately isomorphic to $(X_{\mathbf{b}} \upharpoonright d_{\mathbf{b}}, <_{\mathbf{b}})$, following Definition 2.9.⁷
- (b) **s** is **weakly closed under exponentiation**, or has weak exponentiation, when: for every $\mathbf{a} \in \text{Or}(\mathbf{s})$ there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $(X_{\mathbf{b}}, <_{\mathbf{b}})$ and $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ are internally isomorphic and $(Y_{\mathbf{b}}, <_{\mathbf{b}})$, $(Y_{\mathbf{a}}, <_{\mathbf{a}})$ are internally isomorphic.

Discussion 3.9: The phrase “**s** has exponentiation” clearly covers 3.7, 3.8(a) and 3.8(b). One of the issues raised by 3.8 is whether $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ may map onto $X_{\mathbf{b}}$ in such a way that $d_{\mathbf{b}}$ is not above the range of the map. To distinguish 3.7 from 3.8, we will say “simple exponentiation.”

COROLLARY 3.10: *Let **s** be a weak cofinality spectrum problem which has exponentiation, $\mathbf{a} \in \text{Or}(\mathbf{s})$. Then:*

- (1) $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ is a discrete linear order in which every nonempty definable subset has a first and last element.
- (2) There is $\mathbf{b} \in \text{Or}(\mathbf{s})$ so that $(X_{\mathbf{b}}, \leq_{\mathbf{b}})$ has a $(\mathfrak{t}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}})$ -cut.

Proof. (1) This is inherited from the order-isomorphism to an element of $\text{Or}(\mathbf{s})$.
 (2) Let $\mathcal{T}_{\mathbf{a}}$ witness treetops. By Fact 3.5, the order $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ has a $(\mathfrak{t}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}})$ -cut. Given $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ is order-isomorphic to $X_{\mathbf{b}}$, clearly $(X_{\mathbf{b}}, \leq_{\mathbf{b}})$ has a $(\mathfrak{t}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}})$ -cut. ■

We arrive at a fact which will be useful throughout the paper: if **s** has exponentiation, then $\mathfrak{p}_{\mathbf{s}} = \mathfrak{t}_{\mathbf{s}}$ and ‘the first cut is symmetric’.

THEOREM 3.11:

- (1) *Let **s** be a weak CSP with exponentiation. Then $\mathfrak{p}_{\mathbf{s}} \leq \mathfrak{t}_{\mathbf{s}}$.*
- (2) *Let **s** be a CSP with exponentiation. Then $\mathfrak{p}_{\mathbf{s}} = \mathfrak{t}_{\mathbf{s}}$.*
- (3) *Let **s** be a CSP with exponentiation. Then*

$$\mathfrak{t}_{\mathbf{s}} = \min\{\kappa : (\kappa, \kappa) \in \mathcal{C}^{\text{ct}}(\mathbf{s})\}$$

and the first cut in $\mathcal{C}^{\text{ct}}(\mathbf{s})$ is necessarily symmetric, that is, if

$$\mu = \min\{\kappa + \lambda : (\kappa, \lambda) \in \mathcal{C}^{\text{ct}}(\mathbf{s})\}$$

⁷ In Definition 2.9, accurate isomorphism of two linear orders $\mathbf{b}_0, \mathbf{b}_1$ involves an isomorphism of sets plus an order-isomorphism below the bounds $d_{\mathbf{b}_i}$. For these purposes, consider the d for $\mathcal{S}_{\mathbf{a}}$ to be $\max \mathcal{S}_{\mathbf{a}}$.

then $(\mu, \mu) \in \mathcal{C}^{ct}(\mathbf{s})$.

Proof. First we prove (1). Corollary 3.10(2) shows that if $\mathbf{t}_s = \kappa$, then

$$(\kappa, \kappa) \in \mathcal{C}^{ct}(\mathbf{s}),$$

so $\mathbf{p}_s \leq \kappa + \kappa = \kappa$. Thus $\mathbf{p}_s \leq \mathbf{t}_s$.

When in addition \mathbf{s} is a cofinality spectrum problem the analysis of [9] applies. By Theorem A, §2 above we have that $\mathcal{C}(\mathbf{s}, \mathbf{t}_s) = \emptyset$, thus $\mathbf{t}_s \leq \mathbf{p}_s$, proving (2).

For (3), the proof of 3.11 shows that $\mathbf{p}_s = \min\{\kappa : (\kappa, \kappa) \in \mathcal{C}^{ct}(\mathbf{s})\}$. Since $\mathbf{p}_s = \mathbf{t}_s$, this is sufficient. ■

However, as we will see in Theorem 6.3, the situation for the local versions of these cardinals, Definition 6.1, is more subtle. It would be interesting to explore this further.

4. On bounded arithmetic

In this section, working towards our first main application in §5, we set up CSPs and weak CSPs arising from models of PA or BPA and check when they are closed under exponentiation in the sense just described.

Definition 4.1: A formula is called bounded if all of its quantifiers are bounded. By BPA we mean bounded Peano arithmetic, that is, the restriction of the Peano axioms containing induction only for bounded formulas.

When working with models of PA or BPA, we will use the notation x^y in accordance with:

Fact 4.2 (Gaifman and Dimitracopoulos [3], see [28] §1.1): Let $I\Delta_0$ denote basic arithmetic with bounded induction. There exists a Δ_0 formula $\varphi(x, y, z)$, which we denote by $x^y = z$, that can be shown in $I\Delta_0$ to have all the usual properties of the graph of exponentiation except for the sentence

$$\forall x \forall y \exists z (x^y = z).$$

Regarding Definition 4.3, we will focus on $\mathbf{s}_3[N]$ for N a model of PA or BPA.

Definition 4.3 (Some canonical weak CSPs from models): Let $N \models BPA$ or $N \models PA$, and $\ell = 2, 3$. We define $\mathbf{s} = \mathbf{s}_\ell[N, D]$. Omitting D means $D = N$.

- (0) D is an initial segment of N closed under addition. D is **nontrivial** if it has a nonstandard member. We require that:

- (a) If $\ell = 2$, then $d \in D$ implies that $N \models (\forall x)(x^d \text{ exists})$.
 - (b) The case $\ell = 3$ is covered in (2)(c).
- (1) $\Delta = \Delta_{\mathbf{s}}$ is either:
- (a) in the case of BPA, the set of all bounded formulas, or
 - (b) in the case of PA, the set of all formulas $\varphi(x,y,z)$ with $\ell(x) = \ell(y) = 1$, which satisfy 2.1(1), (2), (4), (7).
- (2) $\mathbf{a} \in \text{Or}(\mathbf{s})$ when the data of $(X_{\mathbf{a}}, <_{\mathbf{a}})$ is internally isomorphic⁸ to some $(X_{\mathbf{a}}, \leq_{\mathbf{a}}, d_{\mathbf{a}}, \mathcal{T}_{\mathbf{a}})$ which satisfies:
- (a) the set of elements of $X_{\mathbf{a}}$ is a Δ -definable bounded subset of N .
 - (b) $<_{\mathbf{a}}$ is a Δ -definable linear order on $X_{\mathbf{a}}$.
 - (c) $d_{\mathbf{a}} \in X_{\mathbf{a}}$, and $(\{d : d < d_{\mathbf{a}}\}, <_{\mathbf{a}})$ is an initial segment of N with the usual order.
 - If $\ell = 2$, we require that $d_{\mathbf{a}} \in D$.
 - If $\ell = 3$, we require that $d_{\mathbf{a}} \in D$ and $|X_{\mathbf{a}}|^{d_{\mathbf{a}}}$ exists.
 - (d) $\mathcal{T}_{\mathbf{a}}$ is defined and satisfies the conditions from 2.1, and its defining formulas depend uniformly on the formula defining $X_{\mathbf{a}}$.
- (3) Write $\mathbf{s}^+[N, D]$ to indicate that we close the set of orders of $\mathbf{s}[N, D]$ under taking Cartesian products, where $d_{\mathbf{a} \times \mathbf{b}}$ is understood to be non-trivial if $d_{\mathbf{a}}, d_{\mathbf{b}}$ are, and that the order on at least one of the pairs is given by the Gödel pairing function.

For use in later papers, we record here:

Definition 4.4: In the context of 4.3, suppose f is a nondecreasing function with $\text{dom}(f) = N$ and $\text{range}(f) \subseteq \{I : I \text{ an initial segment of } N\}$, such that: $d \in f(a)$ implies that a^d exists in N and whenever a is nonstandard, $f(a)$ contains some nonstandard d . Define $\mathbf{s}[N, f]$ by requiring that for each \mathbf{a} ,

- (a) $d_{\mathbf{a}} \leq f(\max(X_{\mathbf{a}}) + 1)$, and
- (b) $f(\max(X_{\mathbf{a}}) + 1) = I$ implies $d_{\mathbf{a}} \in I$.

Let us also give a name to a recurrent assumption.

Definition 4.5 (Reasonable models): We say $N \models \text{BPA}$ is reasonable when $a \in N$ implies that $\{a^n : n \text{ a standard integer}\}$ is a bounded subset of N .

CLAIM 4.6: *Let $N \models \text{BPA}$.*

⁸ Recall that as the set is listed with its order, this means: internally order-isomorphic.

- (1) If N is \aleph_1 -saturated or just recursively saturated, then N is reasonable.
- (2) N is reasonable iff N is a candidate in the sense of 5.2 below.
- (3) If D_1 is a reasonable initial segment of N , then

$D_2 = \text{cl}(D_1, N) := \{a \in N : \text{for some } d \in D_1 \text{ and standard } n, N \models "a < d^n"\}$
 is also reasonable.

- (4) If D is an initial segment of N , then $\text{cl}(D, N)$ is closed under products.

Proof. (1), (2), (4) are immediate.

(3) Let $a_2 \in D_2$ and $n \in N$ be standard. Let $d_1 \in D_2$ and m standard be such that $N \models d_2 < (d_1)^m$. We define x_ℓ by induction on $\ell \leq n \cdot m$ as follows: $x_0 = x$, $x_\ell = (x_{\ell-1})^{d_1}$. Note that $(x_\ell)^{d_\ell}$ exists by the assumption on D_1 . So x_{nm} is well defined. Since

$$x^{d_2} \leq x^{d_1^m} = (x^{d_1})^{d_1} \dots (x^{d_1})^{d_1} \leq x_{nm}$$

where the dots \dots indicate the multiplication has m terms, x^{d_2} exists. ■

Observation 4.7: If $N \models \text{BPA}$ and $\mathbf{s} = \mathbf{s}_\ell[N, D]$ for $\ell \in \{2, 3\}$, then \mathbf{s} is a hereditary weak CSP.

CLAIM 4.8: Let $N \models \text{BPA}$ and suppose that for $n, d \in N$ we have that n^d exists. Then the tree of functions from d to n is definable in N , by bounded formulas. Specifically, the operations from 2.1(7) for the tree of sequences of length d into $([0, n], <^N)$ are all definable by bounded formulas.

Proof. For compatibility with 2.1, we denote $[0, n]$ by $X_{\mathbf{a}}$ and d by $d_{\mathbf{a}}$. Let \mathcal{T} denote the sequences of elements of $X_{\mathbf{a}}$ of length $\leq d_{\mathbf{a}}$. First we show that for each $\eta \in \mathcal{T}$ there is $c \in N$ which is a code for η (and that this is uniformly and boundedly definable). This applies the long known fact that Gödel coding may be carried out in BPA; we sketch a proof for completeness, following the method of Wilkie–Paris [28].

We use B -adic coding (for $B = 2$), representing each element of N as a word in the finite alphabet $\{0, 1\}$. (This ignores the empty word, which could be accommodated by using a different finite alphabet or $B = 3$.) Since for each $\mathcal{T}_{\mathbf{a}}$ or $\mathcal{S}_{\mathbf{a}}$ we will be coding sequences of elements of uniformly bounded length, we don't need a separate symbol to indicate a transition between codes for distinct elements. Let $\text{lgd}(x)$ denote $\max\{\ell : 2^\ell \leq x\}$, the dyadic length.

By 4.3(2), we may assume every element of $\text{Or}(\mathbf{s})$ is isomorphic to a canonical one, i.e., to $\mathbf{a} \in \text{Or}(\mathbf{s})$ where:

- $X_{\mathbf{a}} = [0, \max(X_{\mathbf{a}})]$,
- $\leq_{\mathbf{a}}$ agrees with the order of N restricted to $X_{\mathbf{a}}$
- $d_{\mathbf{a}}$ is such that $(\max X_{\mathbf{a}})^{d_{\mathbf{a}}}$ exists, which is equivalent to:

$$(\exists n)(|X_{\mathbf{a}}| \leq 2^n \leq 2|X_{\mathbf{a}}|) \wedge \text{“}2^{n \cdot d_{\mathbf{a}}}\text{ exists”},$$

- $(\exists n, m)(d_{\mathbf{a}} \leq 2^m \leq 2d_{\mathbf{a}} \wedge |X_{\mathbf{a}}| \leq 2^n \leq 2|X_{\mathbf{a}}| \text{ and } 2^{n \cdot 2^m} \text{ exists}),$
- so w.l.o.g. $d_{\mathbf{a}} = 2^m, |X_{\mathbf{a}}| = 2^n$.

For an element a of $X_{\mathbf{a}}$, let $\text{rep}_B(a)$ denote the B -adic representation of a of length exactly $n + 1$, padded with zeros if necessary (this is possible by the choice of n). In the expression below let “ $2^m + \text{rep}_B(a)$ ” mean in base 2, so this will effectively move $\text{rep}_B(a)$ (which is a sequence of length $\leq n$) over m spaces.

Let $\varphi_{\text{code}}(x, i, b)$ mean:

- $x < 2^{n \cdot 2^{m+1}}$,
- $i < d_{\mathbf{a}}$,
- $b < \max X_{\mathbf{a}}$,
- $(\exists x_1 x_2)(x_1 \hat{\ } x_2 \triangleleft x \wedge \text{lgd}(x_1) = (n+1)i \wedge \text{lgd}(x_2) = n+1 \wedge x_2 = 2^{n+1} + \text{rep}_B(b))$.

Informally, φ_{code} asserts that x is the code for a sequence, thought of as consisting of no more than $d_{\mathbf{a}}$ consecutive blocks of length $n + 1$ (leaving one extra space for the coding of $\mathcal{S}_{\mathbf{a}}$ in the proof of Claim 4.9), the $(i + 1)$ st of which is $\text{rep}_B(b)$.

As written, several values of x may code the same sequence; we may avoid this by restricting to x such that no $y < x$ codes the same sequence.

Then for any given $\mathbf{a} \in \text{Or}(\mathbf{s})$, for our fixed values of n, m , we may naturally represent $\mathcal{T}_{\mathbf{a}}$ by

$$\begin{aligned} \mathcal{T}_{\mathbf{a}} = \{ &x : x < 2^{n \cdot 2^{m+1}}, (\exists i < n)(\text{lgd}(x) = (m + 1) \cdot i) \\ &\text{and } (\forall j \leq i)(\exists x_0, x_1 < x)(x_0 \hat{\ } x_1 \triangleleft x \text{ and } 2^m \leq x_0 < 2^m + 2^n)\}. \end{aligned}$$

Since we have fixed the values of $X_{\mathbf{a}}$ (thus, of $\max(X_{\mathbf{a}})$) and $d_{\mathbf{a}}$, we can easily build on ψ_{code} to find bounded formulas defining: the partial order on elements of \mathcal{T} by initial segment, length, concatenation, and value of the function at a given element of its domain. ■

CLAIM 4.9: *Let $N \models BPA$ be reasonable and suppose that $\mathbf{s} = \mathbf{s}_3[N, D]$. Then \mathbf{s} has simple exponentiation.*

Proof. Recall that to say \mathbf{s} has simple exponentiation means that for every $\mathbf{a} \in \text{Or}(\mathbf{s})$ there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $(X_{\mathbf{b}}, <_{\mathbf{b}})$ and $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ are internally isomorphic. Weak exponentiation adds that also $(Y_{\mathbf{b}}, <_{\mathbf{b}})$, $(Y_{\mathbf{a}}, <_{\mathbf{a}})$ are internally isomorphic.

Let $\mathbf{a} \in \text{Or}(\mathbf{s})$ be nontrivial. By construction of \mathbf{s}_3 , we may identify \mathbf{a} with its “canonical” isomorphic image, so assume $X_{\mathbf{a}}$ is an initial segment of N with the usual order. We will show that the hypothesis that $|X_{\mathbf{a}}|^{d_{\mathbf{a}}}$ exists, i.e., that $(\max(X_{\mathbf{a}}))^{d_{\mathbf{a}}}$ exists, already ensures weak exponentiation.

As $\mathbf{s} = \mathbf{s}_3(N)$, there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ so that $(X_{\mathbf{b}}, \leq_{\mathbf{b}})$ is internally isomorphic to $([0, n_*], <^N)$ and this isomorphism takes $d_{\mathbf{b}}$ to d_* and $\mathcal{T}_{\mathbf{b}}$ to \mathcal{T}_* . Then \mathcal{T}_* is boundedly definable by Claim 4.8. Continuing in the notation of that proof, to code the tree consisting of $\mathcal{S}_{\mathbf{a}} = \mathcal{T}_{\mathbf{a}} \times \{0, 1\}$, define f as follows:

$$\begin{aligned} &\text{if } x = (\sum_{i < \lg(x)} (b_i + 2^m) \cdot 2^{(m+1)^i}, \mathbf{i}), \mathbf{i} \in \{0, 1\}, \\ &\text{then } f(x) = \sum_{i < \lg(x)} (b_i + 2^m) \cdot 2^{(m+1)^i} + \mathbf{i}, \end{aligned}$$

recalling that we have left one unit of space by arranging our coding into blocks of size $n + 1$. Let $\psi_{\text{code}}^+ = \text{range}(f)$, which is definable by a bounded formula. If $x_1, x_2 \in \mathcal{S}_{\mathbf{a}}$ then $x_1 <_{\mathcal{S}_{\mathbf{a}}} x_2$ iff $f(x_1) < f(x_2)$. This proves that $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$ is order-isomorphic to an initial segment of N .

Let $n_* = \max \text{range}(f)$. Since N is reasonable, there is some nonstandard $d_* \in N$ such that $(n_*)^{d_*}$ exists. Applying Claim 4.8 once more, we have a tree \mathcal{T}_* of functions from d_* to $[0, n_*]$ which is definable by bounded formulas. By the definition of $\mathbf{s}_3(N)$, there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ with $X_{\mathbf{b}}$ internally order-isomorphic to $[0, n_*]$ under the usual order so that the image of $d_{\mathbf{b}}$ is d_* .

This proves \mathbf{s} has simple exponentiation.

If in addition $(n_*)^{d_{\mathbf{a}}}$ exists, let $d_{\mathbf{b}} = d_{\mathbf{a}}$. Then the same internal order-isomorphism takes $Y_{\mathbf{b}}$ to $Y_{\mathbf{a}}$, and \mathbf{s} has weak exponentiation. This completes the proof. ■

CLAIM 4.10: *Suppose $N \models \text{BPA}$, N is reasonable and $\mathbf{s} = \mathbf{s}_3[N, D]$. Then \mathbf{s} is a CSP.*

Proof. As \mathbf{s} is a hereditary weak c.s.p. there are two potentially missing conditions, 2.1(5)–(6).

Let nontrivial \mathbf{a}, \mathbf{b} be given. By definition of \mathbf{s}_3 , we may assume that $X_{\mathbf{a}}, X_{\mathbf{b}}$ are internally order-isomorphic to initial segments $[0, n_{\mathbf{a}}], [0, n_{\mathbf{b}}]$ of N respectively (so in what follows we identify $X_{\mathbf{a}}, X_{\mathbf{b}}$ with these images). Without

loss of generality, $n_{\mathbf{a}} \leq n_{\mathbf{b}}$. Let $\text{Pr} : N \times N \rightarrow N$ denote the pairing function $(x, y) \mapsto (x + y + 1)^2 + x$. Consider the set

$$X_{\mathbf{a} \times \mathbf{b}} = \{\text{Pr}(x, y) : x \in X_{\mathbf{a}}, y \in X_{\mathbf{b}}\} \subseteq [0, (n_{\mathbf{a}} + n_{\mathbf{b}} + 1)^2].$$

Then Pr is an isomorphism from $X_{\mathbf{a}} \times X_{\mathbf{b}}$ onto an initial segment of N which we call $X_{\mathbf{c}}$. Let $\leq_{\mathbf{c}} = \leq^N$ be the usual order. Let the order $\leq_{\mathbf{a} \times \mathbf{b}}$ be such that Pr is an order-isomorphism from $(X_{\mathbf{a} \times \mathbf{b}}, \leq_{\mathbf{a} \times \mathbf{b}})$ onto $(X_{\mathbf{c}}, \leq_{\mathbf{c}})$. Note that this pairing function satisfies 2.1(6).

Let $n_* = \max(X_{\mathbf{c}})$. As we assumed N is reasonable, there is some nonstandard d_* such that $(n_*)^{d_*}$ exists. Let $d_{\mathbf{c}} = d_*$. Now existence of the tree $\mathcal{T}_{\mathbf{c}}$ is by Claim 4.8. Recalling the closure under isomorphism from 4.3(2), the product $\mathbf{a} \times \mathbf{b}$ is indeed a nontrivial element of \mathbf{s} . Thus 2.1(5) holds, which completes the proof. ■

CONCLUSION 4.11: Assume $N \models \text{BPA}$, N is reasonable and $\mathbf{s} = \mathbf{s}_3(N, D)$. Then \mathbf{s} is a cofinality spectrum problem with exponentiation, and so Theorem 3.11 applies: $\mathbf{p}_{\mathbf{s}} = \mathbf{t}_{\mathbf{s}}$.

5. Saturated models of Peano arithmetic

We now apply the above analysis to cuts in models of Peano arithmetic. The naturalness of this application comes from two sources. First, it was proved in [9, §5] that csps always allow for the recovery of a certain amount of Peano arithmetic.

Second, Peano arithmetic is a context in which there is a history of interesting work on sufficient conditions for saturation, also related to the history of work on Keisler’s order. The result that models of Peano arithmetic are saturated if and only if the underlying order is saturated is generally known as Pabion’s 1982 theorem [15], quoted below. Later, Kaufman and Schmerl have given an alternate proof. To explain the present results, it is useful to notice that this theorem is already a consequence of the proof of Shelah’s 1978 theorem that any theory of linear order is in the maximal Keisler class [17, Theorem 2.6] (though Pabion’s work was done independently).

THEOREM B (Pabion, 1982 [15]): Let $\lambda > \aleph_0$ and M be a model of PA. Then M is λ -saturated iff $(M, <)$ is λ -saturated.

THEOREM C (Shelah, 1978 [17] Cor. 2.7 pp. 337–341):

- (1) *If \mathcal{D} is a regular ultrafilter on λ such that M^λ/\mathcal{D} is λ^+ -saturated for M some model of linear order, then N^λ/\mathcal{D} is 1-atomically- λ^+ -saturated for any N in a countable language, i.e., λ^+ -saturated for types consisting of atomic formulas, e.g., $x < a, b < x$.*
- (2) *Hence if T has the strict order property it is maximal in Keisler's order.*

In Theorem 5.7 we will weaken the condition to requiring only that the underlying order has no symmetric cuts. A reasonable question, as raised by the referee, is that it could be suspected that these older results could be pushed further to get the result on symmetric cuts (i.e., the analogue of $\mathfrak{p} = \mathfrak{t}$). It seems the answer is no, that the new result is not just an incremental improvement, as discussed below.

We now give a sketch of the proof of Theorem C. It too relies on a connection between orders and trees, though of a somewhat simpler kind. We have a type in some regular ultrapower we would like to realize, and we assume that ultrapower is sufficiently saturated for linear order. Let $\langle \varphi_i : i < \alpha_0 \rangle$ list the type; without loss of generality it is closed under conjunctions. We expand the original model to have a tree of finite sequences of formulas, and in it a downward closed sub-tree of the sequences which are realized. In the ultrapower we choose t_i in the sub-tree, increasing with i in the sub-tree sense, by induction on $i \leq \alpha_0$; this t_i is a possibly nonstandard member of this sub-tree which is consistent with every truly finite subset of our type and includes φ_j for $j < i$. The cases of i a successor or zero are straightforward. For a limit there are two steps, both of which use the saturation of linear orders. First we find an upper bound d_i in the whole (nonstandard) tree in [17, VI, 2.7]; we translate the tree to a linear order and use “no symmetric cut”. This is done by replacing the tree by a linear order, replacing a member (node) of the tree by an interval. Second, below each node s_α the tree is a linear order. We choose by induction on $j < \alpha_0$ a member $d_{i,j}$ above the $c_{i_1}, i_1 < i$ and below the $d_{i,j_1}, j_j < j$. For $j = 0$ we use d_i , for j successor we use an appropriate function, and for j limit we use that there is no $(cf(i), cf(j))$ -cut in the ultrapower of the relevant linear order. So for each formula of the type separately there is t_α as required, and the initial segment defined by $\langle t_\beta : \beta < \alpha \rangle$ has the other cofinality big enough to find t_α good for all formulas of the type. Here also we use that there are no asymmetric cuts.

To see the relation to Peano arithmetic, notice that this proof just says that in some language expanding linear order, the types of the given theory may be

coded in such a way that omission of a type corresponds to a cut in a linear order; this coding argument does not rely on ultrapowers except insofar as ultrapowers commute with reducts, and so may be carried out in any model of Peano arithmetic (as well as in any ultrapower).

So from the maximality of linear order in \trianglelefteq we could obtain a sufficient condition for saturation of models of PA. It is therefore natural to hope that the innovations of [9], which allowed for a substantial weakening of the sufficient condition for maximality in Keisler’s order identified in [17, 2.6], will allow for a parallel weakening of the sufficient condition for saturation of models of (bounded) Peano arithmetic in Theorem 5.7 quoted above. By means of csps, Theorem 5.8 carries this out.

One measure of how surprising it is to weaken “no cuts” to “no symmetric cuts” is via negative results. For example, the reduction to symmetric cuts may fail badly in algebraic contexts without sufficient arithmetic structure, such as real closed fields, as in the theorem quoted at the end of this section. As another example, in §4 of [25] it is shown that when generalizing to reduced powers (and atomic saturation), on the one hand it is true that no cuts imply saturation, but on the other hand it is false that no symmetric cuts imply saturation, i.e., there is a counter-example. A different measure of this distance between ‘no cuts’ and ‘no symmetric cuts’ can be seen in problems around the characterization of good regular ultrafilters as those whose “first cut in the cut spectrum is symmetric,” such as in the forthcoming open problems paper [12].

Observation 5.1: For M a model of linear order ($<$) and uncountable λ , M is 1-atomically λ -saturated iff $\theta_1 + \theta_2 \geq \lambda$ whenever M has a cut of cofinality (θ_1, θ_2) and $\theta_1 + \theta_2 \geq \aleph_0$ (equivalently, > 2 , since we may have a $(1, 1)$ -cut). Usually, we can omit the “1-atomically,” as, e.g., in the order reduct of a model of PA or in $Th(\mathbb{Q}, <)$.

In Observation 5.1, note that if we use $x \leq y$, then $\min\{\theta_1, \theta_2\} \geq \aleph_0$ but then an $(\theta_1, 1)$ -cut does not give incompleteness.

The present results are an improvement in two respects. First, we can restrict to the case $\theta_1 = \theta_2$, i.e., symmetric cuts. Second, our results are for bounded Peano arithmetic, not just PA.

Definition 5.2: Call N a **candidate model** when N is a model of bounded Peano arithmetic (with no last element) which is reasonable, i.e., such that for

any $a \in N$ there is a nonstandard d such that a^d exists. We call \mathbf{s} a **candidate c.s.p.** when $\mathbf{s} = \mathbf{s}_3[N]$ for a reasonable N .

To connect to saturation, we bring in a definition from [9].

Definition 5.3 ([9] Definition 4.1): Let \mathbf{s} be a cofinality spectrum problem and λ a regular cardinal. Let $p = p(x_0, \dots, x_{n-1})$ be a consistent partial type with parameters in M_1^+ . We say that p is a **Or-type over M_1^+** if: p is a consistent partial type in M_1^+ and for some $\mathbf{a}_0, \dots, \mathbf{a}_{n-1} \in \text{Or}(\mathbf{s})$, we have that

$$p \vdash \bigwedge_{i < n} "x_i \in X_{\mathbf{a}_i}"$$

and p is finitely satisfiable in $X_{\mathbf{a}_0} \times \dots \times X_{\mathbf{a}_{n-1}}$. We say simply that M_1^+ is λ -**Or-saturated** if every Or-type over M_1^+ over a set of size $< \lambda$ is realized in M_1^+ . Finally, we say that \mathbf{s} is λ -**Or-saturated** if M_1^+ is.

CLAIM 5.4 ([9] Theorem 4.1): *Let \mathbf{s} be a cofinality spectrum problem. If $\kappa < \min\{\mathbf{p}_s, \mathbf{t}_s\}$ then \mathbf{s} is κ^+ -Or-saturated.*

Remark 5.5: Since by our definition any $\mathbf{a} \in \text{Or}(\mathbf{s})$ has a maximum element, Or-saturation does not a priori guarantee that the cofinality of the model is large.

Observation 5.6: For a model N of PA , the following are equivalent:

- (1) N is λ -saturated.
- (2) $\text{cf}(N) \geq \lambda$ and N is boundedly λ -saturated, that is, $N \upharpoonright_{\leq a}$ is λ -saturated for every $a \in N$.

Proof. It suffices to prove (2) implies (1). Given a type $p(x)$ of cardinality $< \lambda$, write $p(x) = \{\varphi_i(x, \bar{a}_i) : i < \lambda\}$. Since $\text{cf}(N) \geq \lambda$ there is some $a_* \in N$ such that $p(x) \cup \{x < a_*\}$ is finitely satisfiable. Let $b_* = 2^{a_*}$. Now for each i there are $c_i < b_*$ and $\varphi'_i(x, y, b_*)$ which is b_* -bounded [meaning that all quantifiers are of the form $(\exists z < b_*)$ or $(\forall z < b_*)$] and $N \models (\forall x)(\varphi_i(x, \bar{a}_i) \equiv \varphi'_i(x, c_i, b_*))$. Let $p''(x) = \{\varphi'_i(x, c_i, b_*) \wedge x < a_* : i < \lambda\}$. This is a finitely satisfiable type in $N \upharpoonright_{\leq b_*}$. Its realization implies realization of p , and it is realized by hypothesis (2). ■

THEOREM 5.7: *Let N be a model of Peano arithmetic and λ an uncountable cardinal. If the reduct of N to the language of order has cofinality $\geq \lambda$ and no (κ, κ) -cuts for $\kappa < \lambda$, then N is λ -saturated.*

Proof. We may assume N is a nonstandard model. Hence it follows that N is reasonable in the sense of 4.3, i.e., $a \in N$ implies that $\{a^n : n \text{ a standard integer}\}$ is a bounded subset of N . (If not, $\text{cf}(N) < \lambda$.)

Assume $\text{cf}(N) \geq \lambda$. Let $\mathbf{s} = \mathbf{s}_3(N)$ be from 4.3. By Conclusion 4.11, \mathbf{s} is a c.s.p. with exponentiation. Thus Theorem 3.11 applies and $\mathbf{p}_\mathbf{s} = \mathbf{t}_\mathbf{s}$. By Claim 5.4 and Observation 5.6, \mathbf{s} is $\min\{\mathbf{p}_\mathbf{s}, \mathbf{t}_\mathbf{s}, \text{cf}(N, <)\}$ -saturated. If $\mathbf{t}_\mathbf{s} \geq \lambda$, we finish, so assume that $\mathbf{t}_\mathbf{s} < \lambda$. By Theorem 3.11, there is $\mathbf{a} \in \text{Or}(\mathbf{s})$ whose $X_\mathbf{a}$ contains a $(\mathbf{t}_\mathbf{s}, \mathbf{t}_\mathbf{s})$ -cut. Recall from Definition 4.3 that since $\mathbf{s} = \mathbf{s}_3(N)$, for each $\mathbf{a} \in \text{Or}(\mathbf{s})$, $(X_\mathbf{a}, \leq_\mathbf{a})$ is internally order-isomorphic to an initial segment of N with the usual order. Then N has a $(\mathbf{t}_\mathbf{s}, \mathbf{t}_\mathbf{s})$ -cut, which completes the proof. ■

THEOREM 5.8: *Let N be a model of BPA which is reasonable, i.e., for every $a \in N$ the set $\{a^n : n \in N \text{ finite}\}$ is bounded. Then the following are equivalent:*

- (1) *For every $n_* \in N$, the model $N_{<n_*} = N \upharpoonright \{a : N \models a < n_*\}$ is λ -saturated.*
- (2) *For every $n_* \in N$, the model $N_{\leq n_*}$ considered as a linear order has no (κ, κ) -cuts for $\kappa = \text{cf}(\kappa) < \lambda$.*

Proof. (1) implies (2) is obvious, so assume (2) holds. Let $\mathbf{s} = \mathbf{s}_3(N)$ be from 4.3. Then by 4.11, \mathbf{s} is a cofinality spectrum problem with exponentiation, and $\mathbf{p}_\mathbf{s} = \mathbf{t}_\mathbf{s}$.

Let $\kappa = \mathbf{p}_\mathbf{s} = \mathbf{t}_\mathbf{s}$, so κ is regular. By 3.10, some $\mathbf{a} \in \text{Or}(\mathbf{s})$ has a (κ, κ) -cut. By the definition of $\mathbf{s}_3(N)$, any $X_\mathbf{a}$ is internally order-isomorphic to a bounded initial segment of N with the usual order. Thus, some bounded initial segment of N has a (κ, κ) -cut. As we've assumed (2), it must be that $\lambda \leq \kappa = \mathbf{p}_\mathbf{s} = \mathbf{t}_\mathbf{s}$.

Let $n_* \in N$ be given. As we assumed N is reasonable, there is a nonstandard d_* such that $N \models "n_*^{d_*} \text{ exists}"$. Recalling 4.3(2), there is a nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ such that $(X_\mathbf{a}, \leq_\mathbf{a})$ is isomorphic to an initial segment of N , containing $[0, n_*]$, with the usual order. Thus, to prove (1), it will suffice to show that every $\mathbf{a} \in \text{Or}(\mathbf{s})$ is γ^+ -Or-saturated for every $\gamma < \lambda$. By 5.4, if $\gamma < \min\{\mathbf{p}_\mathbf{s}, \mathbf{t}_\mathbf{s}\}$ then \mathbf{s} is γ^+ -Or-saturated. Since $\lambda \leq \mathbf{p}_\mathbf{s} = \mathbf{t}_\mathbf{s}$, this completes the proof. ■

Note that Theorems 5.7 and 5.8 show that the situation in models of Peano arithmetic is very different from that in real closed fields, as shown by the next

quoted theorem. By “asymmetric cut” we mean a cut in which the infinite cofinalities of each side are not equal.

THEOREM D (Theorem 1.1 of Shelah [22]): *Let K be an arbitrary ordered field. Then there is a symmetrically complete⁹ real closed field K^+ containing K such that any asymmetric cut of K is not filled. So if K is not embeddable into \mathbb{R} , then K^+ and K necessarily have an asymmetric cut.*

6. On the local cardinals $\mathfrak{p}_{\mathbf{s},\mathbf{a}}$ and $\mathfrak{t}_{\mathbf{s},\mathbf{a}}$

Returning to CSPs generally, in this section we prove Theorem 6.3, a complementary result to Theorem 3.11. The theorem shows that the local cardinals $\mathfrak{p}_{\mathbf{s},\mathbf{a}}$ and $\mathfrak{t}_{\mathbf{s},\mathbf{a}}$ from Definition 6.1 need not normally agree, even in CSPs arising from models of Peano arithmetic, if the underlying M_1^+ is not uniformly saturated.

Definition 6.1: Let \mathbf{s} be a CSP or weak CSP, and $\mathbf{a} \in \text{Or}(\mathbf{s})$.

(1) Let $\mathfrak{p}_{\mathbf{s},\mathbf{a}}$ be

$$\min\{\kappa : \text{there are regular } \kappa_1, \kappa_2 \text{ such that } \kappa_1 + \kappa_2 = \kappa \text{ and } X_{\mathbf{a}} \text{ has a } (\kappa_1, \kappa_2)\text{-cut}\}.$$

(2) Let $\mathfrak{t}_{\mathbf{s},\mathbf{a}}$ be

$$\min\{\kappa : \kappa \geq \aleph_0 \text{ and there is in the tree } \mathcal{T}_{\mathbf{a}} \text{ a strictly increasing sequence of cofinality } \kappa \text{ with no upper bound}\}.$$

We need a preliminary lemma.

LEMMA 6.2: *Let \mathbf{s} be a cofinality spectrum problem, $\mathbf{a} \in \text{Or}(\mathbf{s})$. Suppose $(\mathcal{S}_{\mathbf{s}}, <_{\mathcal{S}_{\mathbf{s}}})$ has a (κ, κ) -cut, for $\kappa \leq \mathfrak{t}_{\mathbf{s}}$. Then either $X_{\mathbf{a}}$ has an (κ, κ) -cut or else $\mathcal{T}_{\mathbf{a}}$ has a branch of cofinality κ with no upper bound.*

Proof. Let

$$(\bar{a}, \bar{b}) = (\langle (a_i, t_i) : i < \kappa \rangle, \langle (b_i, s_i) : i < \kappa \rangle)$$

witness the cut in $\mathcal{S}_{\mathbf{a}}$, with $a_i, b_i \in \mathcal{T}_{\mathbf{a}}$ and $t_i, s_i \in \{0_{\mathbf{a}}, 1_{\mathbf{a}}\}$ for each $i < \kappa$. By definition of cut, we may assume κ is regular.

STEP 1: Simplifying the presentation of intervals.

⁹ This means that any decreasing sequence of closed bounded intervals, of any ordinal length, has nonempty intersection.

By the pigeonhole principle, we may assume the sequences $\langle t_i : i < \kappa \rangle$ and $\langle s_i : i < \kappa \rangle$ are constant. For each $i < \kappa$, write A_i for the closed interval in the linear order $S_{\mathbf{a}}$ whose endpoints are given by (a_i, t_i) and $(a_i, |t_i - 1|)$, and likewise for B_i . By the construction of $S_{\mathbf{a}}$, any two intervals of this form are either concentric or disjoint.

Let $\bar{A} = \langle A_i : i < \kappa \rangle$ and $\bar{B} = \langle B_i : i < \kappa \rangle$.

The task of this step will be to prove that without loss of generality, \bar{A} consists of either pairwise concentric or pairwise disjoint intervals, and likewise for \bar{B} . Here “concentric” means either concentric decreasing: $j < i < \kappa \implies C_j \supseteq C_i$ [for $C = A$ or B] or concentric increasing: $j < i < \kappa \implies C_i \supseteq C_j$.

If $\kappa = \aleph_0$, then by Ramsey’s theorem,¹⁰ we may assume that \bar{A} consists either of concentric intervals or disjoint intervals moving right, meaning $j < i < \kappa$ implies $A_j \cap A_i = \emptyset$ and $(\forall x \in A_j)(\forall y \in A_i)(x < y)$. Likewise, we may assume that \bar{B} consists either of concentric intervals or of disjoint intervals moving left, meaning $j < i < \kappa$ implies $B_j \cap B_i = \emptyset$ and $(\forall x \in B_j)(\forall y \in B_i)(y < x)$.

If $\kappa > \aleph_0$, let $a \wedge b$ denote the maximal common initial segment of $a, b \in \mathcal{T}_{\mathbf{a}}$. For this argument, we use c to denote either a or b . For each i , the sequence $\langle \text{lg}_{\mathbf{a}}(c_i \wedge c_j) : j \in [i, \kappa) \rangle$ is a sequence of elements of $Y_{\mathbf{a}} \subseteq X_{\mathbf{a}}$ [recall Definition 2.7] bounded by $\text{lg}_{\mathbf{a}}(c_i)$. Thus, for some club E_i of κ with $\min E_i > i$, we have that $\bar{\ell}_i := \langle \text{lg}_{\mathbf{a}}(c_i \wedge c_j) : j \in E_i \rangle$ is either constant or $<_{\mathbf{a}}$ -decreasing. Let $S \subseteq \kappa$ be a stationary set of i on which we get the same outcome (either always constant or always decreasing).

Let $E = \bigcap \{ \epsilon < \kappa : \epsilon \text{ a limit ordinal and } \epsilon \in \bigcap_{i < \epsilon} E_i \}$, so E is a club of κ . There are several cases:

- (1) First case: for all $i \in S$, $\bar{\ell}_i$ is constant.
 - (a) If $i \in S$, $j \in E_i$ implies $c_i \wedge c_j = c_i$, then $\langle c_i : i \in S \cap E \rangle$ is a $<_{\mathcal{T}_{\mathbf{a}}}$ -increasing sequence so we are in the concentric decreasing case.
 - (b) If $i \in S$, $j \in E_i$ implies $\text{lg}_{\mathbf{a}}(c_i \wedge c_j) < \text{lg}(c_i)$, then by Fodor’s lemma there is $\gamma \in \kappa$ and a stationary subset X of κ such that $i \in X$ and $j \in E_i$ implies $\text{lg}_{\mathbf{a}}(c_i \wedge c_j) = \gamma$, so we are in the pairwise disjoint case.

¹⁰ This also holds if κ is weakly compact.

- (2) Second case: for all $i \in S$, $\bar{\ell}_i$ is $<_{\mathbf{a}}$ -decreasing. (Remember that the branches of $\mathcal{T}_{\mathbf{a}}$ are internally pseudofinite, but not necessarily well-ordered from an external point of view.) Let $X = S \cap E$. Then $\langle c_i : i \in X \rangle$ is a $\triangleleft_{\mathcal{T}_{\mathbf{a}}}$ -decreasing sequence, so we are in the concentric increasing case.

STEP 2: The concentric cases.

Suppose both \bar{A} and \bar{B} are concentric. Then $\langle a_i : i < \kappa \rangle$ and $\langle b_i : i < \kappa \rangle$ are both $\triangleleft_{\mathbf{a}}$ -linearly ordered sequences in $\mathcal{T}_{\mathbf{a}}$. There are four cases depending on whether each of these sequences is $\triangleleft_{\mathbf{a}}$ -increasing or decreasing.

(a) Suppose both $\langle a_i : i < \kappa \rangle$ and $\langle b_i : i < \kappa \rangle$ are $\triangleleft_{\mathbf{a}}$ -increasing. If they lie along eventually different branches, the original sequence (\bar{a}, \bar{b}) could describe only a pre-cut and not a cut according to the definition of $\mathcal{S}_{\mathbf{a}}$, so we get a contradiction. If they lie along the same branch, then it must be that $\mathcal{T}_{\mathbf{a}}$ has a branch of cofinality κ , as desired.

(b) If $\langle a_i : i < \kappa \rangle$ is $\triangleleft_{\mathbf{a}}$ -increasing while $\langle b_i : i < \kappa \rangle$ is $\triangleleft_{\mathbf{a}}$ -decreasing, these form a cut $(*)$ in the linearly ordered set

$$(\{c \in \mathcal{T}_{\mathbf{a}} : c \triangleleft b_0\}, \triangleleft).$$

Then the projections $(\langle \lg(a_i) : i < \kappa \rangle, \langle \lg(b_j) : j < \kappa \rangle)$ form a pre-cut $(**)$ in $X_{\mathbf{a}}$. If this pre-cut $(**)$ were realized, say by t , then $b_0 \upharpoonright t$ realizes the cut $(*)$, contradiction. This shows that $X_{\mathbf{a}}$ has a (κ, κ) -cut.

(c) If $\langle a_i : i < \kappa \rangle$ and $\langle b_i : i < \kappa \rangle$ are both $\triangleleft_{\mathcal{T}_{\mathbf{a}}}$ -decreasing, the original sequence (\bar{a}, \bar{b}) will not describe a cut, so we ignore this case.

(d) If $\langle b_i : i < \kappa \rangle$ is $\triangleleft_{\mathcal{T}_{\mathbf{a}}}$ -increasing while $\langle a_i : i < \kappa \rangle$ is $\triangleleft_{\mathcal{T}_{\mathbf{a}}}$ -decreasing, the argument is parallel to case (b).

STEP 3: Not both concentric.

Again, there are several possibilities.

Suppose first that neither \bar{A} nor \bar{B} is concentric, so \bar{A} is a sequence of disjoint intervals moving right and \bar{B} is a sequence of disjoint intervals moving left.

Consider the sequence $\langle c_i : i < \kappa \rangle$ where $c_i := \text{lub}\{a_i, b_i\}$ in the tree $\mathcal{T}_{\mathbf{a}}$. By definition of \bar{a} and \bar{b} , this sequence will be either eventually constant or a path through the tree $\mathcal{T}_{\mathbf{a}}$.

If the sequence is a path through the tree, then $\langle (c_i, 0) : i < \kappa \rangle$ is cofinal in \bar{a} and $\langle (c_i, 1) : i < \kappa \rangle$ is cofinal in \bar{b} . So the path $\langle c_i : i < \kappa \rangle$ cannot have an upper bound, as given any such upper bound d_* , by definition of $(\mathcal{S}_{\mathbf{a}}, <_{\mathcal{S}_{\mathbf{a}}})$, we would have that $(d_*, 0)$ and $(d_*, 1)$ both realize the original cut, contradiction.

So in this case, there is a path through \mathcal{T}_a of length κ with no upper bound.

If the sequence is eventually constant, then there is $i_* < \kappa$ such that all $\{a_i, b_i : i_* < i < \kappa\}$ are immediate successors of the same node, say a_* , in \mathcal{T}_a . So $\text{lg}(a_*) < d_a$. By definition of the order $(\mathcal{S}_a, <_{\mathcal{S}_a})$ and the case we are in, this means

$$(\langle a_i(\text{lg}(a_i) - 1) : i_* < i < \kappa \rangle, \langle b_i(\text{lg}(b_i) - 1) : i_* < i < \kappa \rangle)$$

is a pre-cut in X_a . Suppose for a contradiction it were realized by x ; then $c_* := a_* \wedge \langle x \rangle$ would exist since $\text{lg}(a_*) < d_a$. Then in $(\mathcal{S}_a, <_{\mathcal{S}_a})$, $(c_*, 0)$ and $(c_*, 1)$ would both realize the original cut (\bar{a}, \bar{b}) , contradiction.

So in this case, X_a has a (κ, κ) -cut.

Otherwise, precisely one of \bar{A}, \bar{B} is not concentric. The cases are parallel, so assume the non-concentric side is \bar{B} . Define d_i for $i < \kappa$ by $d_i := \text{lub}\{b_i, b_0\}$ in the tree \mathcal{T}_a . Writing C_i for the interval $((c_i, 0), (c_i, 1))$ and D_i for the interval $((d_i, 0), (d_i, 1))$ in \mathcal{S}_a , we have that both $\langle C_i : i < \kappa \rangle$ and $\langle D_i : i < \kappa \rangle$ are concentric sequences of intervals with $D_i \subseteq C_j$ for all $i \gg j$.

This reduces the problem to Step 2.

STEP 4: Finish.

We have shown that in each case either X_a has a (κ, κ) -cut or else \mathcal{T}_a has a strictly increasing path of length κ with no upper bound, so this completes the proof. ■

THEOREM 6.3: *Let κ be a regular uncountable cardinal.*

- (1) *Suppose we are given M a model of PA which is κ -saturated, and $a_* \in M$ nonstandard. Then we can find a countable set $X \subseteq M$ such that letting N be the Skolem hull of $\{a \in M : M \models a \leq a_*\} \cup X$, we have that the reduct $(N, <)$ to the language of order has an (\aleph_0, \aleph_0) -cut.*
- (2) *There is a cofinality spectrum problem \mathfrak{s} with $M_1^{\mathfrak{s}} = N$ and $\mathfrak{a} \in \text{Or}(\mathfrak{s})$ such that*

$$t_{\mathfrak{s}, \mathfrak{a}} < p_{\mathfrak{s}, \mathfrak{a}};$$

in fact, $t_{\mathfrak{s}, \mathfrak{a}} = \aleph_0$ while $p_{\mathfrak{s}, \mathfrak{a}} \geq \kappa$.

- (3) *If, in the conditions above, we have just that $M \models \text{BPA}$ and for some $r_* < a_*$ nonstandard $M \models \text{“}a_*^{r_*} \text{ exists”}$, this is enough.*

Proof. First we prove (1). Let $A = M \upharpoonright a_*$.

Let $\langle F_n : n < \omega \rangle$ list the Skolem functions of M , each appearing infinitely often (for transparency). Let k_n be the arity of F_n , and without loss of generality $k_n \leq n$.

Let p be the type in the variables $z, x_i (i < \omega), y_i (i < \omega)$ and parameter a_* expressing:

- (a) $z < a_*$ is nonstandard,
- (b) $x_0 = a_*$,
- (c) $y_0 = (a_*)^z$,
- (d) $m < n < \omega$ implies $a_* = x_0 \leq x_m < x_n < y_n < y_m \leq y_0$,
- (e) letting $B_n = \{F_n(e_0, \dots, e_{k_n-1}) : e_\ell \in A \cup \{x_\ell, y_\ell : \ell < n\}\}$, we have that $B_n \cap (x_n, y_n) = \emptyset$.

Let us check that p is consistent. By the κ -saturation of M , this will suffice to show it is realized.

Fix $n < \omega$ and consider a finite fragment of $p \upharpoonright z, x_0, \dots, x_n, y_0, \dots, y_n$. Let ℓ be the maximal exponent appearing in conditions of the form (e). We are looking for $r, a_0, \dots, a_n, b_0, \dots, b_n$ such that

- r is nonstandard, or simply above some given natural number,
- $a_* = a_0 < \dots < a_n < b_n < \dots < b_0 = (a_*)^r$,
- $m \leq n$ implies $B_m \cap (a_m, b_m) = \emptyset$.

For each n , the set B_n is definable in M (since it only involves one function F_n , of arity k_n) and of power c_n , where

$$c_n \approx (|A| + 2n)^{k_n} < a_*^{k_n+1}.$$

Let $B_n(w, v_0, \dots, v_n, v'_0, \dots, v'_n)$ denote the set

$$\{F_n(e_0, \dots, e_{k_n-1}) : e_\ell \leq w \vee e_\ell \in \{v_0, \dots, v_n, v'_0, \dots, v'_n\}\}$$

which will likewise have size $\leq w^{k_n} \cdot (2n + 2)^{k_n} < w^{k_n+1}$. Let

$$\varphi_n(w, v_0, \dots, v_n, v'_0, \dots, v'_n)$$

assert that

$$(\forall t)(\exists s \leq x^{k_n+1} \cdot t)(\exists a \exists b)(a < b \leq s \wedge b - a > t \wedge (a, b) \cap B_n(x) = \emptyset).$$

Then clearly for all $m \leq n$, $M \models \forall w \forall \bar{v} \forall \bar{v}' \varphi_m(w, \bar{v}, \bar{v}')$, recalling that k_n is fixed so the exponential notation abbreviates multiplication.

Choosing r such that

$$(k_0 + 2)(k_1 + 2) \cdots (k_n + 2) \leq r < a_*$$

and

$$b_* = b_0 = (a_*)^r$$

we may then choose $a_1, b_1, \dots, a_n, b_n$ by induction on $\ell \leq n$ such that:

- $k < \ell \implies a_k < a_\ell < b_\ell < b_k,$
- $(a_0, b_0) = (a_*, (a_*)^r),$
- $b_\ell - a_\ell > (a_*)^{n \cdot (n+1-\ell)}.$

For $\ell + 1$, we have

$$b_\ell - a_\ell > (a_*)^{(n+2)(n+1-\ell)}$$

and $B(a_*, a_0, \dots, a_\ell, b_0, \dots, b_\ell)$ is internally a set with $\leq (a_*)^{k_\ell+1} \leq (a_*)^{n+1} < (a_*)^{n+2}$ elements, so there is room.

This completes the verification that p is consistent, therefore (by saturation) realized. For the remainder of the proof, fix realizations $r, a_i (i < \omega), b_i (i < \omega)$ of the type p .

[With a little more care, using $(a_*)^{d_* \cdot (d_*+1-\ell)}$ for $\ell < \omega$, we could alternately have chosen the entire countable sequence by induction, avoiding the appeal to the type and \aleph_1 -saturation.]

Let N be the Skolem hull of $M \upharpoonright a_* \cup \{a_n, b_n : n < \omega\}$, so $N \models PA$. Let $\mathbf{s} = \mathbf{s}_1^+[N]$ be the canonical CSP from Definition 4.3. Then:

- (a) $N \preceq M$.
- (b) $M_{\leq a_*} = N_{\leq a_*}$ so $(N, <^N)$ is saturated below a_* by definition of M .
- (c) $(\langle a_n : n < \omega \rangle, \langle b_n : n < \omega \rangle)$ is an (\aleph_0, \aleph_0) -cut, because by construction the Skolem functions do not fill it.
- (d) Thus, if \mathbf{a} is such that $X_{\mathbf{a}} = M_{\leq a_*} = N_{\leq a_*}$, and, say, $d_{\mathbf{a}} = \max X_{\mathbf{a}}$, then $\mathbf{p}_{\mathbf{s}, \mathbf{a}} \geq \kappa$, whereas if \mathbf{b} is such that $X_{\mathbf{b}} = N_{\leq b_*}$, and, say, $d_{\mathbf{b}} = \max X_{\mathbf{b}}$, then $\mathbf{p}_{\mathbf{s}, \mathbf{b}} = \aleph_0$.

This completes the proof of (1). We continue the argument to prove (2). (3) will follow from the proof.

Let \mathbf{a} be such that $X_{\mathbf{a}} = N_{\leq a_*}$. Recalling the small nonstandard exponent r , let $d_* = r + 1$ and consider the definable subtree $\mathcal{T} \subseteq \mathcal{T}_{\mathbf{a}}$ consisting of sequences of length $< d_*$ of numbers $< a_*$; \mathcal{T} has cardinality a_*^r [the tree $\mathcal{T}_{\mathbf{a}}$ will be at least as large].

Recalling that $N \models PA$, \mathbf{s} is closed under strong exponentiation, so there is an injection from $\mathcal{S}_{\mathbf{a}}$ into some $X_{\mathbf{a}}$. Composing with the Gödel pairing function if needed, we may assume there is a definable injection of $\mathcal{S}_{\mathbf{a}}$ (as a set) into some $X_{\mathbf{b}} \subseteq N$. Applying Conclusion 8.4, we obtain a definable order-isomorphism

between $(\mathcal{S}_a, <_{\mathcal{S}_a})$ onto an initial segment of $M_1^+ = N$. Since \mathcal{T} is the set of all sequences from X_a to itself of length $\leq d_a = a_* \gg r$, the cardinality of \mathcal{T} is $\geq a_*^r = b_*$, so its image must contain the (\aleph_0, \aleph_0) -cut. Because X_b has an (\aleph_0, \aleph_0) -cut, necessarily $(\mathcal{S}, <_{\mathcal{S}})$ has such a cut.

By Lemma 6.2, either X_a has an (\aleph_0, \aleph_0) -cut or else \mathcal{T}_a has a branch of cofinality \aleph_0 with no upper bound. Since X_a is κ -saturated by assumption, we must be in the second case. This shows that $t_{s,a} = \aleph_0$. On the other hand, $p_{s,a} \geq \kappa$, by the hypothesis of saturation. ■

Thus, the local cardinals $p_{s,a}$ and $t_{s,a}$ need not be equal, even in well behaved cofinality spectrum problems, if the model M_1^+ is not uniformly saturated.

7. Characterizing the \triangleleft^* -maximal class

In this section we give the first real evidence that SOP_2 is a dividing line by proving that, under instances of GCH, SOP_2 characterizes maximality in the interpretability order \triangleleft^* which will be defined below. The proof uses cofinality spectrum problems. This answers an open question and also gives evidence for a recent conjecture, as we now explain. The use of GCH comes only from Fact 7.2 below. Recall:

Definition 7.1 (SOP_2 , cf. [26] 1.5): T has SOP_2 if there is a formula $\varphi(\bar{x}, \bar{y})$ which does, meaning that in \mathfrak{C}_T there are parameters $\langle \bar{a}_\eta : \eta \in {}^\omega 2 \rangle$, $\ell(\bar{a}_\eta) = \ell(y)$, such that:

- (1) For each $\eta \in {}^\omega 2$, the set $\{\varphi(\bar{x}, \bar{a}_{\eta \upharpoonright \ell}) : \ell < \omega\}$ is consistent.
- (2) For any two incomparable $\eta, \nu \in {}^\omega 2$, the set $\{\varphi(\bar{x}, \bar{a}_\eta), \varphi(\bar{x}, \bar{a}_\nu)\}$ is inconsistent.

Shelah in [19] had defined an order on theories, a natural weakening of Keisler’s order: $T_1 \triangleleft_* T_2$, which holds, roughly speaking, if there is a third theory T_* which interprets both T_1 and T_2 and whose models M_* have the property that if the reduct to $\tau(T_2)$ is saturated, so is the reduct to $\tau(T_1)$. [See Definition 7.7 below.] This was studied and developed more extensively by Džamonja–Shelah [1] and Shelah–Usvyatsov [26]. In the Shelah–Usvyatsov paper, building on work of Džamonja and Shelah, it was shown that:

Fact 7.2 (Shelah and Usvyatsov [26] 3.15(2), under GCH^{11}): If T is $NSOP_2$ then T is necessarily non-maximal in \triangleleft^* .

In Shelah and Usvyatsov [26] and Džamonja and Shelah [1] it was asked:

Question 7.3 (Question 1.8 of [26]): Does \triangleleft^* -maximality characterize either SOP_3 or SOP_2 , maybe both?

Question 7.4 ([1] Question 3.1): Does SOP_2 imply \triangleleft^* -maximality?

Here we settle the question under GCH, giving a positive answer: it characterizes SOP_2 . (Note that the use of GCH is only in the quoted result 7.2, constituting the known half of the characterization, and not in the present half of the argument. Thus, eliminating GCH from the characterization would involve re-proving 7.2 without the set theoretic hypotheses.) Now let us explain the connection to Keisler's order which motivates this work and our solution. SOP_2 is a property which is not yet well understood and was not known, prior to the present paper, to be a dividing line. Recently, however, we proved the following theorem:

THEOREM E (Malliaris and Shelah [9] Theorem 11.11): *Any theory with SOP_2 is maximal in Keisler's order.*

We conjecture there that SOP_2 characterizes the maximum Keisler class. In the current section, we give strong evidence for this conjecture by proving the result for the order \triangleleft^* , which refines Keisler's order.

We now state the main result of this section: we prove that T is \triangleleft^* -maximal if and only if it has SOP_2 (Theorem 7.13 below). In light of Fact 7.2, it suffices to prove that any theory with SOP_2 is \triangleleft^* -maximal. Since \triangleleft^* refines Keisler's order, even though it is not known whether it is a strict refinement, it is not sufficient to quote Theorem E; rather, we use the technology of cofinality spectrum problems developed for the proof of Theorem E.

Given this result, it is natural to try to say more about the dividing line at SOP_2 , which we do in §9.

Convention 7.5: Throughout this section T denotes a complete countable first-order theory.

¹¹ This hypothesis is missing from the statement in [26], but that proof quotes [1] 3.2, which assumes relevant instances of GCH.

We now review the “interpretability order” \triangleleft^* , introduced in Shelah [19] Definition 2.10 as a natural weakening of Keisler’s order. We first need a definition of “interpretation.”

Definition 7.6 (Interpretations, cf. [1] 1.1): Let T_0 and T_* be complete first-order theories. Suppose that

$$\overline{\varphi} = \langle \varphi_R(\overline{x}_R) : R \text{ a predicate or function symbol of } \tau(T_0), \text{ or } = \rangle$$

is such that each $\varphi_R(\overline{x}_r) \in \tau(T_*)$.

- (1) For any model $M_* \models T_*$, we define the model $N = M_*^{\overline{\varphi}}$ as follows:
 - N is a $\tau(T_0)$ -structure
 - $\text{dom}(N) = \{a : M_* \models \varphi_=(a, a)\} \subseteq M_*$,
 - for each predicate symbol R of $\tau(T_0)$, $R^N = \{\overline{a} : M_* \models \varphi_R[\overline{a}]\}$,
 - for each function symbol f of $\tau(T_0)$ and each $b \in N$, $N \models “f(\overline{a}) = b”$ iff $M_* \models \varphi_f(\overline{a}, b)$, and $M_* \models “\varphi_f(\overline{a}, b) \wedge \varphi_f(\overline{a}, c) \implies b = c”$.

Note that by the last clause, we may restrict to vocabularies with only predicate symbols.

- (2) Say that $\overline{\varphi}$ is an **interpretation** of T_0 in T_* if:
 - each $\varphi_R(\overline{x}_r) \in \tau(T_*)$,
 - for any model $M_* \models T_*$, we have that $M_*^{\overline{\varphi}} \models T_0$.
- (3) If there exists $\overline{\varphi}$ which is an interpretation of T_0 in T_* , say simply that “ T_* interprets T_0 .”

Definition 7.7 (The interpretability order \triangleleft^* , cf. [1] 1.2):

- (1) Let T_0, T_1 be complete first-order theories and λ an infinite regular cardinal. We say that $T_0 \triangleleft_\lambda^* T_1$ if there exists a theory T_* such that:
 - (a) T_* interprets T_0 , witnessed by $\overline{\varphi}_0$, and T_1 , witnessed by $\overline{\varphi}_1$, where “interprets” is in the sense of 7.6.
 - (b) For every model $M_* \models T_*$, if $M_*^{\overline{\varphi}_1}$ is λ -saturated, then $M_*^{\overline{\varphi}_0}$ is λ -saturated.
- (2) We say that $T_0 \triangleleft^* T_1$ if $T_0 \triangleleft_\lambda^* T_1$ for all large enough regular λ .

Discussion 7.8: Definition 7.7 is stated for regular cardinals, but it also makes sense for singular cardinals.

As ultrapowers commute with reducts and the choice of index models is irrelevant¹² clearly \triangleleft^* -equivalence refines \triangleleft -equivalence [equivalence in Keisler’s order]. A priori, one would expect that \triangleleft^* is much weaker but this is not known to be the case. In fact, the known Keisler classes (i.e., of stable theories) coincide with those for \triangleleft^* by [19] 2.11.

We will use a result parallel to that familiar from Keisler’s order:

Fact 7.9 (Shelah [19] Observation 2.11 p. 249): Any theory with the strict order property is \triangleleft^* -maximal [i.e., -maximum].

CLAIM 7.10: Let \mathfrak{s} be a c.s.p. so $\mathfrak{p}_{\mathfrak{s}} = \mathfrak{t}_{\mathfrak{s}}$. Suppose $\mathfrak{a} \in \text{Or}(\mathfrak{s})$ has a $(\mathfrak{t}_{\mathfrak{s}}, \mathfrak{t}_{\mathfrak{s}})$ -cut. Then $\mathcal{T}_{\mathfrak{a}}$ witnesses treetops.

Proof. (Included for completeness, this argument simply adapts the proof of [9] Lemma 6.1, which proved that in this case $\mathcal{T}_{\mathfrak{a} \times \mathfrak{a}}$ witnesses treetops, to show that $\mathcal{T}_{\mathfrak{a}}$ witnesses treetops.) Choose a sequence $(\langle a_{\alpha} : \alpha < \mathfrak{t}_{\mathfrak{s}} \rangle, \langle b_{\alpha} : \alpha < \mathfrak{t}_{\mathfrak{s}} \rangle)$ witnessing the cut. By induction on $\alpha < \mathfrak{t}_{\mathfrak{s}} = \mathfrak{p}_{\mathfrak{s}}$ let us choose a path through $\mathcal{T}_{\mathfrak{a}}$ satisfying the following:

- for each α , c_{α} belongs¹³ to the definable subtree of $\mathcal{T}_{\mathfrak{a}}$ consisting of elements x such that: for each $n < \text{max dom}(x)$, $x(n)$ codes a pair [i.e., is of the form $(a + b)^2 + a$ for elements $a <_{\mathfrak{a}} b \in X_{\mathfrak{a}}$], and if $m < n < \text{max dom}(x)$ and $x(m) = \langle a_1, b_1 \rangle$ and $x(n) = \langle a_2, b_2 \rangle$ then $a_1 <_{\mathfrak{a}} a_2 <_{\mathfrak{a}} b_2 <_{\mathfrak{a}} b_1$.
- $\beta < \alpha \implies c_{\beta} \triangleleft c_{\alpha}$.
- For each α , $n_{\alpha} := \text{max dom}(c_{\alpha})$.
- For each α , $c_{\alpha}(n_{\alpha}) = \langle a_{\alpha}, b_{\alpha} \rangle$.

The construction of this tree follows the template of [9]. At successor steps, we concatenate. At limit steps $\alpha < \mathfrak{t}_{\mathfrak{s}}$, we first choose an upper bound c_* for the sequence built so far, by definition of $\mathfrak{t}_{\mathfrak{s}}$. Let $n_* = \text{max dom}(c_*)$. Then since $X_{\mathfrak{a}}$ is pseudofinite, the nonempty, bounded set

$$\{n < n_* : c_*(n) = \langle a, b \rangle \text{ and } a <_{\mathfrak{a}} a_{\alpha} \text{ and } b_{\alpha} <_{\mathfrak{a}} b\}$$

¹² Keisler [5] proved that if \mathcal{D} is a regular ultrafilter on λ and $M \equiv N$ in a countable language then M^{λ}/\mathcal{D} is λ^+ -saturated iff N^{λ}/\mathcal{D} is λ^+ -saturated.

¹³ Alternately, rather than asking that c_{α} code a pair, ask that if $t_1 < t_2 < t_3$ are successive elements of $\text{lg}(c_{\alpha})$, then either $c_{\alpha}(t_1) < c_{\alpha}(t_3) < c_{\alpha}(t_2)$ or else $c_{\alpha}(t_2) < c_{\alpha}(t_3) < c_{\alpha}(t_1)$. Then at limit steps, the condition is that the interval defined by $(c_{\alpha}(n), c_{\alpha}(n + 1))$ includes a_{α} .

has a maximal element n_{**} . Then let

$$c_\alpha = c_* \upharpoonright_{n_{**}} \hat{\ } \langle a_\alpha, b_\alpha \rangle.$$

Having completed the construction of the sequence, notice that it is unbounded in \mathcal{T}_a , because if it were to have an upper bound c_* , then either of the elements in the pair coded by $c_*(\max \text{dom}(c_*))$ would realize our original cut, contradiction. This completes the proof. ■

We will use one further fact about trees in CSPs, which explains that $\text{Tr}(\mathbf{s})$ is quite robust.

Fact 7.11 ([9] Claim 2.14): If $(\mathcal{T}, \triangleleft_a)$ is a definable subtree of $(\mathcal{T}_a, \triangleleft_a)$ and $\langle c_\alpha : \alpha < \kappa \rangle$ is a \triangleleft_a -increasing sequence of elements of \mathcal{T} , then $\langle c_\alpha : \alpha < \kappa \rangle$ has an upper bound in \mathcal{T} if and only if it has an upper bound in \mathcal{T}_a .

In the language of Definition 3.4, Fact 7.11 shows that if \mathcal{T}' is a definable subtree of \mathcal{T}_a , \mathcal{T}_a witnesses treetops *and* there is an infinite increasing sequence in \mathcal{T}' with no upper bound, then \mathcal{T}' witnesses treetops. But there may not be such a sequence, as, e.g., in the trivial case when \mathcal{T}' consists just of the root.

LEMMA 7.12: *Let T_1 be any theory with SOP_2 . Then T_1 is maximal in \ast .*

Proof. Let T_0 be the theory $Th(\omega + \omega^*)$ of an infinite discrete linear order with a first and last element. As just explained, it will suffice to show that there is a theory T_* with the following properties:

For any $M_* \models T_*$,

- there exists an interpretation of T_1 in M_* , denoted $M_*^{[\varphi_1]}$,
- there exists an interpretation of T_0 in M_* , denoted $M_*^{[\varphi_0]}$,
- for any infinite regular cardinal λ , if $M_*^{[\varphi_1]}$ is λ -saturated then $M_*^{[\varphi_0]}$ is λ -saturated.

We build the theory T_* in several steps. Let $\varphi(x, y)$ be a formula of T_1 which has SOP_2 . Let $n_\varphi = \ell(y)$; here and elsewhere, we may omit overlines. Without loss of generality, $\tau(T_1)$ has only predicates. To begin, fix:

- A model $M_1 \models T_1$ and a set of parameters $\langle a_\eta : \eta \in {}^\omega > \omega \rangle$ witnessing SOP_2 for φ . [Recall that by compactness, we may assume our tree witnessing SOP_2 is infinitely branching, with paths consistent and incomparable nodes inconsistent.]

- A model $M_2 \models (\mathcal{H}(\aleph_1), \epsilon)$. Without loss of generality, the domains of M_1 and M_2 are disjoint, and the relation ϵ does not appear in $\tau(M_1)$.

Let M be the model which is the disjoint union of M_1 and M_2 , expanded by the additional relations and functions $\{N, P, F\}$ as follows and no additional structure. (Without loss of generality, these symbols do not occur in $\tau(M_1)$ or $\tau(M_2)$.)

- N^M names ω in M_2 .
- P^M names $\{a_\eta : \eta \in {}^{>n}\omega\} \subseteq \text{dom}(M_1)$.
- F^M is a unary function with domain ${}^{\omega}>\omega \subseteq M_2$ and range $P^M \subseteq M_1$, such that $F^M(\eta) = a_\eta$.

For each $n < \omega$, let M_n be the model M expanded further by the constant c where $c^{M_n} = n \in \omega = N^M = N^{M_n}$. Let \mathcal{D} be a nonprincipal ultrafilter on ω . Let $N_* = \prod_n M_n / \mathcal{D}$ and let $T_* = Th(N_*)$. This completes the construction of T_* , the theory of the ultrapower as just defined. Note that

$$\tau(T_*) = \tau(N_*) = \tau(M_1) \cup \tau(M_2) \cup \{N, P, F, c\}.$$

In any model M of T_* , the constant c^M is a nonstandard element of N^M , i.e., of the (nonstandard) copy of ω .

In addition to T_* and N_* as just defined, let M_* be an arbitrary but fixed model of T_* which satisfies: $M_* \upharpoonright_{\tau(T_*)}$ is λ -saturated. Note that T_* interprets T_0 and T_1 , that is, there exist $\overline{\varphi}_0, \overline{\varphi}_1$ witnessing 7.6. This is because $M_* \upharpoonright_{\tau(T_*)} \models T_1$ by construction, while

$$N_0 := N^{M_*} \upharpoonright c^{M_*} = (\{a \in M_* : M_* \models a \in c\}, \in^{M_*}) \models T_0.$$

To analyze saturation, we define a cofinality spectrum problem as follows. First, let $\text{Or}^0(\mathfrak{s})$ be the smallest set containing the set of all finite nonempty initial segments of N^{M_*} , in each case letting $d_{\mathfrak{a}} = \max X_{\mathfrak{a}}$. [We may use any non-empty order in $\mathcal{H}(\aleph_0)$.] For each $\mathfrak{a}, \mathfrak{b} \in \text{Or}^0(\mathfrak{s})$, let $\mathfrak{a} \times \mathfrak{b}$ be the definable, pseudofinite linear order on $X_{\mathfrak{a}} \times X_{\mathfrak{b}}$ given by the Gödel pairing function, and let $d_{\mathfrak{a} \times \mathfrak{b}} = (d_{\mathfrak{a}}, d_{\mathfrak{b}})$. Let $\text{Or}(\mathfrak{s})$ be the closure as just described of $\text{Or}^0(\mathfrak{s})$ under Cartesian product, as well as initial segment. [If \mathfrak{b} is an initial segment of \mathfrak{a} , define $d_{\mathfrak{b}} = \min\{\max X_{\mathfrak{b}}, d_{\mathfrak{a}}\}$.]

Suppose for a moment that $M_* = N_*$, i.e., suppose we are really in the case of the ultrapower. Then for each $\mathfrak{a} \in \text{Or}(\mathfrak{s})$, $\varphi_{\mathfrak{a}}$ is a formula $\varphi(x, y, z)$ over the empty set. Because we are in N_* , for each suitable parameter \bar{a} , the set $\varphi(x, x, \bar{a})$ is contained in the ultrapower of finite sets. Let $\psi(w, z)$ be such that for each

suitable $\bar{a} \in {}^{\ell(\bar{z})}M_n$ [though by construction, we can restrict to the case where $\bar{a} \subseteq \mathbb{N}^{M_n}$] we have that $\mathcal{T}_{\varphi, \bar{a}} = \{\eta : M \models \psi(\eta, \bar{a})\}$ is the set of finite sequences of members of the finite set $X_{\varphi, \bar{a}} = \{b : M \models \varphi(b, b, \bar{a})\}$ of length $< \max X_{\varphi, \bar{a}}$. Let $\triangleleft = \{(\eta, \nu) : \eta, \nu \in \mathcal{T}_{\varphi, \bar{a}} \text{ and } \eta \text{ is an initial segment of } \nu\}$. These are definable in M_n by the choice of its theory. The length and evaluation functions lg and val can be defined likewise. By Los' theorem, these formulas will define the appropriate trees in the ultrapower. Thus, we get a cofinality spectrum problem. Moreover, for each given $\varphi_{\mathbf{a}}$, each \bar{a} of length $\ell(z)$, and each $t < \omega$, we have that in each index model, the tree $\mathcal{T} = \mathcal{T}_{\varphi, \bar{a}[t]}$ is finite, so its flattening $(\mathcal{S}, <_{\mathcal{S}})$ [in the sense of 3.2 above] can be injectively mapped into N^{M_n} . That is, $M_n \models$ "there exists an injective order-preserving map of $(\mathcal{S}, <_{\mathcal{S}})$ into N ". As the flattening is uniformly definable from ψ , and we are in a model with sufficient set theory, such an injection exists also in the ultrapower and so \mathbf{s} has (strong) exponentiation.

Returning to the case of arbitrary $M_* \equiv N_*$, as we are in a model with sufficient set theory, note that the existence of such associated trees and injections are elementary properties of each given φ and ψ . Thus, also in this more general case, \mathbf{s} is a cofinality spectrum problem with exponentiation.

Having defined \mathbf{s} , we now prove that $\mathfrak{t}_{\mathbf{s}} \geq \lambda$. First we verify that it is sufficient to look at one-dimensional trees, i.e., that treetops are witnessed by some $\mathcal{T}_{\mathbf{a}}$ where $X_{\mathbf{a}} \subseteq N^{M_*}$. By Theorem 3.11, there is a $(\mathfrak{t}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}})$ -cut in some $X_{\mathbf{a}}$, $\mathbf{a} \in \text{Or}(\mathbf{s})$. If $X_{\mathbf{a}}$ is one-dimensional, then by Claim 7.10, we finish. If not, $X_{\mathbf{a}}$ is contained in some finite Cartesian power of N^{M_*} . Let $f : X_{\mathbf{a}} \rightarrow N^{M_*}$ be a definable injection, given by applying the Gödel pairing function finitely many times. By the definition of \mathbf{s} , this is an order-preserving map. (This appeal to the definition is not necessary: we could simply use that such a map is an injection of sets, so by Conclusion 8.4 of the next section, there is an injective order-preserving map into some one-dimensional $X_{\mathbf{b}}$.) Necessarily $X_{\mathbf{b}}$ will have a $(\mathfrak{t}_{\mathbf{s}}, \mathfrak{t}_{\mathbf{s}})$ -cut, so again by Claim 7.10, we finish.

So to show $\mathfrak{t}_{\mathbf{s}} \geq \lambda$, it will suffice to show that for any one-dimensional \mathbf{a} , and any $\kappa < \lambda$, any κ -indexed strictly increasing sequence $\langle \eta_{\alpha} : \alpha < \kappa \rangle$ in $\mathcal{T}_{\mathbf{a}}$ has an upper bound. Recall the constant c from the signature of N_* and M_* . Since $M_* \equiv N_*$, any one-dimensional $X_{\mathbf{a}}$ is contained in a nonstandard initial segment, i.e., for some m , $\mathbb{N}^{M_*} \models "c \leq m \text{ and } X_{\mathbf{a}} \subseteq M_* \upharpoonright m"$. By Fact 7.11, without loss of generality $X_{\mathbf{a}}$ is the full initial segment below m and $\mathcal{T}_{\mathbf{a}}$ is the

full tree $m > m$. In $(\mathcal{H}(\aleph_0), \epsilon)^{M_*}$ we define $\mathcal{T}_a = ({}^m > m, \triangleleft)$ and let

$$\psi(y, m) := (\exists \eta \in \mathcal{T}_a)(F(\eta) = y)$$

define the subset of P^{M_*} corresponding to the image $F(\mathcal{T}_a)$. Now if

$$\bar{\eta} = \langle \eta_\alpha : \alpha < \kappa \rangle$$

is increasing in \mathcal{T}_a , then recalling the SOP_2 -formula φ from the beginning of the proof,

$$p_{\bar{\eta}} = \{ \varphi(x, F(\eta_\alpha)) : \alpha < \kappa \}$$

is a consistent partial type¹⁴ in $M_* \upharpoonright \tau(T_1)$. By the assumption of λ -saturation in that signature, some $d \in M_*$ realizes p , hence

$$\{ \eta \in \mathcal{T}_a : \varphi[d, F(\eta)] \}$$

is a subset of a branch of \mathcal{T}_a by definition of SOP_2 and of F , and is definable in M_* . Then the set

$$\{ a \in X_a : a < d_a \wedge (\exists \eta)(\eta \in \mathcal{T}_a \wedge \varphi[d, F(\eta)] \wedge a \in \text{dom}(\eta)) \}$$

of lengths of such elements is a definable, nonempty, bounded subset of X_a , so contains a greatest element a_* . Any η along the distinguished branch whose domain contains a_* will be an upper bound for $\bar{\eta}$. This completes the proof that $\mathfrak{t}_s \geq \lambda$.

Recall that we had set $N_0 := N^{M_*} \upharpoonright c^{M_*}$ as the domain for our interpretation of T_0 in M_* . Since \mathfrak{s} is a CSP with exponentiation, $\mathfrak{p}_s = \mathfrak{t}_s = \lambda$ by Theorem 3.11. By definition of \mathfrak{p}_s , there are no (κ_1, κ_2) -cuts in any X_a , for $\mathfrak{a} \in \text{Or}(\mathfrak{s})$ and $\kappa_1, \kappa_2 < \mathfrak{t}_s$. This in particular is true for \mathfrak{a} with $X_a = \{ a : M_* \models a < c \}$ and $d_a = \max X_a$. In fact, for arbitrarily large $a \in N^{M_*}$, $\text{Or}(\mathfrak{s})$ contains \mathfrak{a} such that $X_a \supseteq N \upharpoonright_{\leq a}$, in which, therefore, we have no (κ_1, κ_2) -cuts.

Finally, let us prove that $M_*^{[\varphi_0]}$, i.e., the τ_0 -submodel whose domain is N_0 , is λ -saturated. We know that for models of T_0 , every formula is a Boolean combination of the formulas $x = y$, $x < y$, and $\varphi_k(x, z) = (\exists^{1^k} y)(x < y < z)$. Thus $N_0 \models T_0$ is λ -saturated iff every cut (C_1, C_2) of $N_0 \subseteq N^{M_*}$ of cofinality (κ_1, κ_2) is filled, where κ_1, κ_2 are regular and $\kappa_1 + \kappa_2 \leq \lambda$. This was proved in the previous paragraph.

This completes the proof. ■

¹⁴ The fact that comparability of elements in the domain of F is reflected in the consistency of their images will carry over from the models M_n to N_* by Los' theorem and from there to M_* by elementary equivalence.

THEOREM 7.13 (GCH): T is \triangleleft^* -maximal if and only if it has SOP_2 .

Proof. By Fact 7.2 (which assumes relevant instances of GCH) and Lemma 7.12. ■

8. Useful tools and additional definitions

Before turning to a structure theory for $NSOP_2$, we prove several additional facts about CSPs: Claim 8.3 and Conclusion 8.4, which show that from a suitable bijection of sets, we can recover an order-isomorphism. It follows that the assumption that the order on all pairs was given by the Gödel pairing function in the CSP constructed in Lemma 7.12 could be weakened, as mentioned in that proof. Tying up loose ends, we show that one of the main consequences of 2.1(6) can be recovered in weak hereditary CSPs, Claim 8.5, and discuss the barrier to fully recovering [9] for such CSPs in 8.6. Finally, we include a definition of “strong” CSPs, natural when the underlying model M_1^+ is totally ordered.

Definition 8.1: Let $\mathbf{b} \in \text{Or}(\mathbf{s})$. We say that $Z \subseteq X_{\mathbf{b}}$ is **small** in $X_{\mathbf{b}}$ if:

- (a) there is some definable V , $Z \subseteq V \subset X_{\mathbf{b}}$ such that $M_1^+ \models$ “there does not exist $x \in \mathcal{T}_{\mathbf{b}}$ such that x is a bijection from $X_{\mathbf{b}}$ into V ”,
- (b) if $Z \neq \emptyset$, then $\max(Z) < d_{\mathbf{b}}$.

When $Z \subseteq X_{\mathbf{b}}$ is small in $X_{\mathbf{b}}$ and is an initial segment, we call it a **small initial segment**.

Remark 8.2: Condition 8.1(b) allows for concatenation.

CLAIM 8.3: Suppose we are given a weak cofinality spectrum problem \mathbf{s} , $\mathbf{b} \in \text{Or}(\mathbf{s})$, $h \in M_1^+$, $(W, <_W)$ a definable and pseudofinite linear order, and Z a small initial segment of $Y_{\mathbf{b}}$, such that:

$$M_1^+ \models \text{“} h \text{ is a partial bijection with } W \subseteq \text{dom}(h), \text{ range}(h) \subseteq Z\text{”}.$$

Then $(W, <_W)$ is internally order-isomorphic to an initial segment of $Y_{\mathbf{b}}$.

Proof. Let $(W^*, <_{W^*})$ denote the definable set $h(W) \subseteq X_{\mathbf{b}}$ with the definable linear order given by $x <_{W^*} y \iff h^{-1}(x) <_W h^{-1}(y)$. This allows us to identify $(W^*, <_{W^*})$ definably with $(W, <_W)$, and now we prove that $(W^*, <_{W^*})$ is internally order-isomorphic to an initial segment of $(X_{\mathbf{b}}, \leq_{\mathbf{b}})$.

Let \mathcal{T} be the definable subtree of $\mathcal{T}_{\mathbf{b}}$ given by $\varphi(x)$, which says: “ $x \in \mathcal{T}_{\mathbf{b}}$ is a one-to-one order-preserving function from an initial segment of $(X_{\mathbf{b}}, \leq_{\mathbf{b}})$ onto an initial segment of $(W^*, <_{W^*})$ ”. Thus if $a <_{\mathbf{b}} b$ are in $\text{dom}(x)$ then $x(a) <_{W^*} x(b)$.

First, note that the set $\mathcal{F} = \{x \in \mathcal{T} : \varphi(x)\}$ is linearly ordered. This is because any two elements $x, x' \in \mathcal{T}$ must agree on the first element of their domain; and if neither of x, x' extends the other, then the set in their common domain on which they agree is nonempty and definable. But if the last element of this set is not $\max \text{dom}(x)$, we get a contradiction since the order is pseudofinite.

Consider the subset of $X_{\mathbf{b}}$ given by

$$\{a \in X_{\mathbf{b}} : \text{there is } x \text{ such that } \varphi(x) \text{ and } a \in \text{dom}(x)\}.$$

As this subset is nonempty and definable, by assumption it has a last element $a_* \in X_{\mathbf{b}}$. Let $x_* \in \mathcal{T}$ be a function witnessing this, i.e., such that $a_* \in \text{dom}(x_*)$. Necessarily $x_* = \max \mathcal{F}$.

There are three cases.

CASE 1. The desired case: $\text{range}(x_*) = W^*$.

CASE 2. Not case 1, but $\text{dom}(x_*) = X_{\mathbf{b}}$. Then

$$M_1^+ \models \text{“}x_* \in \mathcal{T}_{\mathbf{b}} \text{ is an injection of } X_{\mathbf{b}} \text{ into } Z\text{”}$$

contradicting Definition 8.1.

CASE 3. Not case 1 or 2, so $\text{dom}(x_*) \subsetneq X_{\mathbf{b}}$ and $\text{range}(x_*) \subsetneq W^*$. By Definition 8.1 and the hypotheses of the Claim, “not case 2” implies $\text{lg}(x_*) < d_{\mathbf{b}}$. Writing S for successor, note that as the orders are pseudofinite, the function $x_* \hat{\ } \langle S(f(a_*)) \rangle$ is well defined (meaning the function extending x_* by the additional condition $S(a_*) \mapsto S(f(a_*))$ is well defined). We may concatenate, so this new function belongs to \mathcal{T} , contradicting the choice of a_* .

As cases 2 and 3 are contradictory, we are necessarily in case 1, which completes the proof. ■

CONCLUSION 8.4: *Suppose that \mathbf{s} is a weak cofinality spectrum problem, and:*

- (1) $N \subseteq M_{1,\mathbf{s}}$ is definable and linearly ordered.
- (2) Arbitrarily large initial segments of N are orders for \mathbf{s} ; more precisely, there is $\psi \in \Delta$ so that for any $a \in N$ there is $\mathbf{a} \in \text{Or}(\mathbf{s})$ with

$$\{b \in N : \models b \leq a\} \subseteq \{b \in M_{1,\mathbf{s}} : \models \psi(b, b, a)\} = X_{\mathbf{a}} \subseteq N$$

and $a < d_{\mathbf{a}}$.

- (3) Cardinality of initial segments of N grows internally; more precisely, letting $\theta = \theta(w, y)$ define the tree associated to $\psi(x_1, x_2, y)$, we have that $M_{1,\mathbf{s}} \models “(\forall y_0 \in N)(\exists y_1 \in N)(\text{there does not exist } z \in \theta(w, y_1) \text{ such that } z \text{ is a bijection from } \psi(x, x, y_1) \text{ into } \psi(x, x, y_0))”$.

Let $(W, <_W)$ be a definable, pseudofinite linear order in the model. If there is an internal injection f of sets from W into some initial segment of N , then for some $\mathbf{b} \in \text{Or}(\mathbf{s})$ with $X_{\mathbf{b}} \subseteq N$ there exists an internal order-isomorphism g from $(W, <_W)$ onto an initial segment of $Y_{\mathbf{b}}$.

Proof. By the hypothesis, we can find \mathbf{b}_0 such that

$$X_{\mathbf{b}_0} \supseteq f(W) \quad \text{and} \quad d_{\mathbf{b}_0} > \max(f(W)).$$

Let $\psi(z)$ be the formula in Δ defining the pseudofinite linear order corresponding to the initial segment below z , and let $\theta(z)$ define its associated tree. Then $M_{1,\mathbf{s}} \models “(\exists z)(\text{there does not exist } x \in \theta(z) \text{ such that } x \text{ is a bijection from } \psi(z) \text{ into } X_{\mathbf{b}_0})”$. Let c be any such z . Again by hypothesis, there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{b}} \supseteq \{x : x \leq c\}$ and $d_{\mathbf{b}} > c$. Now apply Claim 8.3. ■

CLAIM 8.5: *If \mathbf{s} is a nontrivial weak c.s.p. and is hereditarily closed, then there exist nontrivial $\mathbf{a}, \mathbf{a}' \in \text{Or}(\mathbf{s})$ which together satisfy (**) of 2.5.*

Proof. Let \mathbf{b} (so also $\mathcal{T}_{\mathbf{b}}$) be nontrivial. In section 5 of [9], we showed that for any $\mathbf{b} \in \text{Or}(\mathbf{s})$, it is possible to define addition, multiplication, and exponentiation on any element of $\text{Or}(\mathbf{s})$ (that is, to define relations on $X_{\mathbf{b}}$ which have all the same properties as the graphs of these functions, except that they are possibly not total). This does not require any assumptions on pairing. This is done in the proof of [9] Lemma 5.3, essentially as follows. Addition is given by:

$$\varphi_+(x, y, z) = (\exists \eta \in \mathcal{T}_{\mathbf{b}})(\text{lg}(\eta) = y \wedge \eta(0) = x \wedge \eta(y-1) = z \wedge (\forall i)(i < \text{lg}(\eta) \implies \eta(S(i)) = S(\eta(i))).$$

[We omit the parameter $\bar{c}_{\mathbf{b}}$ for readability.] To obtain multiplication, $\varphi_{\times}(x, y, z)$, substitute “ $(i < \text{lg}(\eta) \implies \eta(S(i)) = \eta(i) + x)$ ” as necessary, and define exponentiation $\varphi_{\text{exp}}(x, y, z)$ by substituting in the appropriate place “ $(i < \text{lg}(\eta) \implies \eta(S(i)) = \eta(i) \times x)$,” i.e., requiring that the sequence increment by a factor of x . These are graphs of partial functions, which need not be total. We can therefore define “ x is a prime.” Let $\varphi_4(x, y)$ assert that x is the y th

prime by saying: $y > 0 \wedge (\exists \eta \in \mathcal{T}_{\mathbf{b}})(\lg(\eta) = y \wedge \eta(0) = 2 \wedge \eta(y-1) = x \wedge (\forall i)(i < \lg(\eta) \implies \text{“}\eta(S(i)) \text{ is the } \leq_{\mathbf{b}}\text{-least prime number strictly greater than } \eta(i)\text{”})$). Let $\varphi_5(x, n, m)$ assert that x is divisible by the n th prime precisely m times, by asserting the existence of $\eta \in \mathcal{T}_{\mathbf{b}}$ of length m whose first element is x , whose subsequent elements decrease by a factor of the n th prime and whose last element has no more such factors. Let the formula $\varphi_6(x, \eta)$ assert that $x \in X_{\mathbf{b}}$ is a Gödel code for η by stating: “ $\eta \neq \emptyset, \eta \in \mathcal{T}_{\mathbf{b}}, x > 2$ and for all $i < \lg(\eta)$, writing $m = \eta(i)$, we have that x is divisible by the i th prime precisely $m + 1$ times”. Let $\theta(x)$ assert that

$$(\forall y < x)(\exists z)\varphi_+(x, y, z) \wedge (\forall y < x)(\exists z)\varphi_{\times}(x, y, z) \wedge (\forall y < x)(\exists z)\varphi_{exp}(x, y, z).$$

Let $\psi(y)$ be the formula

$$(\forall x < y)\theta(x) \wedge (\forall \eta \in \mathcal{T}_{\mathbf{b}})((\lg(\eta) < y \wedge (\forall i < \lg(\eta))(\eta(i) < y)) \implies (\exists x)(\varphi_6(x, \eta)))$$

which asserts that Gödel codes exist for all functions from $X_{\mathbf{b}} \upharpoonright_y$ to itself.

For our present case, apply this as follows. Recall that $X_{\mathbf{b}}$ is pseudofinite.

First step: Find nonstandard $n_* \leq d_{\mathbf{b}}$ so that the definable set Z of codes for pairs of elements of $[0, n_*]_{\mathbf{b}}$ is contained in $Y_{\mathbf{b}}$. Second step: Find $n_{**} \in X_{\mathbf{b}}, n_{**} \leq n_*$ still nonstandard, so that $X_{\mathbf{b}}$ contains all codes for functions from $[0, n_{**}]_{\mathbf{b}}$ to itself. Third step: Let m_* be maximal $\leq_{\mathbf{b}} n_{**}$ such that

$$\langle m_*, m_* \rangle \leq n_{**},$$

so necessarily m_* is nonstandard too. Let \mathbf{a}_1 be such that $X_{\mathbf{a}_1} = [0, n_{**}]_{\mathbf{b}}$ and $d_{\mathbf{a}_1} = m_*$.

Now define \mathbf{a} so that $X_{\mathbf{a}} = [0, m_*]_{\mathbf{b}}$ and $d_{\mathbf{a}} = m_*$. What about the desired tree \mathcal{T} of functions from $X_{\mathbf{a}}$ to $X_{\mathbf{a}} \times X_{\mathbf{a}}$? Recalling that $d_{\mathbf{a}_1} = m_*$, this tree is naturally isomorphic to the definable sub-tree $\mathcal{T} \subseteq \mathcal{T}_{\mathbf{a}_1}$ whose elements are functions whose range consists only of codes for pairs of elements each of which are $\leq m_*$. By construction, the codes for elements of $\mathcal{T}_{\mathbf{a}_1}$ and therefore for elements of \mathcal{T} form a definable subset of $X_{\mathbf{b}}$. Finally, let $\mathbf{a}' = \mathbf{b}$.

This completes the proof. ■

Discussion 8.6: By a similar argument, in any weak CSP which is hereditarily closed, for some nontrivial $\mathbf{a} \in \text{Or}(\mathbf{s})$ we have available nontrivial elements of $\text{Or}(\mathbf{s})$ which can be thought of as canonically representing any given one of the finite Cartesian powers $\mathbf{a} \times \dots \times \mathbf{a}$ with the desired ordering, e.g., one derived from repeated applications of the Gödel pairing function. However, as

noted in Discussion 2.5, we don't a priori have Cartesian products of distinct $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$. This prevents us from obtaining the uniformity of functions such as lcf across all $X_{\mathbf{a}}$ which was necessary for the main theorems of [9] to go through.

Definition 8.7 (Strong CSPs): Call \mathbf{s} a **strong** CSP, or **lexicographic** CSP if the demands on Cartesian products from 2.1(5) are replaced by:¹⁵

- (1) if $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$, then internally either $|Y_{\mathbf{a}}| \leq |Y_{\mathbf{b}}|$ or $|Y_{\mathbf{b}}| \leq |Y_{\mathbf{a}}|$,
- (2) if $\mathbf{a} \in \text{Or}(\mathbf{s})$ and $d \leq_{\mathbf{a}} d_{\mathbf{a}}$, then there is $\mathbf{b} \in \text{Or}(\mathbf{s})$ such that:
 - (a) $X_{\mathbf{b}} = X_{\mathbf{a}}, <_{\mathbf{b}} = <_{\mathbf{a}}, d_{\mathbf{b}} = d$,
 - (b) $\mathcal{T}_{\mathbf{b}} = \mathcal{T}_{\mathbf{a}} \upharpoonright \{\eta : (\text{lg}_{\mathbf{a}}(\eta) \leq d) \wedge (\forall n < \max \text{dom}(\eta))(\eta(n) \leq d)\}$,
- (3) if $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$ and h is a definable¹⁶ isomorphism from $(Y_{\mathbf{a}}, <_{\mathbf{a}})$ onto $(Y_{\mathbf{b}}, <_{\mathbf{b}})$ then there is $\mathbf{c} = \mathbf{a} \times \mathbf{b}$ such that $X_{\mathbf{c}} = X_{\mathbf{a}} \times X_{\mathbf{b}}, <_{\mathbf{c}}$ is the lexicographic order, $d_{\mathbf{c}} = (d_{\mathbf{a}}, 0_{\mathbf{b}})$, and $\mathcal{T}_{\mathbf{c}}$ is naturally defined.

Observation 8.8: Any strong CSP is a weak CSP, and the hereditary closure of a strong CSP is a strong CSP.

CLAIM 8.9: If \mathbf{s} is a strong CSP which is hereditarily closed, then:

- (a) \mathbf{s} satisfies Property **(**)** of Discussion 2.5.
- (b) Moreover, for any nontrivial $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$, there is a nontrivial $\mathbf{c} \in \text{Or}(\mathbf{s})$ with either $X_{\mathbf{c}} = X_{\mathbf{a}} \times Y_{\mathbf{b}}$ or $X_{\mathbf{c}} = X_{\mathbf{b}} \times Y_{\mathbf{a}}$.
- (c) Suppose that for every $\mathbf{a} \in \text{Or}(\mathbf{s})$ there is $\mathbf{a}' \in \text{Or}(\mathbf{s})$ such that $X_{\mathbf{a}}$ is internally isomorphic to a subset of $X_{\mathbf{a}'}$ and $d_{\mathbf{a}'} = \max(X_{\mathbf{a}'})$. Then \mathbf{s} is closed under Cartesian products.

Proof. (1) By Claim 8.5 and Observation 8.8.

(2) Fix nontrivial $\mathbf{a}, \mathbf{b} \in \text{Or}(\mathbf{s})$. By condition 8.7(1), without loss of generality, $|Y_{\mathbf{a}}| \leq |Y_{\mathbf{b}}|$ witnessed by an internal partial isomorphism h . Since Condition 8.7(3) requires h to be surjective, let $d = h(d_{\mathbf{a}}) \in Y_{\mathbf{b}}$. Since \mathbf{s} is hereditarily closed and $d \leq d_{\mathbf{b}}$, apply Observation 2.12 to find $\mathbf{b}'' \in \text{Or}(\mathbf{s})$ such that

$$X_{\mathbf{b}''} = Y_{\mathbf{b}} \quad \text{and} \quad d_{\mathbf{b}''} = d.$$

Now $h : Y_{\mathbf{a}} \rightarrow Y_{\mathbf{b}''}$ is onto, so by condition 2.7(c) there is $\mathbf{c} = \mathbf{a} \times \mathbf{b}''$ with $X_{\mathbf{c}} = X_{\mathbf{a}} \times X_{\mathbf{b}''} = X_{\mathbf{a}} \times Y_{\mathbf{b}}$ and $d_{\mathbf{c}}$ is nonstandard, thus \mathbf{c} is nontrivial.

¹⁵ So here Δ is retained.

¹⁶ Not necessarily via Δ .

(3) As $d_{\mathbf{a}'} = \max(X_{\mathbf{a}'})$, $d_{\mathbf{b}'} = \max(X_{\mathbf{b}'})$, we have that $Y_{\mathbf{a}'} = X_{\mathbf{a}'}$, $Y_{\mathbf{b}'} = X_{\mathbf{b}'}$, and the previous condition (2) shows that their Cartesian product exists. The hypothesis of (3) allows us to find Cartesian products for any two elements of $\text{Or}(\mathfrak{s})$ by first isomorphically embedding them in suitable larger elements \mathbf{a}' , \mathbf{b}' , finding $\mathbf{c}' = \mathbf{a}' \times \mathbf{b}'$ and then applying hereditary closure. ■

9. Towards a structure theory for $NSOP_2$

In Section 7, we gave the first real evidence that the strong tree property SOP_2 is a dividing line. Motivated by this result, we now look for the beginnings of a structure theory for $NSOP_2$. The key objects are so-called higher formulas, defined using ultrafilters. The main results are first, Theorem 9.21, which characterizes $NSOP_2$ in terms of few higher formulas; second, the Symmetry Lemma 9.15, which characterizes $NSOP_3$ in terms of symmetric inconsistency for higher formulas; and third, Theorem 9.30, which proves that SOP_2 is sufficient for a certain kind of exact saturation to fail.

Convention 9.1: Throughout this section, T is a complete first order theory and $\mathfrak{C} = \mathfrak{C}_T$ is a monster model for T .

SOP_2 was defined in 7.1 above. SOP_3 was first defined in Shelah [19] as a weakening of the strict order property; note that in Definition 9.2, the case where $\psi = \neg\varphi$ is the strict order property.

Definition 9.2 ([19] 2.20, [26] 1.3): T has SOP_3 if there is an indiscernible sequence $\langle \bar{a}_i : i < \omega \rangle$ and formulas $\varphi(\bar{x}, \bar{y})$, $\psi(\bar{x}, \bar{y})$ such that:

- (1) $\{\varphi(\bar{x}, \bar{y}), \psi(\bar{x}, \bar{y})\}$ is contradictory,
- (2) for each $k < \omega$, the following is a consistent partial type:

$$\{\psi(\bar{x}, \bar{a}_j) : j \leq k\} \cup \{\varphi(\bar{x}, \bar{a}_i) : i > k\},$$

- (3) for $j < i$, the set $\{\varphi(\bar{x}, \bar{a}_i), \psi(\bar{x}, \bar{a}_j)\}$ is contradictory.

It is known that SOP_3 implies SOP_2 but it is open whether, on the level of theories, the converse is true; so it is possible that the Symmetry Lemma below will also characterize SOP_2 . (Still, for pairs (T, Δ) the converse fails since SOP_2 , SOP_3 are known to be distinct at the level of formulas.)

We first look for a useful way to capture the asymmetry of SOP_3 . This approach relates to the idea of “semi-definability” from [17] VII.4.

Definition 9.3: If \mathcal{D} is an ultrafilter on $A \subseteq \mathfrak{C}^m$, then for any set $B \subseteq \mathfrak{C}$ we define

$$\text{Av}(\mathcal{D}, B) = \{\psi(\bar{x}; \bar{b}) : \bar{b} \in B, \psi \in [\mathcal{L}], \{\bar{a} \in A : \models \psi(\bar{a}, \bar{b})\} \in \mathcal{D}\},$$

so this is an element of $\mathbf{S}^m(B)$.

Definition 9.4: Let \mathcal{D} be an ultrafilter on ${}^m A$. (We sometimes write $m = \mathbf{m}(\mathcal{D})$ for this arity.)

- (1) We say that the infinite indiscernible sequence $\bar{b} = \langle \bar{b}_s : s \in I \rangle$ is **based** on \mathcal{D} when

$$\text{tp}(\bar{b}_s, A + \bar{\mathbf{b}}_{>s}) = \text{Av}(D, A + \bar{\mathbf{b}}_{>s})$$

where $\bar{\mathbf{b}}_{>s} = \bigcup \{\bar{b}_t : t \in I, s <_I t\}$.

- (2) For each \mathcal{D} , let $\text{ob}(\mathcal{D}) = \text{ob}(\mathcal{D}, A)$ be the set of such $\bar{\mathbf{b}}$, i.e., the set of all infinite indiscernible sequences based on \mathcal{D} (assuming the monster model \mathfrak{C} is well defined).

In Definition 9.4, the elements approach A ; of course we could have inverted the order. Given such an ultrafilter \mathcal{D} and an infinite indiscernible sequence built from it, we may naturally ask when a given formula instantiated along this sequence is consistent.

Definition 9.5: Let $A \subseteq \mathfrak{C}$ and $\varphi = \varphi(\bar{x}, \bar{y}) = \varphi(\bar{x}, \bar{y}, \bar{c})$ for $\bar{c} \in \mathfrak{C}$.¹⁷

- (1) Let $\text{uf}_\varphi(A)$ be the set of ultrafilters \mathcal{D} on ${}^{\text{lg}(\bar{y})} A$ such that if

$$\bar{\mathbf{b}} = \langle \bar{b}_s : s \in I \rangle \in \text{ob}(\mathcal{D}),$$

then

$$\{\varphi(\bar{x}, \bar{b}_s, \bar{c}) : s \in I\}$$

is a consistent partial type.

- (2) For each $k < \omega$, let $\text{uf}_{\varphi, k}(A)$ be the set of ultrafilters \mathcal{D} on ${}^{\text{lg}(\bar{y})} A$ such that if $\bar{\mathbf{b}} = \langle \bar{b}_s : s \in I \rangle \in \text{ob}(\mathcal{D})$ and $s_0 <_I \dots <_I s_{k-1}$, then

$$\{\varphi(\bar{x}, \bar{b}_{s_\ell}, \bar{c}) : \ell < k\}$$

is a consistent partial type. So when $k = \infty$, we may omit it.

We arrive at a key definition of the section: higher formulas $(\varphi, A, \mathcal{D})$, triples such that φ is indeed consistent when instantiated along any $\bar{b} \in \text{ob}(\mathcal{D}, A)$. Two

¹⁷ Usually, \bar{c} is empty, and in any case, we can just incorporate it into the parameters \bar{y} .

subsequent theorems of the section will characterize SOP_2 and SOP_3 in terms of the interaction of these higher formulas.

Definition 9.6 (Higher formulas): Let $\bar{m} = (m_0, m_1)$. Writing m instead of \bar{m} means $m_0 = 1$ and $m = m_1$, or that m_0 is clear from the context. Let $\mathbf{HF}_k^{\bar{m}}$ be the set of triples $\rho = (\varphi, A, \mathcal{D})$ where $\varphi = \varphi(\bar{x}, \bar{y})$ with $\ell(\bar{x}) = m_0$, $\ell(\bar{y}) = m_1$, and no more parameters, \mathcal{D} is an ultrafilter on ${}^{m_1}A$ and

$$\mathcal{D} \in \text{uf}_{\varphi, k}(A).$$

Definition 9.7: In the context of 9.6:

- (1) Let $\mathbf{HF}^{\bar{m}} = \mathbf{HF}_{\infty}^{\bar{m}} = \bigcap_k \mathbf{HF}_k^{\bar{m}}$.
- (2) We may wish to consider higher formulas over a fixed set A , or using a fixed formula φ , in which case our notation will be:
 - (a) Given A , we may write $\mathbf{HF}_k^{\bar{m}}(A)$, or “ $(\varphi, \mathcal{D}) \in \mathbf{HF}_k^{\bar{m}}(A)$ ”.
 - (b) Let $\mathbf{HF}^{\bar{m}}(A) = \mathbf{HF}_{\infty}^{\bar{m}}(A) = \bigcap_k \mathbf{HF}_k^{\bar{m}}(A)$.
 - (c) Given $\varphi = \varphi(\bar{x}, \bar{y})$, with $\ell(\bar{x}) = m_0$ and $\ell(\bar{y}) = m_1$, we may write $\mathbf{HF}^{\varphi} = \mathbf{HF}_{\varphi}^{\bar{m}}, \mathbf{HF}_{\varphi, k}^{\bar{m}}(A)$, etc., where the subscript φ means we restrict to triples whose first element is φ with the given partition of variables.
 - (d) Call the elements of $\mathbf{HF}_{\varphi}(A)$ “higher φ -formulas over A .”

Convention 9.8: In Definition 9.5(1), we may say “ \mathcal{D} is an ultrafilter **over** A ” without mentioning m_1 when it is clear from the context.

We would like to study pairwise consistency or inconsistency of higher formulas as follows. Suppose we are given $(\varphi_0, A_0, \mathcal{D}_0)$, $(\varphi_1, A_1, \mathcal{D}_1)$, $\langle b_{0,s} : s \in I_0 \rangle \in \text{ob}(A_0, \mathcal{D}_0)$ and $\langle b_{0,t} : t \in I_1 \rangle \in \text{ob}(A_1, \mathcal{D}_1)$. If we choose $s \in I_0$ and $t \in I_1$, will $\varphi_0(\bar{x}, \bar{b}_{0,s})$ and $\varphi_1(\bar{x}, \bar{b}_{1,t})$ be consistent? What if we choose finitely many instances from each list? The specter of SOP_3 suggests that we should first fix an interpolation of I_0 and I_1 into a single linear order I and pay attention to the relative position of the indices s and t . The notation we now introduce in 9.9–9.10 is one way to handle this (most of the time we use $\sigma = 2$).

Definition 9.9 (Partitions of linear orders, [27] Definition 1.39): Let $K_{p, \sigma}$ be the class of triples

$$(I, <_I, (P_i^I)_{i < \sigma})$$

where I is linearly ordered by $<_I$ and $\langle P_i^I : i < \sigma \rangle$ is a partition of I .

Definition 9.10: Fix σ and suppose $\langle (\varphi_i, A_i, D_i) : i < \sigma \rangle$ is a sequence of higher formulas. Let

$$\bar{A} = \langle A_i : i < \sigma \rangle$$

and let

$$\bar{D} = \langle D_i : i < \sigma \rangle.$$

Define

$$\text{ob}(\bar{D}) = \text{ob}(\bar{D}, \bar{A})$$

to be the set of $\bar{\mathbf{b}} = \langle \bar{b}_s : s \in I \rangle$ such that:

- (1) $I \in K_{p,\sigma}$,
- (2) if $s \in P_i^I$ then:
 - $\ell g(\bar{b}_s) = \mathbf{m}(D_i)$,
 - $\text{tp}(\bar{b}_s, \bigcup_{j < \sigma} A_j \cup \bar{\mathbf{b}}_{>s}) \subseteq \text{Av}(D_i) := \text{Av}(D_i, \mathfrak{C}_\tau)$.

We may write A instead of \bar{A} when all the A_i are the same. We may write $i = \mathbf{i}(s) = \mathbf{i}(s, I)$ and we may write $I_{\bar{\mathbf{b}}} = I[\bar{\mathbf{b}}]$ for I .

Note that in Definition 9.10, we do not require that $\bar{\mathbf{b}}$ be indiscernible; in fact, it may consist of sequences of differing lengths, if the $\mathbf{m}(D_i)$ differ.

Definition 9.11 (n -inconsistent higher formulas): Assume that for $\ell = 0, 1$,

$$\rho_\ell = (\varphi_\ell, A_\ell, D_\ell) \in \mathbf{HF}_{k_\ell}^{m_\ell}.$$

- (1) We say (φ_0, A_0, D_0) is **n -contradictory** to (φ_1, A_1, D_1) when: for every $\bar{\mathbf{b}} \in \text{ob}(\langle (D_0, D_1), (A_0, A_1) \rangle)$ and every $s_0 < \dots < s_{2n-1}$ with $s_\ell \in P_0^I[\bar{\mathbf{b}}]$ for $\ell < n$ and $s_\ell \in P_1^I[\bar{\mathbf{b}}]$ for $\ell \in [n, 2n)$, we have that

$$\{\varphi_0(\bar{x}, \bar{b}_{s_\ell}) : \ell < n\} \cup \{\varphi_1(\bar{x}, \bar{b}_{s_\ell}) : \ell \in [n, 2n)\}$$

is contradictory.

- (2) In “ n -contradictory,” if $n = 1$ we may omit it and writing $n = \infty$ means “for some n ”. Of course, “ n -consistent” is the negation.
- (3) We say that (φ_0, A_0, D_0) and (φ_1, A_1, D_1) are **mutually n -contradictory** when $(\varphi_\ell, A_\ell, D_\ell)$ is n -contradictory to $(\varphi_{1-\ell}, A_{1-\ell}, D_{1-\ell})$ for $\ell = 0, 1$. On the symmetry of this notion, see Lemma 9.15 below.
- (4) “The set of $\{\rho_i = (\varphi_i, A_i, D_i) : i \in S\}$ are pairwise n -contradictory” will mean that each pair is mutually n -contradictory.

Discussion 9.12: (1) In 9.11(1) we could have allowed the choice of elements from the two partitions to alternate. However, we will see this is immaterial for $NSOP_3$, and moreover is a little less natural when replacing n -contradictory higher formulas by 1-contradictory derived formulas arising as n -fold conjunctions.

(2) This definition enforces an order between the elements s_i in different partitions; one could give a different definition requiring only that $s_0 < \dots < s_{n-1}$ and $s_n < \dots < s_{2n-1}$.

Definition 9.13: Say that T has **symmetric inconsistency** if, fixing $m = \ell(\bar{x})$, for any¹⁸ $m_1, m_2 < \omega$, $\varphi_\ell = \varphi_\ell(\bar{x}_{[m]}, \bar{y}_{[m_\ell]})$ and $(\varphi_\ell, A_\ell, \mathcal{D}_\ell) \in \mathbf{HF}^{m_\ell}$ for $\ell = 1, 2$, we have that $(\varphi_1, A_1, \mathcal{D}_1)$ is ∞ -contradictory to $(\varphi_2, A_2, \mathcal{D}_2)$ iff $(\varphi_2, A_2, \mathcal{D}_2)$ is ∞ -contradictory to $(\varphi_1, A_1, \mathcal{D}_1)$.

As desired, this definition picks up on the asymmetry of SOP_3 :

CLAIM 9.14: *If T has SOP_3 , then 1-inconsistency is not symmetric, i.e., T has symmetric inconsistency in the sense of 9.13.*

Proof. Let the sequence $\langle \bar{a}_n : n < \omega \rangle$ and the formulas $\varphi(\bar{x}, \bar{y}), \psi(\bar{x}, \bar{y})$ witness SOP_3 , see 9.2. Let $A = \{\bar{a}_n : n < \omega\}$. Let \mathcal{D} be an ultrafilter on ${}^{lg(\bar{y})}A$ such that $\{\bar{a}_n : n > k\} \in \mathcal{D}$ for every $k < \omega$. Let $\rho_1 = (\varphi, A, \mathcal{D}), \rho_2 = (\psi, A, \mathcal{D})$. To see that ρ_2 is 1-consistent with ρ_1 but ρ_1 is not 1-consistent with ρ_2 , let I be the linear order $\omega + \omega^*$. Let (P_0, P_1) be any partition of I into two infinite sets, and let $\langle \bar{b}_s : s \in I \rangle$ be as in 9.10 for $((\mathcal{D}, \mathcal{D}), (A, A))$. ■

LEMMA 9.15 (Symmetry lemma): *For T complete the following are equivalent.*

- (1) T is SOP_3 .
- (2) T has symmetric inconsistency.

Proof. The direction (1) implies (2) is given by Claim 9.14.

For the other direction, we will show how an instance of symmetric inconsistency gives rise to SOP_3 . For notational simplicity, we use $m = \ell(\bar{x}) = 1$, since for SOP_3 the arity does not matter. Suppose we are given $m_0, m_1 < \omega$ and $(\varphi_\ell, A_\ell, \mathcal{D}_\ell) \in \mathbf{HF}^{m_\ell}$ for $\ell = 0, 1$.

¹⁸ The notation $\varphi_\ell(\bar{x}_{[m]}, \bar{y}_{[m_\ell]})$ means that $\ell(\bar{x}) = m$ and $\ell(\bar{y}) = m_\ell$.

Let $A = A_0 \cup A_1$, $m = m_0 + m_1$ and

$$D = D_0 \times D_1$$

$$= \{X \subseteq {}^m A : \{\bar{a}_0 \in {}^{m_0} A_0 : \{\bar{a}_1 \in {}^{m_1} (A_1) : \bar{a}_0 \hat{\ } \bar{a}_1 \in X\} \in D_1\} \in D_0\}.$$

By construction, D is an ultrafilter on ${}^m A$. Suppose I is a linear order and

$$\bar{\mathbf{a}} = \langle \bar{a}_s = \bar{a}_{0,s} \hat{\ } \bar{a}_{1,s} : s \in I \rangle$$

is an indiscernible sequence based on D , with $\ell(a_{\ell,s}) = m_\ell$, thus $\langle \bar{a}_{\ell,s} : s \in I \rangle$ is also an indiscernible sequence based on D_ℓ for $\ell = 0, 1$. For any $h : I \rightarrow \{0, 1\}$, we may consider the partition given by

$$I_h = (I, <_I, (P_i^h)_{i < 2}) \quad \text{where } P_i^h = \{s : h(s) = i\}.$$

Then $I_h \in K_{p,\{0,1\}}$ and $\bar{\mathbf{a}}_h := \langle a_{h(s),s} : s \in I \rangle$ is I_h -indiscernible based on (D_0, D_1) .



Figure 1. A sample partition of I ; I indexes a sequence of $m_0 + m_1$ -tuples which is indiscernible based on D . Thus, restricting to the first m_0 elements of tuples with indices in the black regions gives an indiscernible sequence based on D_0 , whereas restricting to the last m_1 elements of tuples with indices in the grey regions gives an indiscernible sequence based on D_1 .

Moreover, if $\langle s_\alpha : \alpha < \omega + \omega \rangle$ is $<_I$ -increasing, then:

(1) _{n} the following are equivalent:

- (a) $\{\varphi_0(x, \bar{a}_{0,s_\alpha}) : \alpha < n\} \cup \{\varphi_1(x, \bar{a}_{1,s_\omega+\alpha}) : \alpha < n\}$ is contradictory.
- (b) (φ_0, A_0, D_0) is n -contradictory to (φ_1, A_1, D_1) .



Figure 2. Let the image indicate that we instantiate φ_0 n times along the D_0 -indiscernible sequence indexed by the black region and φ_1 n times along the D_1 -indiscernible sequence indexed by the grey region.

(2) _{n} the following are equivalent:

- (a) $\{\varphi_1(x, \bar{a}_{1,s_\alpha}) : \alpha < n\} \cup \{\varphi_0(x, \bar{a}_{0,s_\omega+\alpha}) : \alpha < n\}$ is contradictory.
- (b) (φ_1, A_1, D_1) is n -contradictory to (φ_0, A_0, D_0) .

Figure 3. Let the image indicate that we instantiate φ_1 n times along the D_1 -indiscernible sequence indexed by the grey region and φ_0 n times along the D_0 -indiscernible sequence indexed by the black region. Then for any given n , a disparity in consistency between Figures 2 and 3 leads, by taking conjunctions, to an instance of SOP_3 .

Let $\bar{y} = \bar{y}_0 \widehat{\bar{y}}_1$,

$$\varphi_0^+ = \varphi_0^+(x, \bar{y}) = \varphi_0^+(x, \bar{y}_0, \bar{y}_1) = \varphi_0(x, \bar{y}_0),$$

and

$$\varphi_1^+ = \varphi_1^+(x, \bar{y}) = \varphi_1^+(x, \bar{y}_0, \bar{y}_1) = \varphi_1(x, \bar{y}_1),$$

i.e., these are the given formulas formally considered as having more variables. Then for each n , to $(1)_n$ above, we may add the equivalent condition:

(c) $_n$ $\{\varphi_0^+(x, \bar{a}_{s_\alpha}) : \alpha < n\} \cup \{\varphi_1^+(x, \bar{a}_{s_{\omega+\alpha}}) : \alpha < n\}$ is contradictory.

Likewise, for each n , to $(2)_n$ above, we may add the equivalent condition:

(f) $_n$ $\{\varphi_1^+(x, \bar{a}_{s_\alpha}) : \alpha < n\} \cup \{\varphi_0^+(x, \bar{a}_{s_{\omega+\alpha}}) : \alpha < n\}$ is contradictory.

Now if it is not the case that for all n (c) $_n$ iff for all n (f) $_n$, we have a witness to SOP_3 for T (given by the conjunctions of n copies of φ_0 and of φ_1 , respectively). This completes the proof. ■

As a corollary of the proof of Lemma 9.15, we have:

COROLLARY 9.16: *Assume T is $NSOP_3$. If $p_\ell = (\varphi_\ell, A_\ell, \mathcal{D}_\ell)$ for $\ell = 0, 1$ and p_0 is n -consistent with p_1 for every n (i.e., not ∞ -contradictory), then if $I \in K_\sigma^P$, $\sigma = 2$ and we have $\langle \bar{b}_s : s \in I \rangle$ as usual, then $\{\varphi_\ell(x, \bar{b}_s) : s \in P_\ell^I, \ell < 2\}$ is consistent.*

We now work towards Theorem 9.21, using higher formulas to characterize $NSOP_2$. First, we show that having SOP_2 means many pairwise 1-contradictory higher formulas. Recall that:

Definition 9.17: $\text{Ded}^+(\lambda) = \sup\{|\text{lim}(\mathcal{T})|^+ : \mathcal{T} \subseteq {}^\lambda 2 \text{ is nonempty, closed under initial segments and has no } \triangleleft\text{-maximal members and has cardinality } \leq \lambda\}$.

CLAIM 9.18: Assume $\varphi(\bar{x}, \bar{y})$ has SOP_2 in T and $\lambda < \mu < \text{Ded}(\lambda)$. Then there is $A \subseteq \mathfrak{C}_T$ of cardinality λ and $D_\alpha \in \text{uf}_\varphi(A)$ for $\alpha < \mu$ such that $\langle (\varphi, D_\alpha) : \alpha < \mu \rangle$ are pairwise 1-contradictory.

Proof. Let the tree \mathcal{T} witness that $\mu < \text{Ded}(\lambda)$ and let $\nu_\alpha \in \text{lim}(\mathcal{T})$ (i.e., the “leaves”) for $\alpha < \mu$ be pairwise distinct. Let $\langle \bar{a}_\eta : \eta \in \mathcal{T} \rangle$ be such that:

- (a) $(\varphi(\bar{x}, \bar{a}_\eta), \varphi(\bar{x}, \bar{a}_\nu))$ are contradictory when $\eta \perp \nu$ are from \mathcal{T} ,
- (b) $\{\varphi(x, \bar{a}_{\eta \upharpoonright \alpha}) : \alpha \leq \text{lg}(\eta)\}$ is a consistent partial type for $\eta \in \mathcal{T}$.

Let $A = \cup\{\bar{a}_\eta : \eta \in \mathcal{T}\}$ and for $\alpha < \mu$ let D_α be an ultrafilter on ${}^{\text{lg}(\bar{y})}A$ concentrating on the branch ν_α , i.e., such that:

$$\text{if } \beta < \text{lg}(\nu_\alpha), \alpha < \mu \text{ then } \{\bar{a}_\rho : \nu_\alpha \upharpoonright \beta \trianglelefteq \rho \triangleleft \nu_\alpha\} \in D_\alpha.$$

Clearly these ultrafilters are as desired. ■

We will need notation for finitary approximations to SOP_2 -trees.

Definition 9.19: We say that (T, φ) has $NSOP_{2,n}$ when there are no $\bar{b}_\eta \in {}^{\text{lg}(\bar{y})}\mathfrak{C}$ for $\eta \in {}^{n>2}$ such that

- (1) $\eta \perp \nu \Rightarrow \varphi(\bar{x}, \bar{b}_\eta), \varphi(\bar{x}, \bar{b}_\nu)$ are incompatible,
- (2) for $\eta \in {}^{n2}$, $\{\varphi(\bar{x}, \bar{b}_{\eta \upharpoonright \ell}) : \ell < n\}$ is a type.

Fact 9.20: $(T, \varphi) \models NSOP_2$ iff $\bigvee_n ((T, \varphi) \models NSOP_{2,n})$.

We now arrive at the second theorem of the section, which shows how from many pairwise contradictory higher formulas we may build an SOP_2 -tree, complementing 9.18. Recall that by the Symmetry Lemma 9.15 above, as $NSOP_2$ implies $NSOP_3$, being contradictory is a symmetric notion.

THEOREM 9.21: For a theory T the following are equivalent:

- (1) For every infinite A and formula φ , there are no more than $|A|$ pairwise 1-contradictory higher φ -formulas over A .
- (2) T has $NSOP_2$.

Proof. In Claim 9.18, it was shown that SOP_2 implies many pairwise 1-contradictory higher formulas. So it remains to prove the other direction: many pairwise 1-contradictory higher formulas imply SOP_2 .

STEP 0: Setup. Let $\lambda = |A|^+$ and, by Fact 9.20, let n be such that (T, φ) has $NSOP_{2,n}$.

Assume for a contradiction that:

- (1) $(\varphi, D_\alpha) \in \mathbf{HF}_\varphi^m(A)$ for $\alpha < \lambda$ are pairwise 1-contradictory¹⁹ and
- (2) fixing some infinite linear order I , let $\bar{\mathbf{b}}_\alpha = \langle b_{\alpha,s} : s \in I \rangle \in \text{ob}(D_\alpha)$ for $\alpha < \lambda$.

(Recall that the definition of 1-contradictory is for any such $\bar{\mathbf{b}}$.) We will use just that $\lambda = \text{cf}(\lambda) > |A| \geq \aleph_0$.

STEP 1: Approximations. We define the set AP of approximations (to a full $SOP_{2,n}$ -tree) to be the set of \mathbf{x} consisting of:²⁰

(1.1) $\Lambda \subseteq {}^{n>}2$, \bar{a} where:

- (a) Λ is non-empty downward closed,
- (b) $\bar{a} = \langle \bar{a}_\eta : \eta \in \Lambda \rangle$ with each $\bar{a}_\eta \in {}^{\text{lg}(\bar{y})}A$,
- (c) if $\eta \perp \nu$ are from Λ then $\varphi(\bar{x}, \bar{a}_\eta), \varphi(\bar{x}, \bar{a}_\nu)$ are incompatible,
- (d) if $\eta \in \Lambda$ then $\{\varphi(\bar{x}, \bar{a}_{\eta \upharpoonright \ell}) : \ell \leq \text{lg}(\eta)\}$ is a consistent partial type,
- (e) if $\nu \wedge \langle 1 \rangle \in \Lambda$ then $\nu \wedge \langle 0 \rangle \in \Lambda$.

(1.2) $\bar{\mathcal{U}} = \langle \mathcal{U}_\eta : \eta \in \text{frt}(\Lambda) \rangle$, where:

- (a) $\text{frt}(\Lambda) := \{\eta \in \Lambda : \text{lg}(\eta) < n - 1 \text{ and } \eta \wedge \langle 0 \rangle \notin \Lambda \text{ or } \eta \wedge \langle 1 \rangle \notin \Lambda\}$ (the “frontier” for our inductive construction of a tree, i.e., the nodes without two immediate successors),
- (b) each $\mathcal{U}_\eta \subseteq \lambda$ has cardinality λ (the intention is a set of indices for the $\bar{\mathbf{b}}_\alpha$ from Step 0),
- (c) $\mathcal{U}_\eta \cap \mathcal{U}_\nu = \emptyset$ for $\eta \neq \nu$,
- (d) if $\eta \sqsubseteq \nu$ are from $\text{frt}(\Lambda)$, $k < \omega$, $s_0, \dots, s_{k-1} \in I, k + \text{lg}(\eta) \leq n$ and $\alpha \in \mathcal{U}_\nu$, then

$$\{\varphi(x, \bar{a}_{\eta \upharpoonright \ell}) : \ell \leq \text{lg}(\eta)\} \cup \{\varphi(x, \bar{b}_{\alpha, s_\ell}) : \ell < k\}$$

is a consistent partial type,

- (e) if $\eta \in \text{frt}(\mathbf{x}), s \in I, \alpha \in \mathcal{U}_\eta$ and $\nu \in \Lambda, \neg(\nu \sqsubseteq \eta)$, then $\varphi(\bar{x}, \bar{b}_{\alpha, s}), \varphi(\bar{x}, a_\nu)$ are incompatible.

Note the role of the two kinds of parameters: the $\bar{b}_{\alpha,s}$ from Step 0, and the parameters \bar{a}_η for the tree. Informally, the \mathcal{U}_η tell us in which sequences we can expect to continue our consistent partial type while maintaining inconsistency elsewhere.

¹⁹ Here $\mathbf{HF}_{\varphi,n}^m(A)$ suffices.

²⁰ Alternately, we could consider: in $\Lambda_x, \text{frt}(\mathbf{x})$ is a set of pairwise incomparable elements, e.g. the \triangleleft -maximal $\eta \in \Lambda$ of length $< n - 1$. This is simpler here, but then the induction step would require two steps: add $\eta \wedge \langle 0 \rangle$ or $\eta \wedge \langle 1 \rangle$ for some $\eta \in \text{frt}(\mathbf{x})$.

We define a two-place relation \leq_{AP} on AP in the natural way: $\mathbf{x} \leq_{AP} \mathbf{y}$ iff

- (a) $\mathbf{x}, \mathbf{y} \in AP$,
- (b) $\Lambda_{\mathbf{x}} \subseteq \Lambda_{\mathbf{y}}$,
- (c) $\bar{a}_{\mathbf{x},\eta} = \bar{a}_{\mathbf{y},\eta}$ for $\eta \in \Lambda_{\mathbf{x}}$,
- (d) if $\eta \in \text{frt}(\Lambda_{\mathbf{x}}) \cap \text{frt}(\Lambda_{\mathbf{y}})$, then $\mathcal{U}_{\mathbf{y},\eta} \subseteq \mathcal{U}_{\mathbf{x},\eta}$,
- (e) if $\eta \in \text{frt}(\Lambda_{\mathbf{x}}), \nu \in \text{frt}(\Lambda_{\mathbf{y}})$ and $\eta \triangleleft \nu$ but $\nu \upharpoonright (\ell g(\eta) + 1) \notin \Lambda_{\mathbf{x}}$, then $\mathcal{U}_{\mathbf{y},\nu} \subseteq \mathcal{U}_{\mathbf{x},\eta}$.

STEP 2: Strategy. Clearly \leq_{AP} is a partial order on AP. By choice of n in Step 0, if $\mathbf{x} \in AP$ then $\Lambda_{\mathbf{x}} \not\subseteq^{n>2}$. Thus, to obtain a contradiction (and complete the proof) it will suffice to show that:

- (2.1) there is $\mathbf{z} \in AP$ with $\Lambda_{\mathbf{z}} = \{ \langle \rangle \}$, i.e., $AP \neq \emptyset$,
- (2.2) if $\mathbf{x} \in AP$, then there is $\mathbf{y} \in AP$ such that $|\Lambda_{\mathbf{x}}| < |\Lambda_{\mathbf{y}}|$; in fact, $\mathbf{x} <_{AP} \mathbf{y}$.

STEP 3: Verifying Condition (2.1). Let $s_0 <_I \dots <_I s_n$. For each $\alpha < \lambda$ clearly $\mathfrak{C} \models (\exists \bar{x}) \bigwedge_{\ell \leq n} \varphi(\bar{x}, \bar{b}_{\alpha, s_\ell})$. So by (1) and (2) of Step 0,

$$\mathcal{X}_\alpha = \left\{ \bar{a} \in {}^{\ell g(\bar{y})}A : \mathfrak{C} \models (\exists \bar{x}) \left[\varphi(\bar{x}, \bar{a}) \wedge \bigwedge_{\ell=1}^n \varphi(x, \bar{b}_{\alpha, s_\ell}) \right] \right\} \in D_\alpha.$$

For each $\alpha < \lambda$, choose $\bar{b}_\alpha \in \mathcal{X}_\alpha$. [So \bar{b}_α is “canonically consistent” with the partial φ -type given along the sequence $\bar{\mathbf{b}}_\alpha$ by any n members.] As $|A| < \lambda$, the set ${}^{\ell g(\bar{y})}A$ has cardinality $< \lambda = \text{cf}(\lambda)$, so for some \bar{b} the set

$$\mathcal{U} = \{ \alpha < \lambda : \bar{b}_\alpha = \bar{b} \}$$

has cardinality λ .

Now define \mathbf{z} by:

- $\Lambda_{\mathbf{z}} = \{\langle \rangle\}$,
- $\bar{a}_{\langle \rangle} = \bar{b}$,
- $\mathcal{U}_{\langle \rangle} = \mathcal{U}$,

and \mathbf{z} is as required.

STEP 4: Proving Condition (2.2). Let $\Lambda = \Lambda_{\mathbf{x}}$.

The situation at the inductive step is essentially as follows. We would like the tree to become full, so we choose ϱ which is minimal for the property of not having two successors; say, $\varrho \hat{\ } \mathbf{i}$ is missing (there are minor adjustments at the end of this step depending on whether it has a successor at 0 or 1). We try to find a corresponding $\bar{a}_{\varrho \hat{\ } \mathbf{i}}$ for the missing successor of this node, and its set of compatible indices $\mathcal{U}_{\varrho \hat{\ } \mathbf{i}}$, subject to the following constraints:

- (i) consistency of $\bar{a}_{\varrho \hat{\ } \mathbf{i}}$ with comparable nodes, (1.1)(d),
- (ii) inconsistency of $\bar{a}_{\varrho \hat{\ } \mathbf{i}}$ with incomparable nodes, (1.1)(c),
- (iii) consistency of $\bar{a}_{\varrho \hat{\ } \mathbf{i}}$ with large subsets of \mathcal{U}_{ν} , for compatible ν for (1.2)(d),
- (iv) inconsistency of $\bar{a}_{\varrho \hat{\ } \mathbf{i}}$ with \mathcal{U}_{η} , for incompatible η , for (1.2)(e),
- (v) disjointness of $\mathcal{U}_{\varrho \hat{\ } \mathbf{i}}$ from large subsets of \mathcal{U}_{ν} , for incompatible ν , for (1.2)(c).

In this informal explanation, “large” stands in for the fact that we will also have to refine the other \mathcal{U}_{η} to get actual inconsistency or an actual empty intersection. This completes the description of intent.

Now to begin, choose $\varrho \in \text{frt}(\Lambda)$ to be of minimal length, hence necessarily $\text{lg}(\varrho) < n - 1$ and we can choose $\iota < 2$ such that $\varrho \hat{\ } \langle \iota \rangle \notin \Lambda$. Let $k = n - \text{lg}(\varrho)$.

Fix for a while a choice of a distinguished element of \mathcal{U}_{η} for each $\eta \in \text{frt}(\Lambda)$, that is, $\bar{\alpha} = \langle \alpha_{\eta} : \eta \in \text{frt}(\Lambda) \rangle$ where $\alpha_{\eta} \in \mathcal{U}_{\eta}$, and let $s_0 <_I \dots <_I s_{k-1}$.

By our choice of ϱ and (1.2)(c), for every²¹ $\eta \in \text{frt}(\Lambda_{\mathbf{x}}) \setminus \{\varrho\}$ we have that $\alpha_{\eta} \neq \alpha_{\rho}$, hence the following set belongs to $D_{\alpha_{\varrho}}$:

$$\begin{aligned} \mathcal{X}_{\bar{\alpha}, \eta} &= \mathcal{X}_{\bar{\alpha}, \eta}^1 \cup \mathcal{X}_{\bar{\alpha}, \eta}^2 \text{ where} \\ \mathcal{X}_{\bar{\alpha}, \eta}^1 &= \{\bar{b} \in \text{lg}(\bar{y}) A : \varphi(\bar{x}, \bar{b}), \varphi(\bar{x}, \bar{a}_{\nu}) \text{ are incompatible, where } \nu \in \Lambda \wedge \neg(\nu \leq \rho)\}, \\ \mathcal{X}_{\bar{\alpha}, \eta}^2 &= \{\bar{b} \in \text{lg}(\bar{y}) A : \varphi(\bar{x}, \bar{b}), \varphi(\bar{x}, \bar{b}_{\alpha_{\eta}, s_0}) \text{ are incompatible}\}, \end{aligned}$$

since the (φ, D_{α}) are pairwise 1-contradictory and as $\langle \alpha_{\ell} : \ell \in \text{frt}(\mathbf{x}) \rangle$ is without repetitions by the assumption (1.2)(c) for $\mathcal{X}_{\bar{\alpha}, \eta}^2$ and (1.2)(e) for $\mathcal{X}_{\bar{\alpha}, \eta}^1$.

²¹ If ρ has precisely one successor in the tree, this successor may be from $\text{frt}(\Lambda_{\mathbf{x}})$.

Furthermore, as (1.2)(d) holds for \mathbf{x} , the following set belongs to D_{α_ϱ} :

$$\begin{aligned} \mathcal{Z}_{\bar{\alpha},\varrho} := \{ & \bar{b} \in {}^{\ell g(\bar{y})}A : \{\varphi(\bar{x}, \bar{a}_{\eta|\ell}) : \ell \leq \ell g(\varrho)\} \\ & \cup \{\varphi(\bar{x}, \bar{b})\} \\ & \cup \{\varphi(\bar{x}, \bar{b}_{\alpha_\varrho, s_\ell}) : \ell < n - \ell g(\varrho) - 1\} \text{ is a type} \}. \end{aligned}$$

Let $\mathcal{X}_{\bar{\alpha}} = \cap \{\mathcal{X}_{\bar{\alpha},\eta} : \eta \in \text{frt}(\Lambda) \setminus \{\varrho\} \text{ and not comparable to } \varrho\} \cap \mathcal{Z}_{\bar{\alpha},\varrho}$. Since Λ is finite and D_{α_ϱ} is a filter, $\mathcal{X}_{\bar{\alpha}} \in D_{\alpha_\varrho}$.

By the choice of our sequence of ultrafilters in Step 0, for every $\beta \in \mathcal{U}_{\mathbf{x},\varrho} \setminus \{\alpha_\varrho\}$, we know that $(\varphi, D_{\alpha_\varrho}), (\varphi, D_\beta)$ are 1-contradictory. So

$$\mathcal{Y}_{\bar{\alpha},\beta} = \{\bar{b} \in {}^{\ell g(\bar{y})}A : \varphi(x, \bar{b}), \varphi(x, \bar{b}_{\beta, s_0}) \text{ are incompatible}\}$$

belongs to D_{α_ϱ} . So we may choose $\bar{b}_{\bar{\alpha},\beta} \in {}^{\ell g(\bar{y})}A$ which belongs to $\mathcal{X}_{\bar{\alpha}} \cap \mathcal{Y}_{\bar{\alpha},\beta}$. As $|\mathcal{U}_{\mathbf{x},\varrho}| = \lambda = \text{cf}(\lambda) > |A|$, there is $\bar{b}_{\bar{\alpha}} \in {}^{\ell g(\bar{y})}A$ such that

$$\mathcal{W}_{\bar{\alpha}} = \{\beta \in \mathcal{U}_{\mathbf{x},\varrho} : \bar{b}_{\bar{\alpha},\beta} = \bar{b}_{\bar{\alpha}}\}$$

has cardinality λ .

Recall that all of this is for a fixed $\bar{\alpha}, \langle s_0, \dots, s_{k-1} \rangle$.

Now continue to fix $s_0 <_I \dots <_I s_{k-1}$. For every $\gamma < \lambda$ we let $\bar{\alpha}_\gamma = \langle \alpha_{\gamma,\eta} : \eta \in \text{frt}(\Lambda) \rangle$ be defined by

$$\alpha_{\gamma,\eta} = \min(\mathcal{U}_{\mathbf{x},\eta} \setminus \gamma)$$

which is well defined as $\mathcal{U}_{\mathbf{x},\eta}$ is an unbounded subset of λ . So for every $\gamma < \lambda$, we have that $\bar{\alpha}_\gamma, \bar{b}_{\bar{\alpha}_\gamma}, \mathcal{W}_{\bar{\alpha}_\gamma}$ are well defined. So for some \bar{b}_* we have

$$\mathcal{W}_{\bar{b}_*} = \{\gamma < \lambda : \bar{b}_{\bar{\alpha}_\gamma} = \bar{b}_*\}$$

is an unbounded subset of λ .

We can now define the necessary objects, in two cases. For Case 1 below, the final definitions depend on whether the node we are dealing with has a sibling or not.

CASE 1: $1 = |\Lambda_{\mathbf{x}} \cap \{\varrho^\wedge \langle 0 \rangle, \varrho^\wedge \langle 1 \rangle\}|$

Let ι be such that $\varrho^\wedge \langle \iota \rangle \notin \Lambda_{\mathbf{x}}$. We define \mathbf{y} as follows:

- ⊕ (a) $\Lambda_{\mathbf{y}} = \Lambda_{\mathbf{x}} \cup \{\varrho^\wedge \langle \iota \rangle\}$, hence $\text{frt}(\Lambda_{\mathbf{y}}) = \text{frt}(\Lambda) \cup \{\varrho^\wedge \langle \iota \rangle\} \setminus \{\varrho\}$,
- (b) $\bar{b}_{\mathbf{y},\eta}$ is $\bar{b}_{\mathbf{x},\eta}$ if $\eta \in \Lambda_{\mathbf{x}}$,
- (c) $\bar{b}_{\mathbf{y},\eta}$ is \bar{b}_* if $\eta \in \varrho^\wedge \langle \iota \rangle$,
- (d) if $\eta \in \text{frt}(\Lambda) \setminus \{\varrho\}$, then $\mathcal{U}_{\mathbf{y},\eta}$ is

$$\{\alpha \in \mathcal{U}_{\mathbf{x},\eta} : \text{for some } \gamma < \lambda, \bar{b}_{\bar{\alpha}_\gamma} = \bar{b}_* \text{ and } \alpha = \alpha_{\gamma,\eta}\},$$

(e) if $\eta = \varrho^\wedge \langle \iota \rangle$ then

$$\mathcal{U}_{\mathbf{y},\eta} = \{ \alpha \in \mathcal{U}_{\mathbf{x},\varrho} : \text{for some } \gamma < \lambda, b_{\bar{\alpha}_\gamma} = \bar{b}_* \text{ and } \alpha = \alpha_{\gamma,\varrho} \}.$$

For clause (e) recall $\bar{b}_* \in \mathcal{X}_{\bar{\alpha},\varrho} \subseteq \mathcal{Z}_{\bar{\alpha}_\gamma,\varrho}$.

As $\varrho \in \text{frt}(\mathbf{x})$, we are left with Case 2:

CASE 2: $0 = |\Lambda_{\mathbf{x}} \cap \{ \varrho^\wedge \langle 0 \rangle, \varrho^\wedge \langle 1 \rangle \}|$ and let $\iota = 0$. The only difference is that

- in clause (a), $\text{frt}(\Lambda_{\mathbf{y}}) = \Lambda \cup \{ \varrho^\wedge \langle \iota \rangle \}$

so we have to add

(f) if $\eta = \varrho$, then $\mathcal{U}_{\mathbf{y},\eta} = \mathcal{W}_{\bar{\alpha}_\gamma}$ for some $\gamma < \lambda$ such that $\bar{b}_{\bar{\alpha}_\gamma} = \bar{b}_*$.

STEP 5: Finish. Recalling Step 2, we have shown that the assumption of $|A|^{+}$ -pairwise 1-contradictory higher formulas contradicts $NSOP_2$. This completes the proof. ■

In this theorem, replacing 1-contradictory by n -contradictory would be straightforward as we can replace φ by a conjunction, defined as follows:

Definition 9.22: Given φ and n , let

$$\begin{aligned} \varphi_{[n]} &= \varphi_{[n]}(x, \bar{y}_{[n]}) \\ &= \varphi_{[n]}(\bar{x}, \langle y_i : i < \ell g(\bar{y}) \cdot n \rangle) \\ &= \bigwedge_{\ell < n} \varphi(\bar{x}, \langle y_{\ell g(\bar{y}) \cdot \ell + j} : j < \ell g(\bar{y}) \rangle). \end{aligned}$$

We say $\varphi = \varphi(\bar{x}, \bar{y})$ has $NSOP_2$ **robustly** when no $\varphi_{[n]}$ has SOP_2 .

Observation 9.23: If T is $NSOP_2$, then every $\varphi(\bar{x}, \bar{y}) \in \mathcal{L}(\tau_T)$ has $NSOP_2$ robustly in T .

In the last main result of this section, we apply Theorem 7.13 characterizing the maximal class in \leq^* to prove another property of SOP_2 , related to so-called exact saturation (meaning for which singular κ a given theory T has a model which is κ -saturated but not κ^+ -saturated). Exact saturation was studied in Shelah [23] §2 (pp. 31–37) and in a manuscript of Kaplan and Shelah [4] in preparation, which deal with elementary classes and (\mathbf{D}, κ) -sequence homogeneity.

Definition 9.24 (see [23], [4]): For any theory T , define

$$\text{spec}(T) = \{ \kappa : \kappa \geq |T|, \kappa \text{ singular} \}$$

and there exists a κ -saturated not κ^+ -saturated model of T .

The papers [23] and [4] find sufficient conditions but not necessary and sufficient conditions for exact saturation. In the present paper, we define a pseudo-elementary version²² of this spectrum, which we will connect to SOP_2 via \leq^* .

Definition 9.25: The exact saturation spectrum for T is defined to be the set:

$$\text{ESS}(T) = \{(\kappa, \mu) : \kappa \geq \mu \geq |T|, \kappa \text{ singular and for any } T_1 \supseteq T, |T_1| \leq \mu, \\ \text{there is a } \kappa\text{-saturated not } \kappa^+\text{-saturated} \\ \text{member of } PC(T_1, T)\}.$$

In future work, we hope to be able to get necessary and sufficient conditions for $\text{ESS}(T)$ to be empty (at least restricting ourselves to κ strong limit of large enough cofinality), and it seems plausible that this may be SOP_2 . Here, using the methods of Section 7, we will prove one direction: if T has SOP_2 then $\text{ESS}(T) = \emptyset$, and discuss several open questions. First we recall two known examples.

Fact 9.26 ([23], [4]): Let T be the theory of dense linear order without endpoints. Then for any singular $\kappa \geq |T|$, if $M \models T$ is κ saturated then it is κ^+ -saturated. Thus $\text{spec}(T) = \emptyset$.

Proof. By quantifier elimination, it suffices to show that the cofinality and coinitality of M are at least κ and for any regular κ_1, κ_2 , every (κ_1, κ_2) -pre-cut is filled. Since cofinality and coinitality of the model and of pre-cuts are necessarily regular cardinals, the result is immediate. ■

Fact 9.27 ([23] Example 2.23): There is a theory T with the independence property such that: if T has an exactly κ -saturated model then κ is regular. (In fact it is necessary and sufficient that κ be regular.) Thus $\text{spec}(T) = \emptyset$.

Recall the order \triangleleft^* from Definition 7.7 above. In the following, countability is not essential.

Observation 9.28: Suppose T_0, T_1 are countable theories which are equivalent under the \leq^* order. Then there exists a theory T_* which interprets both T_0 and T_1 , say via $\overline{\varphi_0}$ and $\overline{\varphi_1}$ respectively, such that: for any model $M \models T_*$, and any uncountable κ ,

$$N^{\overline{\varphi_0}} \text{ is } \kappa\text{-saturated if and only if } N^{\overline{\varphi_1}} \text{ is } \kappa\text{-saturated.}$$

²² Compare the results of [17] VI.5 connecting the minimum class in Keisler’s order to saturation properties of a PC -class.

Proof. We may assume T_0 and T_1 have no finite models.

The hypotheses on T_0, T_1 give the following information. Let μ be such that we have T_a witnessing $T_0 \leq^* T_1$ and T_b witnessing $T_1 \leq^* T_0$, and we may assume $|T_a|, |T_b| < \mu$. Let $\overline{\varphi}_\ell^a$ interpret T_ℓ in T_a for $\ell = 0, 1$, and let $\overline{\varphi}_\ell^b$ interpret T_ℓ in T_b for $\ell = 0, 1$. Without loss of generality:

- $\tau(T_0)$ and $\tau(T_1)$ are disjoint.
- $\overline{\varphi}_\ell^a$ says that for any model $M \models T_a$ the universe of $M^{\overline{\varphi}_\ell^a}$ is $(P_\ell^a)^M$, for some unary predicate P_ℓ^a , and T_a implies that P_0^a and P_1^a are disjoint.
- Similarly for T_b .
- $\overline{\varphi}_\ell^a$ is the identity, where this means:
 - $\tau(T_a) \supseteq (\tau(T_0) \cup \tau(T_1))$,
 - $P \in \tau(T_\ell)$ implies $\varphi_{\ell,P}^a = P$, and

$$T_a \vdash (\forall \bar{x})(P(\bar{x}) \implies \forall_{\ell < \lg(\bar{x})} P_{\ell,a}(x)),$$

- similarly for function symbols $F \in \tau(T_\ell)$, adding that they are interpreted as partial functions with domain the predicate P_ℓ^a (alternately, we could have assumed without loss of generality that $\tau(T_\ell)$ has only predicates).
- Each $\overline{\varphi}_\ell^b$ is likewise the identity.
- $\tau(T_a) \cap \tau(T_b) = (\tau(T_0) \cup \tau(T_1))$.

Consider now the statement of the Observation. In order to prove it, it would suffice if we could find a model $M_a \models T_a$ and a model $M_b \models T_b$ such that the interpretations of T_0 in M_a and M_b , respectively, are isomorphic, and likewise the respective interpretations of T_1 in M_a and M_b are isomorphic. Why would this suffice? Because given such M_a and M_b , assuming w.l.o.g. F_0, F_1, A, B are not already used symbols in the language, we may simply consider a third structure M_* whose universe is the disjoint union of a copy of M_a (whose universe is the interpretation of the predicate A) and a copy of M_b (whose universe is the interpretation of the predicate B) and which contains partial function symbols F_0 , interpreted as a bijection between the interpretation of T_0 in the copy of M_a and in the copy of M_b , and F_1 , interpreted as a bijection between the interpretation of T_1 in the copy of M_a and in the copy of M_b . Now $T_* := Th(M_*)$ is the theory we were looking for. That is, in any sufficiently saturated model $M_* \models T_*$, $M_* \upharpoonright A$ behaves like a model of T_a (so saturation transfers from the interpretation of T_0 to that of T_1) and likewise $M_* \upharpoonright B$ behaves like a model of T_b (so saturation transfers from the interpretation of T_1

to that of T_0) and the functions F_0, F_1 identify the two interpretations of T_0, T_1 respectively.

One way to construct such models is to recall that ultrapowers commute with reducts, and that any two elementarily equivalent models have isomorphic ultrapowers. Begin with $M_{0,a} \models T_a$ and $M_{0,b} \models T_b$. Choose the ultrafilter \mathcal{D}_0 so that the \mathcal{D}_0 -ultrapowers of $M_{0,a}^{\overline{\varphi_0^a}}$ and $M_{0,b}^{\overline{\varphi_0^b}}$ are isomorphic. Call these ultrapowers $M_{1,a}$ and $M_{1,b}$ respectively. (As ultrapowers commute with reducts, we may consider $M_{1,a}$ and $M_{1,b}$ as models of the full T_a and T_b respectively.) Consider the model M_c which is the disjoint union of $M_{1,a}$ and $M_{1,b}$ and expand M_c by adding symbols giving the isomorphism between $M_{1,a}^{\overline{\varphi_0^a}}$ and $M_{1,b}^{\overline{\varphi_0^b}}$. Let $M_c^{\overline{\varphi_1^a}}$ have the obvious meaning of $M_{1,a}^{\overline{\varphi_1^a}}$ considered within this model M_c . Next, choose the ultrafilter \mathcal{D}_1 so that $M_c^{\overline{\varphi_1^a}}$ and $M_c^{\overline{\varphi_1^b}}$ are isomorphic. Let M_* be the \mathcal{D}_1 -ultrapower of M_c . In this model, as sketched above, we may expand T_c so as to make the interpretations of T_1 isomorphic. Let T_* be the theory of this expanded model. This completes our construction.

We had fixed an infinite μ so that $|T_a| + |T_b| < \mu$, and $|T_0| + |T_1| = \aleph_0$ by assumption. The construction gives that also $|T_*| < \mu$. ■

COROLLARY 9.29:

- (1) Let T_0 be any theory for which $\text{spec}(T_0) = \emptyset$. Let T_1 be any theory such that T_0 and T_1 are equivalent in the order \leq^* . Then $\text{ESS}(T_1) = \emptyset$.
- (2) Suppose T_0, T_1 are \leq^* -equivalent, as witnessed by T_* with $|T_*| \leq \mu$. Suppose $\kappa \geq \mu$ is singular and $(\kappa, \mu) \notin \text{ESS}(T_0)$. Then $(\kappa, \mu) \notin \text{ESS}(T_1)$.

Proof. (1) Let T_* be a theory witnessing their equivalence, such as that given by Observation 9.28. Let $\overline{\varphi_0}$ and $\overline{\varphi_1}$ witness the interpretations of T_0 and of T_1 , respectively, and let $N^{\overline{\varphi}}$ and $N^{\overline{\varphi^*}}$ denote the respective interpretations in a given model $N \models T_*$. Now let κ be a singular cardinal, $\kappa > |T_1|$ and suppose $M \in PC(T_*, T_1)$ is κ -saturated. Let $N \models T_*$ with $M = N \upharpoonright \tau(T_1)$ witness that $M \in PC(T_*, T_1)$. The fact that M is κ -saturated says precisely that $N^{\overline{\varphi_1}}$ is κ -saturated. By hypothesis of equivalence, $N^{\overline{\varphi_0}}$ is κ -saturated as well. By choice of T_0 , $N^{\overline{\varphi_1}}$ is κ^+ -saturated. Applying the hypothesis of equivalence in the other direction, $N^{\overline{\varphi_0}}$ is κ^+ -saturated. This completes the proof.

(2) As $(\kappa, \mu) \notin \text{ESS}(T_0)$ there is a theory $T_{00} \supseteq T_0$, $|T_{00}| \leq \mu$, and a κ -saturated not κ^+ -saturated member of $PC(T_{00}, T_0)$. In order to make use of this, we would first modify the construction of the theory T_* (by means of a better

choice of T_a, T_b) so that for any $M_* \models T_*$ we have that $M_*^{\overline{\{\varnothing\}}} \in PC(T_{00}, T_0)$. Since $|T_{00}| \leq \mu$, we can do this while preserving $|T_*| \leq \mu$. Then the proof continues as in part (1). ■

THEOREM 9.30: *If T has SOP_2 , then $ESS(T) = \emptyset$. That is, if T has SOP_2 , then for some $T_1 \supseteq T$ of cardinality $|T|$, for every singular $\kappa > |T|$, if $M \in PC(T_1, T)$ is κ -saturated then it is κ^+ -saturated.*

Proof. By Theorem 7.13, Fact 9.26 and Corollary 9.29(1). ■

Question 9.31: Does Theorem 9.30 hold for the theory T_{rg} of the random graph?

In the case of $\text{spec}(T_{rg})$ rather than $ESS(T_{rg})$, note that:

CLAIM 9.32: *If $\kappa > \text{cf}(\kappa) + |T|$ then $\kappa \in \text{spec}(T_{rg})$.*

Proof. Let M be a κ^+ -saturated model of T_{rg} . Let $\langle a_\alpha : \alpha < \kappa \rangle$ be pairwise distinct members of M which form an empty graph, i.e.,

$$\alpha < \beta < \lambda \implies M \models \neg R(a_\alpha, a_\beta).$$

Consider the submodel $N \subseteq M$ whose domain is

$$\{b \in M : (\exists^{<\kappa} \alpha < \kappa)(R(b, a_\alpha))\}.$$

Then N is a model of T_{rg} which is as required: it is κ -saturated but it is not κ^+ -saturated, as $\{R(x, a_\alpha) : \alpha < \kappa\}$ is omitted. ■

Before giving some further evidence, we record here that there are other natural directions these investigations could take.

Discussion 9.33: Instead of asking whether $\kappa \in ESS(T)$ for $\kappa > \text{cf}(\kappa) + |T|$, we may fix a specific way to construct a κ -saturated model and ask if this implies that the constructed model is κ^+ -saturated. For example:

- (a) We may consider T dependent, $|M| = \{a_\alpha : \alpha < \alpha_*\}$, $u_\alpha \in [\alpha]^{<\kappa}$, $\text{tp}(a_\alpha, \{a_\beta : \beta < \alpha\})$ does not split over $\{a_\beta : \beta < \alpha\}$, and M is κ -saturated; see [21].
- (b) As in (a), but we may ask that T has Skolem functions.
- (c) As in (b), and also $\text{tp}(a_\alpha, \{a_\beta : \beta < \alpha\})$ is finitely satisfiable in $\text{Sk}(\{a_\beta : \beta \in u_\alpha\})$; see [17], VII, section 4.
- (d) We may consider models built using $\mathcal{P}^-(n)$ -diagrams in some explicit way.

Returning to ESS, on the positive side we can settle the case of stable T .

CLAIM 9.34: *Let T be a complete theory. If T is stable, $\kappa = \kappa^{|T|} > \text{cf}(\kappa)$, and $\kappa \geq \mu \geq T$, then $(\kappa, \mu) \in \text{ESS}(T)$.*

Proof. Recall that under these cardinal hypotheses there is a model of T of cardinality κ which is κ -saturated; see [17] Chapter III Theorem 3.12. Clearly this model is not κ^+ -saturated. Given any $T_1 \supseteq T$, $|T_1| \leq \mu$, we may build a model of T_1 of cardinality κ in the same way whose restriction to T is κ -saturated but evidently not κ^+ -saturated. ■

Question 9.35: What about simple theories?

Problem 9.36: Determine whether the following is true: for T complete and countable,

$$\text{ESS}(T) = \emptyset \text{ if and only if } T \text{ has } \text{SOP}_2.$$

These investigations suggest some interesting parallel questions for ultrafilters. The reader may recall that if λ is a singular cardinal and the ultrafilter is λ -good, then it is λ^+ -good (since linear order is in the maximal Keisler class, [17].VI.2, a proof similar to that of Fact 9.26 goes through).

Problem 9.37: Suppose λ is singular. Does there exist a regular ultrafilter which λ -saturates but does not λ^+ -saturate ultrapowers of the random graph?

The property of an ultrafilter being “ (λ, \aleph_0) -perfect” was defined in [11].

Problem 9.38: Suppose λ is singular. If \mathcal{D} is a regular ultrafilter which is (κ, \aleph_0) -perfect for every $\kappa < \lambda$, is \mathcal{D} also (λ^+, \aleph_0) -perfect? If not, must it at least produce λ^+ -saturated ultrapowers of the random graph?

10. Balanced pairs of models of PA

Having understood the importance of symmetric cuts of models of PA earlier in the paper, we may wonder what else can be asked about such cuts. In this final section, we consider a question of Kossak, based on work of Schmerl, concerning so-called balanced pairs of models of PA. The analysis applies some work of the second author on EM models.

Definition 10.1: Let $M \preceq_{\text{end}} N$ be models of Peano arithmetic. Call M **balanced** when the cut described by a sequence cofinal in M and a sequence coinital in $N \setminus M$ is symmetric, i.e., the cofinalities of both sides of the cut are equal.

THEOREM F (Schmerl, Theorem 4.6 [16]): *Suppose that $M \models PA$ is saturated. Then there are continuum many different theories of (N, M) where $M \preceq_{\text{end}} N$ and M is balanced.*

Question 10.2 (Kossak [6]): We know of continuum many nonisomorphic balanced (κ, κ) pairs, surely there must be more?

To resolve this we will apply a theorem from Shelah [20]. Let τ be a given signature. For Theorem G which we now quote, there is no restriction on the cardinality of τ , but in our application we will assume τ is countable to get an end extension type. For our purposes, ‘weakly skeleton like’ may be taken as a black box, as will be explained.

THEOREM G ([20] Theorem 3.19 p. 38): *Suppose that $\lambda > \kappa$, \mathcal{K}_λ is a family of τ -models, each of cardinality λ , $\varphi(\bar{x}, \bar{y})$ is an asymmetric formula with vocabulary $\subseteq \tau$, and $\text{lg}(\bar{x}) = \text{lg}(\bar{y}) < \aleph_0$. Suppose that for every linear order J of cardinality λ there are $M \in \mathcal{K}_\lambda$ and $\bar{a}_s \in M$ for $s \in J$ such that $\langle \bar{a}_s : s \in J \rangle$ is weakly $(\kappa, \varphi(\bar{x}, \bar{y}))$ -skeleton-like in M . Then in \mathcal{K}_λ , there are 2^λ pairwise non-isomorphic models.*

COROLLARY 10.3: *Suppose $\lambda > \kappa$, \mathcal{K}_λ is a family of τ -models as in G, each of cardinality λ . Suppose that for every linear order J of cardinality λ there are $M \in \mathcal{K}_\lambda$ and Φ such that $M = EM_\tau(J, \Phi)$. Then in \mathcal{K}_λ , there are 2^λ pairwise non-isomorphic models.*

Proof. The condition “ $\langle \bar{a}_s : s \in J \rangle$ is weakly $(\kappa, \varphi(\bar{x}, \bar{y}))$ -skeleton-like in M ,” [20] Definition 3.1 p. 28, is always satisfied when $M = EM_\tau(J, \Phi)$, [20] 3.2 p. 29. ■

Definition 10.4:

- (1) By FPA_τ we mean τ -**feeble Peano arithmetic**, that is, the vocabulary τ consists of the two-place predicate $\{<\}$ and the axioms say $<$ is a discrete linear order with first and no last element, as well as all the cases of the induction scheme in \mathcal{L}_τ .
- (2) PA_τ is defined similarly but includes $+$, \times .

The “feeble” is because we do not require $+$, \times to be present.

Definition 10.5: Let M be a model which is linearly ordered and let p be a type which is unbounded in M . We call p an **end extension type** of M if every model $N \supset M$, which is generated by M and a single new element realizing p , is an end extension of M .

Example 10.6: The first-order theory of any expansion M of $(\omega, <)$ is a model of $FPA_{\tau(M)}$. The existence of end-extension types in such models is proved by MacDowell–Specker for the case $PA_{\tau M}$, i.e., with plus and times, and without this in [18].

CLAIM 10.7: *Suppose that:*

- (a) $M_* \models PA$ (or just: M_* is a model in a countable signature τ of FPA_{τ}),
- (b) $\aleph_0 \leq \kappa_\ell = \text{cf}(\kappa) \leq \lambda$, for $\ell = 1, 2$,
- (c) $|M_*| \leq \lambda$.

Then the class

$$\{N : N \text{ is an end extension of } M_*, ||N|| = \lambda, \text{cf}(N) = \kappa_2, \\ \text{and the co-initiality of } N \setminus M_* \text{ is } \kappa_1\}$$

has 2^λ pairwise nonisomorphic models.

Proof of 10.7. There are two steps.

STEP 1: *EM* end extensions. Let p be the definition of an end extension type. If we are in a model of PA , this exists by a theorem of Gaifman [2], or MacDowell–Specker [14]; or see [7]. (Gaifman also proved, e.g., that p can be taken to be minimal, which does not follow from [14].) If M_* is just a model of the feeble PA $FPA_{\tau(M_*)}$, then it exists by [18]. [Note: for the rest of the proof, we do not use that the language is countable, just that $|\tau| \leq \lambda$. On eliminating the restriction of countability, see [24].]

Fix a model N so that $M_* \preceq N$ and N is $(2^\lambda)^+$ -saturated, and I a $(< \lambda^+)$ -universal²³ linear order of cardinality $2^{<\lambda} \leq ||N||$, and $I_0 \subseteq I$ of order-type ω . Write $\text{Sk}(X, N)$ for the Skolem hull in N of $X \subseteq N$. By induction on $t \in I$ we may choose $\langle a_t : t \in I_0 \rangle$ such that:

- $a_t \in N$, and
- $tp(a_t, \text{Sk}(M_* \cup \{a_s : s <_I t\}), N)$ is defined by the end extension type p .

Then $\langle a_t : t \in I_0 \rangle$ is an indiscernible sequence over M_* inside N . But N is λ^+ -saturated, hence we can find $a_t \in N$ for $t \in I \setminus I_0$ such that $\langle a_t : t \in I \rangle$ is an indiscernible sequence over M_* in M .

For $J \subseteq I$, let M_J denote $\text{Sk}(M_* \cup \{a_t : t \in J\})$.

Let $\tau_* = \tau(T) \cup \{c_a : a \in M_*\}$, and let $T_* = \text{Th}(M_*, c_a)_{a \in M_*}$, where c_a is interpreted as a .

²³ Recall that a linear order is $< \mu$ -universal if any linear order of cardinality $< \mu$ may be embedded into it.

Then for some Φ , $M_J = EM_{\tau_*}(J, \Phi)$. Moreover, for each such J ,

- M_J is an end extension of M_* ,
- M_J has cardinality $\theta + |J|$, where $\|M_*\| = \theta$,
- if J has no first element then the co-initiality of $M_J \setminus M$ is equal to the coinitiality of J .

STEP 2: Observation. Let \mathcal{K}_λ^0 be the set of models N such that: $M_* \preceq_{\text{end}} N$, $\|N\| = \lambda$. We may already quote 10.3 which, by the analysis just given, shows that \mathcal{K}_λ^0 has 2^λ nonisomorphic models whose coinitiality is determined by the order J which is input to the EM skeleton.

In the next step, we need to show that this count remains true when we restrict to coinitiality κ (since κ was given in advance). As regards symmetric pairs, recall M_* is arbitrary but fixed for the proof, so by the end of this proof we will have the result for all M_* , and in particular for M_* of cofinality κ .

STEP 3: A correspondence. We now translate between \mathcal{K}_λ^0 from Step 2 and the set

$$\mathcal{K}_\lambda = \mathcal{K}_\lambda^1(\kappa, M_*) = \{N : M_* \preceq_{\text{end}} N, \|N\| = \lambda, N \setminus M_* \text{ has coinitiality } \kappa\}$$

For regular $\kappa_1, \kappa_2 \leq \lambda$, whenever $N = EM_{\tau_*}(I, \Phi)$ there is Φ^2 such that:

- (a) $|\tau_\Phi| = \|M_*\| + \kappa_1 + \kappa_2$,
- (b) for any linear order J ,

$$EM_{\tau_\Phi}(J, \Phi^2) = EM_{\tau_*}(\kappa_1^* + J + \kappa_2, \Phi),$$

- (c) thus (by (b) and Step 1) $EM_{\tau_\Phi}(J, \Phi^2)$ is an end extension of M_* and has cofinality κ_2 and co-initiality κ_1 over M_* .

Let $\kappa_1 = \kappa$ and $\kappa_2 = \lambda$, so $\kappa_1 + \kappa_2 + \|M_*\| = \lambda$. So for each linear order J of cardinality λ and each skeleton Φ there is Φ^2 such that $EM_{\tau_\Phi}(J, \Phi^2) \in \mathcal{K}_\lambda$. Quoting 10.3, in \mathcal{K}_λ there are 2^λ pairwise non-isomorphic models. ■

CONCLUSION 10.8: *Let $\kappa \leq \lambda$ be regular. Then there are 2^λ balanced (κ, κ) pairs of models of PA of cardinality λ , i.e., 2^λ many pairs (N, M) where $M \preceq_{\text{end}} N$, the cut described by a sequence cofinal in M and a sequence coinitial in $N \setminus M$ is symmetric of cofinality κ , and $|N| = \lambda$.*

Proof. Apply 10.7 in the case where M_* has cofinality κ . ■

Discussion 10.9: So in fact, to get many pairs of models we do not need to vary M_* as the question might suggest.

Moreover, alternatively, we could use $I = (I_1 + I_2 + I_3)$ where $I_1 \cong \kappa_1^+$, $I_2 = \kappa_2$, I_3 a $(< \lambda^+)$ -universal linear order, and for $J \subseteq I_2$ let M_J be the Skolem hull of $M_* \cup \{a_t : t \in I_1 + J + I_3\}$.

References

- [1] M. Džamonja and S. Shelah, *On \triangleleft^* -maximality*, *Annals of Pure and Applied Logic* **125** (2004), 119–158. References are to the extended edition available at <http://shelah.logic.at/listb.html>, Paper 692.
- [2] H. Gaifman, *Models and types of Peano arithmetic*, *Annals of Mathematical Logic* **9** (1976), 223–306.
- [3] H. Gaifman and C. Dimitracopoulos, *Fragments of Peano's arithmetic and the MRDP theorem*, in *Logic and Algorithmic (Zurich, 1980)*, *Monographies de L'Enseignement Mathématique*, Vol. 30, Université de Genève, Geneva, 1982, pp. 187–206.
- [4] I. Kaplan, S. Shelah and P. Simon, *Exact saturation in simple and NIP theories*, *Journal of Mathematical Logic*, accepted.
- [5] H. J. Keisler, *Ultraproducts which are not saturated*, *Journal of Symbolic Logic* **32** (1967), 23–46.
- [6] R. Kossak, private communication.
- [7] R. Kossak and J. Schmerl, *The Structure of Models of Peano Arithmetic*, *Oxford Logic Guides*, Vol. 50, The Clarendon Press, Oxford University Press, Oxford, 2006.
- [8] M. Malliaris and S. Shelah, *General topology meets model theory, on \mathfrak{p} and \mathfrak{t}* , *Proceedings of the National Academy of Sciences of the United States of America* **110** (2013), 13300–13305.
- [9] M. Malliaris and S. Shelah, *Cofinality spectrum theorems for model theory, set theory and general topology*, *Journal of the American Mathematical Society* **29** (2016), 237–297.
- [10] M. Malliaris and S. Shelah, *Existence of optimal ultrafilters and the fundamental complexity of simple theories*, *Advances in Mathematics* **290** (2016), 614–681
- [11] M. Malliaris and S. Shelah, *Keisler's order has infinitely many classes*, *Israel Journal of Mathematics*, accepted.
- [12] M. Malliaris and S. Shelah, *Open problems on ultrafilters and some connections to the continuum*, in *Foundations of Mathematics*, *Contemporary Mathematics*, Vol. 690, American Mathematical Society, Providence, RI, to appear.
- [13] J. Moore, *Model theory and the cardinals \mathfrak{p} and \mathfrak{t}* , *Proceedings of the National Academy of Sciences of the United States of America* **110** (2013), 13238–13239.
- [14] R. MacDowell and E. Specker, *Modelle der arithmetik*, in *Infinitistic Methods (Proceedings of the Symposium on Foundations of Mathematics, Warsaw, 1959)*, Pergamon Press, Oxford; Państwowe Wydawnictwo Naukowe, Warsaw, 1961, pp. 257–263.
- [15] J.-F. Pabion, *Saturated models of Peano arithmetic*, *Journal of Symbolic Logic* **47** (1982), 625–637.
- [16] J. Schmerl, *Elementary cuts in saturated models of Peano arithmetic*, *Notre Dame Journal of Formal Logic* **53** (2012), 1–13.

- [17] S. Shelah, *Classification Theory and the Number of Nonisomorphic Models*, Studies in Logic and the Foundations of Mathematics, Vol. 92, North-Holland, Amsterdam–New York, 1978.
- [18] S. Shelah, *End extensions and numbers of countable models*, Journal of Symbolic Logic **43** (1978), 550–562.
- [19] S. Shelah, *Toward classifying unstable theories*, Annals of Pure and Applied Logic **80** (1996), 229–255.
- [20] S. Shelah, *General non-structure theory*, Paper E59, available at <http://shelah.logic.at/>
- [21] S. Shelah, *Classification theory for elementary classes with the dependence property—a modest beginning*, Scientiae Mathematicae Japonicae **59** (2004), 265–316.
- [22] S. Shelah, *Quite complete real closed fields*, Israel Journal of Mathematics **142** (2004), 261–272. Extended version available at <http://shelah.logic.at/files/757.pdf>
- [23] S. Shelah, *Dependent theories and the generic pair conjecture*, Communications in Contemporary Mathematics **17** (2015).
- [24] S. Shelah, *Models of expansions of \mathbb{N} with no end extensions*, Mathematical Logic Quarterly **57** (2011), 341–365.
- [25] S. Shelah, *Atomic saturation of reduced powers*, arXiv:1601.04824.
- [26] S. Shelah and A. Usvyatsov, *More on SOP_1 and SOP_2* , Annals of Pure and Applied Logic **155** (2008), 16–31.
- [27] S. Shelah, *Dependent dreams: recounting types*, arXiv: 1202.5795.
- [28] A. J. Wilkie and J. B. Paris, *On the scheme of induction for bounded arithmetic formulas*, Annals of Pure and Applied Logic **35** (1987), 261–302.