

CENTRALITY OF K_2 FOR CHEVALLEY GROUPS: A PRO-GROUP APPROACH

BY

ANDREI LAVRENOV, SERGEY SINCHUK AND EGOR VORONETSKY

St. Petersburg University

7/9 Universitetskaya nab., Saint Petersburg 199034, Russia

e-mail: avlavrenov@gmail.com, sinchukss@gmail.com, voronetckiiigor@yandex.ru

ABSTRACT

We prove the centrality of $K_2(F_4, R)$ for an arbitrary commutative ring R . This completes the proof of the centrality of $K_2(\Phi, R)$ for any root system Φ of rank ≥ 3 . Our proof uses only elementary localization techniques reformulated in terms of pro-groups. Another new result of the paper is the construction of a crossed module on the canonical homomorphism $\text{St}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R)$, which has not been known previously for exceptional Φ .

1. Introduction

The aim of this paper is to give a uniform proof of the centrality of the K_2 -functor modeled on Chevalley groups for an arbitrary commutative ring. Our proof is based on the technique of pro-groups introduced by the third-named author in [32] and [33]. Recall that [32] is dedicated to the centrality of the linear K_2 for not necessarily commutative rings while [33] focuses on the study of the centrality problem in the context of odd-unitary groups, a large class of groups generalizing the usual classical groups; see [17].

Let us briefly explain for the reader how our results fit into the general picture. Recall that Steinberg groups $\text{St}(\Phi, R)$ are certain groups given by generators and relations, that were classically introduced by R. Steinberg and J. Milnor in [23, 15] as combinatorial approximations of the simply-connected Chevalley

Received September 17, 2020 and in revised form October 6, 2021

groups. For example, every element of the special linear group $SL_n(\mathbb{F}_q)$ over a finite field can be presented as a product of elementary transvections $t_{ij}(a)$ ($i \neq j$, $a \in \mathbb{F}_q$). These elements are taken as the generators of the Steinberg group $St_n(\mathbb{F}_q)$. Moreover, any two such presentations of the same element of $SL_n(\mathbb{F}_q)$ can be rewritten one into the other via a sequence of the following “elementary” relations:

- $t_{ij}(a)t_{ij}(b) = t_{ij}(a + b)$;
- $t_{ij}(a)t_{hk}(b) = t_{hk}(b)t_{ij}(a)$ for $j \neq h$ and $i \neq k$;
- $t_{ij}(a)t_{jk}(b)t_{ij}(-a)t_{jk}(-b) = t_{ik}(ab)$ for $i \neq k$.

These relations are called **Steinberg relations** and are taken as the defining relations of the Steinberg group $St_n(\mathbb{F}_q)$. Thus, in the finite field case one has $St_n(\mathbb{F}_q) \cong SL_n(\mathbb{F}_q)$.

For an infinite field F the Steinberg group $St(\Phi, F)$ no longer coincides with the simply-connected Chevalley group $G_{sc}(\Phi, F)$, but rather is the universal central extension of the latter. By the classical theory of central extensions the kernel of the natural homomorphism $St(\Phi, F) \rightarrow G_{sc}(\Phi, F)$ coincides with the Schur multiplier $H_2(G_{sc}(\Phi, F), \mathbb{Z})$. The concrete presentation of this kernel in the field case was obtained by H. Matsumoto in [14]. Similar presentation for the special linear group over a skew field also has been obtained by U. Rehmann in [19].

Recall that the algebraic K_2 -functor was originally defined by J. Milnor as the kernel of the homomorphism $St_\infty(R) \rightarrow GL_\infty(R)$, i.e., as the group of “nontrivial” relations between transvections in the infinite-dimensional general linear group. Also he showed that the Schur multipliers of the linear Steinberg group $St_n(R)$ are trivial for $n \geq 5$ and an arbitrary R . This shows that $St_n(R)$ is a good candidate for the role of the universal central extension of the elementary subgroup $E_n(R) = \text{Im}(St_n(R) \rightarrow SL_n(R))$.

Milnor’s approach was followed by M. Stein who defined Steinberg groups $St(\Phi, R)$ in the context of arbitrary Chevalley groups and, by analogy, defined the functor $K_2(\Phi, R)$ as the kernel of the natural homomorphism

$$(1.1) \quad \text{st}: St(\Phi, R) \rightarrow G_{sc}(\Phi, R).$$

Also M. Stein and W. van der Kallen computed Schur multipliers of $St(\Phi, R)$ for Φ of rank ≥ 3 ; see [21, 7]. The Schur multipliers in the setting of unitary Steinberg groups over noncommutative rings have also been studied in [3, 28, 10, 9].

The first proof of the centrality of the linear functor

$$K_{2,n}(R) = \text{Ker}(\text{St}_n(R) \rightarrow \text{SL}_n(R))$$

for an arbitrary commutative ring R was obtained by W. van der Kallen in [6] for $n \geq 4$. His proof was influenced by the proof of the Suslin normality theorem, which asserts that the elementary subgroup $E_n(R)$ is normal in $\text{SL}_n(R)$ for an arbitrary commutative R and $n \geq 3$. The latter result is one of the ingredients in Suslin's proof of the so-called K_1 -analogue of the Serre problem; see [26]. The result of van der Kallen was further generalized by M. Tulenbaev to almost commutative rings, i.e., algebras module finite over their centers; see [29].

While the centrality of $K_2(\Phi, R)$ has been long known for rings of small Krull dimension, e.g., for semilocal rings (see [22]), the question whether it holds for an arbitrary commutative ring has remained open since [6]. On the other hand, the analogue of Suslin's normality theorem for Chevalley groups of rank ≥ 2 , i.e., the normality of the subgroup

$$E_{\text{sc}}(\Phi, R) := \text{Im}(\text{st})$$

for an arbitrary commutative ring R , was soon obtained by G. Taddei, see [27]. Since then the normality theorem was generalized to even larger classes of groups, e.g., isotropic reductive groups and odd unitary groups; see [17, 16, 25].

The first advancement in the solution of the centrality problem for arbitrary commutative rings since [6] was the counterexample of M. Wendt [35] which showed that the centrality of K_2 may fail for root systems of rank 2 (similarly, there is a counterexample to Suslin's normality theorem for the rank 1 group $\text{SL}_2(R)$). Soon the papers [11, 20, 13] of the first- and second-named authors appeared, in which it was shown that the centrality of K_2 does, indeed, hold for all the Chevalley groups of type C_ℓ , D_ℓ , E_ℓ provided $\ell \geq 3$. In [11] a symplectic analogue of the technique of [6] was developed, while in [20, 13] the proof was based on the amalgamation theorem for relative Steinberg groups which reduced the problem of centrality to the already-known linear case. Nevertheless, neither of these two approaches seemed to work for the root systems B_ℓ and F_4 . Finally, in [32] the third-named author has developed a novel pro-group approach, which allowed him not only to generalize [6, 29] to an even larger class of noncommutative rings, but also to prove the centrality of K_2 for odd unitary groups and, thus, cover all classical groups, including the case B_ℓ ,

see [33]. In the present paper these methods are employed to verify the centrality of K_2 in the case $\Phi = F_4$, the only case which has remained open until now. This is accomplished in Theorem 3 below. Our method of proof also applies to root systems of type $A_{\geq 3}$, $D_{\geq 4}$, $E_{6,7,8}$, and thus gives a proof of the centrality of K_2 independent of the earlier papers [6, 20, 13]. In the course of the proof we also reprove Taddei's normality theorem [27] in the aforementioned cases; see Lemma 5.3. Thus, the main consequence of [33] and our Theorem 3 can be formulated as follows.

THEOREM: *Let R be a commutative ring, Φ be an irreducible root system of rank ≥ 3 . Then the map $\text{st}: \text{St}(\Phi, R) \rightarrow E_{\text{sc}}(\Phi, R)$ is a central extension.*

In fact, in Theorem 3 we prove a stronger result. More precisely, we endow the canonical homomorphism (1.1) with the structure of a crossed module; see Definition 5.1. This is yet another new result of the paper. Notice that no construction of a crossed module on (1.1) was presented in [11, 20, 13]. Some of the material of the present paper (e.g., Section 4.1) is required only for this construction and is not needed for the proof of the centrality of K_2 .

Using the above theorem we obtain a result comparing Stein's K_2 -groups modeled on Chevalley groups with **Quillen's unstable K_2 -groups**. Recall that the latter are defined by means of **Quillen's +-construction**: for a root system Φ of rank ≥ 3 take the perfect normal subgroup $E_{\text{sc}}(\Phi, R) \trianglelefteq G_{\text{sc}}(\Phi, R)$ and set

$$K_2^Q(\Phi, R) := \pi_2 \text{BG}_{\text{sc}}(\Phi, R)_{E_{\text{sc}}(\Phi, R)}^+.$$

The above group also coincides with the homology group $H_2(E_{\text{sc}}(\Phi, R), \mathbb{Z})$; see, e.g., [34, §IV.1]. Thus, combining the above theorem with [21, 7] we obtain the following result.

COROLLARY: *For an arbitrary commutative ring R and a root system Φ of rank ≥ 3 there is an exact sequence*

$$1 \longrightarrow H_2(\text{St}(\Phi, R), \mathbb{Z}) \longrightarrow K_2^Q(\Phi, R) \longrightarrow K_2(\Phi, R) \longrightarrow 1,$$

in which the group $H_2(\text{St}(\Phi, R), \mathbb{Z})$ is trivial in the cases

$$\Phi = A_{\geq 4}, B_{\geq 4}, C_{\geq 4}, D_{\geq 5}, E_{6,7,8},$$

or is one of the groups in the following list:

$$H_2(\mathrm{St}(\mathbf{A}_3, R), \mathbb{Z}) = R/\langle 2, (t^2 - t)(s^2 - s) \mid t, s \in R \rangle,$$

$$H_2(\mathrm{St}(\mathbf{B}_3, R), \mathbb{Z}) = R/\langle 6, 3(t^2 - t)(s^2 - s), 2(t^3 - t) \mid t, s \in R \rangle,$$

$$H_2(\mathrm{St}(\mathbf{C}_3, R), \mathbb{Z}) = R/\langle t^2 - t \mid t \in R \rangle,$$

$$H_2(\mathrm{St}(\mathbf{D}_4, R), \mathbb{Z}) = (R/\langle t^2 - t \mid t \in R \rangle) \times (R/\langle 2, (t^2 - t)(s^2 - s) \mid t, s \in R \rangle),$$

$$H_2(\mathrm{St}(\mathbf{F}_4, R), \mathbb{Z}) = R/\langle t^2 - t \mid t \in R \rangle.$$

In particular, for $\Phi = \mathbf{A}_3, \mathbf{C}_3, \mathbf{D}_4, \mathbf{F}_4$ the equality $K_2(\Phi, R) = K_2^Q(\Phi, R)$ holds iff R does not have residue fields isomorphic to \mathbb{F}_2 and $K_2(\mathbf{B}_3, R) = K_2^Q(\mathbf{B}_3, R)$ iff R does not have residue fields isomorphic to \mathbb{F}_2 or \mathbb{F}_3 .

Since the proof of the main theorem is rather long and technical and the setting of the papers [32, 33] is already rather general, let us briefly sketch the key ideas of our proof assuming, for simplicity, that the root system Φ is simply-laced and the ring R is an integral domain.

The proof is based on a variant of Quillen–Suslin’s local-global principle. Consider an arbitrary element g from $K_2(\Phi, R)$. We want to show that g commutes with $x_\alpha(r)$ for every $r \in R$ and $\alpha \in \Phi$. For $\alpha \in \Phi$ consider the set

$$I_\alpha = \{a \in R \mid [g, x_\alpha(aR)] = 1\},$$

which is easily seen to be, in fact, an ideal. To prove the centrality of $K_2(\Phi, R)$ it suffices to show that $I_\alpha = R$ for all $\alpha \in \Phi$. Assume the contrary and consider an arbitrary maximal ideal $M \trianglelefteq R$ containing I_α for some α . By the centrality of $K_2(\Phi, R_M)$ one has the identity $[\lambda_M(g), x_\alpha(r)] = 1$ for all $r \in R_M$ (we denote by λ_M the localization homomorphism $R \rightarrow R_M$). If we were able to show for some $s_0 \in R \setminus M$ that the identity

$$(1.2) \quad [g, x_\alpha(s_0R)] = 1$$

holds in $\mathrm{St}(\Phi, R)$ or, in other words, that $s_0 \in I_\alpha \cap R \setminus M$, this would lead to a contradiction and hence would complete the proof of the centrality of $K_2(\Phi, R)$. Unfortunately, formally proving this is not easy.

In the earlier papers the proof went as follows; cf. [29, 20, 13, 12]. First, a formal variable was thrown in and the original question was reduced to the question of the triviality of $[g, x_\alpha(t)]$ in $\mathrm{St}(\Phi, R[t])$, which, in turn, was further reduced to proving the Dilation principle, an assertion about principal localizations similar to (1.2) (see e.g., [20, Lemma 15]). The proof of the Dilation

principle was based on the isomorphism of **relative Steinberg groups**

$$\mathrm{St}(\Phi, R_a[t], tR_a[t]) \cong \mathrm{St}(\Phi, R \times tR_a[t], tR_a[t]);$$

cf. [20, Remark 3.12]. Essentially this isomorphism means that it is possible to define a conjugation action of $\mathrm{St}(\Phi, R_a)$ on the group $\mathrm{St}(\Phi, R \times tR_a[t], tR_a[t])$. This last step, however, required finding nontrivial ad hoc presentations of relative Steinberg groups. For example [12] relied on a variant of van der Kallen's "another presentation" [6] for symplectic groups, while [20, 13] were based on the amalgamation theorem for relative Steinberg groups; see [20, Theorem 9]. However, it was not easy to transfer to all Chevalley groups either of these two approaches.

A remarkable idea of the third-named author was to abandon relative Steinberg groups completely and instead work with sequences of **unrelativized** Steinberg groups $\mathrm{St}(\Phi, a^n R)$, $n \in \mathbb{N}$ given by the usual Steinberg presentations. It turns out that it is possible to naturally define the "conjugation" action of the Steinberg group $\mathrm{St}(\Phi, R_a)$ on such a sequence. Naively, for $g \in \mathrm{St}(\Phi, R_a)$ the automorphism of conjugation by g corresponds to a collection of group homomorphisms

$$\{\mathrm{St}(\Phi, a^{c^*(m)} R) \rightarrow \mathrm{St}(\Phi, a^m R)\}_{m \in \mathbb{N}},$$

where $c^* : \mathbb{N} \rightarrow \mathbb{N}$ is some function depending on g .

In fact, it is possible to even further generalize this construction to arbitrary localizations, not just principal ones. Indeed, if $S \subseteq R$ is an arbitrary multiplicative subset of a domain R , then one can consider the following diagram. Its objects are unrelativized Steinberg groups $\mathrm{St}(\Phi, sR)$, $s \in S$, while its arrows are the homomorphisms

$$\begin{aligned} \mathrm{St}(\Phi, s_2 R) &\rightarrow \mathrm{St}(\Phi, s_1 R), \\ x_\alpha(a) &\mapsto x_\alpha(a), \end{aligned}$$

which may exist, obviously, only if s_1 divides s_2 . The above diagram is called the **Steinberg pro-group** and its arrows are called the **structure morphisms**. The key claim is that it is possible to define an action of $\mathrm{St}(\Phi, S^{-1}R)$ on the Steinberg pro-group.

For each $x_\alpha(a/s)$, $a \in R$, $s \in S = R \setminus M$ the automorphism of conjugation by $x_\alpha(a/s)$ corresponds to a collection of group homomorphisms

$$(1.3) \quad \{\text{conj}_{x_\alpha(a/s)}(t): \text{St}(\Phi, c^*(t)R) \rightarrow \text{St}(\Phi, tR)\}_{t \in S},$$

where $c^*: S \rightarrow S$ is the function $t \mapsto ts$, called the **index function**. The construction of the homomorphisms $\text{conj}_{x_\alpha(a/s)}(t)$ will be outlined below.

In fact, a morphism between pro-groups should be defined not just as a collection of group homomorphisms, but as an equivalence class of such collections. By definition, two pro-group morphisms m_1 and m_2 with index functions $m_1^*, m_2^*: S \rightarrow S$ are equivalent if for every t one can choose s divisible by both $m_1^*(t)$ and $m_2^*(t)$ so that the homomorphisms

$$m_i(t): \text{St}(\Phi, m_i^*(t)R) \rightarrow \text{St}(\Phi, tR), \quad i = 1, 2$$

coincide after precomposition with the structure morphisms

$$\text{St}(\Phi, sR) \rightarrow \text{St}(\Phi, m_i^*(t)R), \quad i = 1, 2.$$

We refer the reader to Section 2.1 for a more thorough treatment of the formalism of pro-groups.

After the automorphisms of conjugation by $x_\alpha(a/s)$ are defined, we verify that they satisfy the Steinberg relations defining $\text{St}(\Phi, S^{-1}R)$ and, moreover, that the Steinberg symbols $\{b_1, b_2\}$, $b_1, b_2 \in S^{-1}R^\times$ act trivially on the pro-group. Technically, this is an easier step since it is enough to verify the corresponding equalities on suitable group generators. Now, for $g = \prod x_{\alpha_i}(a_i)$ the pro-group automorphism $\text{conj}_{\lambda_M(g)}$ can be defined as the composition of $\text{conj}_{x_{\alpha_i}(a_i/1)}$. A key point here is that the resulting pro-group automorphism is compatible with the usual conjugation action of $g \in \text{St}(\Phi, R)$, i.e., for some $s_0 \in S$ the group homomorphism

$$\text{conj}_{\lambda_M(g)}(1): \text{St}(\Phi, s_0R) \rightarrow \text{St}(\Phi, R)$$

coincides with the obvious homomorphism $h \mapsto ghg^{-1}$. Since our conjugation action factorizes through $\text{St}(\Phi, S^{-1}R)$ we conclude that the conjugation by $g \in K_2(\Phi, R)$ acts trivially on $x_\alpha(s_0R)$, which finishes the proof of (1.2). This part of the proof is formalized in Sections 4 and 5.

Now let us focus on the construction of the action of $\text{St}(\Phi, S^{-1}R)$ on the Steinberg pro-group. In order to construct the automorphism of conjugation by $x_\alpha(a/s)$ we need to introduce the group $\text{St}(\Phi \setminus \{-\alpha\}, tR)$ whose presentation is obtained from the standard presentation of the Steinberg group $\text{St}(\Phi, tR)$

by omitting all the generators $x_{-\alpha}(tr)$, $r \in R$ and all the relations involving such generators. Now the analogues of the homomorphisms (1.3) between such groups can be constructed fairly easily based on the remaining part of the Steinberg presentation (below N denotes a sufficiently large natural number):

$$(1.4) \quad x_{\beta}(s^N b) \mapsto \begin{cases} x_{\alpha+\beta}(N_{\alpha,\beta}s^{N-1}ab)x_{\beta}(s^N b) & \text{if } \alpha + \beta \in \Phi, \\ x_{\beta}(s^N b) & \text{if } \alpha + \beta \notin \Phi \cup \{0\}. \end{cases}$$

Notice that there is no simple expression similar to the right-hand side of (1.4) in the case $\beta = -\alpha$, which explains our need to use groups $\text{St}(\Phi \setminus \{-\alpha\}, tR)$ as domains for the homomorphisms (1.3). It follows from Lemma 4.2 that (1.4) defines a collection of well-defined homomorphisms:

$$\text{conj}'_{x_{\alpha}(r/s)}(t): \text{St}(\Phi \setminus \{-\alpha\}, s^N tR) \rightarrow \text{St}(\Phi \setminus \{-\alpha\}, tR).$$

The obvious defect of the formula (1.4) is that its correctness depends on the integrality of R . However, it is easy to fix this using the notion of a homotope of a ring introduced in [32, 33]. By definition, for $t \in R$ the t -**homotope** $R^{(t)}$ of R is the ring without unit, multiplication in which is given by the formula $x * y = txy$. The homotope $R^{(t)}$ is isomorphic to tR as a ring without unit if R happens to be a domain. Now, replacing everywhere above the group $\text{St}(\Phi, tR)$ with $\text{St}(\Phi, R^{(t)})$, one can formulate the analogue of (1.4) for not necessarily integral rings. We believe that the proof which works for arbitrary commutative rings has extra elegance as compared to the proof which first reduces to the case of domains. However, the reader not comfortable with the notion of a ring homotope may follow the latter path and think of $R^{(t)}$ as just a synonym for tR ; see Section 5.2 for an outline of such a proof.

Although, the groups $\text{St}(\Phi, tR)$ and $\text{St}(\Phi \setminus \{-\alpha\}, tR)$ need not be isomorphic in general, the pro-groups composed of them turn out to be isomorphic in the category of pro-groups, see Theorem 1. Notice also that the corresponding assertion for unital rings, i.e., the isomorphism

$$\text{St}(\Phi, R) \cong \text{St}(\Phi \setminus \{-\alpha\}, R)$$

is an easy corollary of a much stronger result called Curtis–Tits presentation, which allows one to omit most of the Steinberg generators from the presentation of $\text{St}(\Phi, R)$; see e.g., [2, Corollary 1.3]. In fact, the main difference of the present paper from [32, 33] is that our proof is based on Steinberg presentations omitting roots, while [32, 33] rely on computations with relative root systems.

If written in down-to-earth terms, the aforementioned isomorphism of pro-groups amounts to the following collection of group homomorphisms in the “non-obvious” direction (M is a sufficiently large natural number):

$$\{F_\alpha(t): \text{St}(\Phi, t^M R) \rightarrow \text{St}(\Phi \setminus \{-\alpha\}, tR)\}_{t \in S}.$$

Of course, the natural choice for the image of $x_\alpha(t^M r)$ would be the commutator $[x_{\alpha-\beta}(t^{M-1}b), x_\beta(\pm t)]$ for some $\beta \in \Phi$ such that $\alpha - \beta \in \Phi$. The fact that this definition does not depend on the choice of β can be checked using the Hall–Witt identity (2.3) (notice that we can further decompose $x_{\alpha-\beta}(t^{M-1}b)$ into a commutator in $\text{St}(\Phi, tR)$). Similarly, we can verify that these commutators satisfy the relations missing from the presentation of $\text{St}(\Phi \setminus \{-\alpha\}, t^M R)$. Keeping track of index functions and individual homomorphisms $F_\alpha(t)$ in such computations, however, is not very convenient, so, to simplify the proof of Theorem 1, we prefer to write down all commutator formula calculations inside the category of pro-groups. In addition, to reduce the number of cases that need to be considered, we also assume throughout the paper that Φ is not of type B_ℓ or C_ℓ . The latter assumption makes it possible to decompose any root of Φ into a sum of roots having the same length. Now, composing $F_\alpha(t)$ and $\text{conj}'_{x_\alpha(r/s)}(t)$ in a suitable way we obtain the sought pro-group morphism $\text{conj}_{x_\alpha(r/s)}$.

ACKNOWLEDGEMENTS. The authors of the paper would like to express their sincere gratitude to the anonymous referee for a careful reading of the first version of this paper and numerous helpful suggestions.

The authors of the paper were supported by “Native towns”, a social investment program of PJSC “Gazprom Neft”. The first-named author was also supported by the Ministry of Science and Higher Education of the Russian Federation, agreement No. 075-15-2022-287. The second-named author was also supported by RFBR grant No. 18-31-20044. The second- and third-named authors were also supported by the Foundation for the Advancement of Theoretical Physics and Mathematics “BASIS”. The first-named author is a winner of Young Russian Mathematics contest and would like to thank its sponsors and the jury.

2. Preliminaries

Throughout this paper all commutators are left normed, i.e., $[x, y] = xyx^{-1}y^{-1}$. We denote by x^y and ${}^y x$ the elements xyx^{-1} and $yx y^{-1}$. In this paper we also make use of the following commutator identities:

$$(2.1) \quad [x, yz] = [x, y] \cdot {}^y[x, z],$$

$$(2.2) \quad [xy, z] = {}^x[y, z] \cdot [x, z].$$

Recall that the Hall–Witt identity asserts that

$$(2.3) \quad [[x, y], {}^y z] \cdot [[y, z], {}^z x] \cdot [[z, x], {}^x y] = 1.$$

As a corollary we obtain that

$$(2.4) \quad [x, z] = 1 \text{ implies } [x, [y, z]] = [[x, y], {}^y z].$$

Finally, it is not hard to check that

$$(2.5) \quad [x, y] = 1 \text{ implies } [x, [y, z]] = [y, {}^x z] [z, y].$$

2.1. GENERALITIES ON PRO-OBJECTS. Let \mathcal{C} be an arbitrary category. In this section we recall the construction of the pro-completion of \mathcal{C} (cf. [8, Section 6.1]).

Recall that a nonempty small category \mathcal{I} is called **filtered** if

- for any two objects $i, k \in \text{Ob}(\mathcal{I})$ there is a diagram $i \rightarrow j \leftarrow k$ in \mathcal{I} (i.e., i and k have an upper bound);
- every two parallel morphisms $i \rightrightarrows j$ are equalized by some morphism $j \rightarrow k$ in \mathcal{I} .

A **pro-object** in \mathcal{C} is, by definition, a functor $X^{(\infty)}: \mathcal{I}_X^{\text{op}} \rightarrow \mathcal{C}$, i.e., a contravariant functor from a filtered category \mathcal{I}_X , called the **category of indices of $X^{(\infty)}$** , to the category \mathcal{C} .

We denote by $X^{(i)}$ the value of $X^{(\infty)}$ on an index i . The values of the functor $X^{(\infty)}$ on the arrows of \mathcal{I} are called the **structure morphisms** of $X^{(\infty)}$.

The category of pro-objects is denoted by $\text{Pro}(\mathcal{C})$. The hom-sets in this category are given by the formula

$$(2.6) \quad \text{Pro}(\mathcal{C})(X^{(\infty)}, Y^{(\infty)}) = \varprojlim_{j \in \mathcal{I}_Y} \varinjlim_{i \in \mathcal{I}_X} \mathcal{C}(X^{(i)}, Y^{(j)}).$$

Let us recall a more explicit description of morphisms in $\text{Pro}(\mathcal{C})$. By definition, a **pre-morphism** $f: X^{(\infty)} \rightarrow Y^{(\infty)}$ consists of the following data:

- a set-theoretic function $f^*: \text{Ob}(\mathcal{I}_Y) \rightarrow \text{Ob}(\mathcal{I}_X)$;

- a collection of morphisms $f^{(i)}: X^{(f^*(i))} \rightarrow Y^{(i)}$ in \mathcal{C} parametrized by $i \in Ob(\mathcal{I}_Y)$. These morphisms are required to satisfy the following additional assumption: for every morphism $i \rightarrow j$ in \mathcal{I}_Y there exists a sufficiently large index $k \in Ob(\mathcal{I}_X)$ such that the composite morphisms $X^{(k)} \rightarrow X^{(f^*(i))} \rightarrow Y^{(i)}$ and $X^{(k)} \rightarrow X^{(f^*(j))} \rightarrow Y^{(j)} \rightarrow Y^{(i)}$ are equal.

The composition of two pre-morphisms $f: X^{(\infty)} \rightarrow Y^{(\infty)}$ and $g: Y^{(\infty)} \rightarrow Z^{(\infty)}$ is defined as the pre-morphism $g \circ f$, where

$$(g \circ f)^*(i) = f^*(g^*(i)) \quad \text{and} \quad (g \circ f)^{(i)} = g^{(i)} \circ f^{(g^*(i))}.$$

Two parallel pre-morphisms $f, g: X^{(\infty)} \rightarrow Y^{(\infty)}$ are called equivalent if for every $i \in Ob(\mathcal{I}_Y)$ there exists a sufficiently large index $j \in Ob(\mathcal{I}_X)$ such that the composite morphisms $X^{(j)} \rightarrow X^{(f^*(i))} \rightarrow Y^{(i)}$ and $X^{(j)} \rightarrow X^{(g^*(i))} \rightarrow Y^{(i)}$ are equal. Finally, a **morphism** $X^{(\infty)} \rightarrow Y^{(\infty)}$ is an equivalence class of pre-morphisms. Note that the equivalence relation is preserved by the composition operation.

There is a fully faithful functor $\mathcal{C} \rightarrow \text{Pro}(\mathcal{C})$ sending $X \in Ob(\mathcal{C})$ to the pro-object $X: \mathbf{1}^{\text{op}} \rightarrow \mathcal{C}$. It is clear from (2.6) that

$$(2.7) \quad \text{Pro}(\mathcal{C})(X, Y) \cong \mathcal{C}(X, Y);$$

$$(2.8) \quad \text{Pro}(\mathcal{C})(X^{(\infty)}, Y) \cong \varinjlim_{i \in \mathcal{I}_X} \mathcal{C}(X^{(i)}, Y);$$

$$(2.9) \quad \text{Pro}(\mathcal{C})(X, Y^{(\infty)}) \cong \varprojlim_{i \in \mathcal{I}_Y} \mathcal{C}(X, Y^{(i)}).$$

Moreover, it is also clear from (2.6) and (2.8) that the following assertion holds.

LEMMA 2.1: $X^{(\infty)}$ is the projective limit of $X^{(i)}$ in the category $\text{Pro}(\mathcal{C})$.

The category of pro-sets $\text{Pro}(\mathbf{Set})$ has all finite limits by [8, Prop. 6.1.18] and therefore is a cartesian monoidal category.

Let us describe an explicit construction of limits in one important special case. Let $X^{(\infty)}, Y^{(\infty)}: \mathcal{I}^{\text{op}} \rightarrow \mathbf{Set}$ be a pair of pro-sets with the same index category. Recall that the pointwise product of functors $X^{(\infty)} \times Y^{(\infty)}: \mathcal{I}^{\text{op}} \rightarrow \mathbf{Set}$ is given by $(X^{(\infty)} \times Y^{(\infty)})^{(i)} = X^{(i)} \times Y^{(i)}$. Clearly, $X^{(\infty)} \times Y^{(\infty)}$ is a product of $X^{(\infty)}$ and $Y^{(\infty)}$ in the category $\mathbf{Fun}(\mathcal{I}^{\text{op}}, \mathbf{Set})$.

There is an obvious identity-on-objects functor $\mathbf{Fun}(\mathcal{I}^{\text{op}}, \mathbf{Set}) \rightarrow \text{Pro}(\mathbf{Set})$ sending a natural transformation $\varphi: X^{(\infty)} \rightarrow Y^{(\infty)}$ to the morphism of pro-sets given by the pre-morphism $\varphi^* = \text{id}_{\text{Ob}(\mathcal{I})}$, $\varphi^{(i)} = \varphi_i$. The diagonal morphism $\Delta: X^{(\infty)} \rightarrow X^{(\infty)} \times X^{(\infty)}$ and the canonical projection morphisms $\pi_X: X^{(\infty)} \times Y^{(\infty)} \rightarrow X^{(\infty)}$, $\pi_Y: X^{(\infty)} \times Y^{(\infty)} \rightarrow Y^{(\infty)}$ can be defined in the category $\text{Pro}(\mathbf{Set})$ as the images of the corresponding natural transformations in $\mathbf{Fun}(\mathcal{I}^{\text{op}}, \mathbf{Set})$ under this functor.

We claim that the point-wise product $X^{(\infty)} \times Y^{(\infty)}$ satisfies the universal property of products in the category $\text{Pro}(\mathbf{Set})$. Indeed, this follows from (2.6) and the fact that filtered colimits commute with finite limits. This argument also shows that the functor $\mathbf{Fun}(\mathcal{I}^{\text{op}}, \mathbf{Set}) \rightarrow \text{Pro}(\mathbf{Set})$ preserves finite limits.

2.2. GROUP AND RING OBJECTS IN PRO-SETS. In this paper we consider only commutative rings. To distinguish between unital and non-unital rings we reserve the word “ring” only for unital rings and refer to non-unital rings as **rngs**. We denote the category of rings (resp. rngs) as **Ring** (resp. **Rng**).

Let T be an algebraic theory. Throughout this paper we will be mostly interested in the case where T is the theory of groups or the theory of rngs.

Since the category $\text{Pro}(\mathbf{Set})$ is cartesian monoidal, we may speak of the category $\mathbf{Mod}(T, \text{Pro}(\mathbf{Set}))$ of models of T in $\text{Pro}(\mathbf{Set})$. In the special case where T is the theory of groups (resp. rngs) this category is precisely the category of group (resp. rng) objects in $\text{Pro}(\mathbf{Set})$.

The forgetful functor F from the category of pro-groups $\text{Pro}(\mathbf{Grp})$ to the category of pro-sets $\text{Pro}(\mathbf{Set})$ is faithful. Every pro-group $G^{(\infty)}$ defines the structure of a group object in $\text{Pro}(\mathbf{Set})$ on $F(G^{(\infty)})$. It is easy to see that a morphism $f \in \text{Pro}(\mathbf{Set})(F(G^{(\infty)}), F(H^{(\infty)}))$ comes from $\text{Pro}(\mathbf{Grp})$ if and only if it is a morphism of group objects. Indeed, if $f: F(G^{(\infty)}) \rightarrow F(H^{(\infty)})$ is a pre-morphism and the corresponding morphism preserves the multiplication in the sense of group objects, then the maps $f^{(i)}: G^{(f^*(i))} \rightarrow H^{(i)}$ are group homomorphisms after a restriction to $G^{(j)}$ for sufficiently large j . Thus, $\text{Pro}(\mathbf{Grp})$ is equivalent to a full subcategory of the category of group objects in $\text{Pro}(\mathbf{Set})$.

Definition 2.2: Let n be a natural number and $F_{n,T}$ be the free algebra on n generators in the theory T . If T is the theory of groups, then F_n is the free group $F(t_1, \dots, t_n)$. Similarly, if T is the theory of rngs, $F_{n,T}$ is the sub-rng of the ring $\mathbb{Z}[t_1, \dots, t_n]$ consisting of polynomials over \mathbb{Z} without free terms.

Given a word $w \in F_{n,T}$ and a T -model $X^{(\infty)}$ in $\text{Pro}(\mathbf{Set})$, one can construct the $\text{Pro}(\mathbf{Set})$ -morphism

$$w^{(\infty)}: \underbrace{X^{(\infty)} \times \cdots \times X^{(\infty)}}_{n \text{ times}} \rightarrow X^{(\infty)},$$

which “interprets” the word w . This morphism can be obtained as an appropriate composition of diagonal morphisms $\Delta_{X^{(\infty)}}$, projections π_i (which are part of the structure of a cartesian monoidal category on $\text{Pro}(\mathbf{Set})$) and the morphisms defining the structure of a T -model on $X^{(\infty)}$.

In order to simplify the notation, to denote the interpretation morphism $w^{(\infty)}$ we often use the original term for w , in which every symbol of a free variable t_i is replaced by the symbol $t_i^{(\infty)}$.

Example 2.3: Consider the case where T is the theory of groups. Let $G^{(\infty)}$ be a group object in $\text{Pro}(\mathbf{Set})$ and $w = [t_1, t_2] = t_1 t_2 t_1^{-1} t_2^{-1} \in F(t_1, t_2)$ be the word representing the generic commutator. In this case the interpretation morphism $w^{(\infty)}$ (also denoted $[t_1^{(\infty)}, t_2^{(\infty)}]$) can be defined as the composition

$$\begin{array}{ccc} G^{(\infty)} \times G^{(\infty)} & \xrightarrow{\Delta \times \Delta} & G^{(\infty)} \times G^{(\infty)} \times G^{(\infty)} \times G^{(\infty)} \\ & & \downarrow \langle \pi_1, \pi_3, i\pi_2, i\pi_4 \rangle \\ & & G^{(\infty)} \times G^{(\infty)} \times G^{(\infty)} \times G^{(\infty)} \xrightarrow{m(m \times m)} G^{(\infty)}, \end{array}$$

where the structure of a group object on $G^{(\infty)}$ is given by the triple (m, i, e) .

2.3. HOMOTOPES OF RINGS AND PRO-RINGS. For the rest of the paper R denotes an arbitrary ring and S denotes a fixed multiplicative subset of R containing a unit. Denote by \mathcal{S} the category, whose objects are the elements of S and whose morphisms $\mathcal{S}(s, s')$ are all $s'' \in S$ such that $ss'' = s'$. The composition and the identity morphisms are induced by the ring structure on R . It is clear that \mathcal{S} is a filtered category. Unless stated otherwise, all the pro-sets that we encounter in the sequel have \mathcal{S} as their category of indices.

We introduce the notion of a homotope of a ring inspired by a similar notion from nonassociative algebra.

Definition 2.4: Let s be an element of R . By definition, the **s -homotope** of R is the rng $R^{(s)} = \{a^{(s)} \mid a \in R\}$ with the operations of addition and multiplication given by

$$a^{(s)} + b^{(s)} = (a + b)^{(s)}, \quad a^{(s)}b^{(s)} = (asb)^{(s)}, \quad a, b \in R.$$

Clearly, $R^{(s)}$ has the structure of an R -algebra given by the formula

$$a \cdot b^{(s)} = (ab)^{(s)}, \quad a, b \in R.$$

For $s, s' \in S$ there is a homomorphism of R -algebras

$$R^{(ss')} \rightarrow R^{(s')}, \quad a^{(ss')} \mapsto (as)^{(s')}.$$

Denote by $R^{(\infty)}$ the formal projective limit of the projective system $R^{(s)}$, where $s \in Ob(\mathcal{S})$. Thus, $R^{(\infty)}$ is an object of the category $\text{Pro}(\mathbf{Rng})$. It can also be considered as an rng object in $\text{Pro}(\mathbf{Set})$.

Notice that every pro-rng $R^{(\infty)}$ can be considered as a pro-group by forgetting its multiplicative structure pointwise.

Remark 2.5: Denote by sR the principal ideal of R generated by s . There is an rng homomorphism $R^{(s)} \rightarrow sR$ given by $r^{(s)} \mapsto sr$. In the case where R is an integral domain this homomorphism is easily seen to be an rng isomorphism.

The reason why we use homotopes rather than principal ideals in the definition of $R^{(\infty)}$ is that there is always a “division by s ” homomorphism of R -modules $R^{(ss')} \rightarrow R^{(s')}$ given by $r^{(ss')} \mapsto r^{(s')}$, while the similar map $ss'R \rightarrow s'R$ may not exist if R does not happen to be a domain. The existence of this division homomorphism will be important in Section 4.

Notice also that the projective limit of $R^{(s)}$ in \mathbf{Rng} is often trivial. Indeed, if R is a domain, then the limit of $R^{(s)}$ in \mathbf{Rng} computes the intersection of the principal ideals $\bigcap_{s \in S} sR$, which often coincides with the zero rng. This also shows that the set of **global elements** of $R^{(\infty)}$ (i.e., the hom-set in $\text{Pro}(\mathbf{Set})$ from the terminal object 1 to $R^{(\infty)}$) is often trivial and therefore will be of little interest to us.

Recall that a morphism $f \in \mathcal{C}(X, Y)$ is called a **split epimorphism** (or a **retraction**) if it admits a section, i.e., there exists $g \in \mathcal{C}(Y, X)$ such that $fg = \text{id}_Y$. Retractions are preserved under pullbacks.

LEMMA 2.6: *The rng multiplication morphism $m: R^{(\infty)} \times R^{(\infty)} \rightarrow R^{(\infty)}$ is a split epimorphism of pro-sets.*

Proof. Consider the following pre-morphism of pro-sets:

$$u: R^{(\infty)} \rightarrow R^{(\infty)} \times R^{(\infty)}, \quad u^*(s) = s^2, \quad u^{(s)}: c^{(s^2)} \mapsto (1^{(s)}, c^{(s)}).$$

Clearly, this is indeed a pre-morphism and

$$m^{(s)}(u^{(s)}(c^{(s^2)})) = 1^{(s)}c^{(s)} = (sc)^{(s)},$$

which shows that $mu = \text{id}_{R^{(\infty)}}$. ■

The majority of the calculations encountered in the present paper occur in the category of pro-sets or pro-groups. Most often we need to prove certain equalities between composite pro-set or pro-group morphisms. It turns out that the usual notation for categorical composition makes these calculations too lengthy and hardly readable. In order to remedy this and also make our computations look like the usual computations with root unipotents in Steinberg groups, we need to introduce a certain way to denote pro-set (pro-group) morphisms, specifically composite ones.

Convention 2.7: First of all, notice that the present conventions apply only to algebraic expressions in which the symbol (∞) occurs in the upper index of all free variables (e.g., the identities of Lemma 2.8 or Lemma 3.6, but not the identities (R1)–(R4)). We call such expressions **pro-expressions**.

- Any pro-expression encountered in the sequel is meant to denote a certain morphism of pro-sets. Apart from the variables marked with the index (∞) , a pro-expression may also involve group or rng operations and other pro-set morphisms.
- Whenever a pro-expression does not involve other pro-set morphisms, it should be understood according to Definition 2.2 (the domain and codomain will usually be clear from the context).
- To denote the composition of pro-set morphisms we use the syntax of substituted expressions. For example, if $f, g: X^{(\infty)} \rightarrow Y^{(\infty)}$ are morphisms of pro-groups and $[t_1^{(\infty)}, t_2^{(\infty)}]$ is the pro-group morphism from Example 2.3, then $[f(a^{(\infty)}), g(b^{(\infty)})]$ denotes the composite morphism

$$[t_1^{(\infty)}, t_2^{(\infty)}] \circ (f \times g): X^{(\infty)} \times X^{(\infty)} \rightarrow Y^{(\infty)}.$$

- Any equality of pro-expressions should be understood as the equality of pro-set morphisms defined by these expressions.
- The exact names of the variables occurring in a pro-expression are unimportant and are usually chosen arbitrarily. On the other hand, there is always some natural order on the variables that allows one to read the pro-expression unequivocally. In particular, $[t_1^{(\infty)}, t_2^{(\infty)}]$ and $[a^{(\infty)}, b^{(\infty)}]$ denote the same pro-set morphism $G^{(\infty)} \times G^{(\infty)} \rightarrow G^{(\infty)}$, while $[t_1^{(\infty)}, t_2^{(\infty)}]$ and $[t_2^{(\infty)}, t_1^{(\infty)}]$ are two different morphisms.
- The domain of the pro-set morphism defined by a pro-expression will usually be clear from the context. Most often, it is a power of a single pro-object. In this case to determine the exponent one should count the number of different variables occurring in the expression.
- The notation for the multiplication operation is usually suppressed, i.e., we prefer the notation $a^{(\infty)}b^{(\infty)}$ to $m(a^{(\infty)}, b^{(\infty)})$ or $a^{(\infty)} \cdot b^{(\infty)}$.
- The syntax of tuples is used to denote the product of morphisms. For example, if $f, g: X^{(\infty)} \rightarrow Y^{(\infty)}$ are morphisms of pro-sets, then the notation $(f(x_1^{(\infty)}), g(x_2^{(\infty)}))$ means simply $f \times g$.
- If g is a morphism of pro-sets with $X^{(\infty)} \times Y^{(\infty)}$ as its domain, then we write $g(a^{(\infty)}, b^{(\infty)})$ instead of $g((a^{(\infty)}, b^{(\infty)}))$.
- Notice that the trivial group 1 is a zero object in the category $\text{Pro}(\mathbf{Grp})$, therefore for any pro-groups $G^{(\infty)}, H^{(\infty)}$ there is a unique morphism $G^{(\infty)} \rightarrow H^{(\infty)}$ passing through 1 ; this morphism will also be denoted by 1 .

Now we are ready to formulate our next result.

LEMMA 2.8: *Let $G^{(\infty)}$ be a pro-group, $R^{(\infty)}$ be the pro-rng defined above and $g: R^{(\infty)} \times R^{(\infty)} \rightarrow G^{(\infty)}$ be a morphism of pro-sets. There is a morphism $f: R^{(\infty)} \rightarrow G^{(\infty)}$ of pro-groups such that*

$$g(a^{(\infty)}, b^{(\infty)}) = f(a^{(\infty)}b^{(\infty)})$$

if and only if g satisfies the following identities in $\text{Pro}(\mathbf{Set})$:

- $[g(a_1^{(\infty)}, b_1^{(\infty)}), g(a_2^{(\infty)}, b_2^{(\infty)})] = 1$;
- $g(a_1^{(\infty)} + a_2^{(\infty)}, b^{(\infty)}) = g(a_1^{(\infty)}, b^{(\infty)})g(a_2^{(\infty)}, b^{(\infty)})$;
- $g(a^{(\infty)}, b_1^{(\infty)} + b_2^{(\infty)}) = g(a^{(\infty)}, b_1^{(\infty)})g(a^{(\infty)}, b_2^{(\infty)})$;
- $g(a^{(\infty)}b^{(\infty)}, c^{(\infty)}) = g(a^{(\infty)}, b^{(\infty)}c^{(\infty)})$.

Proof. The necessity of the identities is clear. By Lemma 2.6 the morphism f is unique, so it suffices to show that it exists. By Lemma 2.1 it suffices to consider the case where G is a group. Let g be a morphism satisfying the above identities. By definition, there exists $s \in S$ such that g is given by a homomorphism $g' : R^{(s)} \times R^{(s)} \rightarrow G$ satisfying the first three identities and the identity

$$g'((asb)^{(s)}, c^{(s)}) = g'(a^{(s)}, (bsc)^{(s)}).$$

Consider the map $f' : R^{(s^2)} \rightarrow G$ given by

$$f'(c^{(s^2)}) = g'(1^{(s)}, c^{(s)});$$

it is a homomorphism by the first and the third identities. From the last two identities we conclude that for all $a, b \in R$ one has

$$\begin{aligned} f'(a^{(s^2)}b^{(s^2)}) &= g'(1^{(s)}, (s^2ab)^{(s)}) \\ &= g'((sa)^{(s)}, (sb)^{(s)}) \\ &= g'(a^{(s^2)}, b^{(s^2)}). \end{aligned}$$

It is clear that f' defines the required morphism f of pro-groups. ■

2.4. STEINBERG GROUPS AND STEINBERG PRO-GROUPS. Let Φ be an irreducible root system of rank ≥ 3 . We assume that the root system Φ is contained in a Euclidean space $V = \mathbb{R}^\ell$ whose inner product we denote by $(-, -)$. For a pair of roots $\alpha, \beta \in \Phi$ we denote by $\langle \alpha, \beta \rangle$ the integer

$$\frac{2(\alpha, \beta)}{|\beta|^2}.$$

Recall that a root subset $\Sigma \subseteq \Phi$ is called **closed** if $\alpha, \beta \in \Sigma$, $\alpha + \beta \in \Phi$ imply $\alpha + \beta \in \Sigma$. A closed root subset Σ is called **symmetric** (resp. **special**) if $\Psi = -\Psi$ (resp. $\Psi \cap -\Psi = \emptyset$). By definition, a **root subsystem** $\Psi \subseteq \Phi$ is a symmetric and closed root subset.

We start by recalling the definition of the Steinberg group.

Definition 2.9: Let R be a ring. The **Steinberg group** $\text{St}(\Phi, R)$ is given by generators $x_\alpha(a)$, where $\alpha \in \Phi$ and $a \in R$ and the following list of defining

relations (cf. [18]):

$$(R1) \quad x_\alpha(a) \cdot x_\alpha(b) = x_\alpha(a + b);$$

$$(R2) \quad [x_\alpha(a), x_\beta(b)] = 1,$$

if $\alpha + \beta \notin \Phi \cup \{0\}$;

$$(R3) \quad [x_\alpha(a), x_\beta(b)] = x_{\alpha+\beta}(N_{\alpha,\beta} \cdot ab),$$

if $\alpha + \beta \in \Phi$ but $\alpha + 2\beta, 2\alpha + \beta \notin \Phi$;

$$(R4) \quad [x_\alpha(a), x_\beta(b)] = x_{\alpha+\beta}(N_{\alpha,\beta} \cdot ab) \cdot x_{2\alpha+\beta}(N_{\alpha,\beta}^{2,1} \cdot a^2b),$$

if $\alpha + \beta, 2\alpha + \beta \in \Phi$.

The constants $N_{\alpha,\beta}$ and $N_{\alpha,\beta}^{2,1}$ appearing in the above relations are called the **structure constants** of the Chevalley group of type Φ . Let us take a closer look at them.

First of all, notice that we excluded the case $\Phi = G_2$ so the only possibilities for $N_{\alpha,\beta}$ appearing in the above relations are ± 1 or ± 2 . Notice that $|N_{\alpha,\beta}| = 2$ only when α and β are short but $\alpha + \beta$ is long, in which case we set $\widehat{N}_{\alpha,\beta} = \frac{1}{2}N_{\alpha,\beta}$. In the other cases $|N_{\alpha,\beta}| = 1$. Now, by definition,

$$(2.10) \quad N_{\alpha,\beta}^{2,1} = N_{\alpha,\beta} \cdot \widehat{N}_{\alpha,\alpha+\beta}.$$

It is clear that $|N_{\alpha,\beta}^{2,1}| = 1$.

Many different methods of the choice of signs of the structure constants have been proposed in the literature, see e.g., [31]. Regardless of their concrete choice, however, the structure constants always must satisfy certain relations. First of all, recall from [31, §14] that

$$(2.11) \quad N_{\alpha,\beta} = -N_{\beta,\alpha} = -N_{-\alpha,-\beta} = \frac{|\alpha + \beta|^2}{|\alpha|^2} N_{\beta,-\alpha-\beta} = \frac{|\alpha + \beta|^2}{|\beta|^2} N_{-\alpha-\beta,\alpha}.$$

These identities will be used in the sequel without explicit reference.

We also will need another identity for structure constants. To formulate it succinctly, we extend the domain of the structure constant function $N_{-,-}$ by setting $N_{\alpha,\beta} = 0$ whenever $\alpha + \beta \notin \Phi \setminus \{0\}$. Now if α, β, γ is a triple of pairwise linearly independent roots such that $\alpha + \beta + \gamma \neq 0$, then one has

$$(2.12) \quad N_{\alpha,\beta+\gamma}N_{\beta,\gamma} + N_{\beta,\gamma+\alpha}N_{\gamma,\alpha} + N_{\gamma,\alpha+\beta}N_{\alpha,\beta} = 0.$$

This identity is an equivalent form of (N9) in [31, §14] (cf. also (H4) in [18]).

Sometimes it is convenient to write down the relations (R2)–(R4) as a single relation

$$(2.13) \quad [x_\alpha(a), x_\beta(b)] = \prod_{\substack{i\alpha+j\beta \in \Phi \\ i,j > 0}} x_{i\alpha+j\beta}(N_{\alpha,\beta}^{i,j} \cdot a^i b^j),$$

in which

$$N_{\alpha,\beta}^{1,2} := -N_{\beta,\alpha}^{2,1} \quad \text{and} \quad N_{\alpha,\beta}^{1,1} := N_{\alpha,\beta}.$$

Notice that the multiplicative identity of R is actually never used in the definition of the Steinberg group, which allows one to use it in the situation when R is an rng.

Definition 2.10: Let Φ be a root system of rank ≥ 3 . Applying the Steinberg group functor $\text{St}(\Phi, -)$ to the projective system of rngs $R^{(s)}$ (see Definition 2.4) we obtain a projective system of groups, whose formal projective limit will be denoted by $\text{St}^{(\infty)}(\Phi, R)$ and will be called the **Steinberg pro-group**.

Remark 2.11: There is a group homomorphism $\text{St}(\Phi, R^{(s)}) \rightarrow \text{St}(\Phi, sR)$ given by $x_\alpha^{(s)}(a) \mapsto x_\alpha(sa)$. By Remark 2.5 this homomorphism is an isomorphism in the case where R is an integral domain. Thus, similarly to pro-rngs, Steinberg pro-groups often do not have global elements.

For every root $\alpha \in \Phi$ there is a “root subgroup” morphism in $\text{Pro}(\mathbf{Grp})$,

$$x_\alpha : R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi, R),$$

defined by the pre-morphism

$$x_\alpha^* = \text{id}_S, \quad x_\alpha^{(s)}(a^{(s)}) = x_\alpha(a^{(s)}).$$

We denote by $G_{\text{sc}}(\Phi, -)$ the simply-connected Chevalley–Demazure group scheme corresponding to a root system Φ . We also fix a pinning for $G_{\text{sc}}(\Phi, -)$. By this we mean that a split maximal torus T of $G_{\text{sc}}(\Phi, -)$ and a Borel subgroup B containing T are chosen, and for every $\alpha \in \Phi$ there is an embedding $t_\alpha : \mathbb{G}_a \rightarrow G_{\text{sc}}(\Phi, -)$. For $a \in R$ we call the element $t_\alpha(a)$ of $G_{\text{sc}}(\Phi, R)$ an **elementary root unipotent**, cf. [31]. Notice that sometimes different notation is used for these elements (e.g., $x_\alpha(a)$ or $e_\alpha(a)$, cf. [14, 21, 31]).

Recall that for an ideal $I \trianglelefteq R$ the **congruence subgroup** $G_{\text{sc}}(\Phi, R, I)$ is defined as the kernel of the homomorphism $G_{\text{sc}}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R/I)$.

Recall also that one can define the unitalization of an rng R as the semidirect product $R \rtimes \mathbb{Z}$ (cf., e.g., [20, Definition 3.2]).

Definition 2.12: Consider the projective system of congruence subgroups

$$G(\Phi, R^{(s)}) := G_{\text{sc}}(\Phi, R^{(s)} \rtimes \mathbb{Z}, R^{(s)}) = \text{Ker}(G_{\text{sc}}(\Phi, R^{(s)} \rtimes \mathbb{Z}) \rightarrow G_{\text{sc}}(\Phi, \mathbb{Z}))$$

with the structure morphisms induced by the structure morphisms of $R^{(\infty)}$. Define the **simply-connected Chevalley pro-group** $G^{(\infty)}(\Phi, R)$ as the formal projective limit of this system.

Recall that there is a well-defined homomorphism $\text{st}: \text{St}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R)$ sending each generator $x_\alpha(a)$ to the root unipotent $t_\alpha(a)$. The pro-group analogue $\text{st}: \text{St}^{(\infty)}(\Phi, R) \rightarrow G^{(\infty)}(\Phi, R)$ of this homomorphism is given by the pre-morphism $\text{st}^* = \text{id}_S$, $\text{st}^{(s)}(x_\alpha(a^{(s)})) = t_\alpha(a^{(s)})$.

Remark 2.13: Let $\Sigma \subseteq \Phi$ be a closed special root subset and R be an rng. It is well-known that the restriction of the canonical map $\text{St}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R)$ to the subgroup

$$U(\Sigma, R) = \langle x_\alpha(a) \mid a \in R, \alpha \in \Sigma \rangle \leq \text{St}(\Phi, R)$$

is injective. Moreover, for any chosen linear order on Σ the map $R^\Sigma \rightarrow U(\Sigma, R)$ given by $(r_\alpha)_\alpha \mapsto \prod_\alpha x_\alpha(r_\alpha)$ is a bijection. As a consequence, the subgroup $U(\Sigma, R)$ admits presentation by means of generators $x_\alpha(a)$ for $\alpha \in \Sigma$ and the relations (R1)–(R4) in which $\alpha, \beta \in \Sigma$.

Remark 2.14: The definition of the group $G(\Phi, R^{(s)})$ can be reformulated in terms of Hopf algebras. Denote by H_Φ the Hopf \mathbb{Z} -algebra corresponding to the group scheme $G_{\text{sc}}(\Phi, -)$. Thus,

$$G_{\text{sc}}(\Phi, R) = \mathbf{Ring}(H_\Phi, R).$$

For a pair A, B of abelian groups we define the product $A \widehat{\otimes} B$ as follows:

$$A \widehat{\otimes} B := (A \otimes_{\mathbb{Z}} B) \oplus A \oplus B.$$

Recall that if R_1 and R_2 are rngs then $R_1 \widehat{\otimes} R_2$ realizes their coproduct in the category **Rng**. The product structure is induced from that of $(R_1 \rtimes \mathbb{Z}) \otimes_{\mathbb{Z}} (R_2 \rtimes \mathbb{Z})$. The canonical morphisms $R_i \rightarrow R_1 \widehat{\otimes} R_2$, $i = 1, 2$ are the obvious inclusions $r_1 \mapsto (0, r_1, 0)$, $r_2 \mapsto (0, 0, r_2)$. **Rng** has the structure of a symmetric monoidal category with zero rng as the unit object and $\widehat{\otimes}$ as the tensor product. For an rng R we denote by m the natural homomorphism $R^{\widehat{\otimes} 2} \rightarrow R$ given by

$$(s \otimes t, u, v) \mapsto st + u + v.$$

Notice also that the usual \mathbb{Z} -module tensor product $R_1 \otimes_{\mathbb{Z}} R_2$ has the natural structure of an rng and that $R_1 \otimes R_2 \subseteq R_1 \widehat{\otimes} R_2$. A priori there are no homomorphisms $R_i \rightarrow R_1 \otimes R_2$. However, if R_1 happens to be a ring, there is a canonical homomorphism $R_2 \rightarrow R_1 \otimes R_2$.

Recall that a **Hopf monoid** H in a symmetric monoidal category, by definition, is a bimonoid $(H, m, u, \Delta, \varepsilon)$ with a commutative m such that there is a morphism $\varsigma: H \rightarrow H$, called the **antipode**, satisfying $u \circ \varepsilon = m \circ (1 \otimes \varsigma) \circ \Delta$. We claim that the augmentation ideal $\text{Ker}(\varepsilon)$ of H_{Φ} has the structure of a Hopf monoid in **Rng**. The coproduct Δ and the antipode ς are induced from the corresponding morphisms defining the structure of a Hopf algebra on H_{Φ} . The counit ε is the unique homomorphism $\text{Ker}(\varepsilon) \rightarrow 0$.

Now if R is an arbitrary ring and $s \in R$ then there is a canonical isomorphism $\mathbf{Rng}(\text{Ker}(\varepsilon), R^{(s)}) \cong \mathbf{G}(\Phi, R^{(s)})$ induced by the mapping

$$(h: \text{Ker}(\varepsilon) \rightarrow R^{(s)}) \mapsto (\tilde{h}: H_{\Phi} \rightarrow R^{(s)} \rtimes \mathbb{Z}),$$

where $\tilde{h}(a) = (h(a - \varepsilon(a)), \varepsilon(a))$. The structure of a group on $\mathbf{G}(\Phi, R^{(s)})$ is then induced from the Hopf monoid structure on $\text{Ker}(\varepsilon)$ by duality. For instance, the neutral element in this group is the zero map $0: \text{Ker}(\varepsilon) \rightarrow R^{(s)}$. The product of $g, h \in \mathbf{Rng}(\text{Ker}(\varepsilon), R^{(s)})$ is the composite map

$$\text{Ker}(\varepsilon) \xrightarrow{\Delta} \text{Ker}(\varepsilon) \widehat{\otimes}^2 \xrightarrow{g \widehat{\otimes} h} R^{(s)} \widehat{\otimes} R^{(s)} \xrightarrow{m} R^{(s)}.$$

Remark 2.15: To formulate the next lemma, we need to introduce a way of reinterpreting the usual Steinberg relations (R2)–(R4) as relations between the morphisms x_{α} in the category $\text{Pro}(\mathbf{Grp})$. We call these relations the **pro-analogues** of (R2)–(R4). To obtain these relations one has to add the upper index (∞) to the free variables a, b occurring in (R2)–(R4) and then read the resulting expressions according to Convention 2.7. The fact that these relations hold is an easy consequence of our definitions.

Notice that the pro-analogue of (R1) is the identity

$$x_{\alpha}(a^{(\infty)})x_{\alpha}(b^{(\infty)}) = x_{\alpha}(a^{(\infty)} + b^{(\infty)}),$$

or in the usual notation $m \circ (x_{\alpha} \times x_{\alpha}) = x_{\alpha} \circ +_{R^{(\infty)}}$. It is clear that this relation simply means that x_{α} is a morphism of group objects in $\text{Pro}(\mathbf{Set})$, which is a consequence of the condition that x_{α} is a pro-group morphism.

Let $\mathcal{X} = \{x_i\}_{i \in I}$ be a collection of morphisms in $\text{Pro}(\mathbf{Grp})$ with a common codomain, which we denote by $S^{(\infty)}$. We say that \mathcal{X} **generates** $S^{(\infty)}$ if for every pair of pro-group morphisms $f_1, f_2: S^{(\infty)} \rightarrow G^{(\infty)}$ in order to verify the equality $f_1 = f_2$ it is enough to check that $f_1 x_i = f_2 x_i$ for $x_i \in \mathcal{X}$.

LEMMA 2.16: *The morphisms*

$$x_\alpha: R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi, R), \quad \alpha \in \Phi$$

generate $\text{St}^{(\infty)}(\Phi, R)$. Moreover, for every pro-group $G^{(\infty)}$ to obtain a morphism $f: \text{St}^{(\infty)}(\Phi, R) \rightarrow G$ it suffices to construct a collection of pro-group morphisms $f_\alpha: R^{(\infty)} \rightarrow G$, $\alpha \in \Phi$, satisfying the pro-analogues of (R2)–(R4), in which x_α 's are replaced with f_α 's.

Proof. Notice that by Lemma 2.1 both of the assertions can be verified in the special case where $G^{(\infty)} = G$ is a group.

Let us verify the first assertion. Since there is only a finite number of roots in Φ one can find a sufficiently large index $s \in S$ such that the homomorphisms $f_1^{(s)}, f_2^{(s)}: \text{St}(\Phi, R^{(s)}) \rightarrow G$ become equal after precomposition with x_α for all $\alpha \in \Phi$. Since the root elements $x_\alpha(a^{(s)})$ generate $\text{St}(\Phi, R^{(s)})$ we conclude that the morphisms f_1 and f_2 are equal.

Let us verify the second assertion. By definition, a morphism $f_\alpha: R^{(\infty)} \rightarrow G$ corresponds to a single group homomorphism $f'_\alpha: R^{(s_\alpha)} \rightarrow G$ for some $s_\alpha \in S$. Precomposing each f'_α with the structure morphism $R^{(s)} \rightarrow R^{(s_\alpha)}$ for sufficiently large $s \in S$, we obtain a collection of homomorphisms $\tilde{f}_\alpha: R^{(s)} \rightarrow G$.

Notice that (R2)–(R4) specify only a finite collection of identities in $\text{Pro}(\mathbf{Set})$ to which f_α 's must satisfy. Unwinding the definitions, we find a sufficiently large index $s \in S$ such that \tilde{f}_α satisfy the same identities (with f_α 's replaced by \tilde{f}_α 's). Thus, we obtain a group homomorphism $\text{St}(\Phi, R^{(s)}) \rightarrow G$, which, in turn, determines a morphism $\text{St}^{(\infty)}(\Phi, R) \rightarrow G$. ■

3. Elimination of roots

Throughout this section we always assume that R is an arbitrary commutative ring, $S \subseteq R$ is a multiplicative subset of R and Φ is a root system of rank ≥ 3 different from B_ℓ and C_ℓ .

LEMMA 3.1: *Let $\alpha, \beta, \gamma \in \Phi$ be a triple of roots having the same length such that $\alpha = \beta + \gamma$. Then there exist roots $\beta_1, \gamma_1 \in \Phi \setminus (\mathbb{Z}\beta + \mathbb{Z}\gamma)$ having the same length as β and, moreover, such that $\beta_1 + \gamma_1 = \beta$. The roots $\beta, \gamma, \beta_1, \gamma_1$ are contained in a root subsystem $\Phi_0 \subseteq \Phi$, whose type is either A_3 or C_3 .*

Proof. Denote by Φ' the subset of Φ consisting of roots having the same length as β . It follows from our assumptions that either Φ is simply-laced and $\Phi' = \Phi$ or $\Phi = F_4$, in which case Φ' is a root subset isomorphic to D_4 (notice that Φ' is closed only if β is long). In either case the Dynkin diagram of Φ' is simply laced, connected and has at least 3 vertices. Since all root subsystems of Φ' of type A_2 lie in the same orbit under the action of the Weyl group of Φ' , we may assume β and γ to be basic roots corresponding to adjacent nodes on the Dynkin diagram of Φ' , in which case the first assertion is clear.

For any choice of β_1 and γ_1 notice that the smallest root subsystem Φ_0 containing $\beta, \gamma, \beta_1, \gamma_1$ is irreducible and has rank 3, which implies that it is either of type C_3 (if $\Phi = F_4$ and β is short) or A_3 (otherwise). ■

As an immediate application of the above lemma we prove the following generation result, which is a stronger version of the first claim of Lemma 2.16.

LEMMA 3.2: *Let $\Phi_0 \subseteq \Phi$ be a rank 2 subsystem. The collection of root subgroup morphisms $x_\gamma: R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi, R)$ for $\gamma \in \Phi \setminus \Phi_0$ generates $\text{St}^{(\infty)}(\Phi, R)$.*

Proof. Thanks to Lemma 2.16 it suffices to show that $f_1 x_\delta = f_2 x_\delta$ for all $\delta \in \Phi_0$. Let δ be a root lying in Φ_0 . We claim that δ can be decomposed as a sum $\gamma_1 + \gamma_2$ for some $\gamma_1, \gamma_2 \in \Phi \setminus \Phi_0$ having the same length as δ . We need to consider several cases.

In the case $\Phi_0 \cong A_2$ the claim follows from Lemma 3.1. Consider the case $\Phi_0 \cong B_2$, which is only possible for $\Phi = F_4$. Since all root subsystems of type B_2 lie in the same orbit under the action of $W(F_4)$ (cf., e.g., [5, Table 8]), we may assume that Φ_0 corresponds to the central edge of the Dynkin diagram of F_4 and δ is one of the nodes incident to it, in which case the claim is obvious. Finally, consider the case $\Phi_0 \cong A_1 + A_1$. If either Φ is simply-laced or Φ_0 has roots of different length, acting by a suitable element of $W(\Phi)$ we may achieve that the basis vectors of Φ_0 correspond to a pair of unjoined nodes of the Dynkin diagram of Φ , in which case the claim is also obvious. On the other hand, if $\Phi = F_4$ and the roots of Φ_0 have the same length, the claim can be proved by passing to the root subsystem $\Phi' \cong D_4$ consisting of roots of Φ having the same length as the roots of Φ_0 .

As an immediate consequence of the claim we conclude that

$$\begin{aligned} f_1(x_\delta(a^{(\infty)}b^{(\infty)})) &= [f_1(x_{\gamma_1}(N_{\gamma_1,\gamma_2}a^{(\infty)})), f_1(x_{\gamma_2}(b^{(\infty)}))] \\ &= [f_2(x_{\gamma_1}(N_{\gamma_1,\gamma_2}a^{(\infty)})), f_2(x_{\gamma_2}(b^{(\infty)}))] \\ &= f_2(x_\delta(a^{(\infty)}b^{(\infty)})), \end{aligned}$$

which together with Lemma 2.6 implies the required equality $f_1x_\delta = f_2x_\delta$. ■

Definition 3.3: We denote by $\text{St}(\Phi \setminus \{\alpha\}, R)$ the group given by generators $x_\beta(a)$, $\beta \neq \alpha$, $a \in R$ and the subset of the set of relations (R1)–(R4) consisting of those relations, expressions for which do not contain $x_\alpha(a)$.

Similarly to Definition 2.10 one can define the pro-group $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$ as the formal projective limit of the projective system of groups $\text{St}(\Phi \setminus \{\alpha\}, R^{(s)})$ where $s \in S$.

For $\beta \neq \alpha$ we use the same notation x_β for the “root subgroup” morphisms

$$R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R).$$

Obviously, these morphisms satisfy the pro-analogues of (R2)–(R4) not containing α in their notations.

We can formulate the analogue of Lemma 2.16 for $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$.

LEMMA 3.4: *The morphisms x_β for $\beta \neq \alpha$ generate $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$. To construct a morphism $f: \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R) \rightarrow G^{(\infty)}$ it suffices to construct a collection of pro-group morphisms $f_\beta: R^{(\infty)} \rightarrow G^{(\infty)}$ satisfying the pro-analogues of those relations (R2)–(R4) in which x_α does not appear (with x_β ’s replaced with f_β ’s). Moreover, among these pro-analogues we can omit all relations of the form*

$$[x_\beta(b^{(\infty)}), x_\gamma(c^{(\infty)})] = 1$$

for all pairs of long roots β, γ such that $\beta + \gamma = 2\alpha$.

Proof. The proof of the first two assertions is analogous to Lemma 2.16. Let us verify the last assertion. Notice that Φ must be of type F_4 , otherwise there are no such relations. Acting by a suitable element of $W(F_4)$ we may assume, without loss of generality, that $\alpha = \frac{1}{2}(e_1 + e_2 + e_3 + e_4)$, $\beta = e_1 + e_2$, $\gamma = e_3 + e_4$ (we assume that F_4 is realized in \mathbb{R}^4 as in [4, Ch. VI, §4.9]). Set

$$\gamma_1 = e_4 \quad \text{and} \quad \gamma_2 = e_3 - e_4.$$

Since $\beta + \gamma_1, \beta + \gamma_2, \beta + \gamma_1 + \gamma_2 \notin \Phi \cup \{0, 2\alpha\}$,

$$\begin{aligned} & [x_\beta(b^{(\infty)}), x_\gamma(N_{\gamma_1, \gamma_2}^{2,1} c^{(\infty)2} d^{(\infty)})] \\ &= [x_\beta(b^{(\infty)}), x_{\gamma_1 + \gamma_2}(N_{\gamma_1, \gamma_2} c^{(\infty)} d^{(\infty)}) x_\gamma(N_{\gamma_1, \gamma_2}^{2,1} c^{(\infty)2} d^{(\infty)})] \quad \text{by (R2), (2.1)} \\ &= [x_\beta(b^{(\infty)}), [x_{\gamma_1}(c^{(\infty)}), x_{\gamma_2}(d^{(\infty)})]] = 1 \quad \text{by (R2), (R3), (2.4)}. \end{aligned}$$

The morphism $c^{(\infty)2} d^{(\infty)}: R^{(\infty)} \times R^{(\infty)} \rightarrow R^{(\infty)}$ is a split epimorphism of pro-sets by an argument similar to the proof of Lemma 2.6. Thus, the identity $[x_\beta(b^{(\infty)}), x_\gamma(c^{(\infty)})] = 1$ is a consequence of the other defining relations for $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$. ■

There is a morphism of pro-groups $F_\alpha: \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R) \rightarrow \text{St}^{(\infty)}(\Phi, R)$, given by the pre-morphism $F_\alpha^* = \text{id}_S$ and the homomorphisms

$$F_\alpha^{(s)}: \text{St}(\Phi \setminus \{\alpha\}, R^{(s)}) \rightarrow \text{St}(\Phi, R^{(s)})$$

induced by the obvious embedding of generators.

Notice that, in general, the individual homomorphisms $F_\alpha^{(s)}$ need *not* be isomorphisms (at least if $s \neq 1$). On the other hand, as the following result shows, the morphism of pro-groups F_α is an isomorphism.

THEOREM 1: *For every root $\alpha \in \Phi$ the morphism F_α is an isomorphism of pro-groups.*

The proof of Theorem 1 occupies the rest of this subsection. Our immediate goal is to construct the root subgroup morphism

$$\tilde{x}_\alpha: R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$$

“missing” from the presentation of $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$.

Definition 3.5: Let $\beta, \gamma \in \Phi$ be an arbitrary fixed pair of roots such that $\alpha = \beta + \gamma$ and α, β , and γ have the same length. Denote by $x_{\beta, \gamma}(b^{(\infty)}, c^{(\infty)})$ the morphism

$$[x_\beta(N_{\beta, \gamma} b^{(\infty)}), x_\gamma(c^{(\infty)})]: R^{(\infty)} \times R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R).$$

Below we will construct the morphism \tilde{x}_α based on the morphism $x_{\beta, \gamma}$, see Lemma 3.7. However, in order to be able to do this we first need to prove the following assertion.

LEMMA 3.6: *The morphism $x_{\beta,\gamma}$ satisfies the following relations:*

$$(3.1) \quad [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_\delta(d^{(\infty)})] = 1, \quad \text{if } \alpha + \delta \notin \Phi \cup \{0\}, \delta \neq \alpha;$$

$$(3.2) \quad [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_\delta(d^{(\infty)})] = x_{\alpha+\delta}(N_{\alpha,\delta} \cdot b^{(\infty)} c^{(\infty)} d^{(\infty)}), \\ \text{if } \alpha + \delta \in \Phi \text{ but } \alpha + 2\delta, 2\alpha + \delta \notin \Phi;$$

$$(3.3) \quad [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_\delta(d^{(\infty)})] \\ = x_{\alpha+\delta}(N_{\alpha,\delta} \cdot b^{(\infty)} c^{(\infty)} d^{(\infty)}) \cdot x_{2\alpha+\delta}(N_{\alpha,\delta}^{2,1} \cdot b^{(\infty)^2} c^{(\infty)^2} d^{(\infty)}) \\ \text{if } \alpha + \delta, 2\alpha + \delta \in \Phi;$$

$$(3.4) \quad [x_\delta(d^{(\infty)}), x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)})] \\ = x_{\alpha+\delta}(N_{\delta,\alpha} \cdot b^{(\infty)} c^{(\infty)} d^{(\infty)}) \cdot x_{\alpha+2\delta}(N_{\delta,\alpha}^{2,1} \cdot b^{(\infty)} c^{(\infty)} d^{(\infty)^2}) \\ \text{if } \alpha + \delta, \alpha + 2\delta \in \Phi.$$

Proof. First of all, observe that the above relations are obtained from (R2)–(R4) by replacing $x_\alpha(a^{(\infty)})$ with $x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)})$.

Let us first consider the case $\delta \in \Phi \setminus (\mathbb{Z}\beta + \mathbb{Z}\gamma)$. We claim that the relations between the remaining root subgroup morphisms of $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$ suffice to rewrite the left-hand side of any of the relations (3.1)–(3.4) as follows:

$$(3.5) \quad [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_\delta(d^{(\infty)})] \\ = x_\beta(N_{\beta,\gamma} b^{(\infty)}) x_\gamma(c^{(\infty)}) x_\beta(-N_{\beta,\gamma} b^{(\infty)}) x_\gamma(-c^{(\infty)}) x_\delta(d^{(\infty)}) x_\delta(-d^{(\infty)}) \\ = \prod_{\substack{i\beta+j\gamma+k\delta \in \Phi \\ i,j \geq 0; k > 0}} x_{i\beta+j\gamma+k\delta}(A_{i,j,k} b^{(\infty)^i} c^{(\infty)^j} d^{(\infty)^k}).$$

Indeed, observe that the root subset $\Sigma = \Phi \cap (\mathbb{Z}_{\geq 0}\beta + \mathbb{Z}_{\geq 0}\gamma + \mathbb{Z}_{> 0}\delta)$ is special and does not contain $-\beta, -\gamma$ or α . Thus, we can iteratively simplify the expression for the conjugate of x_δ in the above formula obtaining a product of root subgroup morphisms corresponding to a subset of roots of Σ at each step. The integers $A_{i,j,k}$ in (3.5) depend only on the roots β, γ, δ (a priori they also depend on the chosen order of factors). To determine them one can compute the commutator $[x_\alpha(1), x_\delta(1)]$ in $\text{St}(\Phi, \mathbb{Z})$ via the same procedure as in (3.5):

$$(3.6) \quad [x_\alpha(1), x_\delta(1)] = \prod_{\substack{i\beta+j\gamma+k\delta \in \Phi \\ i,j \geq 0; k > 0}} x_{i\beta+j\gamma+k\delta}(A_{i,j,k}).$$

By Remark 2.13 the integers $A_{i,j,k}$ are uniquely determined by (3.6). It is easy to see that, depending on the relative position of α and δ , the integers $A_{i,j,k}$ coincide with the structure constants in the right-hand sides of (R2), (R3) or (R4), in particular, $A_{i,j,k} = 0$ for $i \neq j$. Thus, (3.1)–(3.4) follow from (3.5) in the specified case.

It remains to verify Steinberg relations in the case $\delta \in \Phi \cap (\mathbb{Z}\beta + \mathbb{Z}\gamma)$. By Lemma 3.1 there exist roots $\beta_1, \beta_2 \in \Phi \setminus (\mathbb{Z}\beta + \mathbb{Z}\gamma)$ such that $|\beta_1| = |\beta_2| = |\beta|$ and $\beta = \beta_1 + \beta_2$. Recall that the root subsystem $\Phi_0 \subseteq \Phi$ containing $\beta, \gamma, \beta_1, \beta_2$ is of type A_3 or C_3 . In both cases we may assume that $\alpha + \beta_2 \notin \Phi$.

Let us first verify the relation (3.1) in the case $\delta \in \{\beta, \gamma\}$. By symmetry we may assume that $\delta = \beta$. Direct computation shows that

$$\begin{aligned}
 & [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_{\beta}(N_{\beta_1,\beta_2}b_1^{(\infty)}b_2^{(\infty)})] \\
 (3.7) \quad &= [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), [x_{\beta_1}(b_1^{(\infty)}), x_{\beta_2}(b_2^{(\infty)})]] \quad \text{by (R3)} \\
 &= [[x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_{\beta_1}(b_1)], x_{\beta_1}(b_1^{(\infty)})x_{\beta_2}(b_2^{(\infty)})] \quad \text{by (2.4),(3.1)}.
 \end{aligned}$$

In the case $\Phi_0 \cong A_3$ the inner commutator in the last expression is trivial by (3.1), therefore so is the outer commutator (notice that α and β_1 form an acute angle). In the case $\Phi_0 \cong C_3$ the last expression can be further simplified using (R3) and the already proved relation (3.2) as follows:

$$\dots = [x_{\alpha+\beta_1}(N_{\alpha,\beta_1}b^{(\infty)}c^{(\infty)}b_1^{(\infty)}), x_{\beta}(N_{\beta_1,\beta_2}b_1^{(\infty)}b_2^{(\infty)})x_{\beta_2}(b_2^{(\infty)})].$$

Since $\alpha + \beta_1$ forms an acute angle with both β and β_2 , the latter commutator is trivial by (2.1) and (3.1). In both cases the right-hand side of (3.7) is trivial. Consequently, from Lemma 2.6 we obtain the equality

$$[x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_{\beta}(d^{(\infty)})] = 1.$$

Thus, we have verified (3.1) in all cases.

Finally, it remains to verify (3.2) in the case $\delta \in \{-\beta, -\gamma\}$. By symmetry it suffices to consider the case $\delta = -\beta$. From the already proved relations (3.1)–(3.2) and the other Steinberg relations we obtain

$$\begin{aligned}
 & [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_{-\beta}(N_{-\beta_1,-\beta_2}b_1^{(\infty)}b_2^{(\infty)})] \\
 &= [x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)})x_{-\beta_1}(b_1^{(\infty)}), x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)})x_{-\beta_2}(b_2^{(\infty)})]x_{-\beta}(-N_{-\beta_1,-\beta_2}b_1^{(\infty)}b_2^{(\infty)}) \\
 &= \prod_{\substack{i\alpha-j\beta_1-k\beta_2 \in \Phi \\ i,j,k \geq 0}} x_{i\alpha-j\beta_1-k\beta_2}(B_{i,j,k}b^{(\infty)^i}c^{(\infty)^j}b_1^{(\infty)^j}b_2^{(\infty)^k}),
 \end{aligned}$$

where $B_{i,j,k}$ are certain integers. Direct computation (or an argument similar to the one used in the first part of the proof) shows that the constants $B_{i,j,k}$ are all zero with the sole exception of $B_{1,1,1}$, which is equal to $N_{\alpha,-\beta}$. Consequently, from Lemma 2.6 we obtain

$$[x_{\beta,\gamma}(b^{(\infty)}, c^{(\infty)}), x_{-\beta}(d^{(\infty)})] = x_\gamma(N_{\alpha,-\beta}b^{(\infty)}c^{(\infty)}d^{(\infty)}),$$

which finishes the proof of (3.2). ■

LEMMA 3.7: *The morphism $x_{\beta,\gamma}$ satisfies the identities listed in Lemma 2.8 and therefore gives rise to a pro-group morphism $R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$, which we denote by \tilde{x}_α . The resulting morphism \tilde{x}_α does not depend on the choice of β and γ .*

Proof. First of all, observe that (3.1) implies the identity

$$[x_{\beta,\gamma}(b_1^{(\infty)}, c_1^{(\infty)}), x_{\beta,\gamma}(b_2^{(\infty)}, c_2^{(\infty)})] = 1.$$

Notice also that (2.2) and (3.1) imply that

$$\begin{aligned} x_{\beta,\gamma}(b_1^{(\infty)} + b_2^{(\infty)}, c^{(\infty)}) &= [x_\beta(N_{\beta,\gamma}b_1^{(\infty)})x_\beta(N_{\beta,\gamma}b_2^{(\infty)}), x_\gamma(c^{(\infty)})] \\ &= x_{\beta,\gamma}(b_1^{(\infty)}, c^{(\infty)})x_{\beta,\gamma}(b_2^{(\infty)}, c^{(\infty)}). \end{aligned}$$

Similarly, one can show the equality

$$x_{\beta,\gamma}(b^{(\infty)}, c_1^{(\infty)} + c_2^{(\infty)}) = x_{\beta,\gamma}(b^{(\infty)}, c_1^{(\infty)})x_{\beta,\gamma}(b^{(\infty)}, c_2^{(\infty)}).$$

Thus, the morphism $x_{\beta,\gamma}$ satisfies the first three requirements of Lemma 2.8.

Now suppose that $\alpha = \beta_1 + \gamma_1 = \beta_2 + \gamma_2$ are two different decompositions for α such that α, β_i and γ_i have the same length and α, β_1, β_2 are linearly independent. By Lemma 3.1 such decompositions exist and the roots β_i, γ_i are contained in a root subsystem Φ_0 of type A_3 or C_3 . Swapping β_1 with γ_1 , if necessary, we also may assume that β_1 and β_2 form an acute angle. Set $\delta = \gamma_1 - \gamma_2 = \beta_2 - \beta_1$ and

$$\varepsilon_1 = N_{\beta_1,\gamma_1}, \quad \varepsilon_2 = N_{\beta_2,\gamma_2}, \quad \varepsilon_3 = N_{\delta,\gamma_2}, \quad \varepsilon_4 = N_{\beta_1,\delta}.$$

In order to simplify keeping track of the relative angles and the length of the roots, the reader may assume that β_i, γ_i are concretely realized in the space \mathbb{R}^3 as follows: $\beta_1 = e_1 - e_2, \gamma_1 = e_2 - e_3$ and

- $\beta_2 = e_1 - e_4, \gamma_2 = e_4 - e_3, \delta = e_2 - e_4$ in the case $\Phi_0 \cong A_3$,
- $\beta_2 = e_1 + e_2, \gamma_2 = -e_2 - e_3, \delta = 2e_2$ in the case $\Phi_0 \cong C_3$.

Here e_i denote the standard basis vectors of \mathbb{R}^3 and we assume that the root systems A_3 and C_3 are realized in \mathbb{R}^3 as in [4, Ch. VI, §§4.6–4.7].

Substituting the triple $(\beta_1, \gamma_1, -\beta_2)$ into (2.12), it is not hard to conclude that in both cases one has $\varepsilon_1\varepsilon_2\varepsilon_3\varepsilon_4 = 1$. Now direct calculation shows that

$$\begin{aligned}
& x_{\beta_1, \gamma_1}(b^{(\infty)}, c^{(\infty)}d^{(\infty)}) \\
&= [x_{\beta_1}(\varepsilon_1 b^{(\infty)}), x_{\gamma_1}(c^{(\infty)}d^{(\infty)})] \\
&= [x_{\beta_1}(\varepsilon_1 b^{(\infty)}), [x_\delta(\varepsilon_3 c^{(\infty)}), x_{\gamma_2}(d^{(\infty)})]] \\
&\quad \text{by (R3) if } \Phi_0 \cong A_3 \text{ and by (R2),(R4),(2.1) if } \Phi_0 \cong C_3 \\
&= [[x_{\beta_1}(\varepsilon_1 b^{(\infty)}), x_\delta(\varepsilon_3 c^{(\infty)})], x_{\gamma_2}(d^{(\infty)})] \quad \text{by (R2),(2.4)} \\
&= [x_{\beta_2}(\varepsilon_2 b^{(\infty)}c^{(\infty)}), x_{\gamma_2}(d^{(\infty)})x_{\gamma_1}(c^{(\infty)}d^{(\infty)})] \quad \text{by (R3)} \\
&= [x_{\beta_2}(\varepsilon_2 b^{(\infty)}c^{(\infty)}), x_{\gamma_2}(d^{(\infty)})] \quad \text{by (R2),(2.1)} \\
&= x_{\beta_2, \gamma_2}(b^{(\infty)}c^{(\infty)}, d^{(\infty)}).
\end{aligned}$$

Thus, in both cases we obtain the equality

$$(3.8) \quad x_{\beta_1, \gamma_1}(b^{(\infty)}, c^{(\infty)}d^{(\infty)}) = x_{\beta_2, \gamma_2}(b^{(\infty)}c^{(\infty)}, d^{(\infty)}).$$

It follows from (3.8) that

$$x_{\beta_1, \gamma_1}(b^{(\infty)}, u_1^{(\infty)}u_2^{(\infty)}d^{(\infty)}) = x_{\beta_1, \gamma_1}(b^{(\infty)}u_1^{(\infty)}u_2^{(\infty)}, d^{(\infty)})$$

therefore by Lemma 2.6, $x_{\beta_1, \gamma_1}(b^{(\infty)}, u^{(\infty)}c^{(\infty)}) = x_{\beta_1, \gamma_1}(b^{(\infty)}u^{(\infty)}, c^{(\infty)})$. Thus, the last requirement of Lemma 2.8 is satisfied. Consequently, there exists a unique morphism of pro-groups $\tilde{x}_\alpha: R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$ satisfying

$$(3.9) \quad x_{\beta_1, \gamma_1}(b^{(\infty)}, c^{(\infty)}) = \tilde{x}_\alpha(b^{(\infty)}c^{(\infty)}) = x_{\beta_2, \gamma_2}(b^{(\infty)}, c^{(\infty)}). \quad \blacksquare$$

The next step in the proof of Theorem 1 is to verify that \tilde{x}_α satisfies the relations (R2)–(R4). Combining Lemmas 3.6 and 3.7 we immediately obtain the following.

COROLLARY 3.8: *The root subgroup morphism \tilde{x}_α satisfies the pro-analogues of Steinberg relations of type (R2)–(R4), in which it occurs on the left-hand side.*

Thus, to finish the proof of Theorem 1 it suffices to consider the case where \tilde{x}_α occurs in the right-hand side of the commutator formula.

LEMMA 3.9: *The root subgroup morphism \tilde{x}_α satisfies the pro-analogues of Steinberg relations of type (R4) in which it occurs on the right-hand side.*

Proof. We only need to consider the case $\Phi = F_4$. We need to verify the relation obtained from

$$(3.10) \quad [x_\beta(b^{(\infty)}), x_\gamma(c^{(\infty)})] = x_{\beta+\gamma}(N_{\beta,\gamma}b^{(\infty)}c^{(\infty)})x_{2\beta+\gamma}(N_{\beta,\gamma}^{2,1}b^{(\infty)^2}c^{(\infty)})$$

by replacing either $x_{\beta+\gamma}$ or $x_{2\beta+\gamma}$ with \tilde{x}_α .

First, let us consider the case $\alpha = \beta + \gamma$, in which α, β are short and γ is long. Consider a decomposition $\beta = \beta_1 + \beta_2$ for some short roots β_i . The smallest root subsystem containing β_i and γ is of type C_3 , so we may assume that $\alpha = e_1 + e_3, \beta = e_1 - e_3, \beta_1 = e_1 - e_2, \beta_2 = e_2 - e_3, \gamma = 2e_3$. Set

$$\begin{aligned} \varepsilon_1 &= N_{\beta_1, \beta_2}, & \varepsilon_2 &= N_{\beta_2, \gamma}^{2,1}, & \varepsilon_3 &= N_{\beta_2, \gamma}, \\ \varepsilon_4 &= N_{\beta_1, 2\beta_2 + \gamma}^{2,1}, & \varepsilon_5 &= N_{\beta_1, 2\beta_2 + \gamma}, & \varepsilon_6 &= N_{\beta_1, \beta_2 + \gamma}, & \varepsilon_7 &= N_{\beta_2 + \gamma, \beta}. \end{aligned}$$

Substituting triples $(\beta_1, \beta_2 + \gamma, \beta_2), (\beta_1, \beta_2, \gamma), (\beta_1, \beta_1 + \beta_2, \beta_2 + \gamma)$ into (2.12) we obtain the equalities

$$\varepsilon_2\varepsilon_3\varepsilon_5 = -\varepsilon_1\varepsilon_7, \quad \varepsilon_3\varepsilon_6 = N_{\beta,\gamma}\varepsilon_1, \quad 2\varepsilon_4\varepsilon_5\varepsilon_7 = -\varepsilon_6N_{\beta,\alpha},$$

which together with (2.10) imply the equality

$$\varepsilon_2\varepsilon_4 = \frac{1}{2}\varepsilon_1\varepsilon_3\varepsilon_5\varepsilon_7 \cdot \varepsilon_5\varepsilon_6\varepsilon_7N_{\beta,\alpha} = \frac{1}{2}\varepsilon_1\varepsilon_3\varepsilon_6N_{\beta,\alpha} = \frac{1}{2}N_{\beta,\gamma}N_{\beta,\alpha} = N_{\beta,\gamma}^{2,1}.$$

Now a direct computation using (R2)–(R4) and Corollary 3.8 shows that

$$\begin{aligned} & [x_\gamma(c^{(\infty)}), x_\beta(\varepsilon_1b_1^{(\infty)}b_2^{(\infty)})] \\ &= [x_{\beta_1}(b_1^{(\infty)}), x_\gamma(c^{(\infty)})x_{\beta_2}(b_2^{(\infty)})] \cdot x_\beta(-\varepsilon_1b_1^{(\infty)}b_2^{(\infty)}) \quad \text{by (2.5)} \\ &= [x_{\beta_1}(b_1^{(\infty)}), x_{2\beta_2+\gamma}(-\varepsilon_2b_2^{(\infty)^2}c^{(\infty)})x_{\beta_2+\gamma}(-\varepsilon_3b_2^{(\infty)}c^{(\infty)})x_{\beta_2}(b_2^{(\infty)})] \\ & \quad \cdot x_\beta(-\varepsilon_1b_1^{(\infty)}b_2^{(\infty)}) \\ &= [x_{\beta_1}(b_1^{(\infty)}), x_{2\beta_2+\gamma}(-\varepsilon_2b_2^{(\infty)^2}c^{(\infty)})] \\ & \quad \cdot x_{2\beta_2+\gamma}(-\varepsilon_2b_2^{(\infty)^2}c^{(\infty)})[x_{\beta_1}(b_1^{(\infty)}), x_{\beta_2+\gamma}(-\varepsilon_3b_2^{(\infty)}c^{(\infty)})] \\ & \quad \cdot [x_{2\beta_2+\gamma}(-\varepsilon_2b_2^{(\infty)^2}c^{(\infty)})x_{\beta_2+\gamma}(-\varepsilon_3b_2^{(\infty)}c^{(\infty)}), x_\beta(\varepsilon_1b_1^{(\infty)}b_2^{(\infty)})] \quad \text{by (2.1)} \\ &= x_{\beta_1+2\beta_2+\gamma}(-\varepsilon_2\varepsilon_5b_1^{(\infty)}b_2^{(\infty)^2}c^{(\infty)}) \cdot x_{2\beta_2+\gamma}(-\varepsilon_2\varepsilon_4b_1^{(\infty)^2}b_2^{(\infty)^2}c^{(\infty)}) \\ & \quad \cdot \tilde{x}_\alpha(-\varepsilon_3\varepsilon_6b_1^{(\infty)}b_2^{(\infty)}c^{(\infty)}) \cdot x_{\beta_1+2\beta_2+\gamma}(-\varepsilon_1\varepsilon_3\varepsilon_7b_1^{(\infty)}b_2^{(\infty)^2}c^{(\infty)}) \quad \text{by (2.2), (3.9)} \\ &= x_{2\beta_2+\gamma}(-N_{\beta,\gamma}^{2,1}b_1^{(\infty)^2}b_2^{(\infty)^2}c^{(\infty)}) \cdot \tilde{x}_\alpha(-N_{\beta,\gamma}\varepsilon_1b_1^{(\infty)}b_2^{(\infty)}c^{(\infty)}). \end{aligned}$$

Thus, by Lemma 2.6

$$[x_\gamma(c^{(\infty)}), x_\beta(b^{(\infty)})] = x_{2\beta+\gamma}(-N_{\beta,\gamma}^{2,1}b^{(\infty)2}c^{(\infty)}) \cdot \tilde{x}_\alpha(-N_{\beta,\gamma}b^{(\infty)}c^{(\infty)}),$$

which is an equivalent form of (3.10).

Now consider the case $\alpha = 2\beta + \gamma$, in which β is short and α, γ are long. We can choose a decomposition $\gamma = \gamma_1 + \gamma_2$ for some long roots γ_i . The smallest root subsystem containing β and γ_i is of type B_3 , so we may assume that $\alpha = e_1 + e_3$, $\beta = e_3$, $\gamma = e_1 - e_3$, $\gamma_1 = e_1 - e_2$, $\gamma_2 = e_2 - e_3$. Set

$$\varepsilon_1 = N_{\gamma_1, \gamma_2}, \quad \varepsilon_2 = N_{\beta, \gamma_2}^{2,1}, \quad \varepsilon_3 = N_{\beta, \gamma_2}, \quad \varepsilon_4 = N_{\gamma_1, 2\beta + \gamma_2}, \quad \varepsilon_5 = N_{\gamma_1, \beta + \gamma_2}.$$

Substituting the triples $(\beta, \gamma_2, \gamma_1)$, $(\beta, \beta + \gamma_2, \gamma_1)$ into (2.12) we obtain the identities

$$\varepsilon_3\varepsilon_5 = \varepsilon_1N_{\beta, \gamma}, \quad 2\varepsilon_2\varepsilon_3\varepsilon_4 = \varepsilon_5N_{\beta, \beta + \gamma}.$$

Notice that these identities imply that $\varepsilon_2\varepsilon_4 = \frac{1}{2}\varepsilon_3\varepsilon_5N_{\beta, \beta + \gamma} = \varepsilon_1N_{\beta, \gamma}^{2,1}$. A direct computation using (R2)–(R4), (2.1), (3.9) and Corollary 3.8 shows that

$$\begin{aligned} & [x_\beta(b^{(\infty)}), x_\gamma(\varepsilon_1c_1^{(\infty)}c_2^{(\infty)})] \\ &= [x_{\gamma_1}(c_1^{(\infty)}), x_{\beta(b^{(\infty)})}x_{\gamma_2}(c_2^{(\infty)})]x_\gamma(-\varepsilon_1c_1^{(\infty)}c_2^{(\infty)}) \\ &= [x_{\gamma_1}(c_1^{(\infty)}), x_{2\beta+\gamma_2}(\varepsilon_2b^{(\infty)2}c_2^{(\infty)})x_{\beta+\gamma_2}(\varepsilon_3b^{(\infty)}c_2^{(\infty)})x_{\gamma_2}(c_2^{(\infty)})]x_\gamma(-\varepsilon_1c_1^{(\infty)}c_2^{(\infty)}) \\ &= \tilde{x}_\alpha(\varepsilon_2\varepsilon_4b^{(\infty)2}c_1^{(\infty)}c_2^{(\infty)}) \cdot x_{\beta+\gamma}(\varepsilon_3\varepsilon_5b^{(\infty)}c_1^{(\infty)}c_2^{(\infty)}). \end{aligned}$$

Now Lemma 2.6 and the above identity imply (3.10). \blacksquare

LEMMA 3.10: *The root subgroup morphism \tilde{x}_α satisfies the pro-analogues of Steinberg relations of type (R3) in which it occurs on the right-hand side.*

Proof. We need to verify the relation

$$(3.11) \quad [x_\beta(b^{(\infty)}), x_\gamma(c^{(\infty)})] = \tilde{x}_\alpha(N_{\beta,\gamma}b^{(\infty)}c^{(\infty)}).$$

In the case where α, β and γ have the same length the assertion follows from Lemma 3.7. Thus, we only need to consider the case where $\alpha = \beta + \gamma$ and α is long while β and γ are short. The smallest root subsystem containing α, β, γ is of type B_2 . Without loss of generality we may assume $\alpha = e_1 + e_2$, $\beta = e_1$,

and $\gamma = e_2$. Set $\varepsilon_1 = N_{\beta, \gamma-\beta}^{2,1}$, $\varepsilon_2 = N_{\beta, \gamma-\beta}$. Direct computation shows that

$$\begin{aligned}
 & x_{\beta(d^{(\infty)})} x_{\gamma}(\varepsilon_2 b^{(\infty)} c^{(\infty)}) \cdot \tilde{x}_{\alpha}(\varepsilon_1 b^{(\infty)2} c^{(\infty)}) \\
 &= x_{\beta(d^{(\infty)})}(x_{\gamma}(\varepsilon_2 b^{(\infty)} c^{(\infty)}) \tilde{x}_{\alpha}(\varepsilon_1 b^{(\infty)2} c^{(\infty)})) && \text{by Corollary 3.8} \\
 &= [x_{\beta}(b^{(\infty)}), x_{\beta(d^{(\infty)})} x_{\gamma-\beta}(c^{(\infty)})] && \text{by (R2), Lemma 3.9} \\
 &= x_{\beta(b^{(\infty)}+d^{(\infty)})} x_{\gamma-\beta}(c^{(\infty)}) \cdot x_{\beta(d^{(\infty)})} x_{\gamma-\beta}(-c^{(\infty)}) && \text{by (R1)} \\
 &= \tilde{x}_{\alpha}(\varepsilon_1 (b^{(\infty)} + d^{(\infty)})^2 c^{(\infty)}) x_{\gamma}(\varepsilon_2 (b^{(\infty)} + d^{(\infty)}) c^{(\infty)}) x_{\gamma-\beta}(c^{(\infty)}) \\
 &\quad \cdot \tilde{x}_{\alpha}(-\varepsilon_1 c^{(\infty)} d^{(\infty)2}) x_{\gamma}(-\varepsilon_2 c^{(\infty)} d^{(\infty)}) x_{\gamma-\beta}(-c^{(\infty)}) && \text{by (R4)} \\
 &= x_{\gamma}(\varepsilon_2 b^{(\infty)} c^{(\infty)}) \tilde{x}_{\alpha}(2\varepsilon_1 b^{(\infty)} c^{(\infty)} d^{(\infty)} + \varepsilon_1 b^{(\infty)2} c^{(\infty)}) && \text{by Lemma 3.9}
 \end{aligned}$$

The relation (3.11) now follows from Lemma 2.6 and (2.10). ■

Proof of Theorem 1. To distinguish between the root subgroup morphisms of $\text{St}^{(\infty)}(\Phi, R)$ and $\text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R)$ we rename the root subgroup morphisms x_{β} , $\beta \neq \alpha$ of $\text{St}(\Phi \setminus \{\alpha\}, R^{(\infty)})$ to \tilde{x}_{β} . For the morphisms x_{β} of $\text{St}^{(\infty)}(\Phi, R)$ we continue to use the usual notation.

By Lemma 2.16 the morphisms \tilde{x}_{β} give rise to a morphism

$$G_{\alpha}: \text{St}^{(\infty)}(\Phi, R) \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{\alpha\}, R).$$

By definition, $G_{\alpha} x_{\beta} = \tilde{x}_{\beta}$ for all β . On the other hand, it is clear that $F_{\alpha} \tilde{x}_{\beta} = x_{\beta}$ for $\beta \neq \alpha$. To see that G_{α} is inverse to F_{α} it remains to invoke the first part of Lemma 2.16 and Lemma 3.4. ■

4. The action of $G_{\text{sc}}(\Phi, S^{-1}R)$ on pro-groups

Throughout this section R is a ring and $S \subseteq R$ is a multiplicative subset.

4.1. THE CASE OF CHEVALLEY GROUPS. Our first goal is to show that the group $G_{\text{sc}}(\Phi, S^{-1}R)$ can be made to act on the Chevalley pro-group $G^{(\infty)}(\Phi, R)$ by conjugation. Denote by H_{Φ} the Hopf \mathbb{Z} -algebra of $G_{\text{sc}}(\Phi, -)$ with the structure of a Hopf algebra given by the triple $(\Delta, \varsigma, \varepsilon)$. For shortness, we denote by K the augmentation ideal $\text{Ker}(\varepsilon)$. Recall that $H_{\Phi} = K \rtimes \mathbb{Z}$ is a finitely presented commutative \mathbb{Z} -algebra, therefore K is a finitely generated rng. Indeed, if x_i are the generators of H_{Φ} , then $y_i := x_i - \varepsilon(x_i)$ generate K as an rng as

seen, e.g., from the identity

$$xy - \varepsilon(xy) = (x - \varepsilon(x))(y - \varepsilon(y)) + (x - \varepsilon(x))\varepsilon(y) + \varepsilon(x)(y - \varepsilon(y)).$$

There is a surjective homomorphism $\mathbb{Z}[X_1, \dots, X_n] \rightarrow K \rtimes \mathbb{Z}$ sending X_i to x_i , whose kernel is generated by a finite collection of polynomials g_1, \dots, g_m . Substituting $X_i \mapsto Y_i + \varepsilon(x_i)$ into g_j 's we obtain a new collection of polynomials $\{f_j(Y_1, \dots, Y_n)\}$ without constant terms. It is clear that f_j 's form a complete system of defining relations between y_i . Denote by N the maximum of the degrees of f_j 's. Consider the ‘‘coconjugation’’ homomorphism (written down in Sweedler notation)

$$\text{Coconj}: K \rightarrow H_\Phi \otimes K, \quad h \mapsto \sum h_{(1)}\varsigma(h_{(3)}) \otimes h_{(2)}.$$

Recall that $S^{-1}R$ is isomorphic as an R -module to the filtered colimit of R -modules $\frac{R}{s}$ over the category \mathcal{S} (cf. Section 2.3), each of these modules being isomorphic to R . For $s, s' \in S$ the homomorphism $\text{can}: \frac{R}{s} \rightarrow \frac{R}{ss'}$ is given by

$$\frac{r}{s} \mapsto \frac{rs'}{ss'}.$$

The elements of modules $\frac{R}{s}$ can be ‘‘multiplied’’ with each other via the natural R -module homomorphisms $\frac{R}{s} \otimes \frac{R}{s'} \rightarrow \frac{R}{ss'}$ compatible with the canonical homomorphisms $\frac{R}{s} \rightarrow S^{-1}R$.

Let g be an element of $G_{\text{sc}}(\Phi, S^{-1}R) \cong \mathbf{Ring}(H_\Phi, S^{-1}R)$. Consider the rng homomorphism $g \otimes \text{id}: H_\Phi \otimes K \rightarrow S^{-1}R \otimes K$. Since tensor products commute with colimits, there exists $s_0 \in S$ such that $(g \otimes \text{id}) \circ \text{Coconj}(y_i)$ are the images of certain elements b_i of $\frac{R}{s_0} \otimes K$ under the canonical R -module homomorphism $\frac{R}{s_0} \otimes K \rightarrow S^{-1}R \otimes K$.

Notice that the elements of $\frac{R}{s_0} \otimes K$ also can be multiplied with each other via the obvious multiplication homomorphisms

$$\frac{R}{s} \otimes K \otimes \frac{R}{s'} \otimes K \rightarrow \frac{R}{ss'} \otimes K, \text{ where } s, s' \in S.$$

Using these homomorphisms we can compute the values $f_j(b_1, \dots, b_n)$ of the polynomials f_j 's on b_i 's. It is clear that these values can be interpreted as elements of $\frac{R}{s_0^N} \otimes K$ and that they are mapped to 0 under the homomorphism $\frac{R}{s_0^N} \otimes K \rightarrow S^{-1}R \otimes K$. Consequently, there is an element $s'_0 \in S$ annihilating all of them simultaneously. Set $s_1 = s_0^N s'_0$. It is clear that in $\frac{R}{s_1} \otimes K$ we have the identities $f_j(b_1, \dots, b_n) = 0$ for all j 's. Although b_i 's are not determined uniquely, their choice is ‘‘almost’’ unique in the sense that for any other

collection of elements $\{b'_i\}$ fitting in the left square of the diagram below there is $s_2 \in S$ making the the whole diagram commute:

$$(4.1) \quad \begin{array}{ccc} \{y_i\} & \xrightarrow[\{b'_i\}]{\{b_i\}} \frac{R}{s_1} \otimes K & \xrightarrow{\text{can} \otimes \text{id}} \frac{R}{s_1 s_2} \otimes K \\ \downarrow & \searrow & \swarrow \\ K & \xrightarrow{(g \otimes \text{id}) \circ \text{Coconj}} S^{-1} R \otimes K. & \end{array}$$

Now we are ready to construct a premorphism θ representing the “conjugation by g ” automorphism of pro-sets

$$(4.2) \quad g(-): G^{(\infty)}(\Phi, R) \rightarrow G^{(\infty)}(\Phi, R).$$

By definition, this amounts to constructing a collection of maps

$$\theta_s : \mathbf{Rng}(K, R^{\theta^*(s)}) \rightarrow \mathbf{Rng}(K, R^{(s)})$$

for some function $\theta^* : S \rightarrow S$. Set $\theta^*(s) = s s_1$.

Observe that for every $s, s' \in S$ the mapping $\frac{r}{s} \otimes r'(s s') \mapsto (r r')^{(s')}$ defines an isomorphism $\frac{R}{s} \otimes_R R^{(s s')} \cong R^{(s')}$ of R -modules. In particular, the R -module $\frac{R}{s} \otimes_R R^{(s s')}$ is equipped with a natural structure of an rng.

Let h be an element of $\mathbf{Rng}(K, R^{(s s_1)})$. Notice that the images of b_i under $\text{id} \otimes h$ satisfy all f_j 's. Now the element $\theta_s(h)$ can be constructed as the composite of Coconj and the unique rng homomorphism $K \rightarrow R^{(s)}$ making the following diagram commute:

$$\begin{array}{ccc} \{y_i\} & \xrightarrow{\{b_i\}} \frac{R}{s_1} \otimes K & \\ \downarrow & & \downarrow \text{id} \otimes h \\ K & \xrightarrow[\theta_s(h)]{\quad} \frac{R}{s_1} \otimes_R R^{(s s_1)} & \equiv R^{(s)}. \end{array}$$

It is clear that the morphism (4.2) defined by θ is independent of the choices of s_0 and s_1 . The diagram (4.1) also shows that it is also independent of the choice of b_i 's.

Now let us check that $g(-)$ is a morphism of pro-groups. By a standard result on Hopf algebras and using the fact that K is flat, we may choose the generators y_i of K in such a way that $Y = \bigoplus_i \mathbb{Z} y_i \leq K$ is a finite free submodule and $\Delta(Y) \leq Y \widehat{\otimes} Y$. Consequently, there is a well-defined group homomorphism $Y \rightarrow \frac{R}{s_1} \otimes K$ given by $y_i \mapsto b_i$. From the identity $g(xy) = g_x g_y$ reformulated in

the language of Hopf algebras we conclude that for sufficiently large $s_2 \in S$ the top pentagon in the following diagram commutes:

$$\begin{array}{ccc}
 & Y & \\
 \Delta \swarrow & & \searrow \\
 Y \widehat{\otimes} Y & & \frac{R}{s_1} \otimes K \\
 \downarrow & & \downarrow \text{can} \otimes \Delta \\
 (\frac{R}{s_1} \otimes K)^{\widehat{\otimes} 2} & \xrightarrow{v} & \frac{R}{s_1^2 s_2} \otimes K^{\widehat{\otimes} 2} \\
 \downarrow (\text{id} \otimes h_1) \widehat{\otimes} (\text{id} \otimes h_2) & & \downarrow \text{id} \otimes (h_1 \widehat{\otimes} h_2) \\
 (R^{(ss_1 s_2)})^{\widehat{\otimes} 2} & & \frac{R}{s_1^2 s_2} \otimes (R^{(ss_1^2 s_2)})^{\widehat{\otimes} 2} \\
 \downarrow m & & \downarrow \text{id} \otimes m \\
 R^{(ss_1 s_2)} & \longrightarrow & R^{(s)} \longleftarrow \frac{R}{s_1^2 s_2} \otimes R^{(ss_1^2 s_2)}.
 \end{array}
 \tag{4.3}$$

The homomorphism v in the above diagram is induced by canonical homomorphisms

$$\frac{R}{s_1} \rightarrow \frac{R}{s_1^2 s_2} \quad \text{and} \quad \frac{R}{s_1} \otimes \frac{R}{s_1} \rightarrow \frac{R}{s_1^2 s_2}.$$

It is clear that for any $h_1, h_2: K \rightarrow R^{(ss_1^2 s_2)}$ the bottom square of the above diagram also commutes. The commutativity of the whole diagram implies that the images of $\theta_{ss_1 s_2}(h_1 h_2)$ and $\theta_{ss_1 s_2}(h_1) \theta_{ss_1 s_2}(h_2)$ in $G_{\text{sc}}(\Phi, R^{(s)})$ coincide. Consequently, the morphism (4.2) is a morphism of a group of objects in $\text{Pro}(\mathbf{Set})$ and hence a morphism of pro-groups (cf. the beginning of Section 2.2).

It is not hard to check that $1(-)$ is the identity automorphism of $\text{id}_{G^{(\infty)}}(\Phi, R)$ and that for all $g, g' \in G_{\text{sc}}(\Phi, S^{-1}R)$ one has

$$gg'(-) = g(g'(-)).$$

4.2. THE CASE OF STEINBERG GROUPS. Clearly, the set $\text{Pro}(\mathbf{Grp})(R^{(\infty)})$ of endomorphisms of the additive pro-group of the pro-rng $R^{(\infty)}$ can be endowed with the structure of an associative ring: the addition is pointwise, while the multiplication is the usual composition of endomorphisms.

For every class of fractions $[\frac{r}{s}] \in S^{-1}R$ there is a well-defined pro-group endomorphism $m_{[\frac{r}{s}]}: R^{(\infty)} \rightarrow R^{(\infty)}$ given by

$$m_{[\frac{r}{s}]}^*(s') = ss' \quad \text{and} \quad m_{[\frac{r}{s}]}^{(s')} (a^{(ss')}) = (ra)^{(s')}.$$

It is easy to see that $m_{[\frac{r}{s}]}$ does not depend on the choice of the representative $\frac{r}{s}$. Moreover, it is easy to see that m is actually a homomorphism of rings $S^{-1}R \rightarrow \text{Pro}(\mathbf{Grp})(R^{(\infty)})$ and that

$$(4.4) \quad m_{[\frac{r}{s}]}(a^{(\infty)}b^{(\infty)}) = m_{[\frac{r}{s}]}(a^{(\infty)})b^{(\infty)} = a^{(\infty)}m_{[\frac{r}{s}]}(b^{(\infty)}).$$

For brevity we write $\frac{r}{s}a^{(\infty)}$ instead of $m_{[\frac{r}{s}]}(a^{(\infty)})$.

Our next step is to construct the action of the Steinberg group $\text{St}(\Phi, S^{-1}R)$ on the Steinberg pro-group $\text{St}^{(\infty)}(\Phi, R)$.

Definition 4.1: Let R be a ring, $S \subseteq R$ be a multiplicative subset, and Φ be a root system of rank ≥ 3 different from B_ℓ and C_ℓ .

For $u \in S^{-1}R$ and a root subgroup morphism x_β of $\text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R)$ we define the morphism

$$x_\alpha(u)x_\beta: R^{(\infty)} \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R)$$

via one of the following identities:

$$(4.5) \quad x_\alpha(u)x_\beta(b^{(\infty)}) := x_\beta(b^{(\infty)}) \quad \text{if } \alpha + \beta \notin \Phi \cup \{0\};$$

$$(4.6) \quad x_\alpha(u)x_\beta(b^{(\infty)}) := x_{\alpha+\beta}(N_{\alpha,\beta}ub^{(\infty)})x_\beta(b^{(\infty)})$$

if $\alpha + \beta \in \Phi$ but $\alpha + 2\beta, 2\alpha + \beta \notin \Phi$;

$$(4.7) \quad x_\alpha(u)x_\beta(b^{(\infty)}) := x_{\alpha+\beta}(N_{\alpha,\beta}ub^{(\infty)})x_{2\alpha+\beta}(N_{\alpha,\beta}^{2,1}u^2b^{(\infty)})x_\beta(b^{(\infty)})$$

if $\alpha + \beta, 2\alpha + \beta \in \Phi$;

$$(4.8) \quad x_\alpha(u)x_\beta(b^{(\infty)}) := x_{\alpha+\beta}(N_{\alpha,\beta}ub^{(\infty)})x_{\alpha+2\beta}(N_{\alpha,\beta}^{1,2}ub^{(\infty)^2})x_\beta(b^{(\infty)})$$

if $\alpha + \beta, \alpha + 2\beta \in \Phi$.

LEMMA 4.2: *The morphisms $x_\alpha(u)x_\beta(b^{(\infty)})$ defined above are pro-group morphisms. They give rise to an endomorphism*

$$x_\alpha(u)(-): \text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R) \rightarrow \text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R).$$

Proof. By Lemma 3.4 we need to verify the identities

$$(4.9) \quad x_\alpha(u)x_\beta(b^{(\infty)}) \cdot x_\alpha(u)x_\beta(c^{(\infty)}) = x_\alpha(u)x_\beta(b^{(\infty)} + c^{(\infty)}),$$

$$(4.10) \quad [x_\alpha(u)x_\beta(b^{(\infty)}), x_\alpha(u)x_\gamma(c^{(\infty)})] = \prod_{\substack{i\beta+j\gamma \in \Phi \\ i,j > 0}} x_\alpha(u)x_{i\beta+j\gamma}(N_{\beta,\gamma}^{i,j}b^{(\infty)^i}c^{(\infty)^j}),$$

in which $\beta, \gamma \in \Phi \setminus \{-\alpha\}$, $\beta \neq -\gamma$, $-\alpha \notin (\mathbb{Z}_{>0}\beta + \mathbb{Z}_{>0}\gamma) \cap \Phi$ and, moreover, $\beta + \gamma \neq -2\alpha$. A direct consideration of root systems A_2 and B_2 shows that under these assumptions the root subset $\Sigma = \Phi \cap (\mathbb{Z}_{\geq 0}\beta + \mathbb{Z}_{\geq 0}\gamma + \mathbb{Z}_{\geq 0}\alpha)$ is special (in particular, it does not contain $-\alpha$).

Set $T = \mathbb{Z}[b, c, u]$, $I := \langle b, c \rangle \trianglelefteq T$. Denote by $\mathcal{X}_{\Sigma, I}$ the set

$$\{x_\delta(d) \mid d \in I, \delta \in \Sigma\}$$

of defining generators of $U(\Sigma, I)$ and by $\mathcal{R}_{\Sigma, I}$ the normal subgroup of the free group $F := F(\mathcal{X}_{\Sigma, I})$ generated by the relations

$$R_\beta(b, c) = x_\beta(b)x_\beta(c)x_\beta(b+c)^{-1}$$

of type (R1) and relations $R_{\beta,\gamma}(b, c)$ of type (R2)–(R4) with $\beta, \gamma \in \Sigma$. By Remark 2.13 the group $U(\Sigma, I)$ is isomorphic to $F(\mathcal{X}_{\Sigma, I})/\mathcal{R}_{\Sigma, I}$.

Notice that there is a well-defined conjugation action of $U(\Sigma, T)$ on $U(\Sigma, I)$. The action of $x_\alpha(u)$ on each $x_\beta(b) \in \mathcal{X}_{\Sigma, I}$ is given precisely by formulas (4.5)–(4.8) (with the superscript (∞) removed). The fact that this endomorphism of F defines an endomorphism of $U(\Sigma, I)$ implies that for any Steinberg relation R (which is either $R_\beta(b, c)$ or $R_{\beta,\gamma}(b, c)$) the word $x_\alpha(u)R$ can be rewritten in F as a product $\prod_i R_i^{g_i}$, where R_i is either $R_{\beta_i}(f_i^1(b, c, u), f_i^2(b, c, u))$ or $R_{\beta_i, \gamma_i}(f_i^1(b, c, u), f_i^2(b, c, u))$, $f_i^1, f_i^2 \in I$, $\beta_i, \gamma_i \in \Sigma$ and $g_i \in F$.

Notice that a polynomial $f(b, c, u) \in I$ and $u_0 = [\frac{r}{s}] \in S^{-1}R$ give rise to a morphism $R^{(\infty)} \times R^{(\infty)} \rightarrow R^{(\infty)}$ of pro-sets. Indeed, this morphism is the result of reading the expression $f(b^{(\infty)}, c^{(\infty)}, u_0)$ according to Convention 2.7. It can be presented as a certain composition of morphisms Δ , $m_{[\frac{r}{s}]}$, m , $+$ and morphisms of component reordering, cf. Example 2.3. It follows from the previous paragraph that by inserting and deleting trivial subexpressions of the form $x_\delta(f)x_\delta(f)^{-1}$, $f = f(b^{(\infty)}, c^{(\infty)}, u_0)$ we can rewrite either of the equations (4.9)–(4.10) into a product of conjugates of words $R_{\beta_i}(f_i^1(b^{(\infty)}, c^{(\infty)}, u_0), f_i^2(b^{(\infty)}, c^{(\infty)}, u_0))$ or $R_{\beta_i, \gamma_i}(f_i^1(b^{(\infty)}, c^{(\infty)}, u_0), f_i^2(b^{(\infty)}, c^{(\infty)}, u_0))$. The assertion of the lemma now follows from the fact that the root subgroup morphisms of $\text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R)$ obviously satisfy all the relations $R_\beta, R_{\beta,\gamma}$ for $\beta, \gamma \in \Sigma$. ■

PROPOSITION 4.3: *Let R and Φ be as in the above definition. The identities (4.5)–(4.8) specify a unique well-defined conjugation action of the group $\text{St}(\Phi, S^{-1}R)$ on the pro-group $\text{St}^{(\infty)}(\Phi, R)$. The morphism*

$$\text{st}: \text{St}^{(\infty)}(\Phi, R) \rightarrow \text{G}^{(\infty)}(\Phi, R)$$

is equivariant with respect to this action.

Proof. By Theorem 1 the pro-group $\text{St}^{(\infty)}(\Phi \setminus \{-\alpha\}, R)$ is isomorphic to $\text{St}^{(\infty)}(\Phi, R)$, therefore from Lemma 4.2 we obtain an endomorphism of $\text{St}^{(\infty)}(\Phi, R)$, for which we use the same notation $x_\alpha(u)(-)$. It is easy to see that the mapping

$$u \mapsto x_\alpha(u)(-),$$

in fact, specifies a group homomorphism $S^{-1}R \rightarrow \text{Aut}(\text{St}^{(\infty)}(\Phi, R))$.

The next step of the proof is to verify that the constructed action satisfies the Steinberg relations (R2)–(R4) defining the group $\text{St}(\Phi, S^{-1}R)$. Let $R_{\alpha,\beta}(a, b)$ be one of these relations (as in the proof of the above lemma we interpret it as an element of the free group on generators $x_\alpha(a)$). Denote by Φ_0 the root subsystem of Φ generated by α and β and let γ be a root of $\Phi \setminus \Phi_0$. Since the root subset

$$\Sigma = (\mathbb{Z}_{\geq 0}\alpha + \mathbb{Z}_{\geq 0}\beta + \mathbb{Z}_{\geq 0}\gamma) \cap \Phi$$

is special, we can use an argument similar to the proof of Lemma 4.2 in order to show that the result of conjugation of x_γ with $R_{\alpha,\beta}(a, b)$ coincides with x_γ . By Lemma 3.2 this implies that the automorphism of conjugation with $R_{\alpha,\beta}(a, b)$ coincides with the trivial automorphism of $\text{St}^{(\infty)}(\Phi, R)$.

Now let us verify the equivariance. It is enough to show that

$$(4.11) \quad \text{st}(x_\alpha(u)x_\beta(b^{(\infty)})) = {}^{t_\alpha(u)}\text{st}(x_\beta(b^{(\infty)})), \quad \text{for all } \alpha, \beta \in \Phi, \beta \neq -\alpha.$$

We will verify the equality in the case when $\alpha + \beta \in \Phi$ but $\alpha + 2\beta, 2\alpha + \beta \notin \Phi$, the proof in the other cases being similar. Consider the elementary root unipotent $t_\beta(-): \mathbb{G}_a \rightarrow \text{G}_{\text{sc}}(\Phi, -)$ as a morphism of algebraic groups. Denote by $\text{Cot}_\beta: K \rightarrow X\mathbb{Z}[X]$ the corresponding homomorphism of augmentation ideals of Hopf algebras (as in Section 4.1 we abbreviate $\text{Ker}(\varepsilon)$ to K). Fix $u \in S^{-1}R$. Since root unipotents t_α satisfy (R3) the outer square of the following diagram

commutes:

$$\begin{array}{ccccc}
 K & \xrightarrow{(t_\alpha(u) \otimes \text{id}) \text{Coconj}} & S^{-1}R \otimes K & \xrightarrow{\text{id} \otimes \text{Cot}_\beta} & S^{-1}R \otimes X\mathbb{Z}[X] \\
 & \swarrow & \uparrow & \nearrow & \uparrow \\
 & & Y & \xrightarrow{\{b_i\}} & \frac{R}{s_1} \otimes K \xrightarrow{\text{can} \otimes \text{Cot}_\beta} \frac{R}{s_1^2 s_2} \otimes X\mathbb{Z}[X] \\
 & & \downarrow & & \uparrow v \\
 & & Y \widehat{\otimes} Y & \dashrightarrow & (\frac{R}{s_1} \otimes X\mathbb{Z}[X])^{\widehat{\otimes} 2} \longrightarrow (S^{-1}R \otimes X\mathbb{Z}[X])^{\widehat{\otimes} 2} \\
 & & & & \uparrow \begin{pmatrix} X_1 \mapsto 1 \otimes X \\ X_2 \mapsto N_{\alpha, \beta} u \otimes X \end{pmatrix} \\
 K \widehat{\otimes} K & \xrightarrow{\text{Cot}_\beta \widehat{\otimes} \text{Cot}_{\alpha+\beta}} & & & X_1\mathbb{Z}[X_1] \widehat{\otimes} X_2\mathbb{Z}[X_2].
 \end{array}$$

Similarly, to the construction of (4.3) we can choose a free abelian subgroup $Y \leq K$ of finite rank generating K multiplicatively so that $\Delta(Y) \leq Y \widehat{\otimes} Y$. Further, following the procedure described in (4.1) we can choose s_1 and $\{b_i\}$ in such a manner that the top left square of the above diagram commutes. Also, there is a dashed arrow making the bottom pentagon commute. Finally, there exists sufficiently large s_2 such that the central square of the diagram commutes. This commutativity implies (4.11) (the top path gives the right-hand side while the bottom one gives the left-hand side; the homomorphism v has the same meaning as in (4.3)). Thus, by Lemma 2.16 the morphism $\text{st} : \text{St}^{(\infty)}(\Phi, S^{-1}R) \rightarrow \text{G}^{(\infty)}(\Phi, R)$ is $\text{St}(\Phi, S^{-1}R)$ -equivariant. ■

THEOREM 2: *Let R be a ring, $M \leq R$ be a maximal ideal and Φ be a root system of rank ≥ 3 different from B_ℓ and C_ℓ . Set $S = R \setminus M$. Then the action of $\text{St}(\Phi, R_M)$ defined in Proposition 4.3 gives rise to an action of the group $\text{G}_{\text{sc}}(\Phi, R_M)$ on $\text{St}^{(\infty)}(\Phi, R)$. The morphism $\text{st} : \text{St}^{(\infty)}(\Phi, R) \rightarrow \text{G}^{(\infty)}(\Phi, R)$ is $\text{G}_{\text{sc}}(\Phi, R_M)$ -equivariant.*

Proof. Recall that for $a, b \in R_M^\times$ and $\alpha \in \Phi$ one can define the following elements of $\text{St}(\Phi, R_M)$:

$$\begin{aligned}
 w_\alpha(a) &= x_\alpha(a) \cdot x_{-\alpha}(-a^{-1}) \cdot x_\alpha(a), \\
 h_\alpha(a) &= w_\alpha(a) \cdot w_\alpha(1)^{-1}, \\
 \{a, b\}_\alpha &= h_\alpha(ab) \cdot h_\alpha^{-1}(a) \cdot h_\alpha^{-1}(b).
 \end{aligned}$$

Since R_M is local, the group $\text{G}_{\text{sc}}(\Phi, R_M)$ coincides with its elementary subgroup $\text{E}_{\text{sc}}(\Phi, R)$. Consequently, by [1, Proposition 1.6] and [22, Theorem 2.13] the group $\text{G}_{\text{sc}}(\Phi, R_M)$ is the quotient of $\text{St}(\Phi, R_M)$ by the relations $\{a, b\}_\alpha = 1$

for $a, b \in R_M^\times$ and some fixed long root $\alpha \in \Phi$. Thus, it remains to check that the Steinberg symbols $\{a, b\}_\alpha$ act trivially on $\text{St}^{(\infty)}(\Phi, R)$, which, in turn, follows from the relations

$$(4.12) \quad h_\alpha(a)x_\beta(b^{(\infty)}) = x_\beta(a^{(\beta, \alpha)}b^{(\infty)}), \quad \beta \neq \pm\alpha.$$

To prove (4.12) take a root $\beta \in \Phi$ linearly independent with α . Direct computation using Definition 4.1 shows that

$$h_\alpha(a)x_\beta(b^{(\infty)}) = \prod_{\substack{i\alpha+j\beta \in \Phi \\ j > 0}} x_{i\alpha+j\beta}(P_{i,j}(a) \cdot b^{(\infty)^j})$$

for some Laurent polynomials $P_{i,j}(t) \in \mathbb{Z}[t, t^{-1}]$ depending only on α and β and the chosen order of factors. Since in the Steinberg group $\text{St}(\Phi, \mathbb{Z}[t, t^{-1}])$ we have a similar identity

$$h_\alpha(t)x_\beta(1) = \prod_{\substack{i\alpha+j\beta \in \Phi \\ j > 0}} x_{i\alpha+j\beta}(P_{i,j}(t)) = x_\beta(t^{(\beta, \alpha)}),$$

we conclude by Remark 2.13 that

$$P_{i,j}(t) = 0$$

with the sole exception of $P_{0,1}(t) = t^{(\beta, \alpha)}$. Thus, the proof of (4.12) is complete. ■

5. Proof of the main result

5.1. THE CONSTRUCTION OF A CROSSED MODULE. In this section we prove the main results of the paper, namely we construct a crossed module structure on the homomorphism st and prove the centrality of K_2 . Let us, first, briefly recall the definition of a crossed module.

Definition 5.1: Let N be a group acting on itself by left conjugation. A group homomorphism $\varphi: M \rightarrow N$ is called a **crossed module** if one can define the action of the group N on M in such a way that φ preserves the action of N and, moreover, the identity $\varphi^{(m)}m' = mm'm^{-1}$, called the **Peiffer identity**, holds for all $m, m' \in M$.

It is an easy exercise to check that the kernel of φ is always a central subgroup of M , while the image of φ is normal in N .

For the rest of this section R is an arbitrary commutative ring and M is a maximal ideal of R . As before, Φ is a root system of rank ≥ 3 different from B_ℓ and C_ℓ .

We denote by S the multiplicative system associated to M , i.e., $S = R \setminus M$. By Theorem 2 the group $G_{\text{sc}}(\Phi, R_M)$ acts on both $\text{St}^{(\infty)}(\Phi, R)$ and $G^{(\infty)}(\Phi, R)$ by conjugation and this action is preserved by the morphism st . Consequently, $G_{\text{sc}}(\Phi, R)$ acts on these pro-groups its images $G_{\text{sc}}(\Phi, R_M)$. Consider the following canonical morphisms:

$$\begin{aligned} \pi_R: R^{(\infty)} &\rightarrow R, \\ \pi_{\text{St}}: \text{St}^{(\infty)}(\Phi, R) &\rightarrow \text{St}(\Phi, R), \\ \pi_G: G^{(\infty)}(\Phi, R) &\rightarrow G_{\text{sc}}(\Phi, R). \end{aligned}$$

Notice that the action of $h \in G_{\text{sc}}(\Phi, R)$ on the pro-group $G^{(\infty)}(\Phi, R)$ is compatible with the conjugation action of $G_{\text{sc}}(\Phi, R)$ on itself, i.e.,

$${}^h\pi_G(g^{(\infty)}) = \pi_G({}^hg^{(\infty)}).$$

Similarly, the conjugation action of $h \in \text{St}(\Phi, R)$ on the pro-group $\text{St}^{(\infty)}(\Phi, R)$ is compatible with the conjugation action of $\text{St}(\Phi, R)$ on itself, i.e., ${}^h\pi_{\text{St}}(g^{(\infty)}) = \pi_{\text{St}}({}^hg^{(\infty)})$. Notice also that $x_\alpha\pi_R = \pi_{\text{St}}x_\alpha$ and $\pi_G\text{st} = \text{st}\pi_{\text{St}}$.

LEMMA 5.2: *The subgroup $K_2(\Phi, R) = \text{Ker}(\text{st}: \text{St}(\Phi, R) \rightarrow G(\Phi, R))$ is contained in the center of $\text{St}(\Phi, R)$.*

Proof. Let g be an element of $K_2(\Phi, R)$ and $\alpha \in \Phi$ be a root. Consider the ideal

$$I_\alpha = \{a \in R \mid {}^gx_\alpha(ra) = x_\alpha(ra) \text{ for all } r \in R\}.$$

We need to show that $I_\alpha = R$. Let $M \trianglelefteq R$ be a maximal ideal. By Theorem 2, the element g acts trivially on the pro-group $\text{St}^{(\infty)}(\Phi, R)$ associated with $S = R \setminus M$. Consequently, we obtain the following equality of morphisms $R^{(\infty)} \rightarrow \text{St}(\Phi, R)$:

$$\begin{aligned} {}^gx_\alpha(\pi_R(a^{(\infty)})) &= {}^g\pi_{\text{St}}(x_\alpha(a^{(\infty)})) = \pi_{\text{St}}({}^gx_\alpha(a^{(\infty)})) \\ &= \pi_{\text{St}}(x_\alpha(a^{(\infty)})) = x_\alpha(\pi_R(a^{(\infty)})), \end{aligned}$$

which implies that $I_\alpha \not\subseteq M$. ■

With a similar technique, we also can re-prove Taddei's normality theorem (see [27]).

LEMMA 5.3: *The subgroup $E(\Phi, R) = \text{Im}(\text{st}: \text{St}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R))$ is normal in $G_{\text{sc}}(\Phi, R)$.*

Proof. Let g be an element of $G_{\text{sc}}(\Phi, R)$ and $\alpha \in \Phi$ be a root. Consider the ideal

$$I_\alpha = \{a \in R \mid {}^g t_\alpha(ra) \in E(\Phi, R) \text{ for all } r \in R\}.$$

Again, we need to show that $I_\alpha = R$. Let $M \trianglelefteq R$ be a maximal ideal. Since g acts on the pro-groups $\text{St}^{(\infty)}(\Phi, R)$ and $G^{(\infty)}(\Phi, R)$ associated with the subset $S = R \setminus M$ and this action is preserved by st , we obtain the following equalities of morphisms $R^{(\infty)} \rightarrow G_{\text{sc}}(\Phi, R)$:

$${}^g t_\alpha(\pi_R(a^{(\infty)})) = {}^g \pi_G(\text{st}(x_\alpha(a^{(\infty)}))) = \pi_G(\text{st}({}^g x_\alpha(a^{(\infty)}))) = \text{st}(\pi_{\text{St}}({}^g x_\alpha(a^{(\infty)}))),$$

which implies that $I_\alpha \not\subseteq M$. ■

THEOREM 3: *The homomorphism $\text{st}: \text{St}(\Phi, R) \rightarrow G_{\text{sc}}(\Phi, R)$ can be turned into a crossed module in a unique way.*

Proof. Notice that by Lemma 5.2 we already know that $K_2(\Phi, R)$ is a central subgroup of $\text{St}(\Phi, R)$. Under our assumptions the group $\text{St}(\Phi, R)$ is a perfect central extension of $E_{\text{sc}}(\Phi, R)$, therefore by [21, Lemma 1.1] for every $g \in G_{\text{sc}}(\Phi, R)$ there may exist at most one endomorphism ${}^g(-)$ of $\text{St}(\Phi, R)$ satisfying

$$(5.1) \quad g \text{st}(h)g^{-1} = \text{st}({}^g h) \quad \text{for all } h \in \text{St}(\Phi, R).$$

Thus, to construct an action of $G_{\text{sc}}(\Phi, R)$ on $\text{St}(\Phi, R)$ it suffices to construct an endomorphism ${}^g(-): \text{St}(\Phi, R) \rightarrow \text{St}(\Phi, R)$ satisfying (5.1). Also, this action would automatically satisfy the Peiffer identity.

Let α be a root of Φ . Since $E(\Phi, R) \trianglelefteq G_{\text{sc}}(\Phi, R)$ by Lemma 5.3, the set $Y_\alpha(a) = \text{st}^{-1}(gt_\alpha(a)g^{-1})$ is nonempty for all $a \in R$. Moreover, $Y_\alpha(a)$ is a coset of the subgroup $K_2(\Phi, R)$ and $Y_\alpha(a + a') = Y_\alpha(a)Y_\alpha(a')$.

First of all, let us show that $[Y_\alpha(a), Y_\beta(b)] = 1$ provided $\alpha + \beta \notin \Phi \cup \{0\}$. Let α, β be roots as above, a be an element of R and h be an element of $Y_\alpha(a)$. Set

$$I = \{b \in R \mid [h, Y_\beta(bR)] = 1\}$$

and let $M \trianglelefteq R$ be a maximal ideal. Since

$$[h, \pi_{\text{St}}({}^g x_\beta(b^{(\infty)}))] = \pi_{\text{St}}({}^{gt_\alpha(a)} x_\beta(b^{(\infty)})) \pi_{\text{St}}({}^g x_\beta(-b^{(\infty)})) = 1,$$

we obtain $I \not\subseteq M$ and, consequently, $I = R$, which proves the assertion.

Now let α be a root of Φ . By our assumptions on Φ there exist roots $\beta, \gamma \in \Phi$ of the same length such that $\alpha = \beta + \gamma$. Denote by $y_\alpha(a)$ the only element of the set $[Y_\beta(N_{\beta,\gamma}), Y_\gamma(a)]$. We define the map $g(-)$ on the set of generators of $\text{St}(\Phi, R)$ by ${}^g x_\alpha(a) = y_\alpha(a)$, and we claim that such a definition induces an endomorphism, i.e., that $y_\alpha(a)$ satisfy the Steinberg relations. Clearly, $y_\alpha(a)$ satisfy (R1). Since $y_\alpha(a) \in Y_\alpha(a)$, it is also clear that they satisfy relations (R2).

It remains to check that $y_\alpha(a)$ satisfy the Steinberg relations (R3) and (R4). Let $M \trianglelefteq R$ be a maximal ideal. Recall that $\pi_{\text{St}}({}^g x_\gamma(a^{(\infty)}))$ denotes a certain pro-group morphism $R^{(\infty)} \rightarrow \text{St}(\Phi, R)$, so $[h, \pi_{\text{St}}({}^g x_\gamma(a^{(\infty)}))]$ is a pro-set morphism $R^{(\infty)} \rightarrow \text{St}(\Phi, R)$ for a fixed $h \in Y_\beta(N_{\beta,\gamma})$. Since

$$\text{st}(\pi_{\text{St}}({}^g x_\gamma(a^{(\infty)}))) = g t_\gamma(\pi_R(a^{(\infty)}))g^{-1},$$

we have

$$\begin{aligned} (5.2) \quad y_\alpha(\pi_R(a^{(\infty)})) &= [h, \pi_{\text{St}}({}^g x_\gamma(a^{(\infty)}))] \\ &= \pi_{\text{St}}(\text{st}(h) {}^g x_\gamma(a^{(\infty)})) \pi_{\text{St}}({}^g x_\gamma(a^{(\infty)}))^{-1} \\ &= \pi_{\text{St}}({}^{gt_\beta(N_{\beta,\gamma})} x_\gamma(a^{(\infty)}) {}^g x_\gamma(-a^{(\infty)})) = \pi_{\text{St}}({}^g x_\alpha(a^{(\infty)})). \end{aligned}$$

Now let a be an element of R and α, β be roots of Φ such that $\alpha + \beta, 2\alpha + \beta \in \Phi$. Consider the ideal

$$J = \{b \in R \mid [y_\alpha(a), y_\beta(br)] = y_{\alpha+\beta}(N_{\alpha,\beta}abr)y_{2\alpha+\beta}(N_{\alpha,\beta}^2 a^2 br) \text{ for all } r \in R\}.$$

From (5.2) we obtain

$$\begin{aligned} [y_\alpha(a), y_\beta(\pi_R(b^{(\infty)}))] &= [y_\alpha(a), \pi_{\text{St}}({}^g x_\beta(b^{(\infty)}))] \\ &= \pi_{\text{St}}({}^{gt_\alpha(a)} x_\beta(b^{(\infty)}) {}^g x_\beta(-b^{(\infty)})) \\ &= \pi_{\text{St}}({}^g x_{\alpha+\beta}(N_{\alpha,\beta}ab^{(\infty)}) {}^g x_{2\alpha+\beta}(N_{\alpha,\beta}^2 a^2 b^{(\infty)})) \\ &= y_{\alpha+\beta}(N_{\alpha,\beta}a\pi_R(b^{(\infty)}))y_{2\alpha+\beta}(N_{\alpha,\beta}^2 a^2 \pi_R(b^{(\infty)})). \end{aligned}$$

It follows that $J \not\subseteq M$, i.e., $J = R$. The verification of (R3) is similar but easier. It is clear that the constructed endomorphism satisfies (5.1). \blacksquare

5.2. CONCLUDING REMARKS. It is possible to prove the centrality of K_2 without using the concept of a ring homotopy introduced in Definition 2.4. Indeed, the idea is to reduce to the case of domains using the following result inspired by [24, Lemma 5.3].

LEMMA 5.4: *Let R be a commutative ring, I be an ideal of R and Φ be a root system of rank ≥ 3 . The centrality of $K_2(\Phi, R)$ implies the centrality of $K_2(\Phi, R/I)$.*

Proof. Consider the following commutative diagram:

$$\begin{CD} \text{St}(\Phi, R) @>\rho_{\text{St}}>> \text{St}(\Phi, R/I) \\ @V\text{st}VV @VVV \\ \text{E}_{\text{sc}}(\Phi, R) @>\rho_{\text{E}}>> \text{E}_{\text{sc}}(\Phi, R/I). \end{CD}$$

We denote by $\text{St}(\Phi, R, I)$ the normal closure in $\text{St}(\Phi, R)$ of the subgroup generated by elements $x_\alpha(s)$, where $s \in I$, $\alpha \in \Phi$. We also denote by $\text{E}_{\text{sc}}(\Phi, R, I)$ the image of this subgroup under st . It is clear that $\text{E}_{\text{sc}}(\Phi, R, I)$ lies in the kernel of ρ_{E} .

Denote by H the preimage of $K_2(\Phi, R/I)$ under ρ_{St} . Also set

$$H' = [H, \text{St}(\Phi, R)].$$

The identity (2.1) shows that H' is a normal subgroup of $\text{St}(\Phi, R)$. Under our assumptions the group $\text{St}(\Phi, R)$ is perfect, therefore using Hall–Witt identity (2.3) we obtain

$$\begin{aligned} H' &= [H, \text{St}(\Phi, R)] = [H, [\text{St}(\Phi, R), \text{St}(\Phi, R)]] \\ &\subseteq [[H, \text{St}(\Phi, R)], \text{St}(\Phi, R)] = [H', \text{St}(\Phi, R)]. \end{aligned}$$

Since $\text{st}(H)$ is contained in the congruence subgroup $\text{G}_{\text{sc}}(\Phi, R, I)$ we obtain from the standard commutator formula [30, Theorem 1] the fact that

$$\text{st}(H') \subseteq \text{E}_{\text{sc}}(\Phi, R, I).$$

Thus, $H' \subseteq \text{St}(\Phi, R, I) \cdot K_2(\Phi, R)$. Finally, from our assumptions we obtain

$$H' \subseteq [H', \text{St}(\Phi, R)] \subseteq [\text{St}(\Phi, R, I) \cdot K_2(\Phi, R), \text{St}(\Phi, R)] \subseteq \text{St}(\Phi, R, I),$$

from which the assertion of the lemma follows. ■

Since every commutative ring R can be presented as a quotient of the polynomial ring over \mathbb{Z} with possibly an infinite number of variables, we may restrict ourselves to considering only integral domains throughout the paper. As noted before, in this case the homotopes $R^{(s)}$ turn into the usual principal ideals $sR \trianglelefteq R$.

References

- [1] E. Abe, *Chevalley groups over local rings*, Tohoku Mathematical Journal **21** (1969), 474–494.
- [2] D. Allcock, *Steinberg groups as amalgams*, Algebra & Number Theory **10** (2016), 1791–1843.
- [3] A. Bak, *K-theory of Forms*, Annals of Mathematics Studies, Vol. 98, Princeton University Press, Princeton, NJ, 1981.
- [4] N. Bourbaki, *Éléments de mathématique. Groupes et algèbres de Lie: Chapitres 4, 5 et 6*, Masson, 1981.
- [5] R. W. Carter, *Conjugacy classes in the Weyl group*, Compositio Mathematica **25** (1972), 1–59.
- [6] W. van der Kallen, *Another presentation for Steinberg groups*, Indagationes Mathematica **39** (1977), 304–312.
- [7] W. van der Kallen and M. Stein, *On the Schur multipliers of Steinberg and Chevalley groups over commutative rings*, Mathematische Zeitschrift **155** (1977), 83–94.
- [8] H. Kashiwara and P. Schapira, *Sheaves and Categories*, Grundlehren der mathematischen Wissenschaften, Vol. 332, Springer, Berlin, 2006.
- [9] A. Lavrenov, *On odd unitary Steinberg group*, <https://arxiv.org/abs/1303.6318>.
- [10] A. Lavrenov, *Unitary Steinberg group is centrally closed*, St. Petersburg Mathematical Journal **24** (2013), 783–794.
- [11] A. Lavrenov, *Another presentation for symplectic Steinberg groups*, Journal of Pure and Applied Algebra **219** (2015), 3755–3780.
- [12] A. Lavrenov, *A local-global principle for symplectic K_2* , Documenta Mathematica **23** (2018), 653–675.
- [13] A. Lavrenov and S. Sinchuk, *On centrality of even orthogonal K_2* , Journal of Pure and Applied Algebra **221** (2017), 1134–1145.
- [14] H. Matsumoto, *Sur les sous-groupes arithmétiques des groupes semi-simple déployés*, Annales Scientifiques de l'École Normale Supérieure **2** (1969), 1–62.
- [15] J. Milnor, *Introduction to Algebraic K-Theory*, Annals of Mathematics Studies, Vol. 72, Princeton University Press, Princeton, NJ, 1971.
- [16] V. Petrov and A. Stavrova, *Elementary subgroups of isotropic reductive groups*, St. Petersburg Mathematical Journal **20** (2009), 625–644.
- [17] V. Petrov, *Odd unitary groups*, Journal of Mathematical Sciences (New York) **130** (2005), 4752–4766.
- [18] U. Rehmann, *Präsentationen von Chevalleygruppen über $k[t]$* , <http://www.math.uni-bielefeld.de/~rehmann/Publ/praesentation.pdf>.
- [19] U. Rehmann, *Zentrale Erweiterungen der speziellen linearen Gruppe eines Schiefkörpers*, Journal für die Reine und Angewandte Mathematik **301** (1978), 77–104.
- [20] S. Sinchuk, *On centrality of K_2 for Chevalley groups of type E_ℓ* , Journal of Pure and Applied Algebra **220** (2016), 857–875.
- [21] M. R. Stein, *Generators, relations and coverings of Chevalley groups over commutative rings*, American Journal of Mathematics **93** (1971), 965–1004.
- [22] M. R. Stein, *Surjective stability in dimension 0 for K_2 and related functors*, Transactions of the American Mathematical Society **178** (1973), 165–191.

- [23] R. Steinberg, *Générateurs, relations et revêtements de groupes algébriques*, in *Colloque sur la Théorie des Groupes Algébriques*, Librairie Universitaire, Louvain, 1962, pp. 113–127.
- [24] A. V. Stepanov, *On the normal structure of the general linear group over a ring*, *Journal of Mathematical Sciences (New York)* **95** (1999), 2146–2155.
- [25] A. Stepanov, *Structure of Chevalley groups over rings via universal localization*, *Journal of Algebra* **450** (2016), 522–548.
- [26] A. Suslin, *On the structure of the special linear group over polynomial rings*, *Mathematics of the USSR - Izvestiya* **11** (1977), 221–238.
- [27] G. Taddei, *Normalité des groupes élémentaires dans les groupes de Chevalley sur un anneau*, in *Applications of Algebraic K-theory to Algebraic Geometry and Number Theory. Part II (Boulder, Colo., 1983)*, *Contemporary Mathematics*, Vol. 55, American Mathematical Society, 1986, pp. 693–710.
- [28] G. Tang, *Hermitian groups and K-theory*, *K-Theory* **13** (1998), 209–267.
- [29] M. Tulenbaev, *Schur multiplier of a group of elementary matrices of finite order*, *Journal of Soviet Mathematics* **17** (1981), 2062–2067.
- [30] L. Vaserstein, *On normal subgroups of Chevalley groups over commutative rings*, *Tohoku Mathematical Journal* **38** (1986), 219–230.
- [31] N. Vavilov and E. Plotkin, *Chevalley groups over commutative rings: I. Elementary calculations*, *Acta Applicandae Mathematica* **45** (1996), 73–113.
- [32] E. Voronetsky, *Centrality of K_2 -functors revisited*, *Journal of Pure and Applied Algebra* **225** (2021), Article no. 106547.
- [33] E. Voronetsky, *Centrality of odd unitary K_2 -functor*, <https://arxiv.org/abs/2005.02926>.
- [34] C. A. Weibel, *The k-book: An Introduction to Algebraic K-theory*, *Graduate Texts in Mathematics*, Vol. 145, American Mathematical Society, providence, RI, 2013.
- [35] M. Wendt, *On homotopy invariance for homology of rank two groups*, *Journal of Pure and Applied Algebra* **216** (2012), 2291–2301.