

# The Roles of Cigarette Smoking and the Lung in the Transitions Between Phases of Preclinical Rheumatoid Arthritis

Jeffrey A. Sparks<sup>1</sup> · Elizabeth W. Karlson<sup>1</sup>

Published online: 7 March 2016  
© Springer Science+Business Media New York 2016

**Abstract** While the etiology of rheumatoid arthritis (RA) remains to be fully elucidated, recent research has advanced the understanding of RA pathogenesis to the point where clinical trials for RA prevention are underway. The current paradigm for RA pathogenesis is that individuals progress through distinct preclinical phases prior to the onset of clinically apparent RA. These preclinical RA phases consist of genetic risk, local inflammation, presence of RA-related autoantibodies, asymptomatic systemic inflammation, and early non-specific symptoms prior to clinical seropositive RA. Epidemiologic studies have been important in forming hypotheses related to the biology occurring in preclinical RA. Specifically, studies associating cigarette smoking with overall RA risk as well as transitions between phases of preclinical RA were vital in helping to establish the lung as a potential important initiating site in the pathogenesis of seropositive RA. Herein, we review the epidemiology associating smoking with transitions in preclinical phases of RA as well as the recent literature supporting the lung as a critical site in RA pathogenesis.

**Keywords** Rheumatoid arthritis (RA) · Lung · Pulmonary · Smoking · Anti-citrullinated protein antibodies (ACPA) · Pre-RA

---

This article is part of the Topical Collection on *Rheumatoid Arthritis*

✉ Jeffrey A. Sparks  
jasparks@partners.org

<sup>1</sup> Department of Medicine, Division of Rheumatology, Immunology and Allergy, Brigham and Women's Hospital and Harvard Medical School, 75 Francis Street, Boston, MA 02115, USA

## Introduction

Rheumatoid arthritis (RA) causes an inflammatory polyarthritis affecting 1 % of the population and is characterized by a chronic erosive arthropathy typically of the small joints [1, 2]. RA is a serious public health problem that results in excess disability, morbidity, and mortality [3•, 4]. While the etiology of RA is currently not established, genetics, lifestyle factors, and biomarkers have been associated with risk of RA development in many research investigations over the past few decades [5•]. The current paradigm for RA pathogenesis is that individuals with increased genetic susceptibility progress through preclinical phases prior to clinically apparent RA [6•, 7]. These preclinical RA phases include genetic risk, immune activation without symptoms (characterized by upregulation of cytokines and autoantibody production), arthralgias and other non-specific symptoms, and inflammatory arthritis prior to the development of classifiable RA [8]. Epidemiologic investigations have been helpful to guide basic and translational studies that aim to elucidate the biology of these pre-clinical RA transition phases.

Local loss of immune tolerance may trigger an inflammatory milieu and dysregulation of the adaptive immune system, causing systemic inflammation that eventually manifests as RA. Several anatomic sites have been posited as possible initiating sites for RA. Mucosal surfaces, in particular, are speculated to be the local sites of initial innate immune system dysregulation [9]. The anatomic sites that may be important in the development of RA include the lung, respiratory tract, periodontum, gut, bladder, and reproductive tract [9]. However, the lung and respiratory tract have the most evidence in support of being an initiating site for RA [10•].

While much remains to be elucidated concerning the biologic mechanisms of RA pathogenesis, RA prevention strategies are actively being pursued. Epidemiologic studies have

identified factors, such as genetics, family history, environment, and biomarkers that allow investigators to target those at highest risk for RA prevention [11•, 12••]. The association of RA with potentially modifiable factors, in particular smoking and obesity, suggests that lifestyle interventions such as smoking cessation or weight loss may prevent or delay RA onset [13•, 14–16]. Since many effective targeted and non-targeted drugs for RA now exist, several clinical trials using pharmacologic interventions for RA prevention are currently underway among those at especially high risk for RA due to seropositivity or early symptoms [13•]. Understanding the pathogenesis of RA remains crucial to the success, interpretation, and implementation of these ongoing as well as future RA prevention studies.

The aim of this article is to review the evidence supporting the lung and respiratory tract as an initiating site for RA pathogenesis, with a focus on how the epidemiology of smoking has informed investigations into the biology of pre-clinical seropositive RA pathogenesis in the lung.


### Cigarette Smoking, Cessation, and Overall Risk for Seropositive RA

Cigarette smoking is the best established lifestyle risk factor for RA and may provide the external trigger for those asymptomatic but at increased genetic risk for RA [17–24]. There is evidence that smoking is important throughout the preclinical phases of RA development (see Fig. 1). The association of smoking is particularly strong for seropositive RA (defined as either rheumatoid factor [RF] or anti-cyclic citrullinated peptide [CCP] positivity). Smoking has a clear association with RA risk and may contribute up to 35 % of the attributable

risk for seropositive RA [25, 26•]. In a large meta-analysis that included 11 studies, 13,885 RA cases among a total of 593,576 individuals, current smokers had an odds ratio (OR) of 1.64 for seropositive RA compared to never smokers [17]. This effect of smoking for increased RA risk was more potent in men compared to women [17]. Current male smokers had an OR of 3.91 for seropositive RA compared to male non-smokers [17]. Both smoking status and intensity are associated with increased RA risk with clinically significant risk most apparent for those with >10 pack-years [27, 28••].

Studies suggest that smoking cessation may decrease RA risk [24, 27, 29••]. Among women followed prospectively for 26 years in the Nurses' Health Study, risk of RA for past smokers was similar to the RA risk of non-smokers  $\geq 20$  years after smoking cessation [27]. Similarly, in the Swedish Epidemiological Investigations of RA that compared incident RA cases to population-based controls, former smokers had similar RA risk compared to never smokers  $\geq 20$  years after cessation [24]. In the prospective Swedish Mammography Cohort, risk of RA was still significantly elevated 15 years after smoking cessation compared to never smokers [29••]. While the risk for RA did not return to the baseline RA risk of never smokers in this study, risk steadily decreased over increasing duration of smoking cessation [29••].

While smoking is clearly associated with seropositive RA, this association is attenuated or null for seronegative RA [17, 28••, 30]. Some studies suggest that seronegative RA may be a heterogeneous collection of diseases without a common pathogenesis or distinct risk factors [31]. A recent study aimed to define a homogeneous population of seronegative RA patients by removing patients with detectable RA-related autoantibodies on sensitive research-only assays for anti-citrullinated protein antibodies (ACPA) as well as patients with *HLA-B27*



**Fig. 1** Schematic of proposed biologic mechanisms linking smoking and the lung to preclinical phases in the pathogenesis of anti-citrullinated protein antibody (ACPA)-positive rheumatoid arthritis (RA). An asymptomatic individual with high genetic risk, such as the *HLA-DRB1* shared epitope, is exposed to environmental triggers such as cigarette smoking. This induces mucosal inflammation in the airways and alveoli of the lung. Activated enzymes, including peptidylarginine deiminase (PAD), stimulate citrullination of proteins at these sites to form

neoantigens. Antigen presenting cells (APCs) in the innate immune system present these neoantigens through the *HLA-DRB1* protein to T cells that results in adaptive immune system dysregulation and leads to systemic inflammation. Stimulated B cells in the respiratory mucosa produce ACPA. Systemic inflammation leads to non-specific symptoms such as fatigue and arthralgias. Autoantibodies exhibit specificity to peripheral joints eventually leading to synovitis and the clinical diagnosis of ACPA-positive RA

where spondyloarthropathy may have masqueraded as seronegative RA [32••]. Using these methods, distinct genetic differences were detected between patients with seropositive and seronegative RA implying separate pathogenesis for each disease phenotype. However, most studies rely on the clinical classification of seronegative RA which has inherent clinical heterogeneity. For this reason, the pathogenesis of seropositive RA, particularly ACPA-positive RA, is best developed and is therefore the focus of this article. We refer to ACPA as the general presence of any research or commercial assay for anti-citrullinated protein antibodies, while we refer to CCP (and particular assays, such as CCP2, CCP3, or CCP3.1) as the specific commercial assay.

### Cigarette Smoking and Transitions Between Preclinical Phases of RA

Smoking has been associated with increased risk for progression among preclinical RA transition states (Table 1). Among patients with arthralgias and detectable serum IgM-RF or CCP2, smoking may hasten the emergence of classifiable RA [37••]. However, when evaluating all patients with arthralgias, there is no clear association of smoking with progression to detectable inflammatory arthritis on physical examination [38•]. Among first-degree relatives without RA in the Studies for the Etiology of RA, smoking >10 pack-years, in addition to current smoking status, was associated with twofold increased risk of developing inflammatory joint signs [36•]. Most of these unaffected relatives were asymptomatic and further away from RA development compared to those with seropositive arthralgias, so the particular timing of smoking in relation to preclinical RA stages is likely important.

Smoking may also be important in the development of RA-related autoantibodies. Patients with chronic obstructive pulmonary disease (COPD) without RA have increased levels of CCP and RF compared to controls. Smoking is associated with development of RA-related autoantibodies (CCP, ACPA, and RF) in patients that later develop RA [39•, 40] and in those at high genetic risk [33]. A Swedish study analyzed banked samples prior to the onset of RA patients and examined the *HLA* shared epitope, CCP, particular ACPA targeting specific antigens, and smoking. Smoking was associated with presence of ACPA antibodies closer to disease onset. Ever smokers with the shared epitope had a fourfold increased odds of being CCP2 positive as well as other ACPA within 10.5 years of RA diagnosis compared to never smokers without the shared epitope [34••]. Since the presence of circulating RA-related autoantibodies is associated with very elevated risk for RA, smoking may propagate the generation of RF, CCP, and other ACPA [41, 42••].

Finally, other inhaled irritants such as silica, air pollution, and textile dust may also increase the risk for RA, particularly

seropositive RA [43–46, 47•]. While the epidemiology of these environmental factors is less developed compared to smoking, the associations of these other environmental factors emphasize that the lung, and potentially other mucosal sites, may provide the common link and micro-environment important for a variety of biologic processes that ultimately present clinically as RA.

### *HLA*–Smoking Statistical Interaction Provides Clues to the Biology of Transitions of Preclinical RA

The *HLA* shared epitope, present in the Major Histocompatibility Complex Class II on chromosome 6, is the strongest genetic risk factor for RA, among many other known genetic risk factors [48, 49, 50••]. Shared epitope alleles are the site of transcription for specific amino acid haplotypes in the HLA-DRβ1 protein that are important in antigen processing and presentation and thus important in immune function [51]. Many epidemiologic studies have detected a statistical interaction between the shared epitope and smoking [52, 53, 54•, 55, 56, 57•, 58•, 59]. This finding provided background for the hypothesis that smoking may induce autoantigens interacting with the innate system that stimulates the adaptive immune system and eventually leads to joint specificity through molecular mimicry of autoantibodies and, finally, clinical symptoms of RA [60–65].

Specific amino acid positions within the peptide binding groove of the HLA-DRβ1 protein are associated with especially high risk of seropositive RA [51]. These sites also have statistical interactions with cigarette smoking for risk of seropositive, especially ACPA-positive, RA [52, 53, 54•, 55, 56, 57•, 58•]. In a trans-ethnic study using data from the Nurses' Health Studies in the US, the Swedish Epidemiological Investigations of RA, and the Korean RA Cohort Study, heavy smoking (defined as >10 pack-years) interacted with HLA-DRβ1 positions 11 and 13 for increased risk for seropositive RA [35••]. Specific amino acid haplotypes conferred increased risk of seropositive RA (RF or CCP in the Nurses' Health Study; CCP in both the Swedish and Korean studies) [35••]. For example, presence of valine at HLA-DRβ1 position 11 or histidine at position 13 significantly increased seropositive RA risk while presence of serine at HLA-DRβ1 positions 11 or 13 significantly decreased RA risk [35••]. These results suggest a physical interaction of a neoantigen, specifically induced by smoking, with the HLA-DRβ1 protein. However, the neoantigen(s) produced by this process are yet to be definitively identified.

### Alveolar and Airway Inflammation in Preclinical RA

While pulmonary involvement of patients with RA is well described, particularly interstitial lung disease (ILD) as an

**Table 1** Selected epidemiologic studies investigating smoking in RA preclinical phase transitions

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
Voigt, 1994 [22]	Overall RA risk	King County, Washington US women N = 349 RA cases and 1457 controls	Case-control	Current smoking: OR 1.33 (95 % CI 1.00–1.77) Ever smoking: OR 1.31 (95 % CI 1.07–1.60) Compared to non-smoking	-One of several early studies suggesting a link between smoking and RA risk
Criswell, 2002 [18]	Overall RA risk	Iowa Women's Health Study US women N = 31,336	Prospective cohort	Current smoking: RR 2.0 (95 % CI 1.3–2.9) for RA Quit smoking $\leq 10$ years: RR 1.8 (95 % CI 1.1–3.1) for RA Quit smoking $> 10$ years: RR 0.9 (95 % CI 0.5–2.6) for RA Compared to non-smoking	-158 RA cases developed during 11 years of follow-up -Results suggest that abstinence from smoking may reduce RA risk -Smoking status was as of baseline in 1986 -Study population was mostly postmenopausal women
Costenbader, 2006 [27]	Overall RA risk	Nurses' Health Study US women N = 103,818	Prospective cohort	Current smoking: RR 1.43 (95 % CI 1.16–1.75) for RA Past smoking: RR 1.47 (95 % CI 1.23–1.76) for RA Current smoking: RR 1.58 (95 % CI 1.21–2.06) for RF+RA Compared to non-smoking	-680 RA cases developed during 26 years of follow-up -Demonstrated dose effect of smoking and RA risk -Showed that $> 10$ pack-years was the threshold for significantly increased RA risk -Smoking was specifically associated with increased risk for RF+ RA -After 20 years of smoking cessation, risk of RA was similar to non-smoking -Smoking status was time-varying during follow-up
Sugiyama, 2010 [17]	Overall RA risk	11 case-control studies (N = 4764 RA cases and 13,647 controls) and 5 cohort studies (9121 cases from 566,044 subjects)	Meta-analysis of case-control and cohort studies	All participants: Ever smokers: OR 1.40 (95 % CI 1.25–1.58) for RA Ever smokers: OR 1.66 (95 % CI 1.42–1.95) for RF+RA Among men: Ever smokers: OR 1.89 (95 % CI 1.56–2.28) for RA Ever smokers: OR 3.02 (95 % CI 2.35–3.88) for RF+RA Compared to non-smoking	-Largest study on smoking and RA risk -Clearly demonstrates an effect of smoking on RA risk with potent risks in men and for RF+RA -Did not examine cumulative smoking or CCP status
Di Giuseppe, 2013 [29••]	Overall RA risk	Swedish Mammography Cohort Swedish women N = 34,101	Prospective cohort	Current smoking: RR 2.20 (95 % CI 1.58–3.04) for RA Quit smoking $\geq 15$ years: RR 1.99 (95 % CI 1.23–3.20) for RA Compared to non-smoking	-219 RA cases identified during 8 years of follow-up -Study focused on whether smoking cessation decreases RA risk -There was a clear trend toward decreasing RA risk with years since smoking cessation

**Table 1** (continued)

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
Bhatia, 2007 [33]	Genetic susceptibility to serum RA-related antibody risk	Studies of the etiology of RA US women N = 304	Cross-sectional	≥20 pack-years smoking: HR 56.38 (95 % CI 4.31–736.98) for RF positivity Compared to non-smoking	-Former smokers did not return to the baseline RA risk of non-smokers -RA serostatus was not examined -All subjects were at elevated risk from RA due to <i>HLA-DR4</i> positivity (all were parents of children with type 1 diabetes) -31 subjects had positive RF (only 3 among those with ≥20 pack-years) -Also associated oral contraceptive use with increased risk of RF positivity -Analyzed biorepository of banked samples of patients who later developed RA -Also associated other ACPA with shared epitope and smoking (anti-CEP-1/Eno5-21, anti-CitC1359-369, anti-Fibβ62-81b, anti-Fibβ36-52, and anti-CCP1/Fil307-324) -Demonstrated <i>HLA</i> -smoking interaction in pre-RA subjects regardless of ACPA status prior to RA diagnosis
Kokkonen, 2015 [34••]	Genetic susceptibility to serum RA-related antibody risk	Medical Biobank of Northern Sweden Sweden N = 370 pre-RA cases, N = 203 RA cases, N = 585 controls	Case-control	Among pre-RA cases within 10.5 years of diagnosis: Ever smoking and shared epitope positivity: OR 3.96 (1.87–8.39) for CCP2 positivity Compared to non-smoking, shared epitope negative	
Kim, 2015 [35••]	Genetic susceptibility to RA risk	Nurses' Health Studies, Epidemiological Investigations of RA, Korean RA Cohort US women, Sweden, Korea N = 229 RA cases and N = 360 controls; N = 1654 RA cases and N = 1934 controls; N = 1390 cases and N = 735 controls	Case-control	>10 pack-years and high genetic risk score: OR ranging from 4.42 to 61.49 for seropositive RA Compared to 0–10 pack-years and low genetic risk score	-Used fine mapping of the Major Histocompatibility Complex Class II to calculate genetic risk scores based on amino acid haplotypes of HLA proteins to investigate the <i>HLA</i> shared epitope-smoking interaction in three diverse study populations -Specific interactions were found at amino acid positions 11 and 13 of HLA-DRβ1 -Valine at position 11 and histidine at position 13 of HLA-DRβ1 interacted with smoking to significantly increase risk of seropositive RA -Results implied a physical interaction of peptide binding sites in HLA-DRβ1 and a neoantigen induced by smoking
Sparks, 2016 [36•]	Genetic susceptibility to inflammatory arthritis risk	Studies of the Etiology of RA/US N = 966	Prospective cohort	Current smoking: OR 2.12 (95 % CI 1.33–3.38) for inflammatory joint signs Smoking > 10 pack-years: OR 1.89 (95 % CI 1.26–2.82) for inflammatory joint signs	-All subjects were unaffected first-degree relatives of patients with RA -Inflammatory joint signs defined as tender or swollen RA-specific joints by physical examination

**Table 1** (continued)

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
De Hair, 2013 [37••]	Seropositive arthralgias to RA risk	Seropositive Arthralgia Cohort Netherlands N = 55 patients with seropositivity and arthralgias but no synovitis	Prospective cohort	Current smoking: HR 9.6 (95 % CI 1.3–73.0) for RA Compared to non-smoking	-222 subjects had inflammatory joint signs at RA-specific joints -Highest risk of inflammatory joint signs for smokers who were <50 years old (OR 4.39, 95 % CI 2.22–8.66) -Subgroup followed prospectively for 2 years had similar findings -No associations of RA genetic risk score, obesity, CCP, or RF with inflammatory joint signs -All patients had positive RF or CCP and joint symptoms but no synovitis on physical examination -15 subjects developed classifiable RA during median of 13 months of follow-up -Smoking and overweight/obesity may have synergistic effects in increasing RA risk

*ACPA* anti-citrullinated protein antibodies, *CCP* cyclic citrullinated peptide, *CI* confidence interval, *HR* hazard ratio, *OR* odds ratio, *RA* rheumatoid arthritis, *RF* rheumatoid factor, *RR* relative risk, *US* United States

extra-articular RA disease manifestation, data suggest that lung inflammation can precede RA diagnosis [66•, 67••]. While rates vary depending on RA duration, smoking exposure, medication history, and sensitivity of the testing modality, 8–36 % of RA patients have pulmonary function tests suggesting airways obstruction, 60–80 % of RA patients have airways disease on high-resolution computed tomography (CT) scans, and parenchymal disease is present in up to 79 % of RA patients [10•].

In a cross-sectional study of newly diagnosed RA patients, Wilsher and colleagues evaluated risk factors for lung imaging abnormalities on 60 patients within 1 year of RA diagnosis [68]. They found that bronchial wall thickening was present in 50 % of patients, bronchiectasis in 35 %, ground glass opacities in 18 %, and reticular changes in 12 % [68]. Additionally, CCP and RF correlated with decreasing diffusion capacity to carbon monoxide, suggesting significant alveolar damage in patients with early seropositive RA [68]. Similarly, Fischer and colleagues described significant disease burden in patients with respiratory symptoms who had CCP positivity without RA [69]. On high-resolution CT scans of the chest, 54 % had isolated airways disease, 14 % had ILD alone, and 26 % had airways disease and ILD [69]. A subset who had bronchial biopsies performed showed significant inflammation in the airways [69].

A study in Sweden compared imaging findings on high-resolution CT scans of the chest among 105 untreated patients with early RA to healthy controls [70••]. Parenchymal lung abnormalities were present in 63 % of CCP-positive RA patients compared to only 37 % of CCP-negative patients and 30 % of controls, suggesting that lung inflammation is particularly relevant to ACPA-positive RA [70••]. Airway changes on CT were detected more frequently in RA patients (66 %) compared to controls (42 %), suggesting that local airways inflammation is already present in patients recently diagnosed with RA [70••]. Another study by the same investigators examined bronchial biopsy in similar patients with untreated early RA and found significant local inflammation in the lung at the alveoli and airways [71••]. Adaptive immune cells (B cells, T cells, and plasma cells) were much more likely to be present and activated in CCP-positive patients compared to the CCP-negative RA patients or controls in both bronchial biopsies as well as bronchioalveolar lavage [71••].

These pulmonary abnormalities may have been due to factors other than preclinical RA disease processes. In particular, cigarette smoking may cause many of these abnormalities. However, similar airways changes are also seen in patients with RA who were non-smokers suggesting that smoking alone does not cause these abnormalities. When adjusted for smoking, differences remained between ACPA-positive RA, ACPA-negative RA, and controls. Since patients in these studies had very early RA and were untreated, it is less likely that other clinical factors such as decreased physical activity or medication use explain these differences.

Demoruelle and colleagues investigated pulmonary function and imaging among subjects with seropositivity (positive CCP2 or CCP3.1  $\pm$   $\geq$  2 RF isotypes) without inflammatory arthritis [72]. This group at very elevated risk of RA was compared to autoantibody-negative controls who were unaffected first-degree relatives of RA patients as well as patients with early RA. Seropositive subjects showed significant burden or pulmonary abnormalities compared to controls [72]. Of those who were seropositive, 76 % had airways disease compared to only 33 % of controls. Fifty percent of seropositive cases without RA had bronchial thickening compared to only 13 % of seronegative controls; 69 % of seropositive cases had air trapping compared to 7 % of seronegative controls [72]. Early RA patients had more bronchial wall thickening and parenchymal abnormalities but were otherwise similar to seropositive cases without RA. Smoking was a matching factor and similar trends were seen when analyzing only never smokers.

Together, these studies in early RA and subjects with seropositivity without RA are compelling to emphasize the inflammation that occurs in airways and alveoli prior to disease. Further, many of these patients with RA had no evidence of ILD on biopsy or imaging arguing that these abnormalities are important for ACPA-positive RA and not only patients with RA who develop ILD.

### Citrullination of Antigens in Respiratory Mucosa

While studies demonstrate excess local inflammation in the alveoli and airways in preclinical RA, the biological processes occurring here are challenging to study since these individuals are difficult to identify and are otherwise often healthy. Citrullination is a process by which arginine residues on proteins, such as fibrinogen and enolase, are converted to citrulline by the enzyme peptidylarginine deiminase (PAD) [73•]. Autoantibodies to citrullinated protein antigens are highly specific to RA and presence of these in preclinical phases of RA greatly increases the risk of progression to clinical RA. Therefore, citrullination has been postulated to be one of the biologic processes occurring locally in the lung that may initiate RA pathogenesis. Citrullination may be the important first step in the loss of immune tolerance, forming self-antigens that are then presented to T cells that eventually result in activated B cells producing ACPA [74, 75•].

Reynisdottir and colleagues analyzed fluid from bronchoalveolar lavage and found increased citrullination of proteins in CCP2-positive early untreated RA patients compared to CCP2-negative RA and healthy controls [71••]. Ytterberg and colleagues performed proteomic screening of bronchial and synovial biopsies of RA patients to find proteins in common at both sites that might link the lung and joints [76••]. They found that two citrullinated vimentin peptides were identified in the majority of these patients at both anatomic sites,

perhaps providing a link between initial breakdown of the innate immune system locally and the lung and inflammation in the joints after adaptive immune system dysregulation [76••].

While these findings are provocative, they require replication and would likely not explain the full spectrum of ACPA-positive RA pathogenesis. Smoking may be the extrinsic factor that causes local mucosal inflammation in the lung, upregulating PAD in susceptible hosts, leading to citrullination of proteins that form neoantigens that stimulate the immune system and induce autoimmunity. However, there may be other parallel biologic processes that also lead to RA. These might include a collection of neoantigens (instead of a single instigator), inflammation at other mucosal sites (in particular the gut and periodontum), or genetic susceptibility factors that are currently undiscovered.

### The Lung as a Site of Initial RA-Related Autoantibody Formation

While the particular immunologic mechanisms occurring locally in the lung prior to RA development remains to be fully elucidated, data support the notion that autoantibody development is initiated specifically in the lung (Table 2). Willis and colleagues investigated individuals at high risk of RA due to family history or detectable serum antibodies [77••]. They evaluated a variety of RA-related autoantibodies in the serum and induced sputum of seronegative relatives, seropositive relatives, and early RA patients. They found that 65 % of seropositive at-risk relatives had evidence for at least one RA-related autoantibody in the sputum (CCP2, CCP3, CCP3.1, IgG-RF, IgA-RF, or IgM-RF) compared to 86 % of early RA subjects, and only 35 % of seronegative at-risk relatives [77••]. In addition, most of those with detectable serum RA-related autoantibodies had the same autoantibody detectable in the sputum [77••]. The authors conclude that these results provide evidence that autoantibodies are initially locally produced in the lung prior to epitope spreading and subsequent detection in the serum. Some argued that sputum may have been contaminated from oral secretions in this study, especially relevant since the periodontum is another mucosal surface that may be important in RA pathogenesis [80]. In particular, the bacterium *Propionibacterium gingivalis* synthesizes a type of PAD that may induce local citrullination in the gingiva [81, 82].

Reynisdottir and colleagues also studied whether RA-related autoantibodies were present locally in the lung of newly diagnosed untreated RA patients by obtaining specimens from bronchoalveolar lavage [70••]. They found that CCP-positive RA patients had higher levels of ACPA autoantibodies in the bronchoalveolar lavage samples than in the serum, suggesting that the autoantibody production may have

started in the lung [70••]. These findings also correlated to patients with structural lung disease on high-resolution CT scans of the chest [70••]. Only one CCP-negative RA patient and no healthy controls had detectable ACPA in the lung [70••]. Since the fluid was collected by bronchoalveolar lavage, it was unlikely that oral flora contaminated these specimens. While preclinical cases were unavailable in this study, the RA patients were all untreated and were recently diagnosed arguing that these changes likely preceded the clinical onset of RA.

A study by Janssen and colleagues evaluated whether other diseases involving inflammation of the mucosal surfaces might have detectable RA-related autoantibodies in the serum. They tested CCP2 in the sera of patients with periodontitis, bronchiectasis, and cystic fibrosis as well as patients with RA and healthy controls [78•]. CCP2 seropositivity was associated with RA, bronchiectasis, and cystic fibrosis but not periodontitis [78•]. The authors conclude that this provides further evidence that local inflammation of the mucosa in the lung may provide conditions necessary for citrullination and ACPA production. Studies of seropositive RA-ILD without joint involvement and the previously described study of patients with pulmonary complaints and CCP positivity without RA argue that there is a group of patients on the spectrum of RA where clinical pulmonary disease precedes or is more severe than joint symptoms [69, 83, 84•, 85, 86].

### Future Directions

While much progress has been made in elucidating the pathogenesis of RA, further work is needed in nearly every phase of preclinical RA. While the shared epitope is the strongest genetic risk factor for RA, currently described genetic factors only explain 18 % of the genetic variance for RA [50••, 87•]. Other potential RA susceptibility factors such as epigenetics, the microbiome, and metabolomics may be useful in identifying high-risk individuals and further bridge genetics with environmental factors for RA development [6•, 88]. Other lifestyle factors, particularly metabolic factors such as diet and obesity, may be important in the development of RA [89•, 90•, 91•]. Future epidemiologic studies of interactions of smoking with other lifestyle factors or biomarkers may offer novel hypotheses for the biology behind preclinical RA phase transitions. While smoking is clearly important in RA etiology, many non-smokers develop RA so studying other environmental factors may provide important alternate hypotheses for RA. Studies concerning the microbiome and periodontitis have already offered other hypotheses for RA development [82, 88].

While structural abnormalities in the alveoli and airways of individuals with preclinical RA are well documented, less is known about the immunologic changes occurring locally in these patients. Most studies investigating this have used early



**Table 2** Selected epidemiologic and translational studies investigating the lung in RA preclinical phase transitions

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
Reynisdottir, 2014 [70••]	Alveolar/airways mucosal inflammation, citrullination to form neoantigens, pulmonary RA-related autoantibodies	Lung Investigation in Early RA Sweden N = 105 untreated early RA patients, N = 43 healthy controls	Translational	63 % of ACPA-positive RA patients had parenchymal lung abnormalities, compared with only 37 % of ACPA-negative RA patients and 30 % of healthy controls ( $P < 0.05$ ) Airway changes detected were more frequent in RA patients than in healthy controls (66 % versus 42 %; $P < 0.05$ )	-High resolution computed tomography and bronchoalveolar lavage performed on early untreated RA patients and compared to healthy controls -Results were adjusted for smoking -Increased citrullination in bronchial tissue of CCP-positive RA patients compared to CCP-negative RA patients -ACPA levels were relatively higher in the BAL fluid as compared with the sera of CCP-positive RA patients -Bronchial biopsy and bronchoalveolar lavage performed on early untreated RA patients and compared to healthy controls -Evidence of local inflammation in the lung (germinal centers, B cells, plasma cells, T cells) for ACPA-positive RA compared to ACPA-negative RA and controls -Authors suggest that inflammation preceded clinical symptoms -Smoking was not adjusted for, but abnormalities were not seen in healthy smoking controls
Reynisdottir, 2015 [71••]	Alveolar/airways mucosal inflammation	Lung Investigation in Early RA Sweden N = 24 untreated early RA patients, N = 79 healthy controls	Translational	Lymphocyte infiltration was more frequently found in ACPA-positive patients (50 %) as compared with ACPA-negative patients (17 %) and controls (13 %) BAL samples of patients with ACPA-positive, but not ACPA-negative, RA had significantly higher relative numbers of lymphocytes and expressed higher levels of activation markers compared with controls	-Defined prevalence of respiratory symptoms, and functional/imaging abnormalities -Authors inferred that most lung abnormalities likely occurred prior to the clinical presentation of RA -Changes were seen in non-smokers suggesting pre-disease RA-specific pulmonary processes
Wilsher, 2012 [68]	Alveolar/airways mucosal inflammation	New Zealand N = 60 early RA patients	Cross-sectional	30 % of early RA patients had respiratory symptoms 50 % had bronchial wall thickening CCP and RF positivity was associated with decreased DLCO	-Cases were at increased RA risk due to family history of RA or positive serum RA-related autoantibodies -Findings suggest that autoantibody generation originates in the lung prior to being detectable in the serum
Willis, 2013 [77••]	Asymptomatic pulmonary RA-related autoantibodies	Studies of the Etiology of RA US N = 49 cases	Case-control	One or more RA-related antibodies (CCP2, CCP3, CCP3.1, IgG-RF, IgA-RF, and IgM-RF) was detected in the sputum of 39 % of at-risk seronegative cases	

**Table 2** (continued)

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
Demoruelle, 2012 [72]	Alveolar/airways mucosal inflammation, Asymptomatic serum RA-related autoantibodies	Studies of the Etiology of RA US N = 42 autoantibody-positive cases, N = 15 autoantibody-negative controls, N = 12 seropositive patients with early RA	Case-control	76 % of seropositive controls without RA had any airways disease compared to 33 % of seronegative controls ( $P = 0.005$ ) 50 % of seropositive controls without RA had bronchial thickening compared to 13 % of seronegative controls ( $P = 0.015$ ) 69 % of seropositive controls without RA had air trapping compared to 7 % of seronegative controls ( $P < 0.001$ )	-35–50 % of at risk subjects were smokers -Seropositive cases without inflammatory arthritis were positive for CCP2 or CCP3.1 and/or $\geq 2$ RF isotypes -Results suggest that pulmonary processes are important in pre-clinical RA pathogenesis -Matching factors included smoking and analyses among never smokers were similar, smoking less likely to have confounded associations -Early RA patients had more bronchial wall thickening and parenchymal abnormalities but were otherwise similar to seropositive cases without RA
Janssen, 2015 [78•]	Asymptomatic serum RA-related autoantibodies	Netherlands N = 86 RA patients, N = 114 periodontitis patients, N = 80 with bronchiectasis, N = 41 cystic fibrosis patients, N = 36 healthy controls	Translational	CCP seropositivity was significantly associated with RA, bronchiectasis, and cystic fibrosis	-Study evaluating whether RA-related autoantibodies may be more prevalent in diseases of the oral and respiratory mucosa -Results were adjusted for smoking status (ever/never) -Patients with diseases characterized by inflammation of the respiratory mucosa had significantly more serum RA-related autoantibodies compared to healthy controls
Fischer, 2012 [69]	Alveolar/airways mucosal inflammation, asymptomatic serum RA-related autoantibodies	US N = 74 patients with respiratory symptoms but no RA	Retrospective cohort	On high-resolution computed tomography scans, 54 % had isolated airways disease, 14 % had isolated interstitial lung disease (ILD), 26 % had airways disease and ILD, and 7 % had pulmonary fibrosis and emphysema	-Characterized CCP-positive patients without RA who presented to a pulmonology center -3/74 later went on to develop classifiable RA -58 % were former smokers -Nine out of ten patients had significant airway inflammation in on bronchial biopsy
Ytterberg, 2015 [76••]	Citrullination to form neoantigens, immune tolerance loss, joint specificity	Epidemiological Investigations of RA Sweden	Translational	Five peptides were shared between synovial and bronchial biopsy specimens in RA patients	-Screened for proteins present in both bronchial and synovial biopsies of patients with RA

**Table 2** (continued)

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
Nannini, 2013 [79••]	Clinical RA	<p><math>N = 393</math> patients with RA, <math>N = 152</math> healthy controls, <math>N = 236</math> disease controls</p> <p>Olmsted County, Minnesota US  <math>N = 594</math> RA patients, <math>N = 596</math> non-RA patients</p>	Prospective cohort	<p>Two citrullinated vimentin (cit-vim) peptides were detected in the majority of synovial and lung tissue biopsies</p> <p>Significantly more RA patients (9.6 %) developed obstructive lung disease than non-RA patients (6.2 %), HR 1.54 (95 % CI 1.01–2.34)</p>	<p>-Two citrullinated vimentin peptides were identified in the majority of both anatomic sites</p> <p>-Antibodies to these peptides were detected in the sera of patients with RA</p> <p>-Patients with RA at diagnosis and a matched comparison cohort of patients without RA were followed for mean of 16.3 and 19.4 years, respectively</p> <p>-Excluded obstructive lung disease diagnosed before RA</p> <p>-52 RA patients and 40 non-RA patients developed obstructive lung disease (majority was chronic obstructive pulmonary disease)</p> <p>-Risk of obstructive lung disease was four-fold increased in smokers</p> <p>-Excess obstructive lung disease accounted for 6.3 % of RA mortality burden</p> <p>-Suggests that preclinical RA disease processes in the lung may contribute to clinical outcomes after RA diagnosis</p> <p>-Women without RA were followed for 36 years, during which 964 developed RA</p> <p>-There were 307 deaths among women with RA during follow-up</p> <p>-14.3 % of deaths were due to respiratory causes among women with RA compared to 7.2 % of deaths in women without RA</p> <p>-Results were adjusted for time-varying smoking pack-years before and after RA diagnosis</p> <p>-Most RA respiratory deaths were due to obstructive lung disease, suggesting that airway disease, in addition to ILD, is a major contributor to respiratory mortality in RA</p>
Sparks, 2015 [3••]	Preclinical to Clinical RA	Nurses' Health Study US $N = 119,209$	Prospective cohort	<p>RA was associated with increased respiratory mortality (HR 2.06, 95 % CI 1.51–2.80)</p> <p>Seropositive RA associated with nearly 3-fold increased respiratory mortality (2.67, 95 % CI 1.89–3.77)</p> <p>Compared to non-RA</p>	

**Table 2** (continued)

First author, year [reference]	Preclinical RA phase	Study Population Number of subjects	Study design	Summary	Comments
					-Since findings were independent of smoking before and after RA diagnosis, airway disease may have been initiated during preclinical RA phases

*ACPA* anti-citrullinated protein antibodies, *BAL* bronchoalveolar lavage, *CCP* cyclic citrullinated peptide, *CI* confidence interval, *DLCO* diffusing capacity of the lungs for carbon monoxide, *ILD* interstitial lung disease, *HR* hazard ratio, *RA* rheumatoid arthritis, *RF* rheumatoid factor, *RR* relative risk, *US* United States

RA patients and inferred that these changes occurred prior to clinical onset [70••, 71••]. Since ACPA-positive RA patients often develop seropositivity within 5 years before clinical onset, these transition states have not been able to be studied directly given the difficulty in identifying these patients. Since many are otherwise healthy, the societal benefit of lung biopsy or even bronchoscopy may not outweigh the risk to the individual subjects. However, if the particular citrullinated neoantigen(s) formed during these processes were identified, targeted approaches to RA prevention would likely have much more potential for success and may also have importance in identifying targets for treatment of early and established RA. Lastly, while much progress has been made concerning the natural history of RA using banked specimens of patients who later developed RA, there are only a few longitudinal studies of at risk subjects available (typically unaffected first-degree relatives or those with arthralgias or undifferentiated inflammatory arthritis) [5••, 11•, 42••, 74, 92]. Larger longitudinal studies with longer follow-up of individuals at increased risk are necessary to understand the natural history, but recruitment and retention are major challenges in this otherwise healthy population.

Preclinical disease processes may have clinical implications for patients with established RA. Several groups have recently identified respiratory mortality as a major contributor to the excess mortality of RA patients, particularly those with seropositive RA [3••, 93••, 94]. Emerging evidence suggests that this increased risk may be due to obstructive lung diseases and that RA patients are at particularly increased risk for asthma and COPD [3••, 79••, 95•, 96]. Since airways disease is particularly common in preclinical RA, these lesions may predispose RA patients to later develop clinical obstructive disease, beyond the effect of smoking. Preclinical pulmonary processes may also further subset patients with RA who later develop ILD which has known excess mortality in RA [97]. Identifying RA patients early in the development of ILD may similarly have major prevention or treatment implications and many of these processes likely are initiated in the preclinical RA phases [98••].

Finally, understanding the pathogenesis of RA may have public health implications beyond RA. Studies of excess CVD outcomes among RA patients have impacted the understanding of how chronic inflammation affects the biology of CVD and has had broad public health significance beyond RA [99–103]. Similar to this framework, inflammation and autoimmunity may also be the common link between RA and increased respiratory disease burden and mortality. Bronchiolar immune tolerance loss with resultant systemic inflammation may bridge RA pathogenesis to clinical outcomes [94]. Evaluating the effect of RA on respiratory outcomes therefore has importance not only for RA patients, but may establish the roles of systemic inflammation and autoimmunity on respiratory outcomes beyond RA, which may have broad biologic, clinical, and public health implications.

## Conclusions

The understanding of the pathogenesis of RA has progressed over the past two decades through both epidemiologic and translational studies. Specifically, epidemiologic investigations of smoking and risk of RA have informed the paradigm for RA development and generated novel hypotheses that have been tested in translational studies to further understand the biology of RA pathogenesis. The association of smoking with RA led investigators to initially consider the lung as a site of RA pathogenesis. Later studies associated smoking with progression to inflammatory arthritis or classifiable RA in high-risk populations. The particular association of smoking with seropositive RA, particularly ACPA-positive RA, led investigators to consider citrullination as a biologic process central to RA pathogenesis. The identification of structural lung abnormalities and local autoantibody production in the lungs add to this hypothesis. Finally, the *HLA* shared epitope-smoking interaction, occurring specifically in the peptide binding groove of HLA-DR $\beta$ 1, offers the possibility of a physical interaction of a neoantigen induced by smoking and the immune cells where dysregulation may be initiated.

## Compliance with Ethical Standards

**Conflict of Interest** Jeffrey A. Sparks and Elizabeth W. Karlson declare that they have no conflicts of interest.

**Human and Animal Rights and Informed Consent** This article does not contain any primary data concerning studies with animals or human subjects.

**Funding** Jeffrey A. Sparks was supported by the Rheumatology Research Foundation Scientist Development Award and the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Numbers K23 AR069688 and L30 AR066953. Elizabeth W. Karlson was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Numbers R01 AR049880, K24 AR052403, and P60 AR047782. The funders had no role in the preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of Harvard University, its affiliated academic health care centers, or the National Institutes of Health.

## References

Papers of particular interest, published recently, have been highlighted as:

- Of importance
  - Of major importance
1. Kirwan JR. Links between radiological change, disability, and pathology in rheumatoid arthritis. *J Rheumatol*. 2001;28(4):881–6.
  2. Crowson CS, Matteson EL, Myasoedova E, et al. The lifetime risk of adult-onset rheumatoid arthritis and other inflammatory autoimmune rheumatic diseases. *Arthritis Rheum*. 2011;63(3):633–9.
  3. Sparks JA, Chang SC, Liao KP, et al. Rheumatoid arthritis and mortality among women during 36 years of prospective follow-up: results from the Nurses' Health Study. *Arthritis Care Res (Hoboken)*. 2015. **Large prospective cohort study that followed women before and after RA diagnosis and found nearly three-fold increased risk of respiratory mortality among those with seropositive RA compared to those without RA that was independent of smoking.**
  4. Dadoun S, Zeboulon-Ktorza N, Combescore C, et al. Mortality in rheumatoid arthritis over the last fifty years: systematic review and meta-analysis. *Joint Bone Spine*. 2013;80(1):29–33.
  5. van Steenberg HW, Huizinga TW, van der Helm-van Mil AH. The preclinical phase of rheumatoid arthritis: what is acknowledged and what needs to be assessed? *Arthritis Rheum*. 2013;65(9):2219–32. **Systematic review of preclinical RA summarizing prior studies and setting a research agenda for future studies.**
  6. Sparks JA, Costenbader KH. Genetics, environment, and gene-environment interactions in the development of systemic rheumatic diseases. *Rheum Dis Clin N Am*. 2014;40(4):637–57. **Review of the framework for epidemiologic studies examining genetics, environment, and gene-environment interaction in rheumatic diseases.**
  7. Karlson EW, Deane K. Environmental and gene-environment interactions and risk of rheumatoid arthritis. *Rheum Dis Clin N Am*. 2012;38(2):405–26.
  8. Gerlag DM, Raza K, van Baarsen LG, et al. EULAR recommendations for terminology and research in individuals at risk of rheumatoid arthritis: report from the Study Group for Risk Factors for Rheumatoid Arthritis. *Ann Rheum Dis*. 2012;71(5):638–41.
  9. Demoruelle MK, Deane KD, Holers VM. When and where does inflammation begin in rheumatoid arthritis? *Curr Opin Rheumatol*. 2014;26(1):64–71.
  10. Demoruelle MK, Solomon JJ, Fischer A, et al. The lung may play a role in the pathogenesis of rheumatoid arthritis. *Int J Clin Rheumatol*. 2014;9(3):295–309. **Thorough review examining the rationale and biology supporting the role of the lung in RA pathogenesis.**
  11. Karlson EW, van Schaardenburg D, van der Helm-van Mil AH. Strategies to predict rheumatoid arthritis development in at-risk populations. *Rheumatology (Oxford)*. 2014. **Summarizes prior epidemiologic research studies examining transitions between preclinical phases of RA.**
  12. Sparks JA, Chen CY, Jiang X, et al. Improved performance of epidemiologic and genetic risk models for rheumatoid arthritis serologic phenotypes using family history. *Ann Rheum Dis*. 2015;74(8):1522–9. **Developed and validated prediction models for RA that included genetics, environmental factors, and family history; among those with positive family history, models were able to accurately identify those who developed either seropositive or seronegative RA.**
  13. Deane KD. Can rheumatoid arthritis be prevented? *Best Pract Res Clin Rheumatol*. 2013;27(4):467–85. **Thorough summary of literature supporting the notion that RA can be prevented.**
  14. Lahiri M, Morgan C, Symmons DP, et al. Modifiable risk factors for RA: prevention, better than cure? *Rheumatology (Oxford)*. 2012;51(3):499–512.
  15. Sparks JA, Iversen MD, Miller Kroouze R, et al. Personalized Risk Estimator for Rheumatoid Arthritis (PRE-RA) Family Study: rationale and design for a randomized controlled trial evaluating rheumatoid arthritis risk education to first-degree relatives. *Contemp Clin Trials*. 2014;39(1):145–57.

16. Gerlag DM, Norris JM, Tak PP. RA: from risk factors and pathogenesis to prevention: towards prevention of autoantibody-positive rheumatoid arthritis: from lifestyle modification to preventive treatment. *Rheumatology (Oxford)*. 2015.
17. Sugiyama D, Nishimura K, Tamaki K, et al. Impact of smoking as a risk factor for developing rheumatoid arthritis: a meta-analysis of observational studies. *Ann Rheum Dis*. 2010;69(1):70–81.
18. Criswell LA, Merlino LA, Cerhan JR, et al. Cigarette smoking and the risk of rheumatoid arthritis among postmenopausal women: results from the Iowa Women's Health Study. *Am J Med*. 2002;112(6):465–71.
19. Heliövaara M, Aho K, Aromaa A, et al. Smoking and risk of rheumatoid arthritis. *J Rheumatol*. 1993;20(11):1830–5.
20. Hazes JM, Dijkmans BA, Vandenbroucke JP, et al. Lifestyle and the risk of rheumatoid arthritis: cigarette smoking and alcohol consumption. *Ann Rheum Dis*. 1990;49(12):980–2.
21. Vessey MP, Villard-Mackintosh L, Yeates D. Oral contraceptives, cigarette smoking and other factors in relation to arthritis. *Contraception*. 1987;35(5):457–64.
22. Voigt LF, Koepsell TD, Nelson JL, et al. Smoking, obesity, alcohol consumption, and the risk of rheumatoid arthritis. *Epidemiology*. 1994;5(5):525–32.
23. Symmons DP, Bankhead CR, Harrison BJ, et al. Blood transfusion, smoking, and obesity as risk factors for the development of rheumatoid arthritis: results from a primary care-based incident case-control study in Norfolk, England. *Arthritis Rheum*. 1997;40(11):1955–61.
24. Stolt P, Bengtsson C, Nordmark B, et al. Quantification of the influence of cigarette smoking on rheumatoid arthritis: results from a population based case-control study, using incident cases. *Ann Rheum Dis*. 2003;62(9):835–41.
25. Kallberg H, Ding B, Padyukov L, et al. Smoking is a major preventable risk factor for rheumatoid arthritis: estimations of risks after various exposures to cigarette smoke. *Ann Rheum Dis*. 2011;70(3):508–11.
26. Sparks JA, Chen CY, Hiraki LT, et al. Contributions of familial rheumatoid arthritis or lupus and environmental factors to risk of rheumatoid arthritis in women: a prospective cohort study. *Arthritis Care Res (Hoboken)*. 2014;66(10):1438–46. **Prospective study examining familial and environmental risk factors for RA; the population attributable risk for RA from known environmental risk factors was 41 % compared to 21 % from family history.**
27. Costenbader KH, Feskanich D, Mandl LA, et al. Smoking intensity, duration, and cessation, and the risk of rheumatoid arthritis in women. *Am J Med*. 2006;119(6):503 e1–9.
28. Di Giuseppe D, Discacciati A, Orsini N, et al. Cigarette smoking and risk of rheumatoid arthritis: a dose-response meta-analysis. *Arthritis Res Ther*. 2014;16(2):R61. **Large meta-analysis examining the effects of smoking status and cumulative pack-years on risk of RA serologic phenotypes.**
29. Di Giuseppe D, Orsini N, Alfredsson L, et al. Cigarette smoking and smoking cessation in relation to risk of rheumatoid arthritis in women. *Arthritis Res Ther*. 2013;15(2):R56. **This large prospective study showed that women who quit smokers were at decreased risk of RA compared to those who continued to smoke, although the risk did not return to the baseline risk of RA of never smokers.**
30. Yahya A, Bengtsson C, Lai TC, et al. Smoking is associated with an increased risk of developing ACPA-positive but not ACPA-negative rheumatoid arthritis in Asian populations: evidence from the Malaysian MyEIRA case-control study. *Mod Rheumatol*. 2012;22(4):524–31.
31. Pedersen M, Jacobsen S, Klarlund M, et al. Environmental risk factors differ between rheumatoid arthritis with and without auto-antibodies against cyclic citrullinated peptides. *Arthritis Res Ther*. 2006;8(4):R133.
32. Han B, Diogo D, Eyre S, et al. Fine mapping seronegative and seropositive rheumatoid arthritis to shared and distinct HLA alleles by adjusting for the effects of heterogeneity. *Am J Hum Genet*. 2014;94(4):522–32. **This large study eliminated heterogeneity of seronegative RA using research assays for RA-related autoantibodies and genetic testing for spondyloarthropathies; distinct genetic risk factors between seronegative and seropositive argue that both are different diseases.**
33. Bhatia SS, Majka DS, Kittelson JM, et al. Rheumatoid factor seropositivity is inversely associated with oral contraceptive use in women without rheumatoid arthritis. *Ann Rheum Dis*. 2007;66(2):267–9.
34. Kokkonen H, Brink M, Hansson M, et al. Associations of antibodies against citrullinated peptides with human leukocyte antigen-shared epitope and smoking prior to the development of rheumatoid arthritis. *Arthritis Res Ther*. 2015;17:125. **Comprehensive study examining the relationship between the shared epitope, smoking, and specific autoantibodies against specific ACPA; the shared epitope and smoking significantly interacted for specific ACPA, particularly those closer to onset of symptoms.**
35. Kim K, Jiang X, Cui J, et al. Interactions between amino acid-defined major histocompatibility complex class II variants and smoking in seropositive rheumatoid arthritis. *Arthritis Rheumatol*. 2015;67(10):2611–23. **This trans-ethnic study used fine mapping of the MHC class II region to investigate gene-smoking interaction using amino acid haplotypes; significant interactions were found at amino acid positions 11 and 13 of HLA-DRβ1 suggesting a physical interaction between citrullinated autoantigens produced by smoking and HLA-DR molecules specifically at the peptide-binding groove.**
36. Sparks JA, Chang SC, Deane KD, et al. Associations of smoking and age with inflammatory joint signs among first-degree relatives without rheumatoid arthritis: Results from the Studies of the Etiology of RA. *Arthritis Rheumatol*. 2016 (In Press). **Among unaffected first-degree relatives without RA followed prospectively, smoking was associated with inflammatory joint signs independent of genetics, CCP2, and other environmental factors.**
37. de Hair MJ, Landewe RB, van de Sande MG, et al. Smoking and overweight determine the likelihood of developing rheumatoid arthritis. *Ann Rheum Dis*. 2013;72(10):1654–8. **Among seropositive patients with arthralgias, those that smoked and were overweight were more likely to develop classifiable RA.**
38. van Steenbergen HW, Mangnus L, Reijnen M, et al. Clinical factors, anticitrullinated peptide antibodies and MRI-detected subclinical inflammation in relation to progression from clinically suspect arthralgia to arthritis. *Ann Rheum Dis*. 2015. **This study examined progression from arthralgias to arthritis and found that inflammation detectable on magnetic resonance imaging predicted frank arthritis independent of ACPA and other clinical factors.**
39. Hensvold AH, Magnusson PK, Joshua V, et al. Environmental and genetic factors in the development of anticitrullinated protein antibodies (ACPAs) and ACPA-positive rheumatoid arthritis: an epidemiological investigation in twins. *Ann Rheum Dis*. 2015;74(2):375–80. **Large study examining the contributions of environmental and genetic factors to ACPA positivity using twins; the authors conclude that environmental factors were relatively more important in ACPA serostatus while the shared**

- epitope may be important in progression to arthritis among those with ACPA positivity.**
40. Mahdi H, Fisher BA, Kallberg H, et al. Specific interaction between genotype, smoking and autoimmunity to citrullinated alpha-enolase in the etiology of rheumatoid arthritis. *Nat Genet.* 2009;41(12):1319–24.
  41. Rantapaa-Dahlqvist S, de Jong BA, Berglin E, et al. Antibodies against cyclic citrullinated peptide and IgA rheumatoid factor predict the development of rheumatoid arthritis. *Arthritis Rheum.* 2003;48(10):2741–9.
  42. Arkema EV, Goldstein BL, Robinson W, et al. Anti-citrullinated peptide autoantibodies, human leukocyte antigen shared epitope and risk of future rheumatoid arthritis: a nested case–control study. *Arthritis Res Ther.* 2013;15(5):R159. **This nested case–control study of banked blood collected prior to RA onset, investigated the timing to RA with ACPA and shared epitope positivity.**
  43. Hart JE, Laden F, Puett RC, et al. Exposure to traffic pollution and increased risk of rheumatoid arthritis. *Environ Health Perspect.* 2009;117(7):1065–9.
  44. Hart JE, Kallberg H, Laden F, et al. Ambient air pollution exposures and risk of rheumatoid arthritis: results from the Swedish EIRA case–control study. *Ann Rheum Dis.* 2013;72(6):888–94.
  45. Stolt P, Kallberg H, Lundberg I, et al. Silica exposure is associated with increased risk of developing rheumatoid arthritis: results from the Swedish EIRA study. *Ann Rheum Dis.* 2005;64(4):582–6.
  46. Stolt P, Yahya A, Bengtsson C, et al. Silica exposure among male current smokers is associated with a high risk of developing ACPA-positive rheumatoid arthritis. *Ann Rheum Dis.* 2010;69(6):1072–6.
  47. Too CL, Muhamad NA, Ilar A, et al. Occupational exposure to textile dust increases the risk of rheumatoid arthritis: results from a Malaysian population-based case–control study. *Ann Rheum Dis.* 2015. **This recent study associated exposure to textile dust with increased risk of RA independent of smoking in a Malaysian population-based study.**
  48. Fernando MM, Stevens CR, Walsh EC, et al. Defining the role of the MHC in autoimmunity: a review and pooled analysis. *PLoS Genet.* 2008;4(4):e1000024.
  49. Gregersen PK, Silver J, Winchester RJ. The shared epitope hypothesis. An approach to understanding the molecular genetics of susceptibility to rheumatoid arthritis. *Arthritis Rheum.* 1987;30(11):1205–13.
  50. Okada Y, Wu D, Trynka G, et al. Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature.* 2014;506(7488):376–81. **Large and comprehensive trans-ethnic genome-wide association study of >100,000 subjects that discovered 42 novel RA risk loci.**
  51. Raychaudhuri S, Sandor C, Stahl EA, et al. Five amino acids in three HLA proteins explain most of the association between MHC and seropositive rheumatoid arthritis. *Nat Genet.* 2012;44(3):291–6.
  52. Karlson EW, Chang SC, Cui J, et al. Gene–environment interaction between HLA-DRB1 shared epitope and heavy cigarette smoking in predicting incident rheumatoid arthritis. *Ann Rheum Dis.* 2010;69(1):54–60.
  53. Lundstrom E, Kallberg H, Alfredsson L, et al. Gene–environment interaction between the DRB1 shared epitope and smoking in the risk of anti-citrullinated protein antibody-positive rheumatoid arthritis: all alleles are important. *Arthritis Rheum.* 2009;60(6):1597–603.
  54. Karlson EW, Ding B, Keenan BT, et al. Association of environmental and genetic factors and gene–environment interactions with risk of developing rheumatoid arthritis. *Arthritis Care Res (Hoboken).* 2013;65(7):1147–56. **Comprehensive study that used genetic risk scores, environmental factors, and gene–environment interactions to develop seropositive RA prediction models.**
  55. Criswell LA, Saag KG, Mikuls TR, et al. Smoking interacts with genetic risk factors in the development of rheumatoid arthritis among older Caucasian women. *Ann Rheum Dis.* 2006;65(9):1163–7.
  56. Kallberg H, Padyukov L, Plenge RM, et al. Gene–gene and gene–environment interactions involving HLA-DRB1, PTPN22, and smoking in two subsets of rheumatoid arthritis. *Am J Hum Genet.* 2007;80(5):867–75.
  57. Lundberg K, Bengtsson C, Kharlamova N, et al. Genetic and environmental determinants for disease risk in subsets of rheumatoid arthritis defined by the anticitrullinated protein/peptide antibody fine specificity profile. *Ann Rheum Dis.* 2013;72(5):652–8. **Examined genetic and environmental factors RA subsets defined by specific ACPA epitopes; the strongest association between the shared epitope and smoking were for those who had autoantibodies to citrullinated  $\alpha$ -enolase and vimentin.**
  58. Jiang X, Kallberg H, Chen Z, et al. An immunochip-based interaction study of contrasting interaction effects with smoking in ACPA-positive versus ACPA-negative rheumatoid arthritis. *Rheumatology (Oxford).* 2016;55(1):149–55. **Comprehensively investigated gene–smoking interactions for RA serologic phenotypes; variants in the shared epitope and other genes in the MHC class II region interacted for risk of seropositive RA while no gene–smoking interactions were found for seronegative RA.**
  59. Too CL, Yahya A, Murad S, et al. Smoking interacts with HLA-DRB1 shared epitope in the development of anti-citrullinated protein antibody-positive rheumatoid arthritis: results from the Malaysian Epidemiological Investigation of Rheumatoid Arthritis (MyEIRA). *Arthritis Res Ther.* 2012;14(2):R89.
  60. Klareskog L, Stolt P, Lundberg K, et al. A new model for an etiology of rheumatoid arthritis: smoking may trigger HLA-DR (shared epitope)-restricted immune reactions to autoantigens modified by citrullination. *Arthritis Rheum.* 2006;54(1):38–46.
  61. Klareskog L, Ronnelid J, Lundberg K, et al. Immunity to citrullinated proteins in rheumatoid arthritis. *Annu Rev Immunol.* 2008;26:651–75.
  62. Lee HS, Irigoyen P, Kern M, et al. Interaction between smoking, the shared epitope, and anti-cyclic citrullinated peptide: a mixed picture in three large North American rheumatoid arthritis cohorts. *Arthritis Rheum.* 2007;56(6):1745–53.
  63. Padyukov L, Silva C, Stolt P, et al. A gene–environment interaction between smoking and shared epitope genes in HLA-DR provides a high risk of seropositive rheumatoid arthritis. *Arthritis Rheum.* 2004;50(10):3085–92.
  64. Bos WH, Wolbink GJ, Boers M, et al. Arthritis development in patients with arthralgia is strongly associated with anti-citrullinated protein antibody status: a prospective cohort study. *Ann Rheum Dis.* 2010;69(3):490–4.
  65. El-Gabalawy HS, Robinson DB, Hart D, et al. Immunogenetic risks of anti-cyclical citrullinated peptide antibodies in a North American Native population with rheumatoid arthritis and their first-degree relatives. *J Rheumatol.* 2009;36(6):1130–5.
  66. Chatzidionisiyou A, Catrina AI. The lung in rheumatoid arthritis, cause or consequence? *Curr Opin Rheumatol.* 2016;28(1):76–82. **Thorough review focusing on the role of the lung in the pre-clinical period as well as established RA.**
  67. Catrina AI, Ytterberg AJ, Reynisdottir G, et al. Lungs, joints and immunity against citrullinated proteins in rheumatoid arthritis. *Nat Rev Rheumatol.* 2014;10(11):645–53. **Comprehensive review focusing on biologic mechanisms linked preclinical biologic processes in the lung to the development of inflammatory arthritis.**

68. Wilsher M, Voight L, Milne D, et al. Prevalence of airway and parenchymal abnormalities in newly diagnosed rheumatoid arthritis. *Respir Med*. 2012;106(10):1441–6.
69. Fischer A, Solomon JJ, du Bois RM, et al. Lung disease with anti-CCP antibodies but not rheumatoid arthritis or connective tissue disease. *Respir Med*. 2012;106(7):1040–7.
70. Reynisdottir G, Karimi R, Joshua V, et al. Structural changes and antibody enrichment in the lungs are early features of anti-citrullinated protein antibody-positive rheumatoid arthritis. *Arthritis Rheumatol*. 2014;66(1):31–9. **Investigators performed high-resolution CT scans and bronchoalveolar lavage on patients with early, untreated RA and compared results to healthy controls; there were a high frequency of parenchymal and airways abnormalities in early ACPA-positive RA patients suggesting that the lung is important in RA pathogenesis.**
71. Reynisdottir G, Olsen H, Joshua V, et al. Signs of immune activation and local inflammation are present in the bronchial tissue of patients with untreated early rheumatoid arthritis. *Ann Rheum Dis*. 2015. doi:10.1136/annrheumdis-2015-208216. **Patients with early, untreated RA had bronchial biopsy and bronchoalveolar lavage performed and were compared to healthy controls; there was evidence of significant local inflammation in the lung in early ACPA-positive RA patients.**
72. Demoruelle MK, Weisman MH, Simonian PL, et al. Brief report: airways abnormalities and rheumatoid arthritis-related autoantibodies in subjects without arthritis: early injury or initiating site of autoimmunity? *Arthritis Rheum*. 2012;64(6):1756–61.
73. Pruijn GJ. Citrullination and carbamylation in the pathophysiology of rheumatoid arthritis. *Front Immunol*. 2015;6:192. **Overview of the importance of citrullination and carbamylation in RA pathogenesis.**
74. Sokolove J, Bromberg R, Deane KD, et al. Autoantibody epitope spreading in the pre-clinical phase predicts progression to rheumatoid arthritis. *PLoS ONE*. 2012;7(5):e35296.
75. Dwivedi N, Radic M. Citrullination of autoantigens implicates NETosis in the induction of autoimmunity. *Ann Rheum Dis*. 2014;73(3):483–91. **Review of the role of a form of neutrophil cell death (NETosis) in the citrullination of autoantigens for the development of autoimmunity.**
76. Ytterberg AJ, Joshua V, Reynisdottir G, et al. Shared immunological targets in the lungs and joints of patients with rheumatoid arthritis: identification and validation. *Ann Rheum Dis*. 2015;74(9):1772–7. **Proteomic screening was performed from bronchial and synovial biopsies from patients with RA in order to find common proteins that may be implicated in molecular mimicry linking these sites; two citrullinated vimentin peptides were identified in the majority of both anatomic sites and antibodies to these peptides were detected in the sera of patients with RA.**
77. Willis VC, Demoruelle MK, Derber LA, et al. Sputum autoantibodies in patients with established rheumatoid arthritis and subjects at risk of future clinically apparent disease. *Arthritis Rheum*. 2013;65(10):2545–54. **Induced sputum was obtained from cases at increased risk of RA due to family history or positive serum RA-related autoantibodies; RA-related autoantibodies were detectable in the sputa of cases who were seronegative suggesting that autoantibodies are initially produced in the lungs prior to detection in the serum.**
78. Janssen KM, de Smit MJ, Brouwer E, et al. Rheumatoid arthritis-associated autoantibodies in non-rheumatoid arthritis patients with mucosal inflammation: a case-control study. *Arthritis Res Ther*. 2015;17:174. **Study evaluating whether RA-related autoantibodies may be more prevalent in diseases of the oral and respiratory mucosa; diseases of respiratory mucosa (bronchiectasis and cystic fibrosis) were more likely to have CCP positivity suggesting that respiratory mucosal inflammation may induce RA-related autoantibodies; notably, no cystic fibrosis patients had smoking history.**
79. Nannini C, Medina-Velasquez YF, Achenbach SJ, et al. Incidence and mortality of obstructive lung disease in rheumatoid arthritis: a population-based study. *Arthritis Care Res (Hoboken)*. 2013;65(8):1243–50. **In this population-based incident cohort, patients with RA were more likely to develop obstructive lung diseases compared to a matched cohort; RA patients with obstructive lung disease had two-fold increased risk of mortality compared to RA patients without obstructive lung disease.**
80. Lundberg K, Wegner N, Yucel-Lindberg T, et al. Periodontitis in RA—the citrullinated enolase connection. *Nat Rev Rheumatol*. 2010;6(12):727–30.
81. Wegner N, Wait R, Sroka A, et al. Peptidylarginine deiminase from *Porphyromonas gingivalis* citrullinates human fibrinogen and alpha-enolase: implications for autoimmunity in rheumatoid arthritis. *Arthritis Rheum*. 2010;62(9):2662–72.
82. Mikuls TR, Thiele GM, Deane KD, et al. *Porphyromonas gingivalis* and disease-related autoantibodies in individuals at increased risk of rheumatoid arthritis. *Arthritis Rheum*. 2012;64(11):3522–30.
83. Wang T, Zheng XJ, Liang BM, et al. Clinical features of rheumatoid arthritis-associated interstitial lung disease. *Sci Rep*. 2015;5:14897.
84. Yunt ZX, Solomon JJ. Lung disease in rheumatoid arthritis. *Rheum Dis Clin N Am*. 2015;41(2):225–36. **Thorough review of lung abnormalities in patients with established RA.**
85. Chen J, Shi Y, Wang X, et al. Asymptomatic preclinical rheumatoid arthritis-associated interstitial lung disease. *Clin Dev Immunol*. 2013;2013:406927.
86. Gochuico BR, Avila NA, Chow CK, et al. Progressive preclinical interstitial lung disease in rheumatoid arthritis. *Arch Intern Med*. 2008;168(2):159–66.
87. Viatte S, Plant D, Raychaudhuri S. Genetics and epigenetics of rheumatoid arthritis. *Nat Rev Rheumatol*. 2013;9(3):141–53. **Thorough review of the genetics of RA prior to the Okada et al. paper.**
88. Scher JU, Sczesnak A, Longman RS, et al. Expansion of intestinal *Prevotella copri* correlates with enhanced susceptibility to arthritis. *eLife*. 2013;2:e01202.
89. Lu B, Solomon DH, Costenbader KH, et al. Alcohol consumption and risk of incident rheumatoid arthritis in women: a prospective study. *Arthritis Rheumatol*. 2014;66(8):1998–2005. **Prospective cohort study associating moderate alcohol intake with decreased risk of RA among women.**
90. Qin B, Yang M, Fu H, et al. Body mass index and the risk of rheumatoid arthritis: a systematic review and dose-response meta-analysis. *Arthritis Res Ther*. 2015;17(1):86. **Large meta-analysis of 11 prior studies associating being overweight or obese with risk of RA.**
91. Di Giuseppe D, Wallin A, Bottai M, et al. Long-term intake of dietary long-chain n-3 polyunsaturated fatty acids and risk of rheumatoid arthritis: a prospective cohort study of women. *Ann Rheum Dis*. 2014;73(11):1949–53. **Prospective study suggesting that consumption of omega-3 polyunsaturated fatty acids may decrease the risk for RA development.**
92. Deane KD, Norris JM, Holers VM. Preclinical rheumatoid arthritis: identification, evaluation, and future directions for investigation. *Rheum Dis Clin N Am*. 2010;36(2):213–41.
93. England BR, Sayles H, Michaud K, et al. Cause-specific mortality in US veteran men with rheumatoid arthritis. *Arthritis Care Res (Hoboken)*. 2015. **RA conferred a three-fold increased risk of respiratory-related death compared to age-matched controls in this study performed among male veterans.**



94. Gonzalez A, Icen M, Kremers HM, et al. Mortality trends in rheumatoid arthritis: the role of rheumatoid factor. *J Rheumatol*. 2008;35(6):1009–14.
95. Shen TC, Lin CL, Chen CH, et al. Increased risk of chronic obstructive pulmonary disease in patients with rheumatoid arthritis: a population-based cohort study. *QJM*. 2014;107(7):537–43. **This large claims-based retrospective cohort in Taiwan associated RA with nearly two-fold increased risk of COPD compared to matched controls.**
96. Shen TC, Lin CL, Wei CC, et al. The risk of asthma in rheumatoid arthritis: a population-based cohort study. *QJM*. 2014;107(6):435–42. **This large claims-based retrospective cohort in Taiwan associated RA with nearly two-fold increased risk of asthma compared to matched controls.**
97. Bongartz T, Nannini C, Medina-Velasquez YF, et al. Incidence and mortality of interstitial lung disease in rheumatoid arthritis: a population-based study. *Arthritis Rheum*. 2010;62(6):1583–91.
98. Doyle TJ, Patel AS, Hatabu H, et al. Detection of rheumatoid arthritis-interstitial lung disease is enhanced by serum biomarkers. *Am J Respir Crit Care Med*. 2015. **Developed and validated prediction models for RA-ILD using clinical factors, RA-related autoantibodies and biomarkers; models using the biomarkers matrix metalloproteinase 7, pulmonary and activation-regulated chemokine, and surfactant protein D were highly accurate in discriminating RA-ILD from controls with RA but no ILD.**
99. Skeoch S, Bruce IN. Atherosclerosis in rheumatoid arthritis: is it all about inflammation? *Nat Rev Rheumatol*. 2015;11(7):390–400.
100. Libby P, Ridker PM, Hansson GK. Progress and challenges in translating the biology of atherosclerosis. *Nature*. 2011;473(7347):317–25.
101. Ridker PM, Luscher TF. Anti-inflammatory therapies for cardiovascular disease. *Eur Heart J*. 2014;35(27):1782–91.
102. Everett BM, Pradhan AD, Solomon DH, et al. Rationale and design of the Cardiovascular Inflammation Reduction Trial: a test of the inflammatory hypothesis of atherothrombosis. *Am Heart J*. 2013;166(2):199–207.e15.
103. Ridker PM, Thuren T, Zalewski A, et al. Interleukin-1beta inhibition and the prevention of recurrent cardiovascular events: rationale and design of the Canakinumab Anti-inflammatory Thrombosis Outcomes Study (CANTOS). *Am Heart J*. 2011;162(4):597–605.