



# Correction to: Eco-Friendly Mycogenic Synthesis of ZnO and CuO Nanoparticles for In Vitro Antibacterial, Antibiofilm and Antifungal Applications

Asem A. Mohamed<sup>1</sup> · Mohammed Abu-Elghait<sup>2</sup> · Nehad E. Ahmed<sup>1</sup> · Salem S. Salem<sup>2</sup> 

Published online: 25 September 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

**Correction to: Biological Trace Element Research**

<https://doi.org/10.1007/s12011-020-02369-4>

The original version of this article unfortunately contained a mistake. Figures 7 and 8 were inadvertently interchanged. The corrected Figs. 7 and 8 are correctly presented here.

The original article has been corrected.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

---

The online version of the original article can be found at <https://doi.org/10.1007/s12011-020-02369-4>


---

✉ Salem S. Salem  
salemsalahsalem@azhar.edu.eg

<sup>1</sup> Chemistry of Natural and Microbial Products Department, Pharmaceutical Industries Research Division, National Research Centre, Dokki, Giza 12622, Egypt

<sup>2</sup> Botany and Microbiology Department, Faculty of Science, Al-Azhar University, Nasr City, Cairo 11884, Egypt

**Fig. 7** Light inverted microscopic images of *S. aureus* biofilms grown with various concentrations of CuO-NPs: **a** 0.0 mg/mL, represent the positive control; **b** negative control; **c, d** 3.0 and 1.5 mg/mL above the MIC value; **e** 0.7 mg/mL; **f** 0.3 mg/mL; **g** 0.15 mg/mL; **h** 0.07 mg/mL; **i** 0.03 mg/mL; and **j** 0.01 mg/mL. At concentrations from 0.01 to 0.3 mg/mL (f–j) bacteria have appeared as scattered cells and cannot aggregated together to perform normal biofilm.



**Fig. 8** Antifungal activity for ZnO-NPs and CuO-NPs at 10mg/mL against different phytopathogenic fungal strains

