

# Pharmacological Interventions to Ameliorate Neuropathological Symptoms in a Mouse Model of Lafora Disease

Arnaud Berthier · Miguel Payá · Ana M. García-Cabrero ·  
María Inmaculada Ballester · Miguel Heredia · José M. Serratosa ·  
Marina P. Sánchez · Pascual Sanz

Received: 17 September 2014 / Accepted: 12 January 2015 / Published online: 28 January 2015  
© Springer Science+Business Media New York 2015

**Abstract** Lafora disease (LD, OMIM 254780) is a rare fatal neurodegenerative disorder that usually occurs during childhood with generalized tonic-clonic seizures, myoclonus, absences, drop attacks, or visual seizures. Unfortunately, at present, available treatments are only palliatives and no curative drugs are available yet. The hallmark of the disease is the accumulation of insoluble polyglucosan inclusions, called Lafora bodies (LBs), within the neurons but also in heart, muscle, and liver cells. Mouse models lacking functional *EPM2A* or *EPM2B* genes (the two major loci related to the disease) recapitulate the Lafora disease phenotype: they accumulate polyglucosan inclusions, show signs of neurodegeneration, and have a dysregulation of protein clearance and endoplasmic reticulum stress response. In this study, we have subjected a mouse model of LD (*Epm2b*<sup>-/-</sup>) to different pharmacological interventions aimed to alleviate protein clearance and endoplasmic reticulum stress. We have used two chemical chaperones, trehalose and 4-phenylbutyric acid. In addition, we have used metformin, an activator of AMP-activated protein kinase (AMPK), as it has a recognized

neuroprotective role in other neurodegenerative diseases. Here, we show that treatment with 4-phenylbutyric acid or metformin decreases the accumulation of Lafora bodies and polyubiquitin protein aggregates in the brain of treated animals. 4-Phenylbutyric acid and metformin also diminish neurodegeneration (measured in terms of neuronal loss and reactive gliosis) and ameliorate neuropsychological tests of *Epm2b*<sup>-/-</sup> mice. As these compounds have good safety records and are already approved for clinical uses on different neurological pathologies, we think that the translation of our results to the clinical practice could be straightforward.

**Keywords** Lafora disease · Polyglucosan accumulation · Metformin · 4-Phenylbutyric acid · Trehalose · Pharmacological intervention

## Abbreviations

AMPK	AMP-activated protein kinase
LB	Lafora bodies
LD	Lafora disease
4-PBA	4-Phenylbutyric acid
SDS-	Sodium dodecyl sulfate polyacrylamide gel
PAGE	electrophoresis

**Electronic supplementary material** The online version of this article (doi:10.1007/s12035-015-9091-8) contains supplementary material, which is available to authorized users.

A. Berthier · M. I. Ballester · M. Heredia · P. Sanz (✉)  
Instituto de Biomedicina de Valencia (CSIC), Centro de  
Investigación Biomédica en Red de Enfermedades Raras  
(CIBERER), Jaime Roig 11, 46010 Valencia, Spain  
e-mail: sanz@ibv.csic.es

M. Payá  
Departamento de Farmacología, Facultad de Farmacia, Universidad  
de Valencia and Centro de Reconocimiento Molecular y Desarrollo  
Tecnológico (IDM), 46100 Valencia, Spain

A. M. García-Cabrero · J. M. Serratosa · M. P. Sánchez  
Fundación Jimenez Diaz and Centro de Investigación Biomédica en  
Red de Enfermedades Raras (CIBERER), 28040 Madrid, Spain

## Introduction

Lafora disease (LD, OMIM 254780) is a rare fatal neurodegenerative disorder that usually occurs during childhood with generalized tonic-clonic seizures, myoclonus, absences, drop attacks, or visual seizures [1, 2]. Unfortunately, at present, available treatments are only palliatives and no curative drugs are available yet. The hallmark of the disease is the accumulation of insoluble polyglucosan inclusions, called Lafora

bodies (LBs), within the neurons but also in heart, muscle, and liver cells. Recessive mutations in two main loci, *EPM2A* and *EPM2B*, are known to produce LD [3–5]. The *EPM2A* gene encodes the glucan phosphatase laforin and the *EPM2B* gene encodes the E3-ubiquitin ligase malin. Although it is known that both proteins form a functional complex [6–8], the underlying molecular mechanisms of this pathology are not fully understood. In this sense, although laforin and malin have been involved in the regulation of glycogen metabolism, recent reports indicate that they also play a role in alternative physiological pathways such as endoplasmic reticulum stress response [9–11] and protein clearance [10, 12–14].

Protein homeostasis (proteostasis) is achieved by the coordinated action of an efficient folding and transport of newly synthesized proteins and an active quality control with degradative mechanisms to reduce the load of unfolded/misfolded proteins thereby to prevent abnormal protein aggregation [15]. When the folding capacity is saturated, unfolded/misfolded proteins are usually tagged with ubiquitin moieties to target them for proteasome and/or lysosomal degradation. If these protein clearance mechanisms are not efficient, then unfolded/misfolded proteins accumulate and tend to form oligomeric structures. Thus, the presence of protein aggregates is an indication of an altered proteostasis [15]. LBs contain, in addition to insoluble glycogen-like polysaccharides (polyglucosans), ubiquitinated proteins, advanced glycation-end products, chaperones, autophagy components, and proteasome subunits [12, 16]. Therefore, LD can be considered as another example of proteostasis dysfunction in which, as indicated above, protein clearance and endoplasmic reticulum stress are defective. So, we reasoned that the use of compounds that could alleviate proteostasis dysfunction could be beneficial in the treatment of LD. In this sense, in this work, we have used 4-phenylbutyric acid (4-PBA), a chemical chaperone that reverses misfolding and aggregation of proteins associated with several human neurodegenerative diseases [17–19], and trehalose, a non-reducing disaccharide that also acts as a chemical chaperone preventing protein denaturation and protecting cell integrity against several stresses [20, 21].

AMP-activated protein kinase (AMPK) is a heterotrimeric enzyme consisting in one catalytic AMPK $\alpha$ , one scaffold AMPK $\beta$ , and one regulatory AMPK $\gamma$  subunit that senses energy status in cells and responds to drops in ATP by inhibiting catabolic and activating anabolic pathways, to restore ATP homeostasis [22, 23]. AMPK also plays a role in promoting autophagy by inhibiting mTOR kinase or by phosphorylating ULK1 [23]. As it has been reported that AMPK activation plays a neuroprotective role in different neurodegenerative diseases [24–27], we checked whether metformin, an activator of AMPK widely used in the treatment of type 2 diabetes, could have a beneficial effect on LD.

Our results indicate that among the drugs mentioned above, 4-PBA and metformin have a clear beneficial effect on all the assayed neuropathological symptoms of *Epm2b*<sup>-/-</sup> mice.

## Material and Methods

### Ethic Statement, Animal Care, Mice, and Husbandry

This study was carried out in strict accordance with the recommendations in the Guide for the Care and Use of Laboratory Animals of the Consejo Superior de Investigaciones Científicas (CSIC, Spain). All mouse procedures were approved by the animal committee of the Instituto de Biomedicina de Valencia—CSIC [Permit Number: INTRA12 (IBV-4)]. All efforts were made to minimize animal suffering. To minimize the effect of differences in the genetic background of the animals, and since LD is a recessive autosomal disorder, we crossed *Epm2b*<sup>-/-</sup> mice (with a mixed background 129sv:C57BL/6) previously described [14] with control C57BL/6JRccHsd mice obtained from Harlan Laboratories (Barcelona, Spain) to obtain heterozygous *Epm2b*<sup>+/-</sup> mice. These were crossed among them to obtain heterozygous *Epm2b*<sup>+/-</sup> (used as control) and homozygous *Epm2b*<sup>-/-</sup> littermates, which were used in this study. Mice were maintained in the IBV-CSIC facility on a 12/12 light/dark cycle under constant temperature (23 °C) with food and water provided ad libitum.

### Pharmacological Interventions

Trehalose, metformin, and 4-PBA, all these compounds having good safety records and already approved for clinical uses on different neurological pathologies, were purchased from Sigma (Madrid, Spain). Trehalose (2 %), metformin (12 mM), and 4-PBA (20 mM) were administered dissolved in drinking water. *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) male mice (groups of 8–12 animals) at 3 months of age (we chose this age because at this time the *Epm2b*<sup>-/-</sup> mice clearly show the presence of Lafora bodies in the brain and they are at the beginning of their neurological impairment) were treated or not with trehalose, 4-PBA, or metformin. Treatments were applied for 1 month, and mice were subjected to the behavioral tests described below. Then, treatment was continued for an extra month. After this period, mice were subjected again to the same behavioral tests and sacrificed by cervical dislocation. Brain was recovered; cut in two equal pieces, one part conserved at -80 °C for processing for Western blot analyses and the other part fixed in 4 % paraformaldehyde in phosphate buffer saline (PBS); and dehydrated for posterior histological analyses.

## Histological and Immunohistofluorescence Analyses

Dehydrated tissues were embedded in paraffin and sectioned at 4  $\mu\text{m}$ . For periodic acid-Schiff (PAS) staining, sections were deparaffined, rehydrated, and incubated with 1 % periodic acid for 7 min, followed by additional 21-min incubation with Schiff reagent (Sigma, Madrid, Spain). Alternatively, for immunohistochemistry analyses (IHC), sections were deparaffined, rehydrated, and warmed at 95 °C for 30 min in 10 mM citrate buffer for antigen retrieval. Sections were blocked in blocking buffer (3 % BSA-3 % fat-free milk proteins, in PBS) and incubated overnight at 4 °C with primary antibody diluted at 1/200 in blocking buffer [anti-NeuN (Millipore, Madrid, Spain)]. After three washes of 10 min in PBS, sections were incubated for 1 h at room temperature with the biotin-conjugated rabbit secondary antibody diluted at 1/250 in blocking buffer, washed three times with PBS for 5 min, incubated with the immunopure ABC kit (Fisher Scientific, Madrid, Spain) for 30 min in darkness, washed three times in PBS for 5 min, and developed using metal-enhanced DAB method (Fisher Scientific). Both PAS staining and IHC sections were finally counterstained with hematoxylin (Sigma, Madrid, Spain), dehydrated, and mounted in DPX (Merck, Germany). For immunohistofluorescence analyses (IHF), sections were deparaffined, rehydrated, incubated for 20 min in sodium borohydrate (1 mg/ml) to lower background, and warmed at 95 °C for 30 min in 10 mM citrate buffer for antigen retrieval. Sections were blocked in blocking buffer (3 % BSA-3 % fat-free milk proteins, in PBS) and incubated overnight at 4 °C with primary antibody diluted at 1/250 in blocking buffer [anti-polyubiquitin (Enzo Life Sciences, Madrid, Spain) or anti-glial fibrillary acidic protein (GFAP) (Sigma, Madrid, Spain)]. After three washes of 10 min in PBS, sections were incubated for 1 h at room temperature with the appropriate secondary antibody diluted at 1/200 in blocking buffer, washed once with PBS, incubated with DAPI (Sigma, Madrid, Spain), washed twice with PBS, and mounted in FluoMount (Polysciences Inc., USA). Images were acquired with a DM6000 Leica microscope, and fluorescence was quantified using ImageJ software (NIH, USA).

## Western Blot Analyses

Mouse brain homogenates were lysed in RIPA buffer [20 mM Tris-HCl, pH 7.5; 150 mM NaCl; 1 mM EDTA; 1 mM EGTA; 1 % NP-40; 1 % sodium deoxycholate; 1 mM  $\text{Na}_3\text{VO}_4$  and complete protease inhibitor cocktail (Roche, Spain)] for 20 min at 4 °C and centrifuged at 10,000 $\times g$  for 15 min. The supernatants were collected and 25  $\mu\text{g}$  of total protein was subjected to SDS-PAGE, transferred into a PVDF membrane, and revealed with the appropriated antibodies using Li-Cor protocol and the Odyssey software (Li-Cor Corporation, USA). The antibodies used in this study were anti-BiP/GRP78, anti-pThr172-AMPKalpha, anti-AMPKbeta1/2, anti-

pS641-glycogen synthase, anti-glycogen synthase kinase 3b, and anti-pSer9-glycogen synthase kinase 3b, from Cell Signaling Technology Inc. (Barcelona, Spain); anti-LC3B and anti-actin from Sigma (Madrid, Spain); anti-p62 from Abcam (Cambridge, UK); and anti-glycogen synthase from Epitomics (Burlingame, USA).

## Behavioral Tests

Mouse neuropsychological behavior was examined using a selection of standard tests. Briefly, the tests were conducted as follows:

**Tail Suspension Test** It was performed as indicated in [28]. Mice were suspended at 15-cm height by the tail for 30 s, and the foot-clasping posture was scored as follows: 0 with both extended feet and 1 with one clasped foot, alternative foot clasping, or with both clasped feet.

**Vertical Pole Test** It was performed as indicated in [29]. Mice were placed head upward near the top of a vertical rough-surfaced pole (diameter 10 mm, height 51 cm). The time taken for the mice to turn completely downward (time to turn) and the total time taken to reach the floor (locomotion activity time) were recorded, with a cutoff limit of 120 s for the combined movements.

**Open Field Test** It was performed as indicated in [30]. Mice were placed in the left-bottom corner of a dark square open field (40 $\times$ 40 $\times$ 40 cm), and exploration was tracked for 20 min. Distance, immobility, spent time, and number of entries into three concentric zones were measured with the Smart software (PanLab, Barcelona, Spain).

**Beam Balance Test** It was performed as indicated in [31]. The mouse's home cage was placed at one end of an 80-cm-long beam of 26, 12, or 5-mm width, and the start point was the other wider end of the beam. The tapered beam had ledges (2 cm wide) that provide "safety" when the mouse's foot slipped off of the beam. The mouse was given a single trial to learn to walk from the start of the beam to its home cage. Once the mouse reached its home cage, it was permitted to remain in the cage to reinforce the target location. The mouse was given three additional trials, each separated by the 60-s reinforcement. Time to cross the beam was recorded with a cutoff limit of 120 s.

## Statistical Analysis

The data were expressed as mean $\pm$ standard error of the mean (SEM). Statistical significance was determined as indicated in each case by Kruskal-Wallis test, followed by Conover-Inman's multiple comparison test, by Fisher exact test, or by Wilcoxon-Mann-Whitney test using R software [32].


## Results

### The Different Treatments Affect Their Corresponding Molecular Targets

In order to find a beneficial treatment for LD, we used different strategies aimed either to alleviate proteostasis, which is impaired in the disease, or to activate AMPK since, as indicated above, activation of this kinase plays a neuroprotective role in different neurodegenerative diseases. Three-month-old male heterozygous *Epm2b*<sup>+/-</sup> (control) and homozygous *Epm2b*<sup>-/-</sup> (KO) mice were subjected to different pharmacological interventions [trehalose and 4-phenylbutyric acid (4-PBA) to alleviate proteostasis dysfunction and metformin to activate AMPK], and after the treatment period, mice were subjected to different behavioral tests and sacrificed by cervical dislocation, and brain was recovered for biochemical and histological analyses.

We first checked by Western blot analyses of total brain extracts the efficiency of the different drugs in affecting their corresponding anticipated molecular

targets. Trehalose and 4-PBA may both act as chemical chaperones to relieve proteostasis dysfunction. In agreement with this assumption, we found that after 2 months of treatment with any of these compounds, there was an increase in the levels of the chaperone BiP/GRP78, in both control (*Epm2b*<sup>+/-</sup>,  $p=0.006$  w/o TTT vs. trehalose and  $p=0.03$  w/o TTT vs. 4-PBA) and KO malin (*Epm2b*<sup>-/-</sup>,  $p=0.002$  w/o TTT vs. trehalose and  $p=0.007$  w/o TTT vs. 4-PBA) mice (Fig. 1a). No significant differences were observed between the untreated samples of control (*Epm2b*<sup>+/-</sup>) and KO (*Epm2b*<sup>-/-</sup>) mice at this age, in agreement with previous results [10]. On the other hand, treatment with metformin for 2 months induced the activation of AMPK in both control ( $p=0.03$ ) and KO malin ( $p=0.03$ ) mice, as indicated by higher levels of the phosphorylated form of the catalytic subunit of the AMPK complex (pThr172-AMPK $\alpha$ ) (Fig. 1b). In this case, we use the levels of AMPK $\beta$  subunits as loading control since this antibody gives better results than the anti-AMPK $\alpha$  antibody when analyzing endogenous levels of the trimeric AMPK complex.



**Fig. 1** Drugs used in this study affect their corresponding targets in mice brain. Western blot analyses of whole brain extracts using the indicated antibodies and densitometric quantification of the corresponding blots were carried out as described in “Material and Methods.” A representative blot is presented. **a** Anti-BiP/GRP78 (BiP) and anti-actin (loading control) from *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose (2 %) or 4-PBA (20 mM) for 2 months. \* $p<0.05$ , \*\* $p<0.01$  [ $n=3$  (control) or  $n=4$  (KO)] comparing the indicated groups with the basal condition according to


Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test. **b** Anti-pThr172-AMPK $\alpha$  (pAMPK $\alpha$ ) and anti-AMPK $\beta$ 1/2 (loading control) from *Epm2b*<sup>+/-</sup> control and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with 12 mM metformin for 2 months. Bars indicate mean values of the relative intensity of each protein of at least three independent samples $\pm$ SEM. Asterisks denote significant differences (\* $p<0.05$ ) [ $n=3$  (control) or  $n=4$  (KO)] comparing the indicated groups with the basal condition according to the Wilcoxon-Mann-Whitney test

#### 4-PBA or Metformin Treatments Decrease the Accumulation of Polyglucosan Inclusions and Polyubiquitinated Protein Aggregates

Next, we investigated the effects of all drugs on the formation of polyglucosan inclusions (that stained positive with the periodic acid-Schiff protocol; PAS+), the hallmark of LD, in

different brain areas: cerebellum, pontine, hippocampus, olfactory bulb, cortex, and midbrain. As observed in Fig. 2, no PAS+ inclusions were detected in the brain areas of control (*Epm2b*<sup>+/-</sup>) mice. However, untreated *Epm2b*<sup>-/-</sup> mice accumulated polyglucosan inclusions in different areas with different extent, with higher levels in the cerebellum, pontine, and hippocampus and lower levels in olfactory bulb, cortex, and midbrain


**Fig. 2** Histological analyses of PAS+ inclusions in the brain of *Epm2b*<sup>+/-</sup> (control) or *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin (2 months each). **a** Representative frontal sections of the cerebellum, pontine, and hippocampus from *Epm2b*<sup>+/-</sup> control or *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated, as indicated in the text, were stained with periodic acid-Schiff (PAS) and counterstained with hematoxylin. Black arrowheads point to Lafora bodies. Scale bars 25  $\mu$ m. **b** PAS+ inclusions were counted by eye in three independent samples of the cerebellum, pontine, hippocampus, olfactory bulb, cerebral cortex, and midbrain of seven mice of each group, and mean quantification of the number of polyglucosan inclusions in an area of 0.09 mm<sup>2</sup> was represented. Bars indicate mean of values corresponding to seven mice  $\pm$  SEM. Asterisks denote significant differences (\* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001) comparing the indicated groups with the basal condition (*Epm2b*<sup>-/-</sup> w/o TTT) according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test



(Fig. 2b, white bars). The treatment with trehalose did not modify the accumulation of PAS+ granules (Fig. 2a, b). However, treatment with 4-PBA or metformin clearly reduced the number of PAS+ aggregates in the cerebellum ( $p < 0.001$  *Epm2b*<sup>-/-</sup> w/o TTT vs. 4-PBA and  $p < 0.001$  w/o TTT vs. metformin), pontine ( $p = 0.011$  *Epm2b*<sup>-/-</sup> w/o TTT vs. 4-PBA and  $p < 0.04$  w/o TTT vs. metformin), and hippocampus ( $p < 0.001$  *Epm2b*<sup>-/-</sup> w/o TTT vs. 4-PBA and  $p < 0.001$  w/o TTT vs. metformin) (Fig. 2a, b).

It is known that polyglucosan inclusions that accumulate in LD are also enriched in polyubiquitinated proteins [1, 2]. We confirmed this fact by immunohistochemistry. As shown in Fig. 3, untreated *Epm2b*<sup>-/-</sup> mice presented high levels of polyubiquitinated protein aggregates in the cerebellum, pontine, and hippocampus, whereas *Epm2b*<sup>+/-</sup> control mice did not present any of these aggregates. Then, we checked whether the treatment with the different drugs reduced the levels of polyubiquitinated protein aggregates present in *Epm2b*<sup>-/-</sup>

**Fig. 3** Immunohistochemistry analyses of polyubiquitinated protein aggregates in brain of *Epm2b*<sup>+/-</sup> (control) or *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin (2 months). **a** Representative frontal sections of the cerebellum, pontine, hippocampus, and cortex from *Epm2b*<sup>+/-</sup> control or *Epm2b*<sup>-/-</sup> KO mice untreated (w/o TTT) or treated, as indicated, were incubated with anti-polyubiquitin antibody and AlexaFluor488-conjugated anti-rabbit antibody. Scale bars 100  $\mu$ m. **b** Polyubiquitin-associated fluorescence was quantified with ImageJ software in two independent pictures of the cerebellum, pontine, hippocampus, and cerebral cortex from five mice per group, and mean fluorescence in an area of 0.4 mm<sup>2</sup> was represented. Bars indicate mean of fluorescence values corresponding to five mice  $\pm$  SEM. Asterisks denote significant differences (\* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$ ) comparing the indicated groups with the basal condition (*Epm2b*<sup>-/-</sup> w/o TTT) according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test




mice. Consistent with the results described above, trehalose did not modify or slightly increased (pontine:  $p=0.02$ ; hippocampus:  $p=0.02$ ) the levels of polyubiquitinated protein aggregates present in the different studied brain areas of the treated animals (Fig. 3). However, treatment with 4-PBA or metformin clearly decreased the levels of polyubiquitinated protein aggregates in all the assayed areas of the brain [cerebellum:  $p=0.002$  (4-PBA),  $p<0.001$  (metformin); pontine:  $p=0.004$  (4-PBA),  $p<0.001$  (metformin); hippocampus:

$p<0.001$  (4-PBA),  $p<0.001$  (metformin); cortex:  $p=0.003$  (4-PBA),  $p<0.001$  (metformin)] (Fig. 3a, b).

#### Neurodegeneration and Hippocampal Gliosis Is Reduced by 4-PBA or Metformin Treatments

Lafora disease is characterized by neuronal loss and the appearance of the concomitant gliosis [2, 33–34]. We confirmed these facts in the untreated *Epm2b*<sup>-/-</sup> mice

**Fig. 4** Immunohistochemistry analyses of neurodegeneration in the hippocampus of *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin (2 months). **a** Representative frontal sections of the dentate gyrus (DG) and the cornu ammonis 1 (CA1) of the hippocampus from *Epm2b*<sup>+/-</sup> control or *Epm2b*<sup>-/-</sup> KO mice untreated (w/o TTT) or treated, as indicated, were incubated with anti-Neuronal Nuclei (NeuN) antibody. *Black arrowheads* point to empty spaces between neurons in DG. *Scale bars* 25  $\mu$ m. NeuN-positive and NeuN-negative cells were counted in DG (**b**) and CA1 (**c**) sections from three independent mice, and the percentage of NeuN-positive cells in an area of 0.09 mm<sup>2</sup> was represented. *Bars* indicate mean values corresponding to three independent mice  $\pm$  SEM. *Asterisks* denote significant differences ( $*p<0.05$ ,  $**p<0.01$ ) comparing the indicated groups with the basal condition (*Epm2b*<sup>-/-</sup> w/o TTT) according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test




where we observed that selected areas of the hippocampus [dentate gyrus (DG) and cornus ammonis 1 (CA1) regions] had reduced Neuronal Nuclei (NeuN) staining, a neuron-specific marker (DG  $p=0.016$  and CA1  $p=0.006$  in untreated control vs. *Epm2b*<sup>-/-</sup>) (Fig. 4a, b). Untreated *Epm2b*<sup>-/-</sup> mice presented a less compacted staining (see black arrowheads) suggesting the loss of neurons in these areas (Fig. 4a). Trehalose treatment did not prevent neuronal loss in DG or CA1 areas of the

hippocampus (Fig. 4a, b). However, when compared with untreated *Epm2b*<sup>-/-</sup> mice, treatment with 4-PBA or metformin presented a significant higher NeuN staining in both analyzed hippocampus regions [DG:  $p=0.005$  (4-PBA),  $p=0.016$  (metformin); CA1:  $p=0.005$  (4-PBA),  $p=0.008$  (metformin)], suggesting that these treatments prevented neuronal loss.

Similar areas of the hippocampus were also stained with an antibody that recognized GFAP, an astrocyte marker (Fig. 5).

**Fig. 5** Immunohistofluorescence analyses of gliosis in the hippocampus of *Epm2b*<sup>+/-</sup> (control) or *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin (2 months). **a** Representative frontal sections of the dentate gyrus (DG) and the cornus ammonis 1 (CA1) of the hippocampus from *Epm2b*<sup>+/-</sup> control or *Epm2b*<sup>-/-</sup> KO mice untreated (w/o TTT) or treated, as indicated, were incubated with anti-gial fibrillary acidic protein (GFAP) antibody and AlexaFluor488-conjugated anti-rabbit antibody and counterstained with DAPI to show the area where the analysis was performed (*insets*). *Scale bars* 100  $\mu\text{m}$ . GFAP-associated fluorescence was quantified with ImageJ software in two pictures from DG (**b**) and CA1 (**c**) sections from five mice per group, and mean fluorescence in an area of 0.26  $\text{mm}^2$  was represented. *Bars* indicate the mean of fluorescence values corresponding to five mice  $\pm$ SEM. *Asterisks* denote significant differences ( $*p<0.05$ ,  $***p<0.001$ ) comparing the indicated groups with the basal condition (*Epm2b*<sup>-/-</sup> w/o TTT) according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test





This staining was higher in preparations from untreated *Epm2b*<sup>-/-</sup> in comparison to control mice (DG:  $p=0.006$  and CA1:  $p=0.03$  in control vs. *Epm2b*<sup>-/-</sup>) (Fig. 5), confirming the presence of reactive gliosis. Treatment of *Epm2b*<sup>-/-</sup> with 4-PBA significantly decreased around 20 % ( $p=0.02$ ) and 40 % ( $p<0.001$ ) the gliosis in DG and CA1, respectively (Fig. 5). Similarly, metformin treatment reduced gliosis around 40 % ( $p<0.001$ ) and 60 % ( $p<0.001$ ) in DG and CA1, respectively (Fig. 5). Surprisingly, treatment with trehalose also significantly diminished around 40 % the astrogliosis in both regions ( $p<0.001$  in both cases) (Fig. 5).


#### Trehalose, 4-PBA, or Metformin Ameliorates Some *Epm2b*<sup>-/-</sup> Neuropsychological Dysfunctions

In addition to the above-described molecular and histological analyses, each group of mice was subjected to a battery of neuropsychological tests after 1 month of treatment and also at the end of the period of 2 months of treatment. One of the tests was the tail suspension test (TST). Originally designed to screen antidepressant drugs [28], it is based on the fact that animals subjected to the short-term, inescapable stress of being suspended by their tail develop an immobile posture with both extended feet. Clasping of the feet is recorded as a sign of neurodegeneration. As observed in Fig. 6, untreated *Epm2b*<sup>-/-</sup> mice presented a significant worse performance than untreated control mice ( $p=0.002$  and  $p=0.007$ , after 1 and 2 months of treatment, respectively). Treatment with 4-PBA or metformin showed a tendency to improve the performance after 1 month of treatment, but the differences became statistically significant after 2 months of treatment [ $p=0.045$  (4-PBA) and  $p=0.017$  (metformin)] (Fig. 6). Trehalose treatment did not improve the performance of *Epm2b*<sup>-/-</sup> mice in any case.

The vertical pole test was originally designed to quantify mouse movement disorders in animal models of Parkinson disease [29]. As observed in Fig. 7, untreated *Epm2b*<sup>-/-</sup> mice presented a significant worse performance than untreated control mice in the time to turn and locomotion activity parameters of the test ( $p=0.018$  and  $p=0.028$  after 1 month of treatment and  $p<0.001$  for both parameters after 2 months of treatment). Treatment with trehalose decreased both parameters of this test (time to turn and locomotion activity) after 1 month of treatment ( $p<0.001$  and  $p=0.002$ , respectively), and the amelioration was even higher after 2 months of treatment ( $p<0.001$  for both parameters) (Fig. 7). 4-PBA also improved both parameters after 1 month of treatment ( $p<0.001$  and  $p=0.03$ , respectively), and we observed a great amelioration of both the time to turn and locomotion activity parameters after the second month of treatment ( $p<0.001$  for both) (Fig. 7). After 1 month, metformin-treated animals showed a significant improvement in the time to turn parameter ( $p=0.049$ ) and a clear tendency to improve the locomotion activity, although


it was not statistically significant. However, after 2 months of treatment, the improvement became clearly observed for both parameters ( $p=0.0038$  and  $p=0.0053$  for time to turn and locomotion activity, respectively) (Fig. 7).

Animals were also subjected to additional behavioral tests (open field, used to assay general locomotion activity levels and anxiety, Fig. 8; beam balance, used to assay fine motor coordination and balance, Fig. 9), but we did not find any significant difference in the parameters of these tests between treated and untreated animals, possibly because the neurological deterioration present in the *Epm2b*<sup>-/-</sup> mice at 5 months of age could have affected some neuropsychological abilities but not others.



**Fig. 6** Tail suspension test of *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin during 1 month (a) or 2 months (b). Percentage of *Epm2b*<sup>+/-</sup> (control) or *Epm2b*<sup>-/-</sup> (KO) mice showing normal posture (0) or altered (1) stereotypical clasping of the hind limb upon tail suspension ( $n=8-11$  in each group). Significant differences between the groups are indicated (\* $p<0.05$ , \*\* $p<0.01$ ) according to the Fisher exact test followed by Wilcoxon-Mann-Whitney test

**Fig. 7** Vertical pole test for evaluation of motor coordination of *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with trehalose, 4-PBA, or metformin during 1 month (a, b) or 2 months (c, d). After 1 month of treatment, the time taken by the mice to turn completely downward (time to turn) (a) and the total time taken to reach the floor (locomotion activity) (b) were recorded. After 2 months of treatment, the time to turn (c) and the locomotion activity (d) were also recorded. Bars indicate the mean of values corresponding to eight to 12 mice  $\pm$ SEM. Asterisks denote significant differences between the groups (\* $p$ <0.05, \*\* $p$ <0.01, \*\*\* $p$ <0.001) according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test




## Discussion

More than a century after its description, the progressive myoclonus epilepsy of Lafora type is still a fatal disease leading to the death of patients usually within the decade after the apparition of the first symptoms. Unfortunately, up to now, available treatments are only palliatives and no curative drugs are available yet. Recent reports indicate that the elimination of the capacity of the cells to synthesize glycogen, either by depleting glycogen synthase or the protein targeting to glycogen (PTG) in animal models of LD [35–38], results not only in the elimination of the accumulation of polyglucosans but also in the amelioration of the neuropathological symptoms. These results have led those authors to suggest the use of glycogen synthase inhibitors as a putative strategy to treat LD. While the field awaits for the beneficial effects that these drugs could have in the future, in this work, we have used an alternative strategy to find a beneficial treatment for LD based on the restoration of proteostasis, which is defective in this disease [9–11]. We describe the use of trehalose and 4-PBA, two chemical chaperones with recognized abilities to ameliorate misfolding of proteins [17–21]. In addition, we have checked the effect of metformin, an activator of AMP-activated protein kinase (AMPK), as activation of this kinase plays a neuroprotective role in different neurodegenerative diseases [24–27].

We first observed that the designed treatments affected the corresponding molecular targets in the brain of a mouse model of LD (*Epm2b*<sup>-/-</sup>, lacking malin): trehalose and 4-PBA increased the levels of the chaperone BiP/Grp78, involved in the amelioration of proteostasis dysfunction, and metformin induced the activation of the AMPK complex. Then, we analyzed the possible beneficial effect of these drugs on LD symptoms. We found that the administration of the chemical chaperone 4-PBA reduced the levels of polyglucosan (PAS+) and polyubiquitinated protein aggregates in different areas of the brain of *Epm2b*<sup>-/-</sup> treated mice. This beneficial effect could be due to its action as a chemical chaperone or to its ability to improve the expression of the endogenous chaperone BiP/GRP78. The improved chaperone content of the cell could diminish the formation of intracellular aggregates. The use of trehalose, another small molecule that acts as a chemical chaperone, also improved the expression of BiP/GRP78 but did not have a beneficial effect in reducing intracellular aggregates, perhaps because its role as a chemical chaperone could not be as potent as 4-PBA. Trehalose and 4-PBA were effective in reducing gliosis in the different parts of the hippocampus, but only 4-PBA had in addition a beneficial effect in preventing neuronal loss. The different histological effect of these two drugs had a reflection in the performance of the treated animals when subjected to different neuropsychological

**Fig. 8** Open field analyses of *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with the described drugs during 1 month (a, c, e, g) or 2 months (b, d, f, h). No significant differences between groups ( $n=8-12$  mice in each group) were observed in all recorded parameters: total distance roamed (a, b), total immobility time (c, d), number of entries in the more central zone (zone 3) (e, f), or the distance roamed in zone 3 (g, h), according to Kruskal-Wallis non-parametric test followed by Conover-Inman post hoc test




tests: whereas 4-PBA-treated animals improved the tail suspension and the pole tests, trehalose had only a beneficial effect on the pole test.

We also describe the beneficial effect of metformin, an activator of AMPK, in the treatment of LD. We observed a clear reduction in the number of polyglucosan (PAS<sup>+</sup>) and polyubiquitinated protein aggregates, a reduction of neuronal loss and gliosis in different hippocampal regions, and an

amelioration of the neuropsychological alterations of *Epm2b*<sup>-/-</sup> treated mice. One possible explanation for the positive effect of metformin on LD could be that it maintained a sustained activation of AMPK, and this resulted in neuroprotection. As AMPK is involved in the activation of the autophagy pathway, we checked whether treatment with metformin could have an effect on autophagy, but as shown in Supplementary Fig. S1, we observed no changes in the levels of two

**Fig. 9** Beam balance analyses of *Epm2b*<sup>+/-</sup> (control) and *Epm2b*<sup>-/-</sup> (KO) mice untreated (w/o TTT) or treated with the described drugs during 1 month (a) or 2 months (b). No significant differences between groups ( $n=8-12$  mice in each group) were observed in the time to cross the beam in any of the beam sizes analyzed (5, 12, and 26 mm), according to Kruskal-Wallis non-parametric test followed by Conover-Imman post hoc test



recognized autophagy markers, namely LC3B-II and p62, upon treatment (trehalose and 4-PBA treatment did not affect these parameters either). So, an enhancement of autophagy is not responsible for the beneficial effects of these compounds. On the other hand, activation of AMPK may inhibit glycogen synthase [39]. However, upon metformin treatment, we did not observe changes in the state of different enzymes related to glycogen homeostasis (protein levels and phosphorylated forms of glycogen synthase and glycogen synthase kinase 3 $\beta$  were similar in all the cases, Supplementary Fig. S2; trehalose and 4-PBA treatment did not affect these parameters either). These results are in agreement with previous reports

which indicate that the specific activity of glycogen synthase does not change in the brain of LD mouse models in comparison to controls, especially at young ages as the ones used in our studies (5-month-old animals at the end of the treatments) [40–42]. Thus, we conclude that the beneficial effect of metformin in LD could be due to alternative mechanisms.

In conclusion, we describe for the first time the beneficial effect of the administration of small-molecule compounds on the neuropathological symptoms of LD. It is encouraging to see that the beneficial effect of these drugs is already present after 1 month of treatment and much better results are obtained if the treatment is prolonged for 2 months. As these compounds

have good safety records and are already approved for clinical use on different neurological pathologies, we think their use in clinical practice based on drug repositioning could be straightforward.

**Acknowledgments** We want to thank Carla Rubio-Villena and Maria Adelaida Garcia-Gimeno for their help in the analyses of glycogen-related enzymes. This work was supported by grants from the Spanish Ministry of Education and Science (SAF2011-27442), Fundació La Marató de TV3 (ref. 100130), and an ACCI2012 action from CIBERER. A.B. holds a postdoctoral fellowship from the Program “Junta para la Ampliación de Estudios” (JAE-Doc) co-funded by the European Social Fund (ESF).

## References

- Delgado-Escueta AV (2007) Advances in Lafora progressive myoclonus epilepsy. *Curr Neurol Neurosci Rep* 7:428–433
- Monaghan TS, Delanty N (2010) Lafora disease: epidemiology, pathophysiology and management. *CNS Drugs* 24:549–561
- Minassian BA, Lee JR, Herbrick JA, Huizenga J, Soder S et al (1998) Mutations in a gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy. *Nat Genet* 20:171–174
- Serratos JM, Gomez-Garre P, Gallardo ME, Anta B, de Bernabe DB et al (1999) A novel protein tyrosine phosphatase gene is mutated in progressive myoclonus epilepsy of the Lafora type (EPM2). *Hum Mol Genet* 8:345–352
- Chan EM, Young EJ, Ianzano L, Munteanu I, Zhao X et al (2003) Mutations in NHLRC1 cause progressive myoclonus epilepsy. *Nat Genet* 35:125–127
- Lohi H, Ianzano L, Zhao XC, Chan EM, Turnbull J et al (2005) Novel glycogen synthase kinase 3 and ubiquitination pathways in progressive myoclonus epilepsy. *Hum Mol Genet* 14:2727–2736
- Vilchez D, Ros S, Cifuentes D, Pujadas L, Valles J et al (2007) Mechanism suppressing glycogen synthesis in neurons and its demise in progressive myoclonus epilepsy. *Nat Neurosci* 10:1407–1413
- Solaz-Fuster MC, Gimeno-Alcaniz JV, Ros S, Fernandez-Sanchez ME, Garcia-Fojeda B et al (2008) Regulation of glycogen synthesis by the laforin-malin complex is modulated by the AMP-activated protein kinase pathway. *Hum Mol Genet* 17:667–678
- Liu Y, Wang Y, Wu C, Liu Y, Zheng P (2009) Deletions and missense mutations of EPM2A exacerbate unfolded protein response and apoptosis of neuronal cells induced by endoplasmic reticulum stress. *Hum Mol Genet* 18:2622–2631
- Vernia S, Rubio T, Heredia M, Rodriguez de Cordoba S, Sanz P (2009) Increased endoplasmic reticulum stress and decreased proteasomal function in Lafora disease models lacking the phosphatase laforin. *PLoS One* 4:e5907
- Zeng L, Wang Y, Baba O, Zheng P, Liu Y (2012) Laforin is required for the functional activation of malin in endoplasmic reticulum stress resistance in neuronal cells. *FEBS J* 279:2467–2478
- Rao SN, Maity R, Sharma J, Dey P, Shankar SK et al (2010) Sequestration of chaperones and proteasome into Lafora bodies and proteasomal dysfunction induced by Lafora disease-associated mutations of malin. *Hum Mol Genet* 19:4726–4734
- Aguado C, Sarkar S, Korolchuk VI, Criado O, Vernia S et al (2010) Laforin, the most common protein mutated in Lafora disease, regulates autophagy. *Hum Mol Genet* 19:2867–2876
- Criado O, Aguado C, Gayarre J, Duran-Trio L, Garcia-Cabrero AM et al (2012) Lafora bodies and neurological defects in malin-deficient mice correlate with impaired autophagy. *Hum Mol Genet* 21:1521–1533
- Hetz C, Mollereau B (2014) Disturbance of endoplasmic reticulum proteostasis in neurodegenerative diseases. *Nat Rev Neurosci* 15:233–249
- Sinadinos C, Valles-Ortega J, Boulan L, Solsona E, Tevy MF et al (2014) Neuronal glycogen synthesis contributes to physiological aging. *Aging Cell* 13:935–945
- Ricobaraza A, Cuadrado-Tejedor M, Perez-Mediavilla A, Frechilla D, Del Rio J et al (2009) Phenylbutyrate ameliorates cognitive deficit and reduces tau pathology in an Alzheimer’s disease mouse model. *Neuropsychopharmacology* 34:1721–1732
- Wiley JC, Meabon JS, Frankowski H, Smith EA, Schecterson LC et al (2010) Phenylbutyric acid rescues endoplasmic reticulum stress-induced suppression of APP proteolysis and prevents apoptosis in neuronal cells. *PLoS One* 5:e9135
- Zhou W, Bercury K, Cummiskey J, Luong N, Lebin J et al (2011) Phenylbutyrate up-regulates the DJ-1 protein and protects neurons in cell culture and in animal models of Parkinson disease. *J Biol Chem* 286:14941–14951
- Tanaka M, Machida Y, Niu S, Ikeda T, Jana NR et al (2004) Trehalose alleviates polyglutamine-mediated pathology in a mouse model of Huntington disease. *Nat Med* 10:148–154
- Chen Q, Haddad GG (2004) Role of trehalose phosphate synthase and trehalose during hypoxia: from flies to mammals. *J Exp Biol* 207:3125–3129
- Hardie DG, Ross FA, Hawley SA (2012) AMPK: a nutrient and energy sensor that maintains energy homeostasis. *Nat Rev Mol Cell Biol* 13:251–262
- Carling D, Thornton C, Woods A, Sanders MJ (2012) AMP-activated protein kinase: new regulation, new roles? *Biochem J* 445:11–27
- Poels J, Spasic MR, Callaerts P, Norga KK (2009) Expanding roles for AMP-activated protein kinase in neuronal survival and autophagy. *Bioessays* 31:944–952
- Han Y, Xie N, Cao L, Zhao X, Liu X et al (2011) Adenosine monophosphate-activated protein kinase and peroxisome proliferator-activated receptor gamma coactivator 1alpha signaling provides neuroprotection in status epilepticus in rats. *Neurosci Lett* 500:133–138
- Dulovic M, Jovanovic M, Xilouri M, Stefanis L, Harhaji-Trajkovic L et al (2014) The protective role of AMP-activated protein kinase in alpha-synuclein neurotoxicity in vitro. *Neurobiol Dis* 63:1–11
- Ashabi G, Khodagholi F, Khalaj L, Goudarzvand M, Nasiri M (2014) Activation of AMP-activated protein kinase by metformin protects against global cerebral ischemia in male rats: interference of AMPK/PGC-1alpha pathway. *Metab Brain Dis* 29:47–58
- Steru L, Chermat R, Thierry B, Simon P (1985) The tail suspension test: a new method for screening antidepressants in mice. *Psychopharmacology (Berl)* 85:367–370
- Ogawa N, Hirose Y, Ohara S, Ono T, Watanabe Y (1985) A simple quantitative bradykinesia test in MPTP-treated mice. *Res Commun Chem Pathol Pharmacol* 50:435–441
- Stanford SC (2007) The Open Field Test: reinventing the wheel. *J Psychopharmacol* 21:134–135
- Luong TN, Carlisle HJ, Southwell A, Patterson PH (2011) Assessment of motor balance and coordination in mice using the balance beam. *J Vis Exp* 49:2376
- R-Core-Team (2014) A language and environment for statistical computing. Foundation for Statistical Computing <http://www.R-project.org/>
- Valles-Ortega J, Duran J, Garcia-Rocha M, Bosch C, Saez I et al (2011) Neurodegeneration and functional impairments associated

- with glycogen synthase accumulation in a mouse model of Lafora disease. *EMBO Mol Med* 3:667–681
34. Ortolano S, Vieitez I, Agis-Balboa RC, Spuch C (2014) Loss of GABAergic cortical neurons underlies the neuropathology of Lafora disease. *Mol Brain* 7:7
  35. Turnbull J, DePaoli-Roach AA, Zhao X, Cortez MA, Pencea N et al (2011) PTG depletion removes Lafora bodies and rescues the fatal epilepsy of Lafora disease. *PLoS Genet* 7:e1002037
  36. Pederson BA, Turnbull J, Epp JR, Weaver SA, Zhao X et al (2013) Inhibiting glycogen synthesis prevents Lafora disease in a mouse model. *Ann Neurol* 74:297–300
  37. Turnbull J, Epp JR, Goldsmith D, Zhao X, Pencea N et al (2014) PTG protein depletion rescues malin-deficient Lafora disease in mouse. *Ann Neurol* 75:442–446
  38. Duran J, Gruart A, Garcia-Rocha M, Delgado-Garcia JM, Guinovart JJ (2014) Glycogen accumulation underlies neurodegeneration and autophagy impairment in Lafora disease. *Hum Mol Genet* 23:3147–3156
  39. Carling D, Hardie DG (1989) The substrate and sequence specificity of the AMP-activated protein kinase. Phosphorylation of glycogen synthase and phosphorylase kinase. *Biochim Biophys Acta* 1012:81–86
  40. Tagliabracci VS, Girard JM, Segvich D, Meyer C, Turnbull J et al (2008) Abnormal metabolism of glycogen phosphate as a cause for Lafora disease. *J Biol Chem* 283:33816–33825
  41. DePaoli-Roach AA, Tagliabracci VS, Segvich DM, Meyer CM, Irimia JM et al (2010) Genetic depletion of the malin E3 ubiquitin ligase in mice leads to Lafora bodies and the accumulation of insoluble laforin. *J Biol Chem* 285:25372–25381
  42. Turnbull J, Wang P, Girard JM, Ruggieri A, Wang TJ et al (2010) Glycogen hyperphosphorylation underlies Lafora body formation. *Ann Neurol* 68:925–933