

Norepinephrine and Glucocorticoids Modulate Chronic Unpredictable Stress-Induced Increase in the Type 2 CRF and Glucocorticoid Receptors in Brain Structures Related to the HPA Axis Activation

Marilia B. Malta¹ · Joelcimar Martins² · Leonardo S. Novaes¹ · Nilton B. dos Santos¹ · Luciane Sita³ · Rosana Camarini¹ · Cristoforo Scavone¹ · Jackson Bittencourt^{3,4} · Carolina D. Munhoz¹

Received: 2 February 2021 / Accepted: 22 June 2021 / Published online: 2 July 2021 © The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

The stress response is multifactorial and enrolls circuitries to build a coordinated reaction, leading to behavioral, endocrine, and autonomic changes. These changes are mainly related to the hypothalamus–pituitary–adrenal (HPA) axis activation and the organism's integrity. However, when self-regulation is ineffective, stress becomes harmful and predisposes the organism to pathologies. The chronic unpredictable stress (CUS) is a widely used experimental model since it induces physiological and behavioral changes and better mimics the stressors variability encountered in daily life. Corticotropin-releasing factor (CRF) and glucocorticoids (GCs) are deeply implicated in the CUS-induced physiological and behavioral changes. Nonetheless, the CUS modulation of CRF receptors and GR and the norepinephrine role in extra-hypothalamic brain areas were not well explored. Here, we show that 14 days of CUS induced a long-lasting HPA axis hyperactivity evidenced by plasmatic corticosterone increase and adrenal gland hypertrophy, which was dependent on both GCs and NE release induced by each stress session. CUS also increased CRF₂ mRNA expression and GR protein levels in fundamental brain structures related to HPA regulation and behavior, such as the lateral septal nucleus intermedia part (LSI), ventromedial hypothalamic nucleus (VMH), and central nucleus of the amygdala (CeA). We also showed that NE participates in the CUS-induced increase in CRF₂ and GR levels in the LSI, reinforcing the *locus coeruleus* (LC) involvement in the HPA axis modulation. Despite the CUS-induced molecular changes in essential areas related to anxiety-like behavior, this phenotype was not observed in CUS animals 24 h after the last stress session.

Keywords Chronic stress \cdot Lateral septum \cdot Glucocorticoids \cdot Type-2 CRF receptor \cdot Norepinephrine \cdot Anxiety-like behavior

Carolina D. Munhoz cdmunhoz@usp.br

- ¹ Department of Pharmacology, Universidade de Sao Paulo Instituto de Ciencias Biomedicas, Av. Prof. Lineu Prestes, 1524, room 323, São Paulo, SP 05508-000, Brazil
- ² Central of Facilities, Universidade de Sao Paulo Instituto de Ciencias Biomedicas, São Paulo 05508-000, Brazil
- ³ Laboratory of Chemical Neuroanatomy, Department of Anatomy, Universidade de Sao Paulo Instituto de Ciencias Biomedicas, São Paulo 05508-000, Brazil
- ⁴ Center for Neurosciences and Behavior, Institute of Psychology, University of São Paulo, São Paulo 05508-000, Brazil

Introduction

Physical and psychological stressors recruit different brain nuclei to build a coordinated stress response that activates the neuroendocrine and autonomic systems and causes behavioral changes [1]. Such response enfolds a conserved and efficient interconnection of several systems to maintain the organism's integrity [2, 3]. However, when this self-regulation is ineffective, stress becomes harmful and predisposes the body to diseases, including cardiovascular, immune, and psychiatric disorders [4, 5].

Several brain regions, such as the paraventricular nucleus of the hypothalamus (PVH), septal-hippocampus complex, amygdalar nuclei, pre-frontal and cingulate cortices, *locus coeruleus* (LC), and parabrachial and raphe nuclei, orchestrate the stress response [6]. The converging signals activate

the hypothalamus–pituitary–adrenal (HPA) axis. The PVH releases the corticotropin-releasing factor (CRF), which results in the synthesis and release of the adrenocortico-tropic hormone (ACTH) by the anterior pituitary, ultimately leading to glucocorticoids (GCs) synthesis and release by the adrenal glands [7–9].

The CRF actions in the central nervous system (CNS) and the periphery are critical to integrating and coordinating several physiological systems. During stress adaptation, these complex responses are fine-tuned by several CRFrelated peptides, including the urocortins (UCNs) that exert complementary or sometimes contrasting actions to CRF itself [10]. The CRF peptides (CRF and urocortins) act by binding to two different G-protein coupled receptors, the CRF type 1 and type 2 receptors (CRF₁ and CRF₂), which play different roles in stress responses. The outcomes of CRF receptor activation depend on the ligands and brain nuclei and decisively dictate the stress-induced behavioral state [reviewed on [11-14]]. For example, CRF₁ activation has been associated with the endocrine effects of CRF and behavioral response to stress, whereas the role of CRF₂ is essential to enable physiological and psychological homeostasis and oppose the initial stress response linked to CRF₁ activation; also, CRF1 is more related to the acute phase of the stress response while CRF2 is related to the recovery phase [11–18]. In addition, data show that the relationship between CRF2 and anxiety or depressive behavior in animals subjected to chronic stress models depends on the intensity of the stressor stimuli and/or when these behaviors are evaluated after the stressor stimulus [19]. Particularly, the CRF₂ receptor's genetic deficiency increases anxiety-like behavior and the sensitivity to stress [15] but attenuates the responses to the stress induced by opiate withdrawal [20]. Thus, the contribution of CRF₂ on responses to different stressful events warrants investigation. Despite the importance of CRF₁ in the mediation of stress-induced anxiety or depressive-like behaviors, they will not be evaluated in this study.

In addition to the HPA axis activation, CRF also modulates the stress-induced sympathetic response through the recruitment of the LC, adrenal medulla, and peripheral sympathetic system and plays a significant role in integrating the central and peripheral norepinephrine (NE) release secondary to stressful events [reviewed on [13]], which is fundamental to mediate the neuroendocrine changes and to bring together the stress perception and physiological and behavioral alterations.

However, acute and chronic stress can use different pathways to change the control of the HPA axis and autonomic responses [1, 6, 21]. In a chronic stress situation, neurochemical evidence suggests a continued increase in the HPA and adreno-medullary axis excitability in addition to an augmented sensibility of the LC to CRF and to the enzyme tyrosine hydroxylase, which elevates NE synthesis (CRF-NE loop) [10]. Some of these changes could be associated with the loss of GC self-regulation in the pituitary and PVH. Furthermore, it was also reported neuronal atrophy in the hippocampus and frontal cortex and an increase in the CRF expression in the central nucleus of the amygdala (CeA) and autonomic nuclei, which resulted in the HPA axis hyperactivity and increased excitability of PVH [22].

The disruption of the GC negative feedback on the HPA axis results in its sustained activation, causing the maintenance of elevated systemic GCs levels. Consequently, the higher availability of these hormones allows them to reach brain structures such as the amygdala and the LC, whose projections reinforce the HPA axis activation and promote changes in behavior and normal physiology [21, 23, 24]. In fact, due to their pleiotropic action, the sustained increase in the systemic levels of GCs can trigger metabolic diseases and immunosuppression and contribute to the development of autoimmune diseases and mood disorders [25].

The chronic unpredictable stress (CUS) paradigm is a well-accepted experimental model of stress-induced brain plasticity and mood disorders, such as depressive-like behaviors [26]. The mixture of psychological and physical stressors and reduced chances of adaptation to the different stress stimuli present significant similarity to stressors encountered in daily life [27, 28]. Considering previous data of loss of HPA axis self-regulation in animals submitted to 14 days of CUS [29] and the few evidence of the NE role in CUS studies, the present work evaluates the changes in the CRF₂ and glucocorticoid receptors (GR) in the central nervous system (CNS) of rats exposed to CUS, the influence of GCs and NE signaling in those molecular changes, and their implications in CUS-induced anxiety-like behavior. We found that CUS increased CRF2 mRNA expression and GR levels in central brain structures related to the HPA axis, and both GCs and NE signaling modulated these effects. Interestingly, CUS animals did not show any anxiety-like behavior in the elevated plus-maze (EPM) and open-field (OF) tests.

Materials and Methods

Animals

Animals (adult male Wistar rats, 60 days old at the beginning of the experiments) from the Facility for SPF Rat Production at the Institute of Biomedical Sciences–Animal Facility Network at the University of São Paulo were used following the standards of the Ethics Committee for Animal Use of the Institute of Biomedical Sciences/University of São Paulo (CEUA- ICB 102/06 and 75/05) and the guidelines of the Brazilian National Council for the Control of Animal Experimentation (CONCEA). All efforts were taken to minimize animal suffering and reduce the number of animals used to the minimum required for detection of significant statistical effects.

Chronic Unpredictable Stress Paradigm

Adult male Wistar rats housed in groups of 3 or 4 with ad libitum rat chow and water were kept under a 12 h light/ dark cycle. The animals were randomly assigned to either control (non-stressed) or CUS group and submitted or not to the different pharmacological treatments. The CUS protocol was based on [29]. Briefly, the animals were subjected to diverse psychological and physical stressors for 14 days, applied as follows: day 1 (2:00 p.m.) restraint, 60 min; day 2 (9:00 a.m.) forced swim, 15 min; day 3 (3:00 p.m.) cold isolation, 90 min; day 4 (7:00 p.m.) lights on, overnight; day 5 (10:00 a.m.) forced swim, 5 min; day 6 (7:00 p.m.) water and food deprivation, overnight; day 7 (2:00 p.m.) restraint, 120 min; day 8 (3:00 p.m.) light off, 120 min; day 9 (9:00 a.m.) forced swim, 5 min; day 10 (7:00 p.m.) lights on, overnight; day 11 (2:00 p.m.) cold isolation, 90 min; day 12 (9:00 a.m.) restraint, 60 min; day 13 (7:00 p.m.) water and food deprivation, overnight; day 14 (9:00 a.m.) restraint, 60 min. In all CUS experiments, stress stimuli were applied in the same order described above. All animals were studied 24 h after the last stressor. The control rats were manipulated every day for 10 min in their home cages to avoid nonspecific handling effects.

Pharmacological Treatments

Metyrapone (2-methyl-1,2-di-3-pyridyl-1-propane; Sigma), a steroid 11 β -inhibitor, was first dissolved in 100% ethanol and then diluted (1:10) in (NaCl 0.9%). Metyrapone (Met; 50 mg/kg) was administered intraperitoneally (i.p.) 30 min before each stress session according to [30]. We attempt to avoid a dose that could interfere with the circadian cycle of glucocorticoid synthesis.

Alpha and beta noradrenergic antagonists, phentolamine (P) and atenolol (A), respectively, were individually dissolved in 0.9% saline and concomitantly (AP) injected i.p. 20 min before each stress section in the doses of 5 mg/kg and 2.5 mg/kg respectively [31]. The treatment with both antagonists was chosen considering the presence, distribution, and action of alpha and beta noradrenergic receptors in the body and brain areas, which allowed us to blunt all noradrenergic influences regardless of whether central or peripheral.

For control of the respective pharmacological treatments, non-stressed animals were administered with Met or AP at the same time point of CUS animals and returned to the animal facility.

Experimental Groups and Design

We randomly divided the animals into the following groups:

- Control: no stressed animals
- CUS: animals submitted to chronic unpredictable stress
- C + Met: control animals submitted to chronic metyrapone treatment
- C + AP: control animals submitted to chronic atenolol and phentolamine treatment
- CUS + Met: animals submitted to CUS and treated chronically with metyrapone
- CUS + AP: animals submitted to CUS and treated chronically with atenolol and phentolamine

Euthanasia and Perfusion

Twenty-four hours after the last stress session, the animals were anesthetized with an excess of isoflurane and perfused via the ascending aorta with $\cong 100 \text{ mL}$ of cold 0.9% saline in 1 min, followed by \cong 900 mL of 4% formaldehyde in dH₂O at 4 °C, pH 9.5, for approximately 25 min. After removed from the skull, the brains were post-fixed for 3 h in the same fixative solution plus 20% sucrose and then transferred to RNAse free 20% sucrose in phosphate buffer solution at 4 °C. Regularly spaced brain series $(5 \times 1 \text{-in-5})$ of 30-µm-thick frozen sections were cut in freezing microtome (SM2000R, Leica, Wetzlar, HE, Germany) in the coronal plane, collected in ethylene glycol-based cryoprotectant solution, and stored at - 20 °C until assayed. To preserve RNA integrity for in situ hybridization, all plastics and solutions were RNAse free. Solutions were prepared fresh with DNAse and RNAse free reagent and miliQ water.

Plasma CORT Concentration

Plasma corticosterone (CORT) was measured using an enzyme-linked immunoassay (Enzo Life Sciences). Before the saline perfusion, cardiac blood was collected, transferred to conical tubes (with heparin 10% v/v), and centrifuged at 3500 rpm for 15 min to obtain plasma samples. CORT titers were assessed following the manufacturer's instructions. Plasma samples were diluted at 1:30 in assay buffer before use.

In Situ Hybridization for the CRF₂ in Rat Brain

The in situ hybridization for CRF_2 was carried out using a ³⁵S-labeled antisense cRNA probe from plasmid kindly provided by Paul Sawchenko (The Salk Institute for Biological Studies, La Jolla, CA, USA). Techniques for probe synthesis, hybridization, and autoradiographic localization of mRNA signal were adapted from [32]. In brief, brain sections were mounted onto electrostatically charged slides (Superfrost plus®, Fisher Scientific, USA) and digested with 10 µg/ml of proteinase K for 30 min at 37 °C. The labeled probes (sense and antisense) were used at concentrations of 10^6 cpm/ml and applied to sections overnight at 56 °C in a solution containing 50% formamide, 0.6 M NaCl, 10 mM Tris, pH 8.0; 1 mM EDTA, 0.05% tRNA, 10 mM dithiothreitol, Denhardt's solution, and 10% dextran sulfate. They were treated with 20 µg/ml of ribonuclease A for 30 min at 37 °C and washed in 15 mM NaCl/1.5 mM sodium citrate at 55-60 °C. Sections were dehydrated and exposed to X-ray films for 1-2 days. The slides were then dipped in Kodak NTB-2 liquid autoradiographic emulsion, dried, and exposed at 4 °C in the dark for 20 days. They were then developed with Kodak D-19 and counterstained with thionine (Nissl method) for anatomical reference purposes. Sections were dehydrated, defatted, and coverslipped with DPX mounting medium (Aldrich, Milwaukee, WI, USA). For CRF₂ mRNA expression, the semiquantitative densitometric analysis was performed on emulsion-coated sections. Slides were coded and analyzed blinded. Densitometric analysis of the images was performed under dark-field illumination with the Image-Pro Plus software. We evaluated the CRF₂ mRNA expression in the lateral septal nucleus intermediate part (LSI), and ventromedial hypothalamic nucleus (VMH) integrated optical density (IOD) in arbitrary units (a.u.). The results are expressed by mean of $IOD \pm SEM$ (a.u., n = 6/group) at medial levels of LSI and VMH (0.20 and - 3.14 mm from Bregma for LSI and VMH, respectively).

Immunohistochemistry for GR

Immunohistochemistry was performed using a conventional avidin-biotin immunoperoxidase protocol [33] and Vectastain Elite reagents. The free-floating sections were pretreated with hydrogen peroxide, blocked in 3% normal donkey serum, and incubated overnight at room temperature with anti-GR polyclonal primary antisera raised in rabbit (1:3000 sc-1004 GR Antibody (M-20) Santa Cruz). Sections were incubated for 1 h in biotin-conjugated IgG donkey anti-rabbit (1:800, Jackson Laboratories) and 1 h in avidin-biotin complex (1:333, kit elite Vector Labs). The peroxidase reaction was performed using 0.05% DAB and 0.025% nickel sulfate as chromogens, and 0.03% hydrogen peroxide. Sections were mounted onto gelatin-coated slides, dehydrated, defatted, and coverslipped with DPX mounting medium (Aldrich, Milwaukee, WI, USA). To evaluate the GR levels, we quantified the stained cells in the LSi, CeA, bed nucleus of the stria terminalis (BST), and PVH. The results are expressed as a mean of GR immunoreactive cells \pm SEM. The total area was defined by the product between the number of sections analyzed and the area in a $20 \times$ magnification for each nucleus (0.278 mm² for LSi and CeA; 0.06 mm² for and 0.04 mm² for PVH).

For the CRF_2 -GR co-localization experiments, the immunohistochemistry for GR was performed as described and after the peroxidase reaction, the sections were mounted onto electrostatically charged slides (Fisher Superfrost Plus®, Fisher Scientific, USA). The in situ hybridization for CRF_2 was performed as described earlier.

Behavioral Studies

To correlate the CUS-induced biochemical changes to the behavioral phenotype, the animals were examined for anxiety-like behavior using the EPM and OF tests 24 h after the last stress session in a non-aversive environment and after the acclimatization period.

The EPM apparatus consisted of two closed and two open arms with a central area (100 cm^2) where the animals were positioned at the beginning of the test. The apparatus was made of dark blue painted wood elevated 65 cm from the floor and located in a soundproof test room under indirect light. A video camera was located above the center of the maze to record the rats. Anxiety-like behavior was assessed as a function of the decreased open arm exploration, measured by the total percentage of time at the open arms.

The OF apparatus consists of a circular arena (60 cm diameter with a central compartment of 30 cm) with opaque walls (50 cm high). The floor was divided into squares, which allowed the discrimination of two compartments, the central and peripheral. Each rat was placed on the periphery of the arena, and its activity was recorded for 5 min. This test was used to evaluate the anxiety-like behavior based on rodents' preference to spend a significantly higher amount of time exploring the periphery of the arena than the unprotected center area.

In both tests, the animals explored the apparatuses for 5 min. The behavior was evaluated by the EthoVision System (Noldus Information Technology). Results are expressed as mean of percentage of the time that each group spent in the arms \pm SEM (%) for the EPM and as a mean of the time in central zone \pm SEM (%) for the OF. Different groups of animals were used for each test, and they were not previously exposed to neither EPM nor OF. All trials were videotaped, and the apparatus was cleaned with 5% (vol/vol) ethanol after each trial to avoid olfactory clues.

Statistical Analysis

The results are expressed as mean \pm SEM and statistical analysis was calculated using the GraphPad Prism Software, version 8.0. For the in situ hybridization and immunohistochemistry, the samples from all groups were assayed simultaneously (n=6 animals/group). For serum

corticosterone measurements, samples were distributed as follows: Corticosterone measurement 24 h after the last stressor stimulus: Control=9; C+AP=6; C+MET=11; CUS=6; CUS+AP=10; CUS+MEt=8; and Corticosterone measurement 72 h after the last stressor stimulus: n=23 per group. For behavioral analysis, we had a n=5-7animals/group.

Under the guidance of the statistical team of the Institute of Mathematics and Statistics (IME-USP), we adopted the Kruskal Wallis test [34] to analyze the results from in situ hybridization (CRF₂ mRNA expression) and immunohistochemistry (GR levels). We used the two-tailed Student t-test to analyze the behavior. The ROUT method (Q = 1%) was performed to identify outliers. However, there was no exclusion, and all data collected were included in the statistical analysis. The Shapiro–Wilk normality test was performed. Values of p < 0.05 were considered statistically significant.

Results

CUS Persistently Increased Plasma CORT Levels Through GCs and NE Signaling

CUS augmented plasma CORT levels 24 h after the last stress session $(14.9 \pm 2.2 \ \mu g/dL, p < 0.05)$ when compared to control rats $(4.9 \pm 0.95 \ \mu g/dL)$ (Fig. 1). This effect was accompanied by the hypertrophy of the adrenal gland $(17.4 \pm 0.8 \ and 14.2 \pm 0.8\%)$ of adrenal gland CUS and control, respectively, p < 0.05) in the CUS animals. Chronic treatment with Met (CUS + Met., $2.6 \pm 0.7 \ \mu g/dL)$ and AP (CUS + AP, $3.9 \pm 1.8 \ \mu g/dL)$ blunted the increase of CORT

levels in CUS animals without altering CORT levels of non-stressed group in the same time point $(4.9 \pm 1.0 \ \mu\text{g/dL}, \ p < 0.05)$. At 72 h post-stress, both CORT levels $(2.5 \pm 0.4 \ \mu\text{g} / \text{dL})$ and adrenal glands $(14.5 \pm 0.5\% \text{ of adrenal gland})$ were no different from non-stressed animals in the same time point $(3.5 \pm 0.5 \ \mu\text{g} / \text{dL})$ and $14.2 \pm 0.8\%$ of adrenal gland).

CUS Persistently Increased CRF₂ mRNA Expression in the LSi and VMH, an Effect Dependent on GCs and NE Signaling

Using the in situ hybridization assay, we observed an increase of the CRF₂ mRNA expression in the LSI and VMH of CUS animals $(1.4 \times 10^6 \pm 1.3 \times 10^5 \text{ a.u.} \text{ and } 2.4 \times 10^6 \pm 2.2 \times 10^5 \text{ a.u.}$ for LSI and VMH, respectively) when compared to the control group $(1.0 \times 10^6 \pm 9.4 \times 10^4 \text{ a.u.} \text{ and } 7.9 \times 10^5 \pm 1.0 \times 10^5 \text{ a.u.}$ LSI and VMH respectively, p < 0.05) (Fig. 2 and Table 1). Because GC levels were elevated 24 h after the last stress session, we tested whether this sustained CORT increase helped to mediate the effects of stress on CRF₂ mRNA expression. As shown in Fig. 2, blunting the stress-induced GCs increase with metyrapone (Met) reduced the CRF₂ mRNA expression in both LSI and VMH compared to CUS animals.

Noradrenergic innervation is widespread in the CNS, and its targets include regions implicated in stress-induced effects. We also observed that NE signaling could modulate plasmatic CORT levels (Fig. 1). Because of that, we tested whether the expected stress-induced NE increase contributed to the stress effects on CRF₂ mRNA expression. We observed that the antagonism of both α - and β -adrenergic receptors (phentolamine and atenolol, respectively) immediately





Fig. 1 CUS-induced persistently increase in plasma CORT levels is modulated by GC release and NE signaling. Plasma CORT from control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol and phentolamine (AP, 5 mg/

kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session) were collected 24 h after the last stress session. Results are presented as mean \pm SEM. Significance differences between groups are indicated as * p<0.05 vs control; # p<0.05 vs CUS, Kruskal Wallis and Mann Whitney tests (n=6 for each group)



√Fig.2 CUS-induced increase in the CRF₂ mRNA expression in hypothalamic and extra-hypothalamic brain areas is modulated by GC release and NE signaling. Representative photomicrography of CRF₂ mRNA expression by in situ hybridization assay and in the lateral septal nucleus intermedia part (A) and the ventromedial hypothalamic nucleus (B) in control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol and phentolamine (AP, 5 mg/kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session) at medial levels of LSI and VMH (0.20 and -3.14 mm from Bregma for LSI and VMH, respectively). The graphics illustrate the quantification of the integrated optical density (IOD) represented in the panels. Results are presented as mean ± SEM. Significance differences between groups are indicated as * p < 0.05 vs control; # p < 0.05 vs CUS, Kruskal Wallis and Mann Whitney tests (n=6 for each group). Abbreviations: LSI, lateral septal nucleus intermedia part; LSD, lateral septal nucleus dorsal part; LV, lateral ventricle; cc, corpus callosum; VMH, ventromedial hypothalamic nucleus; 3 V, 3rd ventricle

before each stress session reduced CRF_2 mRNA expression in both LSI (Fig. 2A) and VMH (Fig. 2B) when compared to the CUS animals, 24 h after the last stress session.

Chronic treatment with AP did not alter CRF_2 mRNA expression in LSI and VMH in control animals, although the administration of Met (C+Met, $1.7 \times 10^6 \pm 4.2 \times 10^4$ a.u.) increased the expression of CRF_2 mRNA in VMH when compared to control untreated animals (p < 0.05) (Table 1).

CUS Persistently Increased GR Expression in Brain Regions Related to GC Release Control, an Effect Dependent on GCs and NE Signaling

We evaluated the GR expression in several brain areas implicated in stress response. Our results showed that CUS increased the number of GR-immunoreactive cells in the LSI, PVH, CeA, and BST compared to the control group (p < 0.05). Unlike CRF₂ mRNA expression, the influence of GC or NE in CUS-induced GR levels varied according to the nucleus analyzed. For the LSI and CeA, both treatments (Met or AP) prevented the CUS-induced increase in GR protein levels (p < 0.05; Figs. 3A and 4B). In the BST, only the AP treatment blunted the CUS-induced GR levels increase, suggesting the dependence of NE for this effect and not GCs (Fig. 3B). In the PVH, the primary hypothalamic nucleus in controlling stress responses, both treatments did not prevent the CUS-induced increase in GR expression (p > 0.05) (Fig. 4A).

Chronic treatment with AP did not alter the GR levels in control animals in any nuclei analyzed. However, MET treatment increased GR levels in PVH when compared to control animals not pharmacologically treated (p < 0.05) (Fig. 3 and Table 2).

We next analyzed if the CRF_2 mRNA expression and GR were present in the same cells in the rat brain. Our results

showed both CRF_2 mRNA and GR in the same cells of LSI (Fig. 5).

CUS Did Not Induce Anxiety-Like Behavior When Tested 24 h After the Last Stress Session

Because of all these CUS-induced molecular changes and the relevance of GR and CRF₂ in modulating some behaviors in chronically stressed animals [18, 19, 25, 35], we analyzed the anxiety-like behaviors in our CUS paradigm using the EPM and OF tests after 24 h of the last stress session. In our CUS protocol, no anxiety-related behaviors were observed in CUS animals in both tests. In the EPM test (Fig. 6A), no difference in the time spent in the closed ($68\pm7\%$, CUS) and open arms ($16\pm6\%$, CUS) were observed when compared to control animals ($62\pm7\%$ and $21\pm7\%$, respectively), evaluated 24 h after the last stressor. Consistently, in the OF, the CUS animals did not present anxiety-like behavior (Fig. 6B) since no difference in the time spent in the central zone was observed in CUS group ($95\pm1.07\%$) when compared to control animals ($96.25\pm0.72\%$).

Discussion

In the present study, we showed that 14 days of CUS increased plasmatic CORT levels that persisted for up to 24 h after the last stress exposure. This long-lasting effect was also accompanied by the hypertrophy of adrenal glands, suggesting hyperactivity of the HPA axis in response to

Table 1CUS-induced increase in the CRF_2 mRNA expression inhypothalamic and extra-hypothalamic brain areas is modulated by GCrelease and NE signaling

CRF ₂ mRNA expression					
	LSI	VMH			
Control	$1.0 \times 10^6 \pm 9.4 \times 10^4$	$7.9 \times 10^5 \pm 1.0 \times 10^5$			
C Met	$6.6 \times 10^5 \pm 1.6 \times 10^5$	$1.7 \times 10^{6} \pm 4.2 \times 10^{4} *$			
C AP	$6.6 \times 10^5 \pm 1.4 \times 10^5$	$5.5 \times 10^5 \pm 1.2 \times 10^5$			
CUS	$1.4 \times 10^6 \pm 1.3 \times 10^5 *$	$2.4 \times 10^{6} \pm 2.2 \times 10^{5} *$			
CUS Met	$6.4 \times 10^5 \pm 1.7 \times 10^5 $ #	$6.4 \times 10^5 \pm 1.1 \times 10^5 \#$			
CUS AP	$6.5 \times 10^5 \pm 8.6 \times 10^4 $ #	$1.3 \times 10^6 \pm 2.7 \times 10^5 $ #			

CRF₂ mRNA expression was measured by in situ hybridization assay the in the lateral septal nucleus intermedia part (LSI) and the ventromedial hypothalamic nucleus (VMH) of control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol and phentolamine (AP, 5 mg/kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session). Results are presented as mean±SEM of the integrated optical density (IOD, arbitrary units). Significance differences between groups are indicated as * p <0.05 vs control; # p <0.05 vs CUS, Kruskal Wallis and Mann Whitney tests (n=6 for each group)

#

ф

-0,0

CUS Met

. .

CUS

#

Δ Δ

A A A

CUS AP

۵

CUS AP

#

CUS Met

cus






Fig. 3 CUS-induced increase in the GR protein levels in hypothalamic and extra-hypothalamic brain areas is modulated by GC release and NE signaling. Representative photomicrography of immunoreactive GR cells in the lateral septal nucleus intermedia part (A), bed nucleus of the stria terminalis (B) in control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol and phentolamine (AP, 5 mg/kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session). The graphics illustrate the mean of GR immunoreactive cells /mm². Significance differences between groups are indicated as * p < 0.05 vs control group; # p < 0.05 vs CUS group Kruskal Wallis and Mann Whitney tests (n=6for each group). Abbreviations: LSI, lateral septal nucleus intermedia part; LSD, lateral septal nucleus dorsal part; LV, lateral ventricle; cc, corpus callosum; BST, bed nucleus of the stria terminalis; ca, anterior commissure

CUS. The HPA axis hyperactivity has been reported in different stress contexts, mostly in the chronic ones, and can be considered a risk factor for developing systemic and neuropsychiatry diseases, such as depression and anxiety [36].

CUS-induced increase of plasmatic CORT levels has been related to a vicious cycle, which maintains the stress brain circuitry activated via GC actions on extra-HPA brain nuclei, due, at least in part, to the loss of the GC-mediated HPA axis negative feedback loop [37]. In this situation, according to [38], other brain nuclei, such as the medial amygdala and pre-frontal cortex, become essential in the HPA axis activation, and the following could be happening: (1) changes in HPA axis signaling, (2) amygdala hyperactivation, and (3) prefrontal cortex hypoactivation. These hypotheses are not mutually excluding, and the control of synthesis and release of GCs could be a multifaceted system, not exclusively dependent on the activation/inhibition of the HPA axis [38]. In addition to GCs, the NE released by LC and NTS could act in brain nuclei such as the amygdala and PVH to maintain HPA activity [12, 18, 39-41]. To our knowledge, for the first time, these results showed that the rise and release of NE are as vital as GCs, during every single stressor, to the establishment of the CUS-induced HPA axis hyperactivation.

The importance of NE signaling in controlling HPA axis activation is well known for acute stress paradigms [42]. However, identifying its participation in maintaining chronic hyperactivation of the HPA axis is relatively recent [43, 44]. Centrally released NE also plays an essential role in arousal and alertness [reviewed on [40]], and the dysregulation of the noradrenergic system could contribute to the susceptibility of stress-related illnesses. NE (systemically or centrally released) can excite CRF-containing cells in the PVH to activate the HPA axis. This effect is thought to be mainly mediated by α 1-adrenergic receptors, although a role for β -adrenergic receptors has not been excluded. It is suspected that NE may activate CRF-containing neurons in extrahypothalamic regions, such as the CeA, LSI, and

hippocampus, during stress. However, this warrants further investigations [11].

The interactions between CRF- and NE-containing neurons within the CNS are well established. Intracerebral administration of NE and adrenergic agonists can change the activity of CRF-containing neurons, and, conversely, administration of CRF can alter the activity of noradrenergic neurons [reviewed on [40]]. Here, we evidence the relevance of NE in this paradigm since our results showed that atenolol and phentolamine modulate the plasmatic CORT, GR levels, and CRF₂ mRNA expression in brain structures related to the HPA control in animals submitted to CUS. Considering that atenolol does not cross the blood-brain barrier [45] and phentolamine does [31], therefore, acting in the α -adrenergic receptors throughout the brain, we could imply the role of NE signaling peripherally and centrally in our study. Nonetheless, it is important to consider that stress can modify the blood-brain barrier permeability, allowing some atenolol to pass through it and to play a role centrally as well [45, 46]. Hence, we can infer that stress-induced NE release could, indeed, modulate GC levels, contributing to hyperactivation of HPA axis and the CUS-induced increase in CRF₂ expression and GR levels in the LSI. CRF acts through the CRF₁ and CRF₂ receptors, in the initial and recovering phases of the stress response, respectively [13, 18, 19]. The CUS-induced increase of CRF2 mRNA expression could be related to the recovery phase in the CUS animals, since 72 h after the last stress session, the plasmatic CORT levels and the adrenal gland hypertrophy were not observed.

Glucocorticoids themselves provide the dominant-negative feedback loop onto the HPA axis acting through MR and GR, at least in part, in the circuitry originated in the forebrain [47]. In the present study, we observed CUS upregulated GR levels in the LSI, PVH, BST, and CeA of the animals. All these nuclei are related to integrating the HPA axis and the limbic system [47]. Our results also suggest that the LSI could be one of the significant brain nuclei involved in this process since the CUS increased both CRF_2 mRNA and GR levels, not only in the same nucleus but also in the same cell. The LSI also directly innervates the hippocampus, the BST, and VMH, which could result in the CUS-induced HPA hyperactivation observed 24 h after the last stress session, contributing to the hypothesis proposed by Kling et al. (2009) regarding HPA axis activation in chronic stress [38].

The LC is one of the essential noradrenergic nuclei in the CNS directly activated by acute stress (reviewed in [48]). Our results suggest, for the first time, that in a chronic stressful situation, NE could act either systemically, by sympathetic activation, or centrally, by LS activation, in brain areas such as BTS, CeA, and PVH (CRF-NE feedback), contributing to the maintenance of the stress vicious cycle [49, 50]. Here, we showed that NE participates in the CUSinduced increase in GR levels in the LSI, reinforcing the




Fig. 4 CUS-induced increase in the GR protein levels in hypothalamic and extra-hypothalamic brain areas is modulated by GC release and NE signaling. Representative photomicrography of immunoreactive GR cells in the paraventricular hypothalamic nucleus (**A**), central amygdaloid nucleus (**B**) in control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol

and phentolamine (AP, 5 mg/kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session). The graphics illustrate the mean of GR immunoreactive cells /mm². Significance differences between groups are indicated as * p < 0.05 vs control group; # p < 0.05 vs CUS group Kruskal Wallis and Mann Whitney tests (n=6 for each group). Abbreviations: CeA, central amygdaloid nucleus; opt, optic tract; PVH, paraventricular hypothalamic nucleus; 3 V, 3rd ventricle

Table 2CUS-induced increasein the GR protein levels inhypothalamic and extra-hypothalamic brain areas aremodulated by GC release andNE signaling

	Glucocorticoid receptor level							
	Control	C+Met	C+AP	CUS	CUS + Met	CUS+AP		
LSI	162±9	235 ± 26	270 ± 30	$433 \pm 40^{*}$	259±25#	$262 \pm 25 \pm$		
CeA	260 ± 27	371 ± 43	331 ± 48	$566 \pm 55^{*}$	$346 \pm 24 \#$	309 <u>±</u> 11#		
BST	529 ± 82	$1008 \pm 53*$	822 ± 147	$1280 \pm 29^*$	$903 \pm 72 \#$	882±130		
PVH	814 ± 96	$1617 \pm 190 *$	1164 ± 259	$1850 \pm 144*$	$1615 \pm 235*$	$1881 \pm 48^{\circ}$		

The results illustrate the mean of GR immunoreactive cells /mm² in the lateral septal nucleus intermedia part (LSI), central amygdaloid nucleus (CeA), the bed nucleus of the stria terminalis (BST), and the paraventricular hypothalamic nucleus (PVH) of control and 14 days of chronic unpredictable stress (CUS) exposed animals treated or not with metyrapone (Met, 50 mg/kg, i.p., 30 min before each stress session) or atenolol and phentolamine (AP, 5 mg/kg and 2.5 mg/kg respectively, i.p., 20 min before each stress session). Significance differences between groups are indicated as * p<0.05 vs control group; # p<0.05 vs CUS group Kruskal Wallis and Mann Whitney tests (n=6 for each group)

LC involvement in the HPA axis modulation as proposed by [37].

Therefore, we hypothesized that the loss of HPA axis regulation observed in the present work could be, in part, due to the loss of inhibitory GABAergic projections from the BST to the hypothalamus, since the CUS-induced alterations in LSI and CeA, as well as in plasmatic CORT levels could block the BST-hypothalamus pathway. However, we cannot neglect the role of the CeA-PVH projection over the HPA axis activity [51, 52]. This projection can be altered since we observed that CUS induced an increase in the GR levels in CeA, which could be via CORT and NE signaling, as our treatment with Met and AP revealed. The molecular mechanism behind this NE participation is not fully elucidated. Nonetheless, our results point to the importance of alpha and beta-adrenergic receptors, as well as the increase of GCs (Fig. 7).

Several studies have shown the participation of GR and CRF_2 in specific brain regions such as the amygdala and lateral septal, in some behavioral alterations, including depression and anxiety-like behavior [13, 18, 19, 25, 35]. Despite the CUS-induced molecular plasticity in GR and CRF_2 , in

the paradigm used here, this chronic stress protocol did not induce anxiety-like behavior in rat 24 h after the last stress session. The anxiety-like behavior depends on CRF₁ and CRF₂ brain location, time, and intensity of stress [18]. Henry et al. (2006) tested the relation between CRF₂ signaling and the LS with the anxiety-like behavior. Their data indicate that this effect depends on stress levels since a high-stress environment, but not lower stress conditions, induced anxiety-like behavior [53]. In the present work, CUS animals did not show anxiety-like behavior that could be associated with a combination of moderate stress conditions, recovery interval between each session, and social support since the animals were grouped in 3 to 4 animals per cage throughout the experiments. The effects of social support in grouped rats are attributed to neural plasticity and some neurotransmitters release such as oxytocin, well known for its anxiolytic properties [54, 55].

Considering the absence of behavioral alterations and the increase of CRF_2 mRNA in the present study, we also suggest a possible protective role of CRF and urocortins in this CUS paradigm. Some studies correlate CRF_2 to the reestablishment of the stress response. They suggest that the

Fig. 5 Representative photomicrography of co-localized CRF₂ mRNA expression (silver grains) and GR (DAB, brown color) in the LSI cells indicated by the arrows. Abbreviations: LSI, lateral septal nucleus intermedia part; LSD, lateral septal nucleus dorsal part; LV, lateral ventricle; cc, corpus callosum. Scale bar: **A** 100 μm; **B** 10 μm





Fig. 6 CUS did not induce anxiety-like behavior in rats measured by the elevated plus maze (EPM) and open-field (OF) tests. Results are presented as mean \pm SEM of the percentage of the time spent in open and closed arms in the EMP test (**A**) and the time spent in the center zone of the OF test (**B**) 24 h after the last stress session. No significant differences between groups were observed. Student's t-test, p > 0.05; n = 12for EPM and n = 8 for OF



 CRF_2 activation is responsible for ensuring physiological and psychological homeostasis and counteracts the initial stress-response-provoking effects and anxiety-like behavior induced by CRF_1 activation [reviewed in [13]].

Prolonged or chronic stress changes the rules under which the body regulates homeostasis, requiring new strategies for successful adaptation. Only when the homeostatic pressure becomes too high, the body enters a state of genuine distress that can lead to morbidity and mortality. Thus, stress alters the physiological milieu over a long-term manner (adaptation through change, or allostasis), and the body's response to these changes lies at the center of both successful stress resilience as well as its transition to pathology [56]. Therefore, while responses to chronic stress can clearly have negative consequences, one must consider the possibility that they may solve the organisms' pressing problems at the expense of future success. The switch from adaptive to maladaptive stress responses will be heavily dependent on




Fig.7 Schematic representation summarizing the main findings. Full lines represent excitatory signaling, while dotted lines represent inhibitory signaling. We hypothesized that CUS-induced increase in GR and CRF_2 expression in LSI could be involved in the loss of HPA axis activity control, leading to sustained high CORT levels up

the individual's constitution, based on genetic and acquired strategies to efficiency and limit overdrive of stress systems.

In conclusion, the present study highlights that CRF_2 and GR, mainly in LSI, play a role in fine-tuning CUS responses, which depend on GC and NE signaling in males rats and future work should determine whether similar effects are observed in females. Although further studies are necessary to explore other CUS mood-related behavior and the mechanisms behind the results described in the present work, we could suggest that CUS-modulated molecular changes are related to animals' ability to cope with stressful events, which could result in a state of resilience for anxiety-like behavior.

Acknowledgements We gratefully thank Guiomar Wiesel and Larissa de Sá Lima for technical assistance.

Author Contribution M.B.M. designed research, performed research, analyzed data, wrote the paper, revised the paper final version; J.M. designed research, performed research, revised the paper final version;

to 48 h after stress. These molecular CUS-induced changes modulate the excitatory LSI projection to VMH, BST, and hypothalamus. In addition, the loss of inhibitory signaling from BST to the hypothalamus could be the result of CUS-induced alterations in the BST projections from LSI, hypothalamus, and CeA

L.S.N. performed research, analyzed data, revised the paper final version; N.B.S. performed research, analyzed data, revised the paper final version; L.S. analyzed data, contributed unpublished reagents/analytic tools, revised the paper final version; R.C: contributed unpublished reagents/analytic tools, revised the paper final version; C.S. contributed unpublished reagents/analytic tools, analyzed data, revised the paper final version; J.B. contributed unpublished reagents/analytic tools, analyzed data, revised the paper final version; C.D.M.: designed research, analyzed data, contributed unpublished reagents/analytic tools, wrote the paper and revised the paper final version.

Funding This article was supported by research grants to J.B. from Fundação de Amparo à Pesquisa do Estado de São Paulo (São Paulo Research Foundation — FAPESP) grants #2010/52068–0, #2016/02224–1. J.B. was also supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Agency for the Advancement of Higher Education) and Comité Français d'Evaluation de la Coopération Universitaire avec le Brésil (French Committee for the Evaluation of Academic and Scientific Cooperation with Brazil) grant CAPES-COFECUB 848/15. J.B. is an Investigator with the Conselho Nacional de Desenvolvimento Científico e Tecnológico (National Council for Scientific and Technological Development— CNPq) with grant #426378/2016–4. This work was supported by research grants to C.D.M. from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP: 2008/55178–0, 2012/24727–4, and 2016/03572–3) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq: 479153/2009–4 and 422523/2016–0). This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—Brasil (CAPES)—Finance Code 001. M.B.M. was supported by FAPESP (2006/52566–4) and CAPES. L.S.N. was supported by FAPESP (2010/13843–8 and 2012/24002–0). N.B.S. was supported by CNPq (160570/2012–3); C.D.M., C.S., and R.C. are research fellows from CNPq.

Data Availability (Data Transparency) The data that support the findings of this study are available from the corresponding author, upon reasonable request.

Code Availability (Software Application or Custom Code) Not applicable.

Declarations

Ethics Approval All animals were used following the standards of the Ethics Committee for Animal Use of the Institute of Biomedical Sciences/University of São Paulo (CEUA- ICB 102/06 and 75/05) and the guidelines of the Brazilian National Council for the Control of Animal Experimentation (CONCEA).

Consent to Participate Not applicable.

Consent for Publication All authors have contributed significantly and agreed with the entire content of this manuscript.

Competing interests The authors declare no competing interests.

References

- Ulrich-Lai YM, Herman JP (2009) Neural regulation of endocrine and autonomic stress responses. Nat Rev Neurosci 10(6):397–409
- de Kloet ER (2008) About stress hormones and resilience to psychopathology. J Neuroendocrinol 20(6):885–892
- Goldstein DS, McEwen B (2002) Allostasis, homeostats, and the nature of stress. Stress 5(1):55–58
- Bejean S, Sultan-Taieb H (2005) Modeling the economic burden of diseases imputable to stress at work. Eur J Health Econ 6(1):16–23
- 5. de Kloet ER, Joels M, Holsboer F (2005) Stress and the brain: from adaptation to disease. Nat Rev Neurosci 6(6):463–475
- Van de Kar LD, Blair ML (1999) Forebrain pathways mediating stress-induced hormone secretion. Front Neuroendocrinol 20(1):1–48
- Akil HA, Morano MI (1995) Stress. In: Bloom FE, Kupfer DJ (eds) psychopharmacology: the fourth generation of progress. Raven Press, New York, pp 773–785
- Romier C et al (1993) Solution of structure of human corticotropin releasing factor by HNMR and distance geometry with restrained molecular dynamics. Prot Eng 6:149–156
- 9. Trainer PJ et al (1995) A comparison of the effects of human and ovine corticotropin-releasing hormone on the pituitary-adrenal axis. J Clin Endocrinol Metab 80(2):412–417
- Hillhouse EW, Grammatopoulos DK (2006) The molecular mechanisms underlying the regulation of the biological activity of corticotropin-releasing hormone receptors: implications for physiology and pathophysiology. Endocr Rev 27(3):260–286

- Dunn AJ, Swiergiel AH, Palamarchouk V (2004) Brain circuits involved in corticotropin-releasing factor-norepinephrine interactions during stress. Ann N Y Acad Sci 1018:25–34
- Gallagher JP et al (2008) Synaptic physiology of central CRH system. Eur J Pharmacol 583(2–3):215–225
- Henckens MJ, Deussing JM, Chen A (2016) Region-specific roles of the corticotropin-releasing factor-urocortin system in stress. Nat Rev Neurosci 17(10):636–651
- Shekhar A et al (2005) Role of stress, corticotrophin releasing factor (CRF) and amygdala plasticity in chronic anxiety. Stress 8(4):209–219
- Bale TL et al (2000) Mice deficient for corticotropin-releasing hormone receptor-2 display anxiety-like behaviour and are hypersensitive to stress. Nat Genet 24(4):410–414
- Coste SC et al (2000) Abnormal adaptations to stress and impaired cardiovascular function in mice lacking corticotropinreleasing hormone receptor-2. Nat Genet 24(4):403–409
- Kozicz T et al (2004) Urocortin expression in the Edinger-Westphal nucleus is down-regulated in transgenic mice overexpressing neuronal corticotropin-releasing factor. Neuroscience 123(3):589–594
- Reul JM, Holsboer F (2002) Corticotropin-releasing factor receptors 1 and 2 in anxiety and depression. Curr Opin Pharmacol 2(1):23–33
- Bale TL, Vale WW (2004) CRF and CRF receptors: role in stress responsivity and other behaviors. Annu Rev Pharmacol Toxicol 44:525–557
- Ingallinesi M et al (2012) CRF2 receptor-deficiency eliminates opiate withdrawal distress without impairing stress coping. Mol Psychiatry 17(12):1283–1294
- Carrasco GA, Van de Kar LD (2003) Neuroendocrine pharmacology of stress. Eur J Pharmacol 463(1–3):235–272
- Keller-Wood M (2015) Hypothalamic-pituitary-adrenal axis feedback control. Compr Physiol 5(3):1161–1182
- McEwen BS (1998) Protective and damaging effects of stress mediators. N Engl J Med 338:171–179
- McEwen BS, Sapolsky RM (1995) Stress and cognitive function. Curr Opin Neurobiol 5(2):205–216
- 25. De Kloet ER, Derijk R (2004) Signaling pathways in brain involved in predisposition and pathogenesis of stress-related disease: genetic and kinetic factors affecting the MR/GR balance. Ann N Y Acad Sci 1032:14–34
- Willner P (2017) The chronic mild stress (CMS) model of depression: history, evaluation and usage. Neurobiol Stress 6:78–93
- Joels M et al (2004) Effects of chronic stress on structure and cell function in rat hippocampus and hypothalamus. Stress 7(4):221–231
- Willner P (2005) Chronic mild stress (CMS) revisited: consistency and behavioural-neurobiological concordance in the effects of CMS. Neuropsychobiology 52(2):90–110
- 29. Munhoz CD et al (2006) Chronic unpredictable stress exacerbates lipopolysaccharide-induced activation of nuclear factor-kappaB in the frontal cortex and hippocampus via glucocorticoid secretion. J Neurosci 26(14):3813–3820
- Tata DA, Yamamoto BK (2008) Chronic stress enhances methamphetamine-induced extracellular glutamate and excitotoxicity in the rat striatum. Synapse 62(5):325–336
- Garcia-Bueno B et al (2008) Stress mediators regulate brain prostaglandin synthesis and peroxisome proliferator-activated receptor-gamma activation after stress in rats. Endocrinology 149(4):1969–1978
- 32. Simmons DM, Arriza JL, Swanson LW (1989) A Complete protocol for in situ hybridization of messenger RNAs in brain and other tissues with radio-labeled single-stranded RNA probes. J Histotechnol 12(3):169–181

- Hsu SM, Raine L, Fanger H (1981) Use of avidin-biotin-peroxidase complex (ABC) in immunoperoxidase techniques: a comparison between ABC and unlabeled antibody (PAP) procedures. J Histochem Cytochem 29(4):577–580
- 34. Conover WJ (1999) Pratical nonparametric statistics, 3rd edn. Wiley, Hoboken
- Bakshi VP et al (2007) Stimulation of lateral septum CRF2 receptors promotes anorexia and stress-like behaviors: functional homology to CRF1 receptors in basolateral amygdala. J Neurosci 27(39):10568–10577
- Jeanneteau FD et al (2012) BDNF and glucocorticoids regulate corticotrophin-releasing hormone (CRH) homeostasis in the hypothalamus. Proc Natl Acad Sci U S A 109(4):1305–1310
- 37. Sapolsky R (2003) Taming stress. Sci Am 289(3):86-95
- Kling MA, Coleman VH, Schulkin J (2009) Glucocorticoid inhibition in the treatment of depression: can we think outside the endocrine hypothalamus? Depress Anxiety 26(7):641–649
- Herman JP et al (2003) Central mechanisms of stress integration: hierarchical circuitry controlling hypothalamo-pituitary-adrenocortical responsiveness. Front Neuroendocrinol 24(3):151–180
- Koob GF (1999) Corticotropin-releasing factor, norepinephrine, and stress. Biol Psychiatry 46(9):1167–1180
- Korosi A, Baram TZ (2008) The central corticotropin releasing factor system during development and adulthood. Eur J Pharmacol 583(2–3):204–214
- 42. Herman JP, Cullinan WE (1997) Neurocircuitry of stress: central control of the hypothalamo-pituitary-adrenocortical axis. Trends Neurosci 20(2):78–84
- 43. Flak JN et al (2014) Role of paraventricular nucleus-projecting norepinephrine/epinephrine neurons in acute and chronic stress. Eur J Neurosci 39(11):1903–1911
- 44. Wang B et al (2017) Effects of alpha2A adrenoceptors on norepinephrine secretion from the locus coeruleus during chronic stress-induced depression. Front Neurosci 11:243
- Agon P et al (1991) Permeability of the blood-brain barrier for atenolol studied by positron emission tomography. J Pharm Pharmacol 43(8):597–600
- Madrigal JL et al (2002) Stress-induced increase in extracellular sucrose space in rats is mediated by nitric oxide. Brain Res 938(1-2):87-91

- Radley JJ, Johnson SB, Sawchenko PE (2017) Stress: neuroendocrinology and neurobiology. In: Fink G (ed) Handbook of Stress Series. Academic Press, New York, pp 17–27
- Morilak DA et al (2005) Role of brain norepinephrine in the behavioral response to stress. Prog Neuropsychopharmacol Biol Psychiatry 29(8):1214–1224
- Herman JP et al (2005) Limbic system mechanisms of stress regulation: hypothalamo-pituitary-adrenocortical axis. Prog Neuropsychopharmacol Biol Psychiatry 29(8):1201–1213
- Singewald GM et al (2011) The modulatory role of the lateral septum on neuroendocrine and behavioral stress responses. Neuropsychopharmacology 36(4):793–804
- 51. Gold PW (2015) The organization of the stress system and its dysregulation in depressive illness. Mol Psychiatry 20(1):32–47
- 52. van Bodegom M, Homberg JR, Henckens M (2017) Modulation of the hypothalamic-pituitary-adrenal axis by early life stress exposure. Front Cell Neurosci 11:87
- Henry B, Vale W, Markou A (2006) The effect of lateral septum corticotropin-releasing factor receptor 2 activation on anxiety is modulated by stress. J Neurosci 26(36):9142–9152
- Heinrichs M et al (2003) Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. Biol Psychiatry 54(12):1389–1398
- Southwick SM, Vythilingam M, Charney DS (2005) The psychobiology of depression and resilience to stress: implications for prevention and treatment. Annu Rev Clin Psychol 1:255–291
- 56. Sterling P, Eyer J (1988) Allostasis: a new paradigm to explain arousal pathology. In: Fisher S, Reason J (eds) Handbook of life stress, cognition and health. Wiley, New York, pp 629–649

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.