

Diverse roles of WDR5-RbBP5-ASH2L-DPY30 (WRAD) complex in the functions of the SET1 histone methyltransferase family

AAMIR ALI[†] and SHWETA TYAGI*[‡]*Laboratory of Cell Cycle Regulation, Centre for DNA Fingerprinting and Diagnostics, Hyderabad, India***Corresponding author (Email, shweta@cdfd.org.in)**†Graduate Studies, Manipal University, Manipal, India*

WD repeat containing protein 5 (WDR5), Retinoblastoma Binding Protein 5 (RbBP5), Absent-Small-Homeotic-2-Like protein (ASH2L), and Dumpy-30 (Dpy30) have been reported to be the integral and shared components of all the SET1 family of histone 3 lysine 4 histone methyltransferase (HMT) complexes. Collectively called the WRAD complex, these proteins are pivotal to the HMT activity of the SET1 complexes. Recent reports highlight the novel non-canonical functions of WRAD in cellular processes other than its well-studied role in histone methylation and gene expression. In this review, we examine the diversity in emerging transcription-independent functions of WRAD.

[Ali A and Tyagi S 2017 Diverse roles of WDR5-RbBP5-ASH2L-DPY30 (WRAD) complex in the functions of the SET1 histone methyltransferase family. *J. Biosci.* **42** 155–159]

1. Introduction

Post-translational histone modifications add another layer of regulation to the ever changing chromatin landscape to activate or repress transcription. N- or C-terminal tails of histones can be methylated, phosphorylated, acetylated, ubiquitinated and, even, sumoylated or ribosylated to impact gene expression. All these histone modifications recruit chromatin remodelling protein complexes or alter the structure of chromatin (Bannister and Kouzarides 2011). In particular, lysine methylation regulates the activation (H3K4, H3K36, and H3K79) as well as repression (H3K9, H3K27, and H4K20) of transcription by methylation of specific residues in the histones (reviewed in Martin and Zhang 2005). Histone 3 Lysine 4 methylation (H3K4me) is closely associated with the activated state of the chromatin (Martin and Zhang 2005; Eissenberg and Shilatifard 2010). In mammals, the SET1 family of histone methyltransferases (HMT) is responsible for depositing the H3K4 methylation mark on promoters of active genes (Ernst and Vakoc 2012; Piunti and Shilatifard 2016).

The SET1 family include at least six multi-protein complexes namely Mixed Lineage Leukemia (MLL1 or MLL), MLL2, MLL3, MLL4, Set1A, and Set1B proteins (Cao 2012; Ernst

and Vakoc 2012). All these enzymes have unique non-redundant roles (Ansari and Mandal 2010; Shilatifard 2012; Shinsky and Cosgrove 2015). While Set1A and Set1B impart H3K4me3 mark to majority of gene promoters in mammals, MLL1 has been shown to regulate a subset of genes important during development, including the *Hox* loci (Wu *et al.* 2008). In contrast, MLL3 and MLL4 are involved in depositing the H3K4 mono-methylation mark at the enhancers (Herz *et al.* 2012; Hu *et al.* 2013). Like other methyltransferases, all members from this family act as multi-protein complexes with common subunits namely, WD repeat containing protein 5 (WDR5), Retinoblastoma Binding Protein 5 (RbBP5), Absent-Small-Homeotic-2-Like protein (ASH2L) and Dumpy-30 protein (DPY30) (Dou *et al.* 2006; Steward *et al.* 2006; Takahashi *et al.* 2011; Ernst and Vakoc 2012; van Nuland *et al.* 2013). Additional context-specific subunits also associate to impart unique properties to the complex (Yokoyama *et al.* 2004; Cho *et al.* 2007; Wu *et al.* 2008; Patel *et al.* 2007, 2014).

2. The conventional functions of WRAD

The WDR5, RbBP5, ASH2L, and DPY30 collectively called WRAD, along with the Set1/MLL Su(var)3-9,

Keywords. Cell cycle regulation; SET1 family; transcription; WRAD

Enhancer-of-zeste, Trithorax (SET) domain form a minimal 'core complex' capable of optimal enzymatic activity (Dou *et al.* 2006; Steward *et al.* 2006; Patel *et al.* 2011; Ernst and Vakoc 2012; Dharmarajan *et al.* 2012; Shinsky and Cosgrove 2015; Zhang *et al.* 2015). *In vitro* reconstitution studies with the purified WRAD proteins and the catalytic SET domain of MLL have provided deeper insights into the role of WRAD in the functions and the assembly of the MLL complex (Steward *et al.* 2006; Odho *et al.* 2010; Dharmarajan *et al.* 2012; Senisterra *et al.* 2013). *In vitro*, MLL by itself is a mono-methyltransferase (Patel *et al.* 2008b; reviewed in Cosgrove and Patel 2010). However in association with WRAD, it is capable of di- and tri-methylation activities (Zhang *et al.* 2015). While WDR5, RbBP5, and ASH2L form the minimal WRA complex required for the methylation activity observed with the WRAD complex, DPY-30 increases the catalytic efficiency and specificity of WRAD for histone H3 peptide (Cosgrove and Patel 2010; Patel *et al.* 2011).

Interestingly, *in vitro* studies also suggest that WRAD, which lacks SET domain, can methylate histone 3 peptide (Cosgrove and Patel 2010; Patel *et al.* 2009, 2011). WRAD can mono-methylate histone 3 peptide by itself but requires the presence of MLL SET domain (even when catalytically inactive) for nucleosome mono-methylation (Patel *et al.* 2009, 2011; Dharmarajan *et al.* 2012).

Two independent studies showed that WDR5 subunit bridges the interaction between MLL and the WRAD complex. WDR5 engages in direct interactions with the conserved arginine-containing WDR5 interacting (Win) motif localized in the C-terminus of MLL protein (Patel *et al.* 2008a, 2008b; Dharmarajan *et al.* 2012). This motif is conserved in SET1 family members and shares sequence homology with histone H3 N-terminus. Correspondingly, Phe133 and Phe263 amino acid residues of WDR5 sandwich the guanidinium of central arginine of the Win motif of MLL (Arg3765). Both *in vitro* and *in vivo* studies suggest that point mutation of either Phe133 or Phe263 in WDR5 or Arg3765 in MLL nearly abolishes the interaction between MLL and WRA sub-complex (Patel *et al.* 2008a; Song and Kingston 2008). However, more recently, it has been suggested that RbBP5-ASH2L heterodimer can bind to MLL directly albeit with weak affinity (Shinsky *et al.* 2014; Li *et al.* 2016).

In addition to the catalytic activity, the components of the WRAD complex are also implicated in the maintenance of the global levels of H3K4 methylation and, stability, product specificity and recruitment of SET1 complexes on the chromatin (Briggs *et al.* 2001; Nagy *et al.* 2002; Dou *et al.* 2006; Steward *et al.* 2006; Smith *et al.* 2011; Ernst and Vakoc 2012; Zhang *et al.* 2015).

3. Non-conventional functions of WRAD

Apart from participation of WRAD in the transcriptional activation of the genes by histone methylation, recent findings suggest transcriptional-independent roles of WRAD in the regulation of the cell cycle progression (figure 1). WRAD along with MLL has recently been shown to regulate the mitotic progression in a transcriptional independent manner (Ali *et al.* 2014). Loss of MLL by RNA interference (RNAi) gave rise to defects in S-phase and M-phase progression. While, after extensive mutational analysis, the regulation of the S-phase progression could be attributed to the activity of Transcriptional activation domain (TAD) of MLL (which acts independently of the SET domain), the domain responsible for influencing mitotic progression was not obvious.

Out of the several mitotic defects observed upon knock-down of MLL, two were quantified: (i) binuclei, arising from defective cytokinesis and (ii) micronuclei, marker of error-prone mitotic segregation (Ali *et al.* 2014). These mitotic defects were phenocopied when each subunit of WRAD was knocked down using RNAi individually, suggesting that the whole MLL 'core complex' was required. Further, we could demonstrate that the WDR5-MLL interaction was crucial for this function of MLL, as a recombinant MLL protein with point mutation in the Win motif was unable to rescue mitotic defects, observed upon the knockdown of endogenous MLL protein. However, further analysis highlighted differences from conventional MLL-WRAD activity, most notable being the non-requirement of MLL SET domain for this function. These observations led us to speculate if WRAD complex also enhances the transcriptional activity of MLL (in addition to its histone methyltransferase activity) using its transcriptional activation domain. We put this hypothesis to test by assessing the Win-motif mutant of MLL for its cell proliferative (S-phase) functions, which is dependent on the TAD domain. Even though, we observed that the mutation in Win motif had no effect on the cell proliferation functions of MLL, later experiments revealed that MLL may not require WRAD for its cell proliferation functions. Hence, we cannot say with certainty if WRAD will affect the TAD activity of MLL or not. But what is clear from our studies is that just the MLL-WRAD interaction is enough for mitotic progression, and neither the transcriptional activity (via the TAD) nor the methyltransferase activity (via the SET domain) of MLL plays any role here (Ali *et al.* 2014). Hence WRAD has a novel function with MLL.

Although this study highlights the emerging non-transcriptional, SET domain- and TAD-independent functions of MLL-WRAD, the underlying molecular mechanism is yet to be investigated. It is likely that WRAD may expand the interactome of MLL, fetching in new specific interaction partners for the proper regulation of mitosis. Alternatively,




Figure 1. Diverse functional roles of WRAD. The figure shows the model of WDR5, RbBP5, ASH2L, and DPY30 (WRAD) components participating in conventional (1) and non-conventional (2) functions. (1) The WRAD complex associates with MLL and methylates the promoters to regulate the transcription of genes during development. (2a-i, ii) Members of WRAD complex participate in cell-cycle regulation controlling S- and M-phase progression. (2a-iii) WDR5 subunit of WRAD associates with diverse interaction partners such as KIF4, MKLP10, and PRC1 at the midbody to regulate Cytokinesis; (2b) DPY30 component of WRAD interacts with BIG1 to modulate endosomal transport. WDR5, W: Red; RbBP5, R: Green; ASH2L, A: Blue, and DPY30, D: Orange.

it may act as an HMT by itself, performing methylation of non-histone proteins required in mitosis. In any case, it will be exciting to find out the sub-cellular localizations, interactions, and most importantly, functions of MLL-WRAD complex during mitotic progression to discern the underlying mechanism of mitotic regulation by this complex.

Recently, another study demonstrated that WDR5 localizes to the midbody and interacts with the midbody-localized proteins such as kinesins (KIF4, and MKLP10) and PRC1 proteins, thereby participating in the proper progression of cytokinesis (Bailey *et al.* 2015). Although this finding could show that central arginine-binding cavity of WDR5 is crucial for its midbody localization, yet the proteins involved in the recruitment of WDR5 to the midbody were not explored further (Bailey *et al.* 2015). In addition, DPY-30 subunit of WRAD has been shown to localize to the trans-Golgi compartments and regulate the recycling of the endosomes (Xu *et al.* 2009). The same study demonstrated that the depletion of RAD (RbBP5, ASH2L, and DPY30) perturbed the recycling of the endosomes (Xu *et al.* 2009). Even though these studies have not tested all subunits of WRAD complex; it is very likely that all of them are involved. Supporting this hypothesis, our studies observed cytokinesis defects with all

WRAD subunits, even though only WDR5 has been shown to localize to the midbody (Ali *et al.* 2014; Bailey *et al.* 2015). However, whether MLL is involved in all the above mentioned functions of WRAD is difficult to predict. Interestingly, in size exclusion chromatography of mammalian nuclear extracts, WRA complex in addition to eluting with MLL, also elutes as a distinct ~150 kDa complex lacking MLL, indicating that it may exist as an independent entity in the cell (Steward *et al.* 2006). Therefore, there is a possibility that WRAD may function far beyond the previously studied nuclear HMT function.

4. Future directions

An ever-increasing body of literature has correlated the activity of WRAD with the SET1 family for the histone methyltransferase function. While all these functions of WRAD well connect with its propensity to bind to the gene promoters within the nucleus, recently WRAD complex subunits have been suggested to play roles beyond its nuclear functions. It will be interesting to observe if contribution of WRAD supersedes its thoroughly studied partners to regulate diverse cellular functions.

Acknowledgements

We thank members of the Laboratory of Cell Cycle Regulation for valuable feedback. AA is the recipient of Junior and Senior Research Fellowships of the Council of Scientific and Industrial Research (CSIR), India, towards the pursuit of a Ph.D. degree of Manipal University. This work was supported in part by a grant from DBT (to ST; BT/BR15453/BRB/10/927/2011), DST (to ST; SB/SO/BB-069/2013) and CDFD core funds.

References

- Ali A, Veeranki SN and Tyagi S 2014 A SET-domain-independent role of WRAD complex in cell-cycle regulatory function of mixed lineage leukemia. *Nucleic Acids Res.* **42** 7611–7624
- Ansari KI and Mandal SS 2010 Mixed lineage leukemia: roles in gene expression, hormone signaling and mRNA processing. *FEBS J.* **277** 1790–1804
- Bailey JK, Fields AT, Cheng K, Lee A, Wagenaar E, Lagrois R, Schmidt B, Xia B, *et al.* 2015 WD repeat-containing protein 5 (WDR5) localizes to the midbody and regulates abscission. *J. Biol. Chem.* **290** 8987–9001
- Bannister AJ and Kouzarides T 2011 Regulation of chromatin by histone modifications. *Cell Res.* **21** 381–395
- Briggs SD, Bryk M, Strahl BD, Cheung WL, Davie JK, Dent SYR, Winston F and David Allis C 2001 Histone H3 lysine 4 methylation is mediated by Set1 and required for cell growth and rDNA silencing in *Saccharomyces cerevisiae*. *Genes Dev.* **15** 3286–3295
- Cao F 2012 MLL/SET1 complex: from yeast to human. *Fungal Genomics Biol.* **2** 1–2
- Cho YW, Hong T, Hong SH, Guo H, Yu H, Kim D, Guszczynski T, Dressler GR, *et al.* 2007 PTIP associates with MLL3- and MLL4-containing histone H3 lysine 4 methyltransferase complex. *J. Biol. Chem.* **282** 20395–20406
- Cosgrove MS and Patel A 2010 Mixed lineage leukemia: a structure-function perspective of the MLL1 protein. *FEBS J.* **277** 1832–1842
- Dharmarajan V, Lee JH, Patel A, Skalnik DG and Cosgrove MS 2012 Structural basis for WDR5 interaction (Win) motif recognition in human SET1 family histone methyltransferases. *J. Biol. Chem.* **287** 27275–27289
- Dou Y, Milne TA, Ruthenburg AJ, Lee S, Lee JW, Verdine GL, Allis CD and Roeder RG 2006 Regulation of MLL1 H3K4 methyltransferase activity by its core components. *Nat. Struct. Mol. Biol.* **13** 713–719
- Eissenberg JC and Shilatifard A 2010 Histone H3 lysine 4 (H3K4) methylation in development and differentiation. *Dev. Biol.* **339** 240–249
- Ernst P and Vakoc CR 2012 WRAD: enabler of the SET1-family of H3K4 methyltransferases. *Brief. Funct. Genomics.* **11** 217–226
- Herz HM, Mohan M, Garruss AS, Liang K, Takahashi Y-h, Mickey K, Voets O, Verrijzer CP, *et al.* 2012 Enhancer-associated H3K4 monomethylation by trithorax-related, the drosophila homolog of mammalian MLL3/MLL4. *Genes Dev.* **26** 2604–2620
- Hu D, Gao X, Morgan MA, Herz H-M, Smith ER and Shilatifard A 2013 The MLL3/MLL4 branches of the COMPASS family function as major histone H3K4 monomethylases at enhancers. *Mol. Cell. Biol.* **33** 4745–4754
- Li Y, Han J, Zhang Y, Cao F, Liu Z, Li S, Wu J, Hu C, *et al.* 2016 Structural basis for activity regulation of MLL family methyltransferases. *Nature* **530** 447–452
- Martin C and Zhang Y 2005 The diverse functions of histone lysine methylation. *Nat. Rev. Mol. Cell Biol.* **6** 838–849
- Nagy PL, Griesenbeck J, Kornberg RD and Cleary ML 2002 A trithorax-group complex purified from *Saccharomyces cerevisiae* is required for methylation of histone H3. *Proc. Natl. Acad. Sci. USA* **99** 90–94
- Odho Z, Southall SM and Wilson JR 2010 Characterization of a novel WDR5-binding site that recruits RbBP5 through a conserved motif to enhance methylation of histone H3 lysine 4 by mixed lineage leukemia protein-1. *J. Biol. Chem.* **285** 32967–32976
- Patel SR, Kim D, Levitan I and Dressler GR 2007 The BRCT-domain containing protein PTIP links PAX2 to a histone H3, lysine 4 methyltransferase complex. *Dev. Cell* **13** 580–592
- Patel A, Dharmarajan V and Cosgrove MS 2008a Structure of WDR5 bound to mixed lineage leukemia protein-1 peptide. *J. Biol. Chem.* **283** 32158–32161
- Patel A, Vought VE, Dharmarajan V and Cosgrove MS 2008b A conserved arginine-containing motif crucial for the assembly and enzymatic activity of the mixed lineage leukemia protein-1 core complex. *J. Biol. Chem.* **283** 32162–32175
- Patel A, Dharmarajan V, Vought VE and Cosgrove MS 2009 On the mechanism of multiple lysine methylation by the human mixed lineage leukemia protein-1 (MLL1) core complex. *J. Biol. Chem.* **284** 24242–24256
- Patel A, Vought VE, Dharmarajan V and Cosgrove MS 2011 A novel non-SET domain multi-subunit methyltransferase required for sequential nucleosomal histone H3 methylation by the mixed lineage leukemia protein-1 (MLL1) core complex. *J. Biol. Chem.* **286** 3359–3369
- Patel A, Vought VE, Swatkoski S, Viggiano S, Howard B, Dharmarajan V, Monteith KE, Kupakuwana G, *et al.* 2014 Autome-thylation activities within the mixed lineage leukemia-1 (MLL1) core complex reveal evidence supporting a “two-active site” model for multiple histone H3 lysine 4 methylation. *J. Biol. Chem.* **289** 868–884
- Piunti A and Shilatifard A 2016 Epigenetic balance of gene expression by Polycomb and COMPASS families. *Science* **352** aad9780
- Senisterra G, Wu H, Allali-Hassani A, Wasney GA, Barsyte-Lovejoy D, Dombrowski L, Dong A, Nguyen KT, *et al.* 2013 Small-molecule inhibition of MLL activity by disruption of its interaction with WDR5. *Biochem. J.* **449** 151–159
- Shilatifard A 2012 The COMPASS family of histone H3K4 methylases: mechanisms of regulation in development and disease pathogenesis. *Annu. Rev. Biochem.* **81** 65–95
- Shinsky SA and Cosgrove MS 2015 Unique role of the WD-40 repeat protein 5 (WDR5) subunit within the mixed lineage leukemia 3 (MLL3) histone methyltransferase complex. *J. Biol. Chem.* **290** 25819–25833

- Shinsky SA, Hu M, Vought VE, Ng SB, Bamshad MJ, Shendure J and Cosgrove MS 2014 A non-active-site SET domain surface crucial for the interaction of MLL1 and the RbBP5/Ash2L heterodimer within MLL family core complexes. *J. Mol. Biol.* **426** 2283–2299
- Smith E, Lin C and Shilatifard A 2011 The super elongation complex (SEC) and MLL in development and disease. *Genes Dev.* **25** 661–672
- Song JJ and Kingston RE 2008 WDR5 interacts with mixed lineage leukemia (MLL) protein via the histone H3-binding pocket. *J. Biol. Chem.* **283** 35258–35264
- Steward MM, Lee J, O'Donovan A, Wyatt M, Bernstein BE, Shilatifard A, Donovan AO, Wyatt M, *et al.* 2006 Molecular regulation of H3K4 trimethylation by ASH2L, a shared subunit of MLL complexes. *Nat. Struct. Mol. Biol.* **13** 852–854
- Takahashi Y-h, Westfield GH, Oleskie AN, Trievel RC, Shilatifard A and Skiniotis G 2011 Structural analysis of the core COMPASS family of histone H3K4 methylases from yeast to human. *Proc. Natl. Acad. Sci.* **108** 20526–20531
- van Nuland R, Smits AH, Pallaki P, Jansen PWTC, Vermeulen M and Timmers HTM 2013 Quantitative dissection and stoichiometry determination of the human SET1/MLL histone methyltransferase complexes. *Mol. Cell. Biol.* **33** 2067–2077
- Wu M, Wang PF, Lee JS, Martin-Brown S, Florens L, Washburn M and Shilatifard A 2008 Molecular regulation of H3K4 trimethylation by Wdr82, a component of human Set1/COMPASS. *Mol. Cell. Biol.* **28** 7337–7344
- Xu Z, Gong Q, Xia B, Groves B, Zimmermann M, Mugler C, Mu D, Matsumoto B, *et al.* 2009 A role of histone H3 lysine 4 methyltransferase components in endosomal trafficking. *J. Cell Biol.* **186** 343–353
- Yokoyama A, Wang Z, Wysocka J, Sanyal M, Aufiero DJ, Kitabayashi I, Herr W and Cleary ML 2004 Leukemia proto-oncoprotein MLL forms a SET1-like histone methyltransferase complex with menin to regulate Hox gene expression. *Mol. Cell. Biol.* **24** 5639–5649
- Zhang Y, Mittal A, Reid J, Reich S, Gamblin SJ and Wilson JR 2015 Evolving catalytic properties of the MLL family SET domain. *Structure* **23** 1921–1933

MS received 10 December 2016; accepted 12 January 2017

Corresponding editor: VEENA K PARNAIK