



An efficient one-pot synthesis of carbazole fused benzoquinolines and pyridocarbazoles

KRIPALAYA RATHEESH ARYA^a, HAZEL A SPARKES^b and
KARNAM JAYARAM PILLAI RAJENDRA PRASAD^{a,*}

^aDepartment of Chemistry, Bharathiar University, Coimbatore, Tamilnadu 641 046, India

^bSchool of Chemistry, University of Bristol, Cantock's Close, Bristol BS8 1TS, UK

E-mail: prasad_125@yahoo.com

MS received 1 November 2017; revised 3 January 2018; accepted 23 January 2018; published online 28 March 2018

Abstract. A one-pot, solvent-free protocol for the synthesis of chloro-substituted benzoquinoline-carbazole derivatives *via* a modified Friedländer hetero-annulation reaction between 2, 3, 4, 9-tetrahydrocarbazol-1-one and 3-amino-2-naphthoic acid in the presence of POCl₃ is described. In addition, the direct pseudo multicomponent transformation of 2, 3, 4, 9-tetrahydrocarbazol-1-one, malononitrile and 9-ethyl-3-carbazolecarboxaldehyde results in the formation of a multifunctionalized carbazole through a Knoevenagel–Michael addition-cyclization reaction has also been reported. All the newly synthesized molecules were deduced by spectral and analytical methods.

Keywords. Modified Friedländer reaction; benzoquinoline-carbazoles; pseudo multicomponent reaction; carbazole substituted pyridocarbazole.

1. Introduction

The structural diversity and biological importance of nitrogen containing heterocycles such as carbazole and quinoline derivatives have made them attractive targets in both medicinal and organic chemistry. The heteroaryl-carbazole derivatives have been found to display a diverse array of important functions and are abundant in bioactive natural products.^{1–4} For example, the pyridocarbazole type alkaloids, ellipticine (extracted from the leaves of *Ochrosia elliptica*) and Olivacine (isolated from *Aspidosperma olivaceum*) exhibit a wide spectrum of biological and medicinal activities.^{5–9} The benzoquinoline core structure is also found in a wide variety of biologically active natural products and pharmaceuticals with anti-Parkinson, antipsychotic, antibacterial, UDP (Uridine diphosphate)-glucuronosyl transferase, antimalarial, agonistic and antipsychotic activities.^{10–14} Kantevari *et al.*, reported that the coupling of 9-methyl-9H-carbazole to tetrahydroquinoline nucleus could deliver a new scaffold with better antimycobacterial activity than its individual reactants.¹⁵ Recently our

research group reported carbazole and quinoline-based hybrid moieties for its pharmacological interest.¹⁶

As an extension to our efforts in developing heterocycles of biological interest and also considering the significant role of carbazoles and benzoquinolines in biological applications, we were inspired to synthesize a new series of benzoquinoline-carbazole and carbazole dimer from 2,3,4,9-tetrahydro-1H-carbazol-1-one as the precursor *via* a one-pot protocol.

2. Experimental

2.1 General

All the chemicals were bought from Sigma-Aldrich and Merck and were utilized for the process without further purification. Melting points (M.p.) were determined on a Mettler FP 51 apparatus (Mettler Instruments, Switzerland) and are uncorrected. They are expressed in degree centigrade (°C). FT-IR spectra were recorded on Avatar Model FT-IR (4000–400 cm⁻¹) spectrophotometer. ¹H NMR and ¹³C NMR spectra were recorded on an Agilent- 400 MHz

*For correspondence

(^1H) and 100 MHz (^{13}C) spectrometers respectively in CDCl_3 using TMS (tetramethylsilane) as internal reference; chemical shifts are expressed in parts per million (ppm); coupling constants (J) are reported in hertz (Hz) and the terms J_o and J_m refer to ortho coupling constant and meta coupling constant. The signals were characterized as s (singlet), d (doublet), t (triplet), m (multiplet), bs (broad singlet) and dd (doublet of doublet). Microanalyses were carried out using Vario EL III model CHNS analyzer (Vario, Germany). Absorption spectral measurements were carried out using JASCO V-630 UV-Visible spectrophotometer. Quartz cuvettes of path length 1 cm were used to record the absorption spectra. The emission spectral studies were performed with JASCO FP-6600 spectrofluorometer equipped with a 1 cm quartz cuvette at the Department of Chemistry, Bharathiar University. When known compounds had to be prepared according to literature procedures pertinent references are given. The purity of the products was tested by TLC plates coated with silica gel-G using petroleum ether and ethyl acetate in the ratio of 1:1 as developing solvents.

2.2 General procedure for the synthesis of 7-chloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole **3**

A mixture of 2,3,4,9-tetrahydrocarbazol-1-one **1** (1.0 mmol) and 3-amino-2-naphthoic acid **2** (1.0 mmol) was refluxed with phosphorus oxychloride (5 mL) for 5 h at 120 °C. The completeness of the reaction was monitored by TLC. After completion, the reaction mixture was poured into ice water and extracted with ethyl acetate. Combined organic layers were dried over anhydrous magnesium sulphate. It was then purified on a silica-gel column chromatography (eluent: petroleum ether/ethyl acetate, 99:1).

2.3 The spectral and analytical data of all the compounds **3(a-d)**

2.3a 7-Chloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole (3a): White solid; yield: 283 mg (80%); M.p. 201–203 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3436(NH), 1592 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 9.47 (br s, 1H, NH), 8.66 (s, 1H, ArH), 8.52 (s, 1H, ArH), 8.04–7.98 (m, 2H, ArH), 7.62 (d, 1H, ArH, $J_o = 7.60$ Hz), 7.52–7.47 (m, 2H, ArH), 7.39 (d, 1H, ArH, $J_o = 8.40$ Hz), 7.29–7.27 (m, 1H, ArH), 7.13 (t, 1H, ArH, $J_o = 7.20$ Hz), 3.51–3.48 (m, 2H, CH_2), 3.21–3.17 (m, 2H, CH_2); ^{13}C NMR(100 MHz, CDCl_3) (ppm) δ_{C} : 149.2, 148.4, 143.8, 138.6, 138.1, 134.0, 132.4, 131.6, 128.5, 127.9, 126.8, 126.6, 126.0, 124.7, 124.5, 123.5, 120.0, 119.8, 111.8, 26.7, 19.0; Anal. calcd. for $\text{C}_{23}\text{H}_{15}\text{ClN}_2$: C, 77.85; H, 4.26; N, 7.89; found: C, 77.94; H, 4.22; N, 7.83 %.

2.3b 7-Chloro-3-methyl-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole (3b): White solid; yield: 253 mg (69%); M.p. 198–200 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3462 (NH), 1592 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 9.34 (br s, 1H, NH), 8.64 (s, 1H, ArH), 8.49 (s, 1H, ArH),

8.03–7.97 (m, 2H, ArH), 7.51–7.46 (m, 2H, ArH), 7.38 (s, 1H, ArH), 7.25 (d, 1H, ArH, $J_o = 8.80$ Hz), 7.08 (d, 1H, ArH, $J_o = 8.80$ Hz), 3.49–3.45 (m, 2H, CH_2), 3.17–3.13 (m, 2H, CH_2), 2.45 (s, 3H, CH_3); ^{13}C NMR(100 MHz, CDCl_3) (ppm) δ_{C} : 148.5, 143.9, 138.4, 136.5, 134.0, 132.7, 131.6, 129.3, 128.5, 127.9, 127.0, 126.5, 126.1, 125.9, 124.5, 123.5, 119.3, 111.4, 26.7, 21.4, 19.0 (CH_3); Anal. calcd. for $\text{C}_{24}\text{H}_{17}\text{ClN}_2$: C, 78.15; H, 4.65; N, 7.59; Found: C, 78.24; H, 4.63; N, 7.65%.

2.3c 7-Chloro-1-methyl-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole (3c): White solid; yield: 246 mg (67%); M.p. 199–201 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3281 (NH), 1590 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 9.13 (br s, 1H, NH), 8.68 (s, 1H, ArH), 8.55 (s, 1H, ArH), 8.05–8.12 (m, 2H, ArH), 7.54–7.47 (m, 3H, ArH), 7.12–7.05 (m, 2H, ArH), 3.52–3.49 (m, 2H, CH_2), 3.22–3.19 (m, 2H, CH_2), 2.59 (s, 3H, CH_3); Anal. calcd. for $\text{C}_{24}\text{H}_{17}\text{ClN}_2$: C, 78.15; H, 4.65; N, 7.59; Found: C, 78.22; H, 4.69; N, 7.63%.

2.3d 3,7-Dichloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole (3d): White solid; yield: 213 mg (55%); M.p. 195–197 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3263 (NH), 1549 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.69 (br s, 1H, NH), 8.60 (s, 1H, ArH), 8.06–8.01 (m, 2H, ArH), 7.58 (m, 1H, ArH), 7.55–7.53 (m, 2H, ArH), 7.39 (d, 2H, ArH, $J_o = 8.40$ Hz), 7.25 (s, 1H, ArH), 3.53–3.49 (m, 2H, CH_2), 3.19–3.15 (m, 2H, CH_2), Anal. calcd. for $\text{C}_{23}\text{H}_{14}\text{Cl}_2\text{N}_2$: C, 70.96; H, 3.62; N, 7.20; Found: C, 70.87; H, 3.66; N, 7.27%.

2.4 General procedure for the preparation of carbazole substituted pyrido[2,3-a]carbazoles **6** and **7**

A mixture of 2,3,4,9-tetrahydrocarbazol-1-one **1** (1.0 mmol), malononitrile **4**, (1.0 mmol), 9-ethyl-3-carbazole carboxaldehyde **5** (1.0 mmol) and lithium ethoxide (3 equiv.) in 15 mL of ethanol/methanol was heated to reflux for 3 h. The reaction was monitored by TLC which indicated the formation of the product. The excess of solvent was removed by distillation and the mixture was poured into ice-water. The reaction mixture was then neutralized with 5N HCl and extracted with ethyl acetate. The organic layer was thoroughly washed with water and dried over anhydrous Na_2SO_4 . Upon removal of the solvent, a brown crude mixture was obtained. It was purified by column chromatography over silica gel using petroleum ether: ethyl acetate (97:3) mixture as eluent to afford the corresponding product, 2-ethoxy/methoxy-4-aryl/heteroaryl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile **6** & **7**.

2.5 The spectral and analytical data of all the compounds **6** & **7**

2.5a 2-Ethoxy-4-(9'-ethyl-9H-carbazol-3'-yl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6a): Yellow solid; yield: 351 mg (73%); M.p. 272–274 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3259 (NH), 2216 (CN), 1544 (C=N);

^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.81 (br s, 1H, NH), 8.10 (d, 1H, ArH, $J = 8.00$ Hz), 8.06 (s, 1H, ArH), 7.58–7.44 (m, 6H, ArH), 7.29–7.27 (m, 1H, ArH), 7.15–7.11 (m, 2H, ArH), 4.64 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 4.42 (q, 2H, $\text{N}_9'\text{-CH}_2\text{CH}_3$, $J = 7.20$ Hz), 2.97–2.94 (m, 4H, CH_2), 1.54–1.48 (m, 6H, $\text{C}_2\text{-OCH}_2\text{CH}_3$ & $\text{N}_9'\text{-CH}_2\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3) (ppm) δ_{C} : 163.5, 155.4, 148.4, 140.4, 140.0, 137.8, 132.4, 126.8, 126.1, 126.0, 125.8, 124.4, 122.9, 122.7, 121.9, 120.7, 120.6, 120.2, 119.8, 119.2, 118.6 (CN), 116.1, 111.7, 108.7, 108.6, 93.9, 63.1 ($\text{C}_2\text{-OCH}_2\text{CH}_3$), 37.7 ($\text{N}_9'\text{-CH}_2\text{CH}_3$), 25.4, 19.4, 14.6 ($\text{C}_2\text{-OCH}_2\text{CH}_3$), 13.8 ($\text{N}_9'\text{-CH}_2\text{CH}_3$); Anal. calcd. for $\text{C}_{32}\text{H}_{26}\text{N}_4\text{O}$: C, 79.64; H, 5.43; N, 11.61; Found: C, 79.74; H, 5.47; N, 11.57%.

2.5b *2-Ethoxy-4-(9'-ethyl-9H-carbazol-3'-yl)-10-methyl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6b)*: Yellow solid; yield: 347 mg (70%); M.p. 271–273 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3449 (NH), 2217 (CN), 1551 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.62 (br s, 1H, NH), 8.10 (d, 1H, ArH, $J = 8.00$ Hz), 8.06 (s, 1H, ArH), 7.54–7.41 (m, 5H, ArH), 7.26–7.25 (m, 1H, ArH), 7.07 (d, 2H, ArH, $J = 8.00$ Hz), 4.66 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 4.42 (q, 2H, $\text{N}_9'\text{-CH}_2\text{CH}_3$, $J = 6.80$ Hz), 2.94–2.93 (m, 4H, CH_2), 2.59 (s, 3H, CH_3), 1.54–1.47 (m, 6H, $\text{C}_2\text{-OCH}_2\text{CH}_3$ & $\text{N}_9'\text{-CH}_2\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3) (ppm) δ_{C} : 163.5, 155.4, 148.5, 140.4, 140.0, 137.4, 132.1, 126.1, 126.1, 126.0, 125.9, 125.0, 122.9, 122.7, 121.9, 120.8, 120.7, 120.6, 120.4, 119.3, 119.2, 117.5 (CN), 116.2, 108.7, 108.6, 93.9, 62.9 ($\text{C}_2\text{-OCH}_2\text{CH}_3$), 37.7 ($\text{N}_9'\text{-CH}_2\text{CH}_3$), 25.5, 19.5, 16.7 (CH_3), 14.6 ($\text{C}_2\text{-OCH}_2\text{CH}_3$), 13.8 ($\text{N}_9'\text{-CH}_2\text{CH}_3$); Anal. calcd. for $\text{C}_{33}\text{H}_{28}\text{N}_4\text{O}$: C, 79.81; H, 5.68; N, 11.28; Found: C, 79.90; H, 5.64; N, 11.21%.

2.5c *2-Ethoxy-4-phenyl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6c)*: Yellow solid; yield: 292 mg (80%); M.p. 261–263 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3331 (NH), 2216 (CN), 1552 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.80 (br s, 1H, NH), 7.57 (d, 1H, ArH, $J_o = 8.00$ Hz), 7.53–7.47 (m, 3H, ArH), 7.44 (d, 1H, ArH, $J_o = 8.00$ Hz), 7.34 (d d, 2H, ArH, $J_m = 1.80$ Hz & $J_o = 7.80$ Hz), 7.28 (t, 1H, ArH, $J = 7.40$ Hz), 7.14 (t, 1H, ArH, $J = 7.40$ Hz), 4.62 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 2.96–2.92 (m, 2H, CH_2), 2.87–2.83 (m, 2H, CH_2), 1.51 (t, 3H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz); ^{13}C NMR (100 MHz, CDCl_3) (ppm) δ_{C} : 163.4, 154.2, 148.6, 137.8, 135.5, 132.2, 129.2, 129.0, 128.7, 128.4, 126.9, 126.7, 124.6, 121.3, 120.2, 119.9, 118.8 (CN), 115.7, 111.7, 93.2, 63.0 (OCH_2CH_3), 25.2, 19.4, 14.5 (OCH_2CH_3); Anal. calcd. for $\text{C}_{24}\text{H}_{19}\text{N}_3\text{O}$: C, 78.88; H, 5.24; N, 11.50; Found: C, 78.79; H, 5.28; N, 11.57%.

2.5d *2-Ethoxy-10-methyl-4-phenyl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6d)*: Yellow solid; yield: 299 mg (79%); M.p. 259–261 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3345 (NH), 2218 (CN), 1553 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.59 (br s, 1H, NH), 7.54–7.47 (m, 3H, ArH), 7.42 (d, 1H, ArH, $J_o = 6.80$ Hz), 7.35 (d d, 2H, ArH, $J_m = 1.60$ Hz & $J_o = 6.40$ Hz), 7.10–7.04 (m, 2H,


ArH), 4.64 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 2.95–2.90 (m, 2H, CH_2), 2.87–2.83 (m, 2H, CH_2), 2.58 (s, 3H, CH_3), 1.52 (t, 3H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz); Anal. calcd. for $\text{C}_{25}\text{H}_{21}\text{N}_3\text{O}$: C, 79.13; H, 5.58; N, 11.07; Found: C, 79.20; H, 5.54; N, 11.01%.

2.5e *2-Ethoxy-4-(thiophen-2-yl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6e)*: Yellow solid; yield: 255 mg (69%); M.p. 255–257 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3316 (NH), 2216 (CN), 1557 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.77 (br s, 1H, NH), 7.58 (d, 1H, ArH, $J_o = 8.00$ Hz), 7.53 (d d, 1H, ArH, $J_m = 1.20$ Hz & $J_o = 4.80$ Hz), 7.43 (d, 1H, ArH, $J_o = 8.00$ Hz), 7.29 (d d, 1H, ArH, $J_m = 1.20$ Hz & $J_o = 6.80$ Hz), 7.21–7.12 (m, 3H, ArH), 4.61 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 3.03–2.96 (m, 4H, CH_2), 1.50 (t, 3H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz); ^{13}C NMR (100 MHz, CDCl_3) (ppm) δ_{C} : 163.5, 148.6, 147.0, 137.9, 134.9, 132.1, 129.0, 127.7, 127.4, 126.7, 124.7, 122.6, 120.3, 119.9, 119.1 (CN), 115.5, 111.7, 94.0, 63.1 (OCH_2CH_3), 25.4, 19.3, 14.5 (OCH_2CH_3); Anal. calcd. for $\text{C}_{22}\text{H}_{17}\text{N}_3\text{OS}$: C, 71.14; H, 4.61; N, 11.31; Found: C, 71.23; H, 4.57; N, 11.36%.

2.5f *2-Ethoxy-10-methyl-4-(thiophen-2-yl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6f)*: Yellow solid; yield: 250 mg (65%); M.p. 253–255 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3349 (NH), 2214 (CN), 1552 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.56 (br s, 1H, NH), 7.53 (d d, 1H, ArH, $J_m = 1.20$ Hz & $J_o = 4.80$ Hz), 7.43 (d, 1H, ArH, $J_o = 7.60$ Hz), 7.21–7.14 (m, 2H, ArH), 7.08–7.04 (m, 2H, ArH), 4.63 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 3.03–2.95 (m, 4H, CH_2), 2.57 (s, 3H, CH_3), 1.50 (t, 3H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz); Anal. calcd. for $\text{C}_{23}\text{H}_{19}\text{N}_3\text{OS}$: C, 71.66; H, 4.97; N, 10.90; Found: C, 71.72; H, 4.93; N, 10.97%.

2.5g *2-ethoxy-4-(4'-chlorophenyl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6g)*: Yellow solid; yield: 231 mg (58%); M.p. 249–250 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3286 (NH), 2220 (CN), 1644 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.79 (br s, 1H, NH), 7.57 (d, 1H, ArH, $J_o = 7.20$ Hz), 7.50–7.47 (m, 2H, ArH), 7.44 (d, 1H, ArH, $J_o = 8.80$ Hz), 7.39–7.38 (m, 1H, ArH), 7.31–7.27 (m, 2H, ArH), 7.16–7.12 (m, 1H, ArH), 4.62 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 6.80$ Hz), 2.97–2.93 (m, 2H, CH_2), 2.85–2.81 (m, 2H, CH_2), 1.50 (t, 3H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 6.80$ Hz); Anal. calcd. for $\text{C}_{24}\text{H}_{18}\text{ClN}_3\text{O}$: C, 72.09; H, 4.54; N, 10.51; Found: C, 72.09; H, 4.54; N, 10.51%.

2.5h *2-ethoxy-10-methyl-4-(4'-chlorophenyl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (6h)*: Yellow solid; yield: 231 mg (56%); M.p. 250–252 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3267 (NH), 2214 (CN), 1552 (C=N); ^1H NMR (400 MHz, CDCl_3) (ppm) δ_{H} : 8.59 (br s, 1H, NH), 7.51–7.47 (m, 2H, ArH), 7.43 (d, 1H, ArH, $J_o = 6.40$ Hz), 7.30–7.27 (m, 2H, ArH), 7.10–7.05 (m, 2H, ArH), 4.64 (q, 2H, $\text{C}_2\text{-OCH}_2\text{CH}_3$, $J = 7.20$ Hz), 2.96–2.92 (m, 2H, CH_2),



Scheme 1. Synthesis of 7-chloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazoles **3**.

Table 1. Scope of the modified Friedlander synthesis of Benzoquinoline-carbazole derivatives.

Entry	R ₁	R ₂	R ₃	Product	Time (h)	Yield (%)
1.	H	H	H	3a	5	80
2.	CH ₃	H	H	3b	5	69
3.	H	H	CH ₃	3c	5.5	67
4.	Cl	H	H	3d	5.5	55

2.85–2.81 (m, 2H, CH₂), 2.57 (s, 3H, CH₃), 1.50 (t, 3H, C₂-OCH₂CH₃, $J = 7.20$ Hz); Anal. calcd. for C₂₅H₂₀ClN₃O: C, 72.55; H, 4.87; N, 10.15; Found: C, 72.55; H, 4.87; N, 10.15%.

2.5i 2-ethoxy-4-(4'-methylphenyl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (**6i**): Yellow solid; yield: 333 mg (88%); M.p. 263–265 °C; FT-IR (KBr, cm⁻¹) ν_{\max} : 3336 (NH), 2216 (CN), 1555 (C=N); ¹H NMR (400 MHz, CDCl₃) (ppm) δ_H : 8.79 (br s, 1H, NH), 7.57 (d, 1H, ArH, $J_o = 8.40$ Hz), 7.44 (d, 1H, ArH, $J_o = 8.40$ Hz), 7.31 (d, 2H, ArH, $J_o = 8.00$ Hz), 7.26–7.22 (m, 2H, ArH), 7.13 (t, 1H, ArH, $J = 8.00$ Hz), 4.61 (q, 2H, C₂-OCH₂CH₃, $J = 7.20$ Hz), 2.96–2.91 (m, 2H, CH₂), 2.89–2.84 (m, 2H, CH₂), 2.43 (s, 3H, C₄'-CH₃), 1.50 (t, 3H, C₂-OCH₂CH₃, $J = 7.20$ Hz); ¹³C NMR(100 MHz, CDCl₃) (ppm) δ_C : 163.4, 154.4, 148.5, 138.9, 137.8, 132.5, 132.3, 129.3, 128.3, 126.7, 124.5, 121.4, 120.2, 119.8, 118.7 (CN), 115.8, 111.7, 93.4, 62.9 (OCH₂CH₃), 25.2, 21.36, 19.4 (C₄'-CH₃), 14.5 (OCH₂CH₃); Anal. calcd. for C₂₅H₂₁N₃O: C, 79.13; H, 5.58; N, 11.07; Found: C, 79.22; H, 5.54; N, 11.14%.



2.5j 2-ethoxy-10-methyl-4-(4'-methylphenyl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (**6j**): Yellow solid; yield: 337 mg (86%); M.p. 261–263 °C; FT-IR (KBr, cm⁻¹) ν_{\max} : 3355 (NH), 2214 (CN), 1554 (C=N); ¹H NMR (400 MHz, CDCl₃) (ppm) δ_H : 8.58 (br s, 1H, NH), 7.43–7.40 (m, 1H, ArH), 7.31 (d, 2H, ArH, $J_o = 7.60$ Hz), 7.25–7.22 (m, 2H, ArH), 7.08–7.04 (m, 2H, ArH), 4.63 (q, 2H, C₂-OCH₂CH₃, $J = 6.80$ Hz), 2.94–2.92 (m, 2H, CH₂),

2.90–2.84 (m, 2H, CH₂), 2.57 (s, 3H, CH₃), 2.42 (s, 3H, C₄'-CH₃), 1.50 (t, 3H, C₂-OCH₂CH₃, $J = 6.80$ Hz); Anal. calcd. for C₂₆H₂₃N₃O: C, 79.36; H, 5.89; N, 10.68; Found: C, 79.25; H, 5.84; N, 10.72%.

2.5k 2-ethoxy-10-methyl-4-(4'-methoxyphenyl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (**6k**): Yellow solid; yield: 290 mg (71%); M.p. 256–258 °C; FT-IR (KBr, cm⁻¹) ν_{\max} : 3299 (NH), 2215 (CN), 1640 (C=N); ¹H NMR (400 MHz, CDCl₃) (ppm) δ_H : 8.58 (br s, 1H, NH), 7.43–7.41 (m, 1H, ArH), 7.28 (d d, 2H, ArH, $J_m = 2.00$ Hz & $J_o = 6.80$ Hz), 7.12–7.06 (m, 2H, ArH), 7.03 (d d, 2H, ArH, $J_m = 2.00$ Hz & $J_o = 6.80$ Hz), 4.63 (q, 2H, C₂-OCH₂CH₃, $J = 6.80$ Hz), 3.87 (s, 3H, C₄'-OCH₃), 2.93–2.89 (m, 4H, CH₂), 2.57 (s, 3H, CH₃), 1.51 (t, 3H, C₂-OCH₂CH₃, $J = 6.80$ Hz); Anal. calcd. for C₂₆H₂₃N₃O₂: C, 76.26; H, 5.66; N, 10.26; Found: C, 76.34; H, 5.64; N, 10.33%.

2.5l 2-Methoxy-4-(9-ethyl-9H-carbazol-3-yl)-10-methyl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (**7a**): Yellow solid; yield: 318 mg (68%); M.p. 263–265 °C; FT-IR (KBr, cm⁻¹) ν_{\max} : 3375 (NH), 2216 (CN), 1557 (C=N); ¹H NMR (400 MHz, CDCl₃) (ppm) δ_H : 8.86 (br s, 1H, NH), 8.10 (d, 1H, ArH, $J_o = 7.20$ Hz), 8.06 (s, 1H, ArH, $J_m = 1.20$ Hz), 7.57 (d, 1H, ArH, $J_o = 8.00$ Hz), 7.54–7.48 (m, 2H, ArH), 7.46–7.42 (m, 3H, ArH), 7.30–7.26 (m, 2H, ArH), 7.13 (t, 1H, ArH, $J_o = 7.60$ Hz), 4.62 (q, 2H, N₉'-CH₂CH₃, $J = 7.20$ Hz), 4.18 (s, 3H, C₂-OCH₃), 2.98–2.93 (m, 4H, CH₂), 1.49 (t, 3H, N₉'-CH₂CH₃, $J = 7.20$ Hz); ¹³C NMR(100 MHz, CDCl₃) (ppm) δ_C : 163.7, 155.4, 148.4, 140.4, 140.0, 137.9, 132.3, 126.1, 126.7, 126.2, 125.7, 124.5, 123.6, 122.7, 122.1, 120.7, 120.6, 120.2, 119.3 (CN), 119.2, 118.8, 111.7, 108.7, 108.6, 54.3 (OCH₃), 37.7 (N₉'-CH₂CH₃), 25.4, 19.4, 13.8 (N₉'-CH₂CH₃); Anal. calcd. for C₃₁H₂₄N₄O: C, 79.46; H, 5.16; N, 11.96; Found: C, 79.55; H, 5.12; N, 11.89%.

2.5m 2-Methoxy-4-(9'-ethyl-9H-carbazol-3'-yl)-10-methyl-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile (**7b**): Yellow solid; yield: 313 mg (65%); M.p.



260–262 °C; FT-IR (KBr, cm^{-1}) ν_{max} : 3391 (NH), 2215 (CN), 1556 (C=N); $^1\text{H NMR}$ (400 MHz, CDCl_3) (ppm) δ_{H} : 8.75 (br s, 1H, NH), 8.10 (d, 1H, ArH, $J_o = 8.60$ Hz), 8.58 (d, 1H, ArH, $J_m = 1.60$ Hz), 7.54–7.48 (m, 2H, ArH), 7.46

(m, 2H, ArH), 7.35–7.33 (m, 2H, ArH), 7.27–7.26 (m, 1H, ArH), 7.11 (d d, 1H, ArH, $J_m = 1.60$ Hz & $J_o = 7.60$ Hz), 4.42 (q, 2H, $\text{N}_9\text{'-CH}_2\text{CH}_3$, $J = 7.20$ Hz), 4.17 (s, 3H, $\text{C}_2\text{-OCH}_3$), 2.96–2.90 (m, 4H, CH_2), 2.44 (s, 3H, CH_3),

1.49 (m, 3H, N₉'-CH₂CH₃); Anal. calcd. for C₃₂H₂₆N₄O: C, 79.64; H, 5.43; N, 11.61; Found: C, 79.73; H, 5.47; N, 11.54%.

3. Results and Discussion

The two-component Friedländer reaction of 2-aminoaryl ketones with carbonyl compounds containing a reactive α -methylene group in order to obtain benzoquinoline compounds has been known for a long time.^{17–22} However, to the best of our knowledge, none of these report the synthesis of benzoquinoline-carbazole derivatives via a POCl₃ promoted Friedländer reaction.

Prompted by the encouraging importance of benzoquinoline-carbazole derivatives via Friedländer reaction, we embarked on the synthesis of 7-chloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole **3** by a modified Friedländer hetero-annulation reaction of 2,3,4,9-tetrahydrocarbazol-1-one **1** with 3-amino-2-naphthoic acid **2** using catalytic amounts of POCl₃ under solvent-free condition (Scheme 1). All these reactions were carried out by refluxing an equimolar ratio of the reactants at 120 °C.

Next, we examined the scope of the transformation utilizing the reactivity of different substituted 2,3,4,9-tetrahydrocarbazol-1-ones **1** (a–d) as substrates; all of the reactions afforded the corresponding benzoquinoline-carbazole products **3** (a–d) in moderate to good yields (Table 1).

The proposed structures of the synthesized compounds were consistent with their FT-IR, ¹H NMR, ¹³C NMR spectra and elemental analyses. The FT-IR spectral data of **3a** displayed prominent absorption peaks at 3436 and 1592 cm⁻¹ due to indole NH and C=N stretchings respectively. The ¹H NMR spectrum of **3a** exhibited a broad singlet for the indole NH at δ 9.47 ppm. Two singlets at δ 8.66 and δ 8.52 ppm assigned to C₈ and C₁₃ protons. The two multiplet signals in the region of δ 8.04–7.98 ppm due to C₁₂ & C₉ protons and δ 7.52–7.47 ppm were attributed to the C₁₁ & C₁₀ aromatic protons respectively. The signal due to C₄ proton occurred as a doublet at δ 7.62 ($J_o = 7.60$ Hz) ppm and a doublet at δ 7.39 ($J_o = 8.40$ Hz) arising from the C₁ proton. A multiplet at δ 7.29–7.27 ppm was assigned to a C₃ proton, a triplet at δ 7.13 ($J_o = 7.20$ Hz) was assigned to the C₂ proton while the aliphatic protons (C₆ and C₅) resonated as multiplets centered at δ 3.50 ppm and 3.19 ppm respectively. The ¹³C NMR spectrum of **3a** displayed 23 resonances in agreement with the proposed structure.

The plausible mechanism for the formation of compound **3** has been shown in Scheme 2 as reported earlier by our research group.²³ Initially, 2,3,4,9-tetrahydrocar-

Table 2. Screening of catalyst for the multicomponent synthesis of **6a**^a.


Entry	Solvent	Catalyst (equiv)	Time	Yield ^b
1.	EtOH	Catalyst-free	12	— ^c
2.	EtOH	NaOH	8	20
3.	EtOH	K ₂ CO ₃	8	29
4.	EtOH	Et ₃ N	6	57
5.	EtOH	DABCO	6	31
6.	EtOH	Morpholine	6	27
7.	EtOH	Piperidine	5	38
8.	EtOH	NaOEt (1)	5	47
9.	EtOH	LiOEt (1)	3	58
10.	EtOH	LiOEt (1.5)	3	61
11.	EtOH	LiOEt (2)	3	68
12.	EtOH	LiOEt (3)	3	73
13.	EtOH	LiOEt (4)	6	67

^aReaction of **2**, **3**, **4**, 9-tetrahydrocarbazol-1-one **1a** (1.0 mmol), malononitrile **4** (1.0 mmol) and 9-ethyl-3-carbazolecarboxaldehyde **5a** (1.0 mmol) using LiOEt (3 equiv) under EtOH.

^bIsolated yields, ^cTrace product.

bazol-1-one **1** is condensed with 3-amino-2-naphthoic acid **2** in the presence of an acid POCl₃ as a catalyst to afford the stable geometrical isomer **E** as intermediate **I**. Upon tautomerisation the more stable imine intermediate **I** gave an enamine intermediate **II**. The addition of POCl₃ to the carboxylic acid intermediate **II** causes the formation of a mixed anhydride intermediate **III**, which then underwent an intramolecular electrophilic substitution reaction to yield the intermediate **IV** and this intermediate on subsequent 1,4-prototropic shift followed by an S_N1 reaction *cum* PO₂Cl elimination afforded the intermediate **VI**. The trans dehydration of intermediate **VI** gave the hybrid compound, 7-chloro-benzo[6',7'-a']quino[2',3'-a]-5,6-dihydrocarbazole **3**.

9-Ethyl-3-carbazolecarboxaldehyde derivatives have been utilized as versatile coupling components in the preparation of a number of nitrogen-containing heteroaromatics.²⁴ The simple carbazoles reported so far from our laboratory were found to have anticancer, antibacterial, antioxidant and photophysical activities.¹⁶ However, the dimerised carbazoles/carbazolyl carbazole derivatives have an electron-rich structure, high thermal stability and unique electrical and optical properties^{25,26} compared to its simple carbazoles. The conjugated carbazole dimer enhanced greatly the fluorescence of carbazole.²⁷ The importance of pyridocarbazole and carbazole dimer derivatives prompted us to design a practical and efficient multicomponent reaction for the preparation of highly functionalized carbazole substituted pyridocarbazole derivatives. Recently, our research group envisaged the possibility of pseudo

Table 3. Scope of aryl/heteroaryl substituted pyrido [2, 3-*a*] carbazoles **6** (a–k).

multicomponent reaction for the preparation of pyrido-carbazoles.²⁸

The synthetic pathways employed to prepare the targeted derivatives are depicted in Scheme 3. In a one-pot, pseudo three-component heterocyclo condensation process, 2-ethoxy-4-(9'-ethyl-9*H*-carbazol-3'-yl)-5,6-dihydro-11*H*-pyrido[2,3-*a*]carbazole-3-carbonitrile **6a** was obtained *via* a base mediated Michael addition-cyclization of 2,3,4,9-tetrahydrocarbazol-1-one **1a** with malononitrile **4** and 9-ethyl-3-carbazolecarboxaldehyde

5a in refluxing EtOH. In this reaction, the solvent ethanol could act as both a reactant and a solvent.

Table 2 shows the results obtained from using different reaction conditions for the synthesis of **6a**. In order to justify the significance of base in this multicomponent process, the reaction was first performed in the absence of base wherein the reaction failed to occur even at prolonged reaction time (Table 2, entry 1). In the presence of sodium hydroxide or potassium carbonate, the reaction occurred with isolated yields of **6a** after 8 h of 20% and





Figure 1. Illustration of the structures of **6c** (left) and **6h** (right) (only one unique molecule shown). Anisotropic displacement parameters are depicted at the 50% probability level.




Scheme 4. Synthesis of 2-methoxy-4-(9'-ethyl-9H-carbazol-3'-yl)-5,6-dihydro-11H-pyrido[2,3-a]carbazole-3-carbonitrile **7a**.

29% respectively (Table 2, entries 2 and 3). Carrying out the reaction in the presence of triethylamine (Table 2, entry 4) for 6 h led to a significant increase in the yield of **6a**. The reaction was then performed with bases such as DABCO, morpholine and piperidine (Table 2, entries 5–7), however, these bases were not as effective and resulted in low yields (< 38%) after 5–6 h. Our research group has previously reported the use of NaOEt as a base to afford simple pyridocarbazoles.²⁸ However, in our present work, the use of NaOEt as a catalyst yielded only 47% of carbazole compound (Table 2, entry 8). Interestingly, the use of one equivalent of LiOEt instead of NaOEt increases the product yield up to 58% after 3 h (Table 2, entry 9). The optimum quantity of LiOEt required was screened and it was found that on increasing the amount of catalyst from 1.0 to 3.0 equiv, the yield of the reaction increased gradually but beyond 3.0 equiv, there was no significant improvement in the rate or yield of the reaction even at prolonged time (Table 2, entries 9–13). From the results, LiOEt was found to be the optimum catalyst for this transformation, wherein 73% of product **6a** was obtained in 3h. The literature survey revealed the catalytic potential of LiOEt in many MCRs.²⁹

Based on the appropriate reaction conditions, a series of 2-ethoxy-4-aryl/heteroaryl-5, 11-dihydro-6H-pyrido [2, 3-a] carbazole-3-carbonitrile derivatives **6** (**a–k**) were synthesized. The results are summarized in Table 3.

The structures of compounds **6** (**a–k**) were established on the basis of their elemental analyses and spectral data. The important diagnostic bands in the FT-IR spectrum of **6a** were assigned, with the stretching vibrations at 3259 and 1544 cm^{-1} corresponding to indole NH and C=N groups respectively and the cyano group stretching vibration assigned to a sharp band at 2216 cm^{-1} . The ^1H NMR spectrum displayed a broad singlet at δ 8.81 ppm attributed to the indole NH proton while the signal from C_{5'}-H was visible as a doublet at δ 8.10 ($J_o = 8.00$ Hz) and a sharp singlet at δ 8.06 ppm was assigned to C_{4'} proton. The six aromatic protons at C_{8'}, C_{2'}, C_{1'}, C₇, C₁₀ and C_{7'} positions resonated as multiplets in the region of δ 7.58–7.44 ppm and the C_{6'} aromatic proton appeared as a multiplet at δ 7.29–7.27 ppm. The C₉ and C₈ protons were visible as a multiplet at δ 7.15–7.11 ppm, the OCH₂CH₃ protons appeared as a quartet at δ 4.64 ($J = 7.20$ Hz). The two protons of N_{9'}-CH₂ appeared as a quartet at δ 4.42 ppm ($J = 7.20$ Hz)




Scheme 5. Mechanistic rationalization for the formation of **6** & **7**.

while the methylene protons of C_5 and C_6 resonated as a multiplet at δ 2.97–2.94 ppm. Six methyl protons of OCH_2CH_3 and N_9' - CH_2CH_3 appeared as a multiplet at δ 1.54–1.48 ppm. The ^{13}C NMR spectrum of **6a** displayed 32 resonances in agreement with the proposed structure. The resonance signals at δ 63.1, 37.7, 14.6 and 13.8 were attributed to OCH_2CH_3 , N_9' - CH_2CH_3 , OCH_2CH_3 and N_9' - CH_2CH_3 carbons. The structures of **6c** and **6h** were further confirmed by single X-ray diffraction studies (Figure 1).

After the successful synthesis of 2-ethoxy-4-(9'-ethyl-9*H*-carbazol-3'-yl)-5,6-dihydro-11*H*-pyrido[2,3-*a*]carbazole-3-carbonitrile **6a**, this catalytic system was used for the synthesis of 2-methoxy-4-(9'-ethyl-9*H*-carbazol-3'-yl)-5,6-dihydro-11*H*-pyrido[2,3-*a*]carbazole-3-carbonitrile **7a** by the condensation of same reactants in the presence of MeOH instead of EtOH as solvent (Scheme 4).

The analysis of the ^1H NMR and ^{13}C NMR data of the products led us to conclude that, when MeOH or EtOH was used as a solvent the reaction mechanism changed from a three-component to a four-component pathway, as depicted in Scheme 5. Initially, the intermediate **I** was formed *via* Knoevenagel condensation of 9-ethyl-3-carbazolecarboxaldehyde with malononitrile in the presence of a base. On subsequent base promoted the 1,4-Michael addition of electron deficient Knoevenagel adduct to the carbanion, derived from the synthon, 2,3,4,9-tetrahydro-1*H*-carbazol-1-one affords the dinitrile intermediate **II**, which undergoes prototropic shift and alcoholic addition facilitated by the base, LiOEt to form an intermediate **IV** through the intermediate **III**. An intramolecular cyclization of intermediate **IV** could furnish the intermediate **V**, which upon dehydration and aerial oxidation could give rise to the product **6/7**.

Table 4. The scope of various 9-ethyl-3-carbazolecarboxaldehyde substituted pyrido [2, 3-*a*] carbazole derivatives.

The scope and general applicability of this methodology have been investigated by using different 2, 3, 4, 9-tetrahydrocarbazol-1-ones **1** (**a**, **b**) with 9-ethyl-3-carbazolecarboxaldehyde **5a** in different solvent conditions (Table 4).

4. Conclusion

In conclusion, we have demonstrated that differently substituted 2, 3, 4, 9-tetrahydrocarbazol-1-ones and 3-amino-2-naphthoic acid undergo an efficient Friedlander condensation reaction in the presence of POCl₃ to yield benzoquinoline-carbazoles. The synthon, 2, 3, 4, 9-tetrahydrocarbazol-1-ones further undergo a pseudo three-component reaction with malononitrile and 9-ethyl-3-carbazole carboxaldehyde to afford a carbazole engrafted pyridocarbazole derivatives. The investigation on solvent effect on this reaction procedure revealed that solvent could change reaction mechanism from three-component to four-component pathway. On the whole, the methods presented here are significant in terms

of good yields, short reaction time, cost-effectiveness, readily available substrates and wide scope for production of a diversity of the products and potentially bioactive compounds.

Supplementary Information (SI)

All additional information pertaining to characterization of the compounds using ¹H NMR and ¹³C NMR spectra (Figures S1–S21) and table of Crystal data are given in the supporting information.

Acknowledgements

We would like to thank the University of Mysore, for NMR data. X-ray diffraction was funded by University of Bristol, United Kingdom. Financial support from “UGC-Emeritus fellowship” [Award NO.F.6-6/2015-17/EMERITUS-2015-17-OBC-7410/ (SAII)] for research, is gratefully acknowledged by Prof. K. J. Rajendra Prasad. Arya. K. R acknowledges the award of BSR-Senior Research Fellowship by University Grant Commission (UGC), New Delhi.

References

- Suresh J R, Syam Kumar U K, Ila H and Junjappa H 2001 Aromatic and heteroaromatic annelation studies on 3-[bis(methylthio)methylene]-1-methyloxindole: synthesis of carbazoles and an efficient route to pyrido[2,3-*b*]indoles *Tetrahedron* **57** 781
- Hu L X, Li Z R, Wang Y M, Wu Y B, Jiang J D and Boykin D W 2007 Novel pyridinyl and pyrimidinylcarbazole sulfonamides as antiproliferative agents *Bioorg. Med. Chem. Lett.* **17** 1193
- Dračinský M, Sejbál J, Rygerová B and Stiborová M 2007 An efficient modification of ellipticine synthesis and preparation of 13-hydroxyellipticine *Tetrahedron Lett.* **48** 6893
- Barbieri V and Ferlin M G 2006 Microwave-assisted one-pot synthesis of substituted tetrahydrocarbazole and 8,9,10,11-tetrahydro-7H-pyrido[a]carbazoles *Tetrahedron Lett.* **47** 8289
- Haider N, Jabara R, Khadami F and Wanko R 1998 Synthesis of Pyridazino [4, 5-*b*] carbazoles as potential antitumor agents *Heterocycles* **48** 1609
- Hedin V M, Tabka T, Poulain L, Godard T, Lachevrel M, Saturnino C, Lancelot J C, Le Talaer J Y and Gauduchon P 2000 Biological properties of 5,11-dimethyl-6H-pyrido [3, 2-*b*] carbazoles: a new class of potent antitumor drugs *Anticancer Drug Des.* **15** 109
- de Koning C B, Michael J P and Rosseau A L 2000 A versatile and convenient method for the synthesis of substituted benzo[a]carbazoles and pyrido[2,3-*a*]carbazoles *J. Chem. Soc. Perkin Trans.* **1** 1705
- Hirata K, Ito C, Furukawa H, Itogiawa M, Mark Cosentino L and Lee K H 1999 Substituted 7H-pyrido[4,3-*c*]carbazoles with potent anti-HIV activity *Bioorg. Med. Chem. Lett.* **9** 119
- Ohashi M, Shudo T, Nishijima K, Notso T, Kikuchi A, Yanagibashi K and Nishida H 2000 *Pyridocarbazole derivatives having cGMP-PDE inhibitory activity, United States Patent 6018046* United States Mochida Pharmaceutical Co., Ltd, Tokyo, JP
- Selvi G and Rajendran S P 2004 Synthesis of some new 2-[3-(2-chloro quinolinyl)]-3-aryl-4-thiazolidinones as potent antibacterial agents *J. Asian Chem.* **16** 1017
- Carr B A and Franklin M R 1998 Drug-metabolizing enzyme induction by 2,2'-dipyridyl, 1,7-phenanthroline, 7,8-benzoquinoline and oltipraz in mouse *Xenobiotica* **28** 949
- Mikhailitsyn F S, Kozyreva N P, Rabinovich S A, Maksakovskaya Y V, Kulikovskaya I M, Dadasheva N R, Lebedeva M N, Bekhli A F, Lychko N D and Uvarova N A 1992 Search for new antiparasitic agents, Synthesis, toxicity, and antimalarial effect of some nitrogen-containing heterocycles with 4-(4-alkylpiperazin-1-yl)phenylamino substituents *Med. Parazitol. Parazit. Bolezni.* **50**
- Nozulak J, Vigouret J M, Jaton A L, Hofmann A, Dravid A R, Weber H P, Kalkman H O and Walkinshaw M D 1992 Centrally acting α_1 -adrenoceptor agonists based on hexahydronaphth[2,3-*b*]-1,4-oxazines and octahydrobenzo[g]quinolines *J. Med. Chem.* **35** 480
- Szmuszkowicz J, Darlington W H and Von Voigtlander P F 1988 Preparation and formulation of antipsychotic aminopolyhydrobenz(iso)quinolines and intermediates. WO 8804292 A1 1988 *Chem. Abstr.* **110** 75335
- Kantevari S, Yempala T, Surineni G, Sridhar B, Yogeewari P and Sriram D 2011 Synthesis and antitubercular evaluation of novel dibenzo[b,d]furan and 9-methyl-9H-carbazole derived hexahydro-2H-pyrano[3,2-*c*]quinolines via Povarov reaction *Eur. J. Med. Chem.* **46** 4827
- (a) Prabha K and Rajendra Prasad K J 2016 Synthesis and Cytotoxic Distinction of Benzo[h]naphtho[1,2-*b*] [1,6] Naphthyridine and its Isomeric Benzo[b]naphtho[1,2-*h*][1,6] Naphthyridines *Med. Chem.* **62** 062; (b) Indumathi T, Muthusankar A, Shanmughavel P and Rajendra Prasad K J 2013 Synthesis of hetero annulated carbazoles: exploration of in vitro cytotoxicity and molecular docking studies *Med. Chem. Commun.* **4** 450; (c) Indumathi T, Jamal Ahamed V S, Surk-Sik Moon, Fronczek F R and Rajendra Prasad K J 2011 L-Proline anchored multicomponent synthesis of novel pyrido[2,3-*a*]carbazoles; investigation of in vitro antimicrobial, antioxidant, cytotoxicity and structure activity relationship studies *Eur. J. Med. Chem.* **46** 5580; (d) Murali K, Sparkes H A and Rajendra Prasad K J 2017 Synthesis of hetero annulated isoxazolo-, pyrido- and pyrimido carbazoles: Screened for in vitro antitumor activity and structure activity relationships, a novel 2-amino-4-(3'-bromo-4'-methoxyphenyl)-8-chloro-11H-pyrimido[4,5-*a*]carbazole as an antitumor agent *Eur. J. Med. Chem.* **128** 319; (e) Yamuna E, Yurcho A, Sovesky R J, Smith P M, Zeller M and Rajendra Prasad K J 2011 Elegant one-pot synthesis of quinolino[2', 3' : 7, 6]-cyclohept[1,2-*b*]indole through Friedländer and Pfizinger annulation reaction *Synth. Commun.* **41** 3351
- (a) Ryabukhin S V, Naumchik V S, Plaskon A S, Grygorenko O O and Tolmachev A A 2011 3-Haloquinolines by Friedländer reaction of α -haloketones *J. Org. Chem.* **76** 5774; (b) Le Z-G, Liang M, Chen Z-S, Zhang S-H and Xie Z-B 2017 Ionic liquid as an efficient medium for the synthesis of quinoline derivatives via α -chymotrypsin catalysed Friedländer condensation *Molecules* **22** 762; (c) Tufail F, Saquib M, Singh S, Tiwari J, Singh M, Singh J and Singh J 2017 Bioorganopromoted green Friedländer synthesis: a versatile new malic acid promoted solvent free approach to multisubstituted quinoline *New J. Chem.* **41** 1618
- (a) Zhang X-L, Wang Q-Y, Sheng S-R, Wang Q and Liu X-L 2009 Efficient Friedländer Synthesis of quinoline derivatives from 2-aminoarylketones and carbonyl compounds mediated by recyclable PEG-supported sulfonic acid *Synth. Commun.* **39** 3293; (b) Ramann G A and Cowen B J 2016 Recent advances in metal-free quinoline synthesis *Molecules* **21** 986; (c) Reddy B P, Iniyavan P, Sarveswari S and Vijayakumar V 2014 Nickel oxide nanoparticles catalyzed synthesis of poly-substituted quinolines via Friedlander hetero-annulation reaction *Chin. Chem. Lett.* **25** 1595
- Luo W, Mu Q, Qiu W, Liu T, Yang F, Liu X and Tang J 2011 A novel Friedlander-type synthesis of 3-aryl quinolines from 3-oxo-2,3-diaryl-propionaldehydes *Tetrahedron* **67** 7090

20. Genovese S, Epifano F, Marcotullio M C, Pelucchini C and Curini M 2011 An alternative quinoline synthesis by via Friedländer reaction catalyzed by Yb(OTf)₃ *Tetrahedron Lett.* **52** 3474
21. McNaughton B R and Miller B L 2003 A mild and efficient one-step synthesis of quinolines *Org. Lett.* **5** 4257
22. Satheeshkumar R, Shanker R, Kaminsky W and Rajendra Prasad K J 2016 Novel synthetic and mechanistic approach of TFA catalyzed Friedlander synthesis of 2-acylquinolines from symmetrical and unsymmetrical 1,2-diketones with o-aminoarylketone *ChemistrySelect.* **1** 6823
23. Prabakaran K and Rajendra Prasad K J 2010 Simple and convenient methods for the synthesis of indolo[3,2-c]acridines *Synth. Commun.* **40** 3528
24. Ashok D, Ravi S, Vijaya Lakshmi B and Ganesh A 2015 One-pot synthesis of carbazole based 3-hydroxy-4H-chromen-4-ones by modified Algar–Flynn–Oyamada reaction and their antimicrobial activity *J. Serb. Chem. Soc.* **80** 1361
25. (a) Song Y, Di C-A, Wei Z, Zhao T, Xu W, Liu Y, Zhang D and Zhu D 2008 Synthesis, characterization, and field-effect transistor properties of carbazolevinylene oligomers: from linear to cyclic architectures *Chem. Eur. J.* **14** 4731; (b) Simokaitiene J, Grigalevicius S, Grazulevicius J V, Rutkaite R, Kazlauskas K, Jursenas S, Jankauskas V and Sidaravicius J 2006 Synthesis, photophysical and photoelectrical properties of glass-forming phenothiazinyl- and carbazoly-substituted ethylenes *J. Optoelectron Adv. Mater.* **8** 876; (c) Jin Y, Zhang A, Huang Y and Zhang W 2010 Shape-persistent arylenevinylene macrocycles (AVMs) prepared via acyclic diene metathesis macrocyclization (ADMAC) *Chem. Commun.* **46** 8258
26. Chen C H, Lin J T and Yeh M C P 2006 Stilbene like carbazole dimer-based electroluminescent materials *Tetrahedron* **62** 8564
27. Jiao C-X, Shen Q, Huan S-Y, Shen G-L and Yu R-Q 2005 Conjugated carbazole dimer as fluorescence carrier for preparation of iodine-sensitive chemical sensor *Anal. Chim. Acta* **528** 229
28. Murali K, Sparkes H A, Pandiyan B V and Rajendra Prasad K J 2017 Synthesis, photophysical properties and DFT analysis of highly substituted pyrido carbazole-based “push pull” chromophores *New J. Chem.* **41** 8242
29. Maharani S and Ranjith Kumar R 2015 Domino four-component synthesis of novel cycloocta[b]pyridines *Tetrahedron Lett.* **56** 179