

RESEARCH ARTICLE

Genetic mapping and QTL analysis of agronomic traits in Indian *Mucuna pruriens* using an intraspecific F₂ population

S. MAHESH¹, M. LEELAMBIKA¹, MD JAHEER¹, A. M. ANITHAKUMARI² and N. SATHYANARAYANA^{3*}

¹Department of Biotechnology, Sir M. Visvesvaraya Institute of Technology, Hunasamarahalli, Bengaluru 562 157, India

²Department of Plant Breeding, Wageningen University and Research Centre, 6708 PB Wageningen, The Netherlands

³Department of Botany, Sikkim University, Gangtok, Tadong 737 102, India

Abstract

Mucuna pruriens is a well-recognized agricultural and horticultural crop with important medicinal use. However, antinutritional factors in seed and adverse morphological characters have negatively affected its cultivation. To elucidate the genetic control of agronomic traits, an intraspecific genetic linkage map of Indian *M. pruriens* has been developed based on amplified fragment length polymorphism (AFLP) markers using 200 F₂ progenies derived from a cross between wild and cultivated genotypes. The resulting linkage map comprised 129 AFLP markers dispersed over 13 linkage groups spanning a total distance of 618.88 cM with an average marker interval of 4.79 cM. For the first time, three QTLs explaining about 6.05–14.77% of the corresponding total phenotypic variation for three quantitative (seed) traits and, eight QTLs explaining about 25.96% of the corresponding total phenotypic variation for three qualitative traits have been detected on four linkage groups. The map presented here will pave a way for mapping of genes/QTLs for the important agronomic and horticultural traits contrasting between the parents used in this study.

[Mahesh S., Leelambika M., Jaheer M., Anithakumari A. M. and Sathyanarayana N. 2016 Genetic mapping and QTL analysis of agronomic traits in Indian *Mucuna pruriens* using an intraspecific F₂ population. *J. Genet.* **95**, 35–44]

Introduction

There are many indigenous food legumes whose potential is underexploited and untapped. *Mucuna pruriens*, described as a self-pollinated species, is a tropical legume classified within the phaseoloid clade of Leguminosae, also includes soybean, common bean, mung bean and its relatives. It is reported to be native of China and eastern India (Buckles 1995) and has chromosome number of $2n = 2x = 22$ (Sastrapradja *et al.* 1972; Bairiganjan and Patnaik 1989) and an estimated genome size of 1361 Mbp. The plant exhibits climbing habit, trifoliolate leaves, hairy aerial parts and a long inflorescence of white or purple flowers. The pods containing 4–6 seeds mostly green or brown in colour with thickly covered soft or stiff orange bristles, of which the latter causes intense allergic irritation to human skin.

The proximate nutritional composition, total protein content and *in vitro* protein digestibility of *M. pruriens* seeds is analogous to other edible legumes (Bressani 2002). It is therefore used as a minor food crop by many Indian and African tribes and its use as livestock feed—a common use of it in the early 1900s in the USA and elsewhere—is prevailing (Jorge *et al.* 2007). From an agronomic point, it is known to produce seed yield of 2000 kg/ha (Buckles 1995); perform well under dry farming and low soil fertility conditions (Siddhuraju *et al.* 2000), shows resistance against wide range of diseases (Eilittä *et al.* 2002); exhibits allelopathic properties (Fujii *et al.* 1991) and is effective in lowering nematode population (Carsky and Ndikawa 1998). Its positive impacts as green manure cover crop are documented in earlier studies (Tarawali *et al.* 1999; Jorge *et al.* 2007).

Seeds of *M. pruriens* contain high levels (1–6%) of L-Dopa (L-3,4-dihydroxyphenylalanine; Kavitha and Thangamani 2014), a precursor of dopamine used in the treatment of Parkinson's disease (Haq 1983). However, toxic properties of the drug and resultant side effects such as nausea, anorexia and vomiting in human beings and intestinal ailments in livestock, render the plants, particularly seeds inedible (Szabo

*For correspondence. E-mail: nsathyanarayana@cus.ac.in.
MS and LM raised mapping population, carried out marker analysis and QTL mapping. MJ assisted in field and laboratory works. AAM reviewed the manuscript. NS conceptualized the work and obtained funding.

Keywords. velvet bean; intraspecific linkage map; quantitative trait loci mapping; *Mucuna pruriens*.

and Tebbett 2002). Efforts are therefore underway to breed varieties with safe levels of L-Dopa to make it fit for consumption. On the other hand, even those plants with elevated levels of L-Dopa may be desired as a high value medicinal plant by the pharmaceutical industries.

Genetic linkage maps have emerged as valuable resources to be used as framework for a number of plant breeding applications such as marker-assisted selection (MAS), map-based cloning, physical and comparative mapping (Staub et al. 1996) etc. They are proved to be useful in precise detection of chromosomal locations and to study individual and interactive effects of genes for complex traits in several important legume species such as *Lotus japonicus* (Hayashi et al. 2001), *Medicago truncatula* (Thoquet et al. 2002); *M. sativa* (Julier et al. 2003), *Phaseolus vulgaris* (Yuste-Lisbona et al. 2012) and *Pisum sativum* (Sun et al. 2014). Recently, genetic maps have been successfully developed for even lesser known legume species such as azuki bean (Han et al. 2005); bambara groundnut (Ahmad 2012); lima bean (Bonifácio et al. 2012), yardlong bean (Kongjaimun et al. 2012) etc.

However, in *M. pruriens* except for a few studies involving molecular markers (Capo-Chichi et al. 2001; Padmesh et al. 2006; Sathyanarayana et al. 2010, 2011, 2012; Leelambika et al. 2010) and a lone published linkage map based on amplified fragment length polymorphism (AFLP) markers (Capo-Chichi et al. 2004), little effort has been made towards development and deployment of genomic resources to augment the breeding programmes.

Therefore, in the present study, an F₂ mapping population derived from a cross between two botanical varieties of *M. pruriens* and a set of AFLP markers has been used to construct first intraspecific genetic linkage map. Also, localization of the genetic loci controlling three qualitative and four quantitative traits, of which six are economically important is attempted.

Materials and methods

Plant material

The parents of the mapping population were selected based on the information in germplasm catalogue developed by the present authors. The female parent belonged to botanical variety *utilis* (velvet bean), a cultivated variety. Male parent is a wild genotype belonging to variety *pruriens* (itching bean). The crossing experiments were carried out during May–June 2009 following regular procedures of hand emasculation, dusting of donor pollens and bagging for protected pollination. True F₁ hybrids from the above interbotanical varietal cross were screened based on the dominant flower colour of the male parent. Two hundred and seventy random seeds obtained after selfing single F₁ plant were sown in the field and a F₂ population of 200 surviving individuals was obtained. All F₂ plants were grown in an open field in *Mucuna* Field Gene Bank (figure 1) established at Sir M. Visvesvaraya Institute of Technology, Bengaluru (12.58°N, 77.38°E; sandy loamy soil; 867 mm rainfall per year; channel irrigation) during January–June 2010.



Figure 1. Depiction of (a) phenotypic characters of flower, pod and seeds of F₁ hybrid and F₂ segregating phenotypes, (b) 200 F₂ segregating individuals raised in the field.

DNA isolation and AFLP marker analysis

Genomic DNA was extracted from the young leaf tissue (2–3 week old) of the F₂ individuals using modified CTAB protocol (Doyle and Doyle 1990). AFLP marker analysis was performed following the method described by the authors earlier (Sathyanarayana *et al.* 2011). The details of the adapters and oligonucleotides used as primers are provided in table 1.

The AFLP fragments were scored as dominant, i.e. presence versus absence of bands, and therefore for the parents, F₂ generation markers were assigned to either of the parental allele for map construction. Markers originating from each parent were scored according to standard coding system using A, B and H of Mapmaker (Lander *et al.* 1987) to the individual locus for map construction. The missing data was recorded as ‘-’. The AFLP markers were designated by the primer pair used followed by a number reflecting the fragment position on the gel (e.g. AACCTA4 = *Eco*RI + AAC/*Mse*I + CTA, band 4). The numbers given for the markers are in ascending order by molecular weight. A total of 805 AFLP markers were generated using 12 primer combinations of which 347 marker were polymorphic. Chi-square tests of goodness-of-fit were performed on the segregation data for all the markers.

Phenotyping of the traits

Fully matured pods were harvested from each F₂ plant for collection of seeds. Phenotyping data on length (mm), width (mm), thickness (mm) of seeds and 100-seed weight (g) were recorded as per the method described earlier. The data was recorded in five replicates for random seeds and then averaged. Qualitative traits including flower colour, fruit colour, presence of trichome on the pod, trichome colour and pod itchiness were recorded visually. Flower colour was scored as presence or absence of purple pigment, pod itchiness was

scored as presence of itchiness, and the trichome colour was scored based on ranking of colours (0–3 ranks).

Linkage mapping

Each marker was tested for the segregation ratios. The alterations in the map distances of adjacent markers caused by the removal of markers with a skewed segregation were analysed. Such highly distorted markers were excluded and the linkage map was constructed based on 129 AFLP markers using Antmap ver. 1.2 (Iwata and Ninomiya 2004) software followed by generation of a pictorial depiction of linkage groups using MapChart 2.2 (Voorrips 2002).

Initial grouping of the linkage groups was performed using nearest-neighbouring locus method following the grouping criterion of recombination with a grouping threshold at 0.3. Segregation distortion of markers was determined by chi-square goodness-of-fit analysis, while the locus ordering was done based on LL (log of likelihood) criteria. Marker distances were calculated based on the Kosambi mapping function (Kosambi 1944). The bootstrap test was also performed for the linkage groups with about 1000 iterations and the Antmap colony optimization for better linkage grouping was accomplished with 5000 iterations.

Mapping quantitative trait loci (QTL)

QTL analysis was performed using winQTLcartographer 2.5 (Wang *et al.* 2012) based on three levels namely, single marker analysis (SMA), simple interval mapping (SIM) and composite interval mapping (CIM). In SMA, the test of association between trait values and the marker loci of the genotypes was carried out at a significance levels of 5, 1, 0.1 and 0.01% indicated by *, **, *** and ****, respectively. In SIM, the association between the trait values and

Table 1. Sequence information on the AFLP adapters and primers used.

| Name of the oligonucleotide | Code | Sequence |
|-----------------------------|-------|---|
| <i>Eco</i> RI adapter | E-0 | 5'-AAT TGG TAC GCA GTC TAC-3' 3'-CC ATG CGT CAG ATG CTC-5' |
| <i>Mse</i> I adapter | M-0 | 5'-TAC TCA GGA CTC AT-3' 3'-G AGT CCT GAG TAG CAG-5' |
| <i>Eco</i> RI primer | E-A00 | 5'-GAC TGC GTA CCA ATT C A-3' |
| <i>Mse</i> I primer | M-C00 | 5'-GAT GAG TCC TGA GTA A C-3' |
| <i>Eco</i> RI primer | E-C00 | 5'-GAC TGC GTA CCA ATT C C-3' |
| <i>Mse</i> I primer | M-G00 | 5'-GAT GAG TCC TGA GTA A G-3' |
| <i>Eco</i> RI + 3-CAC | E-CAC | 5'- GAC TGC GTA CCA ATT C CAC-3' |
| <i>Eco</i> RI + 3-CAA | E-CAA | 5'-GAC TGC GTA CCA ATT C CAA-3' |
| <i>Eco</i> RI + 3-ACT | E-ACT | 5'-GAC TGC GTA CCA ATT C ACT-3' |
| <i>Eco</i> RI + 3-AAC | E-ACC | 5'-GAC TGC GTA CCA ATT C AAC-3' |
| <i>Mse</i> I + 3-GCT | M-GCT | 5'-GAT GAG TCC TGA GTA A GCT-3' |
| <i>Mse</i> I + 3-GCA | M-GCA | 5'-GAT GAG TCC TGA GTA A GCA-3' |
| <i>Mse</i> I + 3-CAT | M-CAT | 5'-GAT GAG TCC TGA GTA A CAT-3' |
| <i>Mse</i> I + 3-CAG | M-CAG | 5'-GAT GAG TCC TGA GTA A CAG-3' |
| <i>Mse</i> I + 3-CTA | M-CTA | 5'-GAT GAG TCC TGA GTA A CTA-3' |

Table 2. Comparison of morpho-agronomic characters between the parents, F₁ hybrid and F₂ population.

| Plant ID Parameters | <i>M. pruriens</i> | | F ₁ (hybrid) | F ₂ (200 random individuals) |
|------------------------|------------------------------|-----------------------------|--|---|
| | var. <i>utilis</i> ♀ | var. <i>pruriens</i> ♂ | | |
| Place of collection | Hunasamaranahalli, Karnataka | Triambakeshwar, Maharashtra | <i>Mucuna</i> field Gene Bank Bengaluru, Karnataka | |
| Latitude | 13°14'N | 20°00'N | 13°14'N | 13°14'N |
| Longitude | 77°62'E | 73°77'E | 77°62'E | 77°62'E |
| Days to emergence | 7 | 5 | 11 | 6–10 |
| TLL (cm) | 12.837 ± 2.134 | 7.81 ± 0.378 | 12.00 ± 0.238 | * |
| ALL (cm) | 12.475 ± 6.44 | 7.58 ± 0.577 | 10.00 ± 5.44 | * |
| TLW (cm) | 8.562 ± 1.53 | 3.46 ± 0.374 | 8.00 ± 0.389 | * |
| ALW (cm) | 7.825 ± 1.418 | 3.476 ± 0.447 | 6.70 ± 0.347 | * |
| TLS | Ovate | Ovate–lanceolate | Ovate | Both parental types |
| ALS | Ovate | Ovate–lanceolate | Ovate | Both parental types |
| PL (cm) | 10.925 ± 2.65 | 3.287 ± 0.652 | 11.2 ± 0.532 | * |
| LT | Membranous | Membranous | Membranous | Membranous |
| LC | Green RHS 137C | Dark green, RHS 137 A | Green RHS 137C | Both parental types |
| DF | 127 | 109 | 193 | 125–220 |
| IL (cm) | 34.33 ± 4.99 | 93.00 ± 6.82 | 23.25 ± 5.36 | * |
| NBPC | 31.00 ± 3.74 | 86.75 ± 9.73 | 22.83 ± 4.87 | * |
| FC | Yellow green RHS 145C | Dark violet, RHS 43A | Dark violet, RHS 43A | Both parental types |
| FL (cm) | 5.17 ± 0.287 | 4.68 ± 0.083 | 4.8 ± 0.073 | * |
| PTC | Creamish white | Golden orange | Dense silvery white | Both parental, F ₁ types & black velvety |
| DFMP | 140 | 131 | 260 | 150–280 |
| POL (cm) | 11.25 ± 0.45 | 8.7 ± 0.79 | 10.73 ± 0.228 | * |
| POW (cm) | 2.13 ± 0.126 | 2.15 ± 0.11 | 1.88 ± 0.083 | * |
| POC | Curved | S-shaped | Curved | Both parental types |
| PI | Nonitching | Highly itching | Highly itching | Both parental, F ₁ & extreme phenotypes |
| NPPC | 20.00 ± 5.02 | 15.00 ± 5.00 | 17 ± 1.22 | * |
| NPPP | 203 | 340 | 460 | * |
| DM | 210 | 170 | 310 | 180–320 |
| SL (mm) | 15.17 ± 1.592 | 10.52 ± 1.524 | 12.98 ± 1.301 | 11.95–18.7 |
| SW (mm) | 11.61 ± 0.662 | 7.63 ± 1.132 | 9.29 ± 0.976 | 9.78–14.57 |
| ST (mm) | 7.32 ± 0.308 | 3.83 ± 0.616 | 6.86 ± 0.924 | 5.75–9.71 |
| SC | White | Brown RHS 199C | Black | Both parental & F ₁ types, light brown, black, dark brown, black/brown mottled |
| SP | – | Black mottled | Brown mottled | |
| SS | Rhomboid | Rhomboid | Ovoid | Both parental and F ₁ types |
| NSPP | 5–6 | 4–6 | 5–6 | 5–6 |
| HSW (g) | 93.05 ± 5.84 | 27.75 ± 2.598 | 73.80 ± 7.158 | 42.98 ± 143.8 |

TLL, terminal leaflet length; ALL, adjacent leaflet length; TLW, terminal leaflet width; ALW, adjacent leaflet width; TLS, terminal leaflet shape; ALS, adjacent leaflet shape; PL, petiole length; LT, leaf texture; LC, leaf colour; DF, days to flowering; IL, inflorescence length; NBPC, no. of flower buds per cluster; FC, flower colour; FL, flower length; PTC, pod trichome colour; DFMP, days to first mature pods; POL, pod length; POW, pod width; POC, pod curvature; PI, pod itchiness; NPPC, no. of pods per cluster; NPPP, no. of pods per plant; DM, days to maturity; SL, seed length; SW, seed width; ST, seed thickness; SC, seed colour; SP, seed coat pattern; SS, seed shape; NSPP, no. of seeds per plant; HSW, hundred seeds weight; *not scored; the scale measurements are average value ± standard error; RHS, Royal Horticultural Society colour code chart.

the expected contribution of hypothetical QTL at multiple analysis points between each pair of adjacent marker loci was calculated. A manual threshold value of LR 14 (LOD 3.0) was tested through 1000 permutations scanning of the LOD curves with walk speed of 1 cM at 0.05 significance levels. Like SIM, in CIM, the possibility of a target QTL at multiple analyses points across each marker interval was

evaluated but including the effect of one or more background markers at each point. Similar statistical model like SIM was deployed for CIM.

A LOD threshold of 3.0 and above was used to declare putative QTLs since empirically computed LOD thresholds for the traits were slightly lower than 3.0. The phenotypic variance explained by each QTL (R^2) was calculated

Table 3. Statistical measures on four quantitative seed traits in F₂ population.

| Traits | Range | Mean | Standard | | Skewness |
|---------|--------|--------|--------------|---------------|----------|
| | | | Variance (n) | deviation (n) | |
| SL (mm) | 6.750 | 15.598 | 1.617 | 1.272 | 0.082 |
| SW (mm) | 4.790 | 11.445 | 0.662 | 0.814 | 0.071 |
| ST (mm) | 3.960 | 7.714 | 0.480 | 0.693 | 0.090 |
| HSW (g) | 100.82 | 96.243 | 307.396 | 17.533 | 0.182 |

SL, seed length; SW, seed width; ST, seed thickness; HSW, hundred seeds weight.

inclusive of additive and dominance effects of the markers in the latter two methods of mapping.

Results

Mapping population

The summary of phenotyping data of the parents, F₁ hybrid and the F₂ population are given in table 2. Of the 270 selfed seeds produced from an F₁ hybrid, 200 survived after planting in the field and were used as F₂ mapping population for the present work. The F₁ hybrid as well as some of the F₂ segregants fashioned an un predicted behaviour and produced variations which were not present in either of the parent (figure 1). For instance, F₁ hybrid flowered very late, in comparison to parents; likewise the flowering and maturity dates of few of the F₂ individuals were as late as F₁ hybrid and in few cases even later (table 2). Besides, the F₂ plants also produced black-velvety pods, brown to light brown seeds with both mottled as well nonmottled forms, all of which were absent in parents.

Inheritance of traits

The parental varieties differed with regard to seven qualitative traits such as flower colour, fruit colour, pod itchiness, pod trichome colour, seed shape, seed colour as well as pattern. The F₂ plants exhibited monogenic (3:1) segregation for flower colour ($\chi^2 = 0.41$ at 0.05 probability levels), while pod itchiness showed polygenic inheritance ($\chi^2 = 5.78$ at 0.05 probability levels). Also, the pod trichome colour did not fit Mendelian segregation ratios.

Correlation between the traits

The values for the mean, variance, standard deviation, coefficient of variation and range of quantitative seed traits recorded on the F₂ population are provided in table 3. Pearson correlation analysis revealed significant correlation for all the seed traits (table 4). However, high correlation coefficient value was observed between seed length and seed width (0.620).

Linkage mapping

Twelve AFLP primer combinations (table 5) used for genotyping 200 F₂ individuals produced 805 scorable markers of

Table 4. Matrix of Pearson's correlation coefficient for the four quantitative seed traits.

| Variables | SL | SW | ST | HSW |
|-----------|----|--------------|--------------|--------------|
| SL | 1 | 0.620 | 0.274 | 0.586 |
| SW | | 1 | 0.439 | 0.607 |
| ST | | | 1 | 0.582 |
| HSW | | | | 1 |

Values in bold are different from 0 with a significance level alpha = 0.01.

Table 5. Number of polymorphic markers generated between the parents using 12 AFLP primer combinations.

| Primer combination | Total no. of bands | Total no. of polymorphic bands |
|--------------------|---|--------------------------------|
| 1 E-AAC + M-CAT | 8 | 8 |
| 2 E-AAC + M-CAG | 25 | 22 |
| 3 E-AAC + M-CTA | 38 | 30 |
| 4 E-AAC + M-CTC | 52 | 34 |
| 5 E-ACT + M-CAT | 60 | 28 |
| 6 E-ACT + M-CTA | 69 | 29 |
| 7 E-ACT + M-CTC | 82 | 43 |
| 8 E-ACT + M-CAG | Not considered for analysis as shear was seen throughout wells and no polymorphic bands were seen between parents | |
| 9 E-CAA + M-GCA | 93 | 21 |
| 10 E-CAA + M-GCT | 105 | 42 |
| 11 E-CAC + M-GCA | 126 | 51 |
| 12 E-CAC + M-GCT | 147 | 39 |
| Total | 805 | 347 |

which 347 were polymorphic. The genetic linkage map constructed included 129 AFLP markers after excluding all the distorted ones distributed over 13 linkage groups spanning a total distance of 618.88 cM with an average marker interval of 4.79 cM (table 6; figure 2). Based on the number of markers and the distance coverage, the linkage groups can be classified into three main classes. First one with two linkage groups containing 14–30 markers spanning the highest distance range of all, 120.5–189.96 cM; second with 10 linkage groups with 7–11 markers covering a distance of 3.52–56.51cM and the third one with one linkage group with four markers with a distance of 7.01 cM. Segregation distortion was observed for 218 of the 347 markers leading to smaller linkage groups with 1–2 markers and large linkage groups with wide marker intervals at 0.001 probability levels and these were excluded from the analysis.

QTL mapping

Mapping of both qualitative and quantitative traits were carried out using three different methods which showed increased levels of accuracy and reliability. Firstly, single marker analysis was done for all the seven traits. A statistic measure pr (F) was calculated which basically supports

Table 6. Summary of information on genetic linkage map.

| Linkage group | Number of loci | Map length (cM) | Average marker interval (cM) |
|---------------|----------------|-----------------|------------------------------|
| 1 | 30 | 189.96 | 6.33 |
| 2 | 8 | 37.35 | 4.67 |
| 3 | 9 | 21.04 | 2.34 |
| 4 | 9 | 20.02 | 2.22 |
| 5 | 14 | 120.5 | 8.61 |
| 6 | 9 | 26.05 | 2.89 |
| 7 | 4 | 3.52 | 0.88 |
| 8 | 7 | 9.51 | 1.36 |
| 9 | 4 | 7.01 | 1.75 |
| 10 | 7 | 51.49 | 7.36 |
| 11 | 11 | 23.77 | 2.16 |
| 12 | 8 | 52.15 | 6.52 |
| 13 | 9 | 56.51 | 6.28 |
| Total | 129 | 618.88 | 4.79 |

the model of single marker associations. Based on this, the extent of association of marker with the trait was determined (table 7). Secondly, simple interval mapping was done at LOD of 3.0 with a walk speed of 1.0 cM spanning the linkage groups. Unlike SMA, SIM depicted the hypothetical QTLs for the studied traits by testing the association of marker and the trait at multiple points with additive and dominance effects of genes/markers being considered for

calculation of percentage of effective phenotypic variance. It is represented by the measure denoted as R^2 . The summary of SIM for all the traits across different groups of phenotypic variance is detailed in table 8. One QTL was identified for 100-seed weight depicting a per cent phenotypic variance of 6.82% and five QTLs for qualitative traits with trichome colour showing the highest per cent of phenotypic variance (25.96%) explained by them. Thirdly, composite interval mapping was also done at LOD of 3.0 with a walk speed of 1.0 cM spanning the linkage groups. Like simple interval mapping, CIM depicted the putative QTLs including the effect of one or more background markers at multiple analysis points across each intermarker interval. This helps in one of the two ways, depending on whether the background marker and the target interval are linked. If they are not, inclusion of the background marker makes the analysis more sensitive to the presence of a QTL in the target interval. If they are linked, it may help to separate the target QTL from other linked QTLs on the far side of the background marker. Doing so, this method has resulted in more reliable QTL identification with good illustration of phenotypic variance.

In this model again, additive and dominance effects of genes/markers are considered for calculation of percentage of effective phenotypic variance, R^2 . The summary of CIM for all the traits across different groups which explains




Figure 2. Location of QTLs for three qualitative and three quantitative traits based on combined results of SIM and CIM. The scale on the left side is the genetic distance in centimorgan (cM), marker designations are given on the left side with distance and marker names spanning on both sides of the linkage group. QTLs are shown at the right side in vertical bars with trait names in different colours for different traits. SW, black; ST, dark green; HSW, blue; FC, pink; PI, orange; TC, red. The maps were drawn by the Map Chart 2.2 program. The vertical bar shows the LOD support interval and the line LOD interval of the QTL.

Table 7. Overview of strength of single marker associations with the traits investigated.

| LG | Marker no. | Marker names | Trait association |
|----|------------|--------------|-------------------|
| 1 | 4 | CAAGCA54 | SL** |
| 8 | 2 | ACTCAT9 | SL** |
| 10 | 7 | CACGCT83 | SL** |
| 1 | 4 | CAAGCA54 | SW** |
| 1 | 6 | ACTCTC74 | SW** |
| 2 | 6 | CAAGCA79 | SW** |
| 6 | 4 | ACTCTC39 | SW** |
| 10 | 4 | ACTCTA39 | SW** |
| 5 | 4 | AACCTC25 | ST** |
| 7 | 1 | ACTCTC54 | ST** |
| 8 | 1 | CAAGCA30 | ST** |
| 11 | 5 | ACTCTC55 | ST** |
| 11 | 8 | CAAGCA8 | ST** |
| 11 | 10 | AACCAT8 | ST** |
| 11 | 11 | ACTCTA9 | ST** |
| 12 | 1 | AACCAG2 | ST** |
| 1 | 6 | ACTCTC74 | HSW** |
| 1 | 23 | ACTCTA2 | HSW** |
| 3 | 3 | AACAG18 | HSW** |
| 11 | 5 | ACTCTC55 | HSW** |
| 12 | 7 | AACCTA33 | FC** |
| 13 | 2 | ACTCAT57 | FC** |
| 13 | 7 | CAAGCA92 | FC** |
| 13 | 2 | ACTCAT57 | PI** |
| 13 | 5 | ACTCAT58 | PI*** |
| 13 | 6 | ACTCAT2 | PI** |
| 13 | 7 | CAAGCA92 | PI** |
| 13 | 2 | ACTCAT57 | TC** |
| 13 | 5 | ACTCAT58 | TC** |
| 13 | 6 | ACTCAT2 | TC** |
| 13 | 7 | CAAGCA92 | TC** |

Significance levels of 5, 1 and 0.1% are indicated by *, ** and ***, respectively.

the phenotypic variation is detailed in table 8. About two QTLs were identified for the quantitative traits depicting a

maximum phenotypic variance of 14.77% for seed thickness and six QTLs for qualitative traits with pod itchiness showing the highest per cent of phenotypic variance (21.12%) explained by them.

Generally, if we closely study both simple interval mapping and composite interval mapping results, we can conclude that CIM has identified QTLs with sharp peaks and some of the QTLs that were identified in interval mapping disappeared in CIM and some new QTLs were identified which were not identified in SIM. In our case, it was observed that CIM identified QTLs with comparatively lesser %R² recorded as against SIM; but the resolution of the peak was much sharper than SIM. Interestingly, both SIM and CIM identified three similar QTLs for pod traits with slightly varying %R² confirming the localization of QTLs for the same. Highest %R² of 25.96 and 25.79 were recorded for the QTLs identified for the traits trichome colour and pod itchiness, respectively.

The map positions for both qualitative and quantitative traits based on combined results of SIM and CIM are shown in figure 2.

Discussion

Little is done to develop genomic resources for underutilized plant species in general and *M. pruriens* in particular. Alone linkage map using US core collection (Capo-Chichi *et al.* 2004), published earlier, demonstrated prospects of good genome coverage for linkage studies with AFLP markers, in addition to segregation of pod colour and pod pubescence in F₂ population. Beyond this, there was no effort in the direction of trait-based mapping or QTL studies from this or any other work from any part of the world till date on this crop. Therefore, we present here the first report on the map positions for floral, pod and seed traits in *M. pruriens* with genome coverage analogous to earlier published map.

Table 8. QTLs illustrating the per cent phenotypic variance of traits at different LOD thresholds based on SIM and CIM.

| Trait | LG | Marker | Position | Marker interval | LOD | ADD | DOM | %R ² | Method |
|-------|----|--------|----------|--------------------|-------|-------|-------|-----------------|--------|
| SW-1 | 1 | 4 | 10.5 | CACGCT8–CAAGCA58 | 3.55 | 0.26 | 2.14 | 6.05 | CIM |
| ST-1 | 5 | 3 | 39.2 | CACGCA118–AACCTC25 | 4.22 | –0.37 | 0.33 | 14.77 | CIM |
| HSW-1 | 3 | 2 | 7 | CAAGCA9–AACAG18 | 3.17 | 9.44 | 9.92 | 6.82 | SIM |
| FC-1 | 1 | 3 | 8 | ACTCTC40–CAAGCA18 | 3.76 | –0.25 | 0.16 | 7.88 | CIM |
| PI-1 | 13 | 1 | 3 | ACTCTC10–ACTCAT57 | 3.98 | –0.30 | 0.43 | 14.29 | SIM |
| PI-2* | 13 | 4 | 35.6 | ACTCAT54–ACTCAT58 | 5.98 | –0.37 | 0.57 | 25.79 | SIM |
| | 13 | 4 | 37.6 | ACTCAT54–ACTCAT58 | 6.48 | –0.34 | 0.55 | 21.12 | CIM |
| PI-3* | 13 | 5 | 52 | ACTCAT58–ACTCAT2 | 3.16 | –0.13 | –0.61 | 5.31 | SIM |
| | 13 | 5 | 52 | ACTCAT58–ACTCAT2 | 3.75 | –0.12 | –0.70 | 4.66 | CIM |
| TC-1 | 1 | 7 | 44 | CAAGCT8–CACGCA39 | 12.98 | 0.14 | –1.85 | 2.69 | CIM |
| TC-2 | 5 | 13 | 117 | CAAGCA90–CACGCA79 | 3.24 | 0.48 | –1.74 | 6.00 | CIM |
| TC-3 | 13 | 1 | 4 | ACTCTC10–ACTCAT57 | 8.64 | –0.10 | –2.04 | 5.19 | SIM |
| TC-4* | 13 | 4 | 34.6 | ACTCAT54–ACTCAT58 | 7.04 | –0.95 | 0.91 | 25.96 | SIM |
| | 13 | 4 | 34.6 | ACTCAT54–ACTCAT58 | 5.47 | –0.93 | 0.85 | 16.66 | CIM |

The positive and negative effects indicate that the allele which increases the trait values is in paternal and maternal parents, respectively. *Similar QTLs identified in both SIM and CIM.

The F₁ intraspecific hybrid as well F₂ segregating population showed surprising behaviour and thrown up a few variations which was not present in either parent in this study. This is plausible as the two genotypes used for developing mapping population belong to different botanical varieties, and consequently might have behaved analogous to interspecific hybrid owing to large genetic variations between them. Further, this must have resulted in transgressive type of segregation inducing novel phenotypes in F₂ individuals which is common observation in the progenies involving interspecific mating (Lotsy 1916; Brainerd 1924; Darlington and Mather 1949; Vega and Frey 1980). It is necessary, however, to raise different subsets of F₂ individuals from the same F₁ hybrid at multienvironments to conform this.

High percentage of segregation distortion (62.8%) was observed for the polymorphic markers between the parents of F₂ population. The fact that markers with segregation distortion are clustered around particular region in the linkage groups indicates that segregation distortion in the F₂ population is most likely caused by the genetic factors. At the same time even distribution across a few groups are also observed indicating statistical bias (Plomion *et al.* 1995). Moreover, since statistical bias likely emerging out of genotyping and/or scoring errors are eliminated beforehand in this study by analysing each marker by altering the position and distance between the adjacent markers of different linkage groups, and complete exclusion of such skewed markers from the analysis, these distortions are majorly attributed to genetic factors.

The linkage map realized in this work serves as workable genetic map for *M. pruriens*. It is necessary that such a map should have mean genetic distance of 20 and 10 cM between adjacent markers for mapping of major genes and QTLs, respectively (Xu and Zhu 1994; Fang *et al.* 2000). With the maximum marker interval of 8.61 cM and an average marker interval of 4.79 cM, the present map essentially meets these prerequisites. However, much is still awaited in terms of developing saturated map to facilitate detection of closely linked or functional markers for marker-assisted selection and map-based cloning of genes of interest.

In this study, relatively higher centimorgan distances were observed (in LG's: 1, 5, 10, 12 and 13). Literature suggests that they might have caused due to higher, but <50.0%, total recombination percentages due to higher expected double crosses in addition to the single crossover events (Haldane 1919). Nonetheless, this remains to be verified. Also, the existence of minor linkage groups and several unlinked markers indicates that there are many large gaps with a few markers. This might be due to relatively limiting population size, coupled with low map saturation. However, they may merge into larger groups as and when more markers are assigned to map. Deployment of superior and codominant marker types such as SSRs, SNPs, besides additional AFLP markers generated from primer sets other than *EcoRI/MseI* will be valuable for this purpose.

All the seed traits and pod trichome showed expected segregation in F₂ population. But, the trichome colour did not segregate in Mendelian ratio. Also intermediates were observed for this trait in both F₁ the F₂ population implying the codominant nature of the genes involved. The range of variability both for trichome colour as well as its effect, itching intensity observed in the population along with the three QTLs identified suggests that the trait is polygenic. The two overlapping QTLs identified for these two traits suggest that they are also highly correlated. The trichomes in *M. pruriens* contain chemical substances, mucunain and serotonin, which are known to be responsible for itching, blisters and dermatitis caused by its contact (Sastry and Kavathekar 1990). The fact that the intensity of trichome colour is directly proportional to severity of pod itchiness points to some interactions between genes / biochemical pathways responsible for both the traits underlying these QTLs. Further investigations are needed on molecular and biochemical basis of mucunain and serotonin variations along with dense covering of linkage groups with genomewide markers to unveil the precise loci and/or genes linked to these traits.

In case of quantitative traits, highest correlation was observed between the seed traits SL and SW (0.62). Besides, good correlation was also observed between SL and HSW (0.586); SW and HSW (0.607) and ST and HSW (0.582). Correlation percentage of >50% observed between the seed size traits (seed length, width and thickness) with seed mass suggest high genetic relatedness between them implying their usefulness in breeding programmes.

In conclusion, this is the first report on the development of workable linkage map and identification of QTLs for floral, pod and seed characteristics in *M. pruriens* which form the basis for future genetic studies in this promising species. However, low genome coverage and use of dominant marker (AFLP) are the major limitations. Besides, variation among the F₂ plants which might have included environmental variation is also not accounted. Thus, further experimentations by growing different F₂ subsets at varying environments or developing RIL population besides saturation of linkage map with codominant markers such as SSRs and SNPs will reinforce the findings of the above work and help appropriate it for molecular breeding applications.

Acknowledgements

The authors acknowledge financial support from the Department of Science and Technology (DST) and Department of Biotechnology (DBT), Govt. of India for this project. NS thank Sikkim University, Gangtok and Sri Krishnadevaraya Educational Trust (Sri KET), Bangalore, India for the laboratory and field facilities.

References

- Ahmad N. S. 2012 Genetic analysis of plant morphology in Bambara groundnut (*Vigna subterranea* (L.) Verdc.). BSc Thesis. University of Nottingham, Nottinghamshire, UK.

- Bairiganjan G. C. and Patnaik S. N. 1989 Chromosome evolution in Fabaceae. *Cytologia* **54**, 51–64.
- Brainerd E. 1924 Some natural violet hybrids of North America. *Vermont Agr. Exp. Sta. Bull.* No. 239.
- Bressani R. 2002 Factors influencing nutritive value in food grain legumes: *Mucuna* compared to other grain legumes. In *Food and feed from Mucuna: current uses and the way forward* (ed. B. M. Flores, M. Eilittä, R. Myhrman, L. B. Carew and R. J. Carsky) pp. 164–188, Workshop, CIDICCO, CIEPCA and World Hunger Research Center, Tegucigalpa Honduras, USA.
- Bonifácio E. M., Fonsêca A., Almeida C., Dos Santos K. G. B. and Pedrosa-Harand A. 2012 Comparative cytogenetic mapping between the lima bean (*Phaseolus lunatus* L.) and the common bean (*P. vulgaris* L.). *Theor. Appl. Genet.* **124**, 1513–1520.
- Buckles D. 1995 Velvetbean: a “new” plant with a history. *Econ. Bot.* **49**, 13–25.
- Capo-Chichi L. J. A., Weaver D. B. and Morton C. M. 2001 AFLP assessment of genetic variability among Velvetbean (*Mucuna* sp.) accessions. *Theor. Appl. Genet.* **103**, 1180–1188.
- Capo-Chichi L. J. A., Weaver D. B. and Morton C. M. 2004 An intraspecific genetic map of velvetbean (*Mucuna* sp.) based on AFLP markers. *Theor. Appl. Genet.* **108**, 814–821.
- Carsky R. J. and Ndikawa R. 1998 Identification of cover crops for the semi-arid savanna zone of west Africa. In *Cover crops in west Africa—contributing to sustainable agriculture* (ed. D. Buckles, A. Eteka, M. Osiname, M. Galiba and G. Galiano) pp. 179–187, IDRC, IITA, Sasakawa Global 2000, Ottawa, Canada, Ibadan, Nigeria, Cotonou, Benin.
- Darlington C. D. and Mather K. 1949 *The elements of genetics*. Allen & Unwin, London, UK.
- Doyle J. J. and Doyle J. 1990 Isolation of DNA from plant tissue. *Focus* **12**, 13–15.
- Eilittä M., Bressani R., Carew L. B., Carsky R. J., Flores M., Gilbert R. et al. 2002 *Mucuna* as a food and feed crop: an overview. In *Food and feed from Mucuna: current uses and the way forward*, (ed. B. M. Flores, M. Eilittä, R. Myhrman, L. B. Carew and R. J. Carsky) pp 18–47, workshop CIDICCO, CIEPCA and World Hunger Research Center, Tegucigalpa, Honduras USA.
- Fang X., Wu W. and Tang J. 2000 *DNA marker-assisted breeding in crops*. Scientific Press, Beijing, China.
- Fujii Y., Shibuya T. and Yasuda T. 1991 L-3,4-dihydroxyphenylalanine as an allelochemical candidate from *Mucuna pruriens* (L.) DC. var. *utilis*. *Agric. Biol. Chem.* **55**, 617–618.
- Haldane J. B. S. 1919 The combination of linkage values and the calculation of distances between the loci of linked factors. *J. Genet.* **8**, 299–309.
- Han O. K., Kaga A., Isemura T., Wang X. W., Tomooka N. and Vaughan D. A. 2005 A genetic linkage map for azuki bean (*Vigna angularis* (Wild.) Ohwi & Ohashi). *Theor. Appl. Genet.* **111**, 1278–1287.
- Haq N. 1983 New food legume crops for the tropics, In *Better crops for the food* (ed. J. Nugent and M. O. Cormor) pp. 144–160. Pitman Books, London (Cuba Foundation Symposium, 97), UK.
- Hayashi M., Miyahara A., Sato S., Kato T., Yoshikawa M., Taketa M. et al. 2001 Construction of a genetic linkage map of the model legume *Lotus japonicus* using an intraspecific F₂ population. *DNA Res.* **8**, 301–310.
- Iwata H. and Ninomiya S. 2004 *Ant colony optimization for linkage grouping and locus ordering in genome mapping*. Plant and Animal Genome XII, San Diego, USA.
- Jorge M. A., Eilittä M., Proud F. J., Maasdorp B. V., Beksissa H., Sarial A. K. and Hanson 2007 *Mucuna* species: recent advances in application of biotechnology. *Fruit Veg. Cer. Sci. Biotech.* **2**, 80–94.
- Julier B., Flajoulot S., Barre P., Cardinet G., Santoni S., Huguet T. and Huyghe H. 2003 Construction of two genetic linkage maps in cultivated tetraploid alfalfa (*Medicago sativa*) using microsatellite and AFLP markers. *BMC Plant Biol.* **3**, 9.
- Kavitha C. and Thangamani C. 2014 Amazing bean “*Mucuna pruriens*”—a comprehensive review. *J. Med. Plants Res.* **8**, 138–143.
- Kongjaimun A., Kaga A., Tomooka N., Somta P., Shimizu T., Shu Y. et al. 2012 An SSR-based linkage map of yardlong bean (*Vigna unguiculata* (L.) Walp. sub-sp. *unguiculata* Sesquipedalis group) and QTL analysis of pod length. *Genome* **55**, 81–92.
- Kosambi D. D. 1944 The estimation of map distances from recombination values. *Ann. Eugen.* **12**, 172–175.
- Lander E. S., Green P., Abrahamson J., Barlow A., Daly M. J., Lincoln S. E. and Newburg L. 1987 MAPMAKER: an interactive computer package of constructing primary genetic linkage maps of experimental and natural populations. *Genomics* **1**, 174–181.
- Leelambika M., Mahesh S., Jaheer M. and Sathyanarayana N. 2010 Comparative evaluation of genetic diversity among Indian *Mucuna* species using morphometric, biochemical and molecular approaches. *World J. Agric. Sci.* **6**, 568–578.
- Lotsy J. P. 1916 *Evolution by means of hybridization*. Martinees Nijhoff, The Hague, The Netherlands.
- Padmesh P., Reji J. V., Jinish Dhar M. and Seenii D. 2006 Estimation of genetic diversity in varieties of *M. pruriens* using RAPD. *Biol. Plant.* **50**, 367–372.
- Plomion C., O’Malley D. M. and Durel C. E. 1995 Genomic analysis in maritime pine (*Pinus pinaster*): comparison of two RAPD maps using selfed and open-pollinated seeds of the same individual. *Theor. Appl. Genet.* **90**, 1028–1034.
- Sastrapradja S., Sastrapradja D., Aminah S. H., Lubis I. and Idris S. 1972 Comparative seedling morphology of *Mucuna pruriens* group. *Ann. Bogoriendes.* **5**, 131–136.
- Sastry C. S. T. and Kavathekar Y. Y. 1990 Plants for reclamation of wastelands. Publications and Information Directorate, New Delhi, India.
- Sathyanarayana N., Vikas P. B., Bharath Kumar T. N. and Rajesha R. 2010 RAPD markers for genetic characterization of *Mucuna* species. *Ind. J. Genet.* **70**, 296–298.
- Sathyanarayana N., Leelambika M., Mahesh S. and Jaheer M. 2011 AFLP assessment of genetic diversity among Indian *Mucuna* accessions. *Physiol. Mol. Biol. Plants* **17**, 171–180.
- Sathyanarayana N., Mahesh S., Jaheer M. and Leelambika M. 2012 Genetic diversity of wild and cultivated *M. pruriens* (L.). DC accessions analyzed using thirty morpho-agronomical characters. *Trop. Subtrop. Agroecosyst.* **15**, 249–259.
- Siddhuraju P., Becker K. and Makkar H. P. 2000 Studies on the nutritional composition and antinutritional factors of three different germplasm seed materials of an under-utilized tropical legume, *Mucuna pruriens* var. *utilis*. *J. Agric. Food Chem.* **48**, 6048–6060.
- Staub J. E., Serquen F. C. and Gupta M. 1996 Genetic markers, map construction, and their application in plant breeding. *Hort. Sci.* **31**, 729–740.
- Szabo N. J. and Tebbett I. R. 2002 The chemistry and toxicity of *Mucuna* species. In *Food and feed from Mucuna: current uses and the way forward* (ed. M. Flores, M. Eilittä, R. Myhrman, L. B. Carew, R. J. Carsky) pp. 120–141. Proceedings of an International Workshop, CIDICCO, CIEPCA, World Hunger Research Center, Tegucigalpa, Honduras, USA.
- Sun X., Yang T., Hao J., Xiaoyan Z., Ford R., Jiang J. et al. 2014 SSR genetic linkage map construction of pea (*Pisum sativum* L.) based on Chinese native varieties. *Crop J.*, 170–174.
- Tarawali G., Manyong V. M., Carsky R. J., Vissoh P. V., Osei-Bonsu P. and Galiba M. 1999 Adoption of improved fallows in west Africa: lessons from *Mucuna* and *Stylo* case studies. *Agroforestry Syst.* **47**, 93–122.
- Thoquet P., Ghérardi M., Journet E.-P., Kereszt A., Ané J.-M., Prosperii J.-M. and Hugust T. 2002 The molecular genetic

- linkage map of the model legume *Medicago truncatula*: an essential tool for comparative legume genomics and the isolation of agronomically important genes. *BMC Plant Biol.* **2**, 1.
- Vega U. and Frey K. J. 1980 Transgressive segregation in inter and intraspecific crosses of barley. *Euphytica* **2**, 585–594.
- Voorrips R. E. 2002 MapChart: software for the graphical presentation of linkage maps and QTLs. *J. Hered.* **93**, 77–78.
- Wang S., Basten C. J. and Zeng Z. B. 2012 Windows QTL Cartographer 2.5. Department of Statistics, North Carolina State University, Raleigh, NC (<http://statgen.ncsu.edu/qtlcart/WQTLCart.htm>).
- Xu Y. and Zhu L. 1994 *Molecular quantitative genetics*. Chinese Agricultural Press, Beijing, China.
- Yuste-Lisbona F. J., Santalla M., Capel C., García-Alcázar M., De La Fuente M., Capel J. *et al.* 2012 Marker-based linkage map of Andean common bean (*Phaseolus vulgaris* L.) and mapping of QTLs underlying popping ability traits. *BMC Plant Biol.* **12**, 136.

Received 18 January 2015, in revised form 21 May 2015; accepted 26 June 2015

Unedited version published online: 6 July 2015

Final version published online: 4 February 2016