

# Iran Smell Identification Test (Iran-SIT): a Modified Version of the University of Pennsylvania Smell Identification Test (UPSIT) for Iranian Population

Safa Taherkhani<sup>1</sup>  · Fathollah Moztarzadeh<sup>1</sup> · Jalal Mehdizadeh Seraj<sup>2</sup> · Seyed Saeed Hashemi Nazari<sup>3,4</sup> · Farzad Taherkhani<sup>5</sup> · Jaber Gharehdaghi<sup>6</sup> · Arash Okazi<sup>7</sup> · Sevda Pouraghaei<sup>1</sup>

Received: 25 October 2014 / Accepted: 20 July 2015 / Published online: 5 August 2015  
© Springer Science+Business Media New York 2015

## Abstract

**Introduction** Based on cultural adaption, various types of olfactory tests have been designed in different countries. The aim of this study was to develop a standardized smell identification test to evaluate the olfactory function of Iranian population.

**Methods** For designing Iran smell identification test (Iran-SIT), we selected the most familiar odors for Iranians, and prepared a 24-item smell identification test in four-alternative forced-choice paradigm. We tested 577 healthy subjects aged 6 to 68 years by Iran-SIT. In order to assess

the reliability and stability of Iran-SIT over time, 96 subjects participated in the retest study after 5 months.

**Results** All odors of Iran-SIT were identified by more than 70 % of subjects. Identification scores were significantly changed by age. Children and elderly subjects represented significantly lower identification scores than adult aged 20–50 years. There were no significant differences between adult aged 20–50 years. Test-retest study revealed that Iran-SIT is a highly reliable and valid test (Pearson's correlation coefficient between test-retest identification scores:  $r=0.93$ ). In order to classify adult patients into four olfactory disorder levels, we determined diagnostic criterion of olfactory disorder using Iran-SIT based on the identification score obtained from subjects aged 20–50 years.

**Conclusions** Iran-SIT with high reliability and validity has adequacy to distinguish among normosmia, mild microsmia, severe microsmia, and anosmia.

✉ Safa Taherkhani  
safa.taherkhani@gmail.com

✉ Seyed Saeed Hashemi Nazari  
saeedh\_1999@yahoo.com

<sup>1</sup> Department of Biomedical Engineering (Center of Excellence), Amirkabir University of Technology, P. O. Box 15875-4413, Tehran, Iran

<sup>2</sup> Department of Otolaryngology, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup> Safety Promotion and Injury Prevention Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>4</sup> Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, P. O. Box 14155-6153, Tehran, Iran

<sup>5</sup> Department of Mechanical Engineering, Tehran University, Tehran, Iran

<sup>6</sup> Department of Legal Medicine Research Center, Tehran University of Medical Sciences, Tehran, Iran

<sup>7</sup> Forensic Medicine Department, Tehran University of Medical Sciences, Tehran, Iran

**Keywords** Smell identification test (SIT) · Iranian population · Chemical sense · Olfactory function · Anosmia · Age

## Introduction

The sense of smell widely affects our quality of life through determining the palatability of foods and beverages, enjoyment of flowers and perfumes, reproduction of memory and warning for dangerous situations like spoiled food, fire, and gas leakage (Murphy 1985; Landis et al. 2005). Olfactory disorder may be caused by a number of factors such as sinus or nasal disease, head trauma, toxic chemical exposure, using some drugs, neurodegenerative disease, aging etc. (Murphy 1985; Landis et al. 2005; Doty 2001). Clinical evaluation of

olfactory performance is an essential step to the diagnosis and treatment of olfactory dysfunction. Psychophysical assessment methods consist of odor detection and recognition threshold, odor detection, odor identification, odor discrimination, and odor memory tests, providing an effective and practical way to rapid assessment of olfactory function (Harper et al. 1968; Cain et al. 1992; Doty 1995; Eibenstein et al. 2005).

The most famous olfactory test, University of Pennsylvania smell identification test (UPSIT) is widely used as a diagnostic tool in researches and clinical settings in the USA (Doty et al. 1984a, b). UPSIT focuses on the comparative ability of individuals to identify various odors. It has efficiency for detecting malingering as well as some olfactory disorders (Doty 1995). As the sense of smell highly depends on social/cultural lifestyle of population, familiarity with the odors in the smell test is a considerable matter. Sniffin' Sticks is another olfactory performance assessment test adjusted for European people with a combination of odor identification, odor discrimination, and olfactory threshold tests (Kobal et al. 1996; Hummel et al. 1997). However, indeed, there are some odors in Sniffin' Sticks, which are unfamiliar for the people in some European countries (CĂtanĂ et al. 2012). In recent years, many efforts have been made to standardize olfactory tests according to cultural features (Nordin et al. 2002; Thomas-Danguin et al. 2003; Cardesín et al. 2006; Saito et al. 2006; Cho et al. 2009; Silveira-Moriyama et al. 2010; CĂtanĂ et al. 2012; Oniz et al. 2013). In other words, there is no universal gold standard test for the assessment of olfactory function.

The major issue of this study was the absence of an accepted smell test in Iran. Iranian otolaryngologists have difficulties in determining the degree of olfactory disorders. Although most of otolaryngologists use handmade traditional bottles filled with different fragrances, some of them uses the original version of the University of Pennsylvania smell identification test (UPSIT) or Sniffin' Sticks. Despite this situation, neither the original tests nor the traditional bottles are validated for Iranian population.

In this study, we tried to design a reliable and valid test for clinical assessment of olfactory function based on familiar odors in Iranian population named Iran smell identification test (Iran-SIT). Here, we report the results of pilot, main, and test-retest studies.

## Methods

### Determining Familiar Odors for Iranian Population

Two famous tests, University of Pennsylvania smell identification test (UPSIT) and Sniffin' Sticks, are comprehensively used as the main reference for the smell identification tests developed in different countries (Doty et al. 1984a, b; Kobal

et al. 1996; Hummel et al. 1997). In this study, we considered the 40 odors of the UPSIT as the base of our test. Iran is a big country with different ethnicities, so the odors must be familiar for all Iranians. Thousands of students from all around the country live in Tehran (mostly in dormitories). They represent the cultural diversity of Iran very well. Ninety students were asked to list the familiar odors of the UPSIT. For better linguistic perception, we prepared a translation of the original version of UPSIT in Farsi. In order to replace unfamiliar odors, they were also asked to propose some odors which people from different regions of Iran are commonly encountered in their daily life. By considering the odor categories proposed by Castro et al. (2013) and feasibility of supplying the odorants, the first version of Iran-SIT was designed using 40 items.

### Preparing the First Version of Iran-SIT

In order to prepare the first version of Iran-SIT, the natural or synthetic odorants were provided and fragrance microcapsules were produced unless they were commercially available. Afterwards, to prepare the scratch and sniff stickers, microcapsules were mixed with varnish ink and printed on sticker papers using silk screen printer machine. Finally, a questionnaire containing these stickers was designed in a four-alternative forced-choice (4-AFC) test format.

### Pilot Study

Before the main experiment, a pilot study was carried out by 43 subjects (23 female and 20 male) with ages ranging from 20 to 40 years using the first version of Iran-SIT. The aim of pilot study was to reveal deficiencies in the procedure and select the best odors from 40 items and their alternatives.

**Table 1** The number of subjects (female, male, and total) of each age group

Age Group	Female	Male	Total
5–9	24	23	47
10–14	24	24	48
15–19	28	25	53
20–24	35	31	66
25–29	34	36	70
30–34	30	31	61
35–39	27	24	51
40–44	24	22	46
45–49	18	18	36
50–54	16	15	31
55–59	16	12	28
60–64	12	9	21
65–69	10	9	19
Sum	298	279	577

The subjects were asked to scratch the stickers, sniff them, and choose one of the four alternatives. The data obtained from this experiment was analyzed, and 16 odors with lowest identification score were omitted and some alternatives were switched.

## Main Study

Five hundred seventy-seven healthy subjects (mean age 32.46, SEM (standard error of the mean) 0.681), 298 female (mean age 32.76, SEM 0.97) and 279 male

**Table 2** Forty odors and alternatives of each odor used in UPSIT and the first version of Iran-SIT

UPSIT		The first version of Iran-SIT	
Odors	Alternatives	Odors	Alternatives
Pizza	(Gasoline, peanuts, lilac)	Pizza	(Gasoline, peanuts, <b>jasmine</b> )
Bubble gum	(Dill pickle, wintergreen, watermelon)	Bubble gum	( <b>Cucumber, pizza</b> , watermelon)
Menthol	(Tomato, licorice, strawberry)	<i>Cola</i>	( <b>Basil, fish, pear</b> )
Cherry	(Whiskey, honey, lime)	Cherry	( <b>Vinegar</b> , honey, <b>lemon</b> )
Motor oil	(Grass, pizza, pineapple)	<i>Fish</i>	( <b>Smoke</b> , pizza, pineapple)
Mint	(Skunk, fruit punch, cola)	Mint	( <b>Black pepper</b> , fruit punch, cola)
Banana	(Garlic, cherry, motor oil)	Banana	(Garlic, cherry, <b>gasoline</b> )
Clove	(Licorice, chili, banana)	<i>Tuberose</i>	( <b>Saffron</b> , chili, banana)
Leather	(Clove, lilac, apple)	<i>Honey</i>	( <b>Olive oil, jasmine</b> , apple)
Coconut	(Skunk, cedar, honey)	Coconut	( <b>Smoke, cardamom</b> , honey)
Onion	(Chocolate, banana, fruit punch)	Onion	(Chocolate, banana, fruit punch)
Fruit punch	(Soap, menthol, pumpkin pie)	<i>Kiwifruit</i>	(Soap, <b>mint, honey</b> )
Licorice	(Pineapple, cheddar cheese, cherry)	<i>Vanilla</i>	(Pineapple, <b>old cheese</b> , cherry)
Cheddar cheese	(Paint thinner, cherry, coconut)	<i>Fried chicken</i>	(Paint thinner, cherry, coconut)
Cinnamon	(Cola, pine, coconut)	Cinnamon	(Cola, <b>strawberry</b> , coconut)
Gasoline	(Rose, lemon, peach)	<i>Butter</i>	(Rose, lemon, peach)
Strawberry	(Dill pickle, chocolate, cedar)	Strawberry	( <b>Olive oil</b> , chocolate, <b>thymus</b> )
Cedar	(Gasoline, lemon, root beer)	<i>Biscuit</i>	( <b>Tuberose, cologne, orange</b> )
Chocolate	(Lemon, root beer, black pepper)	Chocolate	(Lemon, <b>vinegar</b> , black pepper)
Gingerbread	(Menthol, apple, cheddar cheese)	<i>Cake</i>	( <b>Mint</b> , apple, <b>old cheese</b> )
Lilac	(Chili, coconut, whiskey)	<i>Jasmine</i>	( <b>Smoke, fish, lemon</b> )
Turpentine	(Soap, skunk, chili)	<i>Eucalyptus</i>	( <b>Coconut, saffron, peach</b> )
Peach	(Chocolate, leather, pizza)	Peach	(Chocolate, <b>coffee</b> , pizza)
Root beer	(Watermelon, banana, smoke)	<i>Hami melon</i>	( <b>Cinnamon, vinegar, kebab</b> )
Dill pickle	(Pineapple, root beer, black pepper)	<i>Mango</i>	( <b>Persil, saffron, fried chicken</b> )
Pineapple	(Smoke, whiskey, onion)	Pineapple	(Smoke, <b>natural gas</b> , onion)
Lime	(Musk, garlic, turpentine)	<i>Dried lim</i>	( <b>Rose water</b> , garlic, peanut)
Orange	(Cheddar cheese, bubble gum, turpentine)	Orange	( <b>Old cheese</b> , bubble gum, <b>garlic</b> )
Wintergreen	(Lime, pumpkin pie, leather)	Cucumber	( <b>Cherry, honey</b> , leather)
Watermelon	(Chili, menthol, orange)	Watermelon	(Chili, <b>smoke</b> , orange)
Paint thinner	(Watermelon, peanut, rose)	<i>Cologne</i>	(Watermelon, peanut, <b>onion</b> )
Grass	(Mint, gingerbread, strawberry)	<i>Coffee</i>	(Mint, <b>bread</b> , strawberry)
Smoke	(Dill pickle, grass, peach)	Smoke	( <b>Mint, jasmine</b> , peach)
Pine	(Smoke, lilac, orange)	<i>Apple</i>	(Smoke, <b>jasmine</b> , orange)
Grape	(Pizza, turpentine, clove)	Grape	(Pizza, <b>garlic, tuberose</b> )
Lemon	(Motor oil, pumpkin pie, rose)	<i>Tangerine</i>	( <b>Gasoline, honey</b> , rose)
Soap	(Black pepper, licorice, peanut)	Soap	(Black pepper, <b>onion</b> , peanut)
Natural gas	(Orange, musk, cola)	<i>Garlic</i>	( <b>Chocolate, cucumber</b> , cola)
Rose	(Lime, mint, bubble gum)	<i>Rose water</i>	( <b>Orange</b> , mint, bubble gum)
Peanut	(Lemon, apple, root beer)	Peanut	(Lemon, apple, rose water)

Replaced odors are in italics and changed alternatives are in bold

(mean age 32.13, SEM 0.96) from different regions of Iran with ages ranging from 6 to 68 years were selected to participate in main study. They were classified in thirteen 5-year age groups. The number of people in

each age group was based on Iran demographics pattern (Asia-Pacific Population Journal 2006). The number of subjects (female, male, and total) of each age group is shown in Table 1. All subjects gave their informed

**Table 3** Identification rate of each odor used in the first version of Iran-SIT

No	Odor	Identification rate (%)	Female Identification (%)	Male Identification (%)
1	Banana	100.0	100.0	100.0
2	Garlic	100.0	100.0	100.0
3	Cologne	100.0	100.0	100.0
4	Rose water	97.7 (84.2–99.6)	100.0	95.0 (67.7–99.4)
5	Mint	97.7 (84.2–99.6)	95.7 (71.5–99.4)	100.0
6	Hami melon	97.7 (84.2–99.6)	100.0	95.0 (67.7–99.4)
7	Chocolate	97.7 (84.2–99.6)	95.7 (71.5–99.4)	100.0
8	Watermelon	97.7 (84.2–99.6)	95.7 (71.5–99.4)	100.0
9	Bubble gum	95.3 (82.3–98.9)	95.7 (71.5–99.4)	95.0 (67.7–99.4)
10	Cola	95.3 (82.3–98.9)	91.3 (68.6–98.0)	100.0
11	Biscuit	95.3 (82.3–98.9)	91.3 (68.6–98.0)	100.0
12	Cinnamon	93.0 (79.7–97.8)	95.7 (71.5–99.4)	90.0 (64.4–97.8)
13	Coconut	93.0 (79.7–97.8)	91.3 (68.6–98.0)	95.0 (67.7–99.4)
14	Pineapple	93.0 (79.7–97.8)	87.0 (64.2–96.1)	100.0
15	Smoke	93.0 (79.7–97.8)	87.0 (64.2–96.1)	100.0
16	Peach	90.7 (76.9–96.6)	91.3 (68.6–98.0)	90.0 (64.4–97.8)
17	Onion	90.7 (76.9–96.6)	87.0 (64.2–96.1)	95.0 (67.7–99.4)
18	Grape	88.4 (74.2–95.2)	82.6 (59.6–93.8)	95.0 (67.7–99.4)
19	Peanut	86.0 (71.5–93.8)	91.3 (68.6–98.0)	80.0 (54.6–93.0)
20	Strawberry	86.0 (71.5–93.8)	95.7 (71.5–99.4)	75.0 (49.7–90.0)
21	Apple	83.7 (68.8–92.2)	78.3 (55.2–91.3)	90.0 (64.4–97.8)
22	Dried lime	79.1 (63.7–89.0)	87.0 (64.2–96.1)	70.0 (45.0–86.9)
23	Soap	74.4 (58.7–85.5)	69.6 (46.6–85.6)	80.0 (54.6–93.0)
24	Jasmine	74.4 (58.7–85.5)	73.9 (50.8–88.5)	75.0 (49.7–90.0)
25	Pizza	69.8 (53.9–81.9)	69.6 (46.6–85.6)	70.0 (45.0–86.9)
26	Fried chicken	69.8 (53.9–81.9)	73.9 (50.8–88.5)	65.0 (40.4–83.5)
27	Vanilla	67.4 (51.5–80.1)	60.9 (38.6–79.3)	75.0 (49.7–90.0)
28	Mango	62.8 (46.9–76.2)	60.9 (38.6–79.3)	65.0 (40.4–83.5)
29	Honey	60.5 (44.7–74.3)	65.2 (42.5–82.5)	55.0 (31.7–76.2)
30	Eucalyptus	55.8 (40.2–70.2)	47.8 (27.4–68.9)	65.0 (40.4–83.5)
31	Tuberose	53.5 (38.1–68.2)	52.2 (31.0–72.5)	55.0 (31.7–76.2)
32	Orange	51.2 (35.9–66.1)	43.5 (23.9–65.2)	60.0 (36.0–79.9)
33	Kiwifruit	51.2 (35.9–66.1)	47.8 (27.4–68.9)	55.0 (31.7–76.2)
34	Cake	48.8 (33.8–64.0)	52.2 (31.0–72.5)	45.0 (23.7–68.2)
35	Cherry	46.5 (31.7–61.8)	43.5 (23.9–65.2)	50.0 (27.6–72.3)
36	Fish	46.5 (31.7–61.8)	52.2 (31.0–72.5)	40.0 (20.0–63.9)
37	Cucumber	44.2 (29.7–59.7)	47.8 (27.4–68.9)	40.0 (20.0–63.9)
38	Tangerine	41.9 (27.6–57.5)	43.5 (23.9–65.2)	40.0 (20.0–63.9)
39	Butter	39.5 (25.6–55.2)	34.8 (17.4–57.4)	45.0 (23.7–68.2)
40	Coffee	34.9 (21.7–50.7)	21.7 (8.6–44.7)	50.0 (27.6–72.3)
	Mean	76.1	75.0	77.4

Values of 95 % confidence interval are shown in parentheses

**Table 4** Identification rate of each odor used in the final version of Iran-SIT

No	Odor	Identification rate (%)	Female Identification (%)	Male Identification (%)
1	Banana	98.8 (97.4–99.4)	99.3 (97.3–99.8)	98.2 (95.7–99.2)
2	Rose water	94.1 (91.8–95.7)	94.3 (90.9–96.4)	93.9 (90.3–96.1)
3	Garlic	92.9 (90.4–94.7)	93.0 (89.4–95.3)	92.8 (89.1–95.3)
4	Mint	92.0 (89.5–93.9)	93.3 (89.8–95.6)	90.7 (86.6–93.5)
5	Hami melon	91.3 (88.7–93.3)	91.6 (87.8–94.2)	91.0 (87.0–93.8)
6	Cologne	90.5 (87.7–92.6)	89.9 (85.9–92.8)	91.0 (87.0–93.8)
7	Cinnamon	89.4 (86.6–91.6)	90.9 (87.0–93.7)	87.8 (83.3–91.1)
8	Coconut	88.6 (85.6–90.9)	90.3 (86.3–93.1)	86.7 (82.1–90.2)
9	Peach	87.0 (83.9–89.5)	90.3 (86.3–93.1)	83.5 (78.6–87.4)
10	Chocolate	85.8 (82.6–88.4)	84.6 (79.9–88.2)	87.1 (82.5–90.5)
11	Bubble gum	85.4 (82.3–88.0)	83.9 (79.2–87.6)	87.1 (82.5–90.5)
12	Pineapple	83.9 (80.6–86.6)	83.2 (78.5–87.0)	84.6 (79.8–88.3)
13	Grape	83.2 (79.9–86.0)	83.6 (78.8–87.3)	82.8 (77.8–86.8)
14	Onion	82.0 (78.6–84.9)	81.5 (76.6–85.5)	82.4 (77.4–86.4)
15	Smoke	81.8 (78.4–84.7)	82.2 (77.4–86.1)	81.4 (76.3–85.5)
16	Biscuit	80.4 (76.9–83.4)	80.9 (75.9–84.9)	79.9 (74.7–84.2)
17	Peanut	79.4 (75.8–82.4)	77.9 (72.7–82.2)	81.0 (75.9–85.2)
18	Cola	76.8 (73.1–80.0)	74.2 (68.8–78.8)	79.6 (74.3–83.9)
19	Watermelon	76.1 (72.4–79.3)	75.8 (70.6–80.3)	76.3 (70.9–80.9)
20	Apple	74.9 (71.1–78.2)	78.9 (73.8–83.1)	70.6 (64.9–75.6)
21	Dried lime	74.2 (70.4–77.5)	79.2 (74.1–83.4)	68.8 (63.1–74.0)
22	Strawberry	73.7 (69.8–77.1)	76.5 (71.3–81.0)	70.6 (64.9–75.6)
23	Soap	72.6 (68.8–76.1)	73.5 (68.1–78.2)	71.7 (66.0–76.6)
24	Jasmine	71.2 (67.3–74.7)	73.2 (67.8–77.9)	69.2 (63.4–74.3)
	Mean	83.6	84.2	82.9

Values of 95 % confidence interval are shown in parentheses

**Fig. 1** Iran smell identification test

**Table 5** Classification of 24 odors used in the final version of Iran-SIT

Category	Specifications	Iran-SIT, odors
Floral	Fragrant, light, natural	Jasmine
		Rose water
		Cologne
		Soap
Fruity	Fresh, smooth, pleasant	Banana
		Hami melon
		Grape
		Pineapple
		Apple
		Peach
		Watermelon
		Strawberry
Pungent	Sharp, nasty, caustic	Bubble gum
		Onion
Minty	Cool, fresh, exhilarating	Garlic
		Mint
Woody	Natural, strong	Cola
		Cinnamon
Sweet	Warm, light, creamy, rich	Chocolate
		Coconut
Baked	Burnt, nutty, heavy, warm	Biscuit
		Peanut
		Smoke
Citrus	Acidic, sharp	Dried lime

consent, and this project was approved by the Ethics Committee for research in Tehran University of Medical Sciences.

In order to select subjects who qualified to participate in the main study, we determined the inclusion/exclusion


criteria. All subjects underwent physical examination which consisted of items like deviated nasal septum (DNS), tight nasal valve, dried nasal mucus, and nasal adhesion (Snow et al. 1991). They were also interviewed about their past medical history. The medical history was evaluated using the questionnaire developed in the University of Pennsylvania Smell and Taste Center (Deems et al. 1991). The questions included sinus or nasal disease, history of pre and post-operative radiotherapy and/or chemotherapy, history of head trauma, toxic chemical exposure, serious upper respiratory problems, history of head and neck surgery, nasal allergies, and family history of smell problems. Based on the results of physical examination and medical history, we selected subjects to participate in the main study. Moreover, people with smoking habit (Doty et al. 1984a, b; Frye et al. 1990; Ishimaru and Fujii 2007) and/or taking medicines affecting olfaction (Mair and Harrison 1991; Henkin 1994) were excluded from the study. Finally, 577 healthy subjects were selected to participate in main study.

By omitting 16 problematic odors, the main experiment was carried out using the modified 24-items test named Iran smell identification test (Iran-SIT). It was designed as a four-alternative test in a forced-choice paradigm. The procedure was fully explained to the all 577 subjects as the following lines. The subjects were asked to scratch the stickers by means of a pencil tip to release the odors. They were encouraged to sniff the scraped sticker immediately and choose one of four alternatives. If they claimed that the odor they smelled was not presented in the alternatives, they were asked to mark the answer closest to their experience. The time interval between each sniff was 30 s. In some cases, the examiner helped administer the test to subjects who could not read or who had impaired eyesight. Iran-SIT score was considered as the number of the items that were correctly answered.

**Table 6** Identification score of each age group

Age group	Minimum	Maximum	Mean	SE	Unadjusted (95 % confident interval)
5–9	6	17	13.47	0.358	12.76–14.17
10–14	9	21	16.71 A	0.445	15.83–17.58
15–19	14	24	20.79 DEF	0.336	20.13–21.45
20–24	16	24	22.00 F	0.262	21.48–22.51
25–29	16	24	22.16 F	0.239	21.68–22.62
30–34	15	24	21.44 EF	0.263	20.92–21.96
35–39	17	24	21.69 EF	0.263	21.16–22.20
40–44	17	24	21.13 DEF	0.275	20.58–21.67
45–49	15	24	21.08 DEF	0.405	20.28–21.87
50–54	15	24	20.19 CDE	0.493	19.22–21.16
55–59	14	23	19.54 BCD	0.475	18.60–20.46
60–64	14	22	18.19 ABC	0.509	17.18–19.19
65–69	14	21	17.58 AB	0.473	16.64–18.50

Means sharing the same letter are not significantly different at the 5 % level. SE standard error



**Fig. 2** Bubble chart of the relation between test and retest study with 5 months interval

### Test-Retest Study

Reliability and stability of Iran-SIT over time was assessed by administrating the Iran-SIT 5 months after main study. Ninety-six (44 female and 52 male) of 577 subjects with ages ranging from 10 to 60 years, who had different identification scores, were selected to participate in the retest study. The retest study was administered using the same procedure as main study.

## Results and Discussion

All the analyses were performed by STATA software version 12 and  $p$  value less than 0.05 was considered statistically significant.

### Determining Familiar Odors for Iranian Population

The aim of this study was to develop a standardized olfactory test for Iranian population considering their cultural background. For designing a reliable and valid test, it was important to choose the familiar odors covering the different smell categories. Because 21 of 40 odors of UPSIT were mostly unfamiliar for the subjects, the first

version of Iran-SIT was provided by replacement in some odors of UPSIT, which were unfamiliar for Iranian population. It was tried to preserve the main categories of odors in all replacement; for instance, jasmine and tuberose was replaced for lilac and clove, cake was replaced for gingerbread, and vanilla was replaced for licorice. Furthermore, we had to change some alternatives related to odors. Finally, our 40-item list was obtained as shown in Table 2.

### Pilot Study

To assess the quality of the odors and to choose the suitable alternatives, we analyzed the data obtained from 43 subjects in the pilot study. The identification percentage (95 % confidence interval) for each odor of the first version of Iran-SIT is presented in Table 3. Results indicated that 16 odors had identification percentages less than 70 %. Most of them were difficult to identify correctly due to manufacturing difficulties, so we had to omit them for the main experiment. Moreover, we switched some alternatives because they misled participants.

### Main Study

We confirmed that all of the 24 odors used in the final version of Iran-SIT obtained an identification percentage of more than 70 % (Table 4). We photographed the final version of Iran-SIT which comprised 24 odors (Fig. 1). There are thousands of aromas that humans can smell. Castro et al. (2013) have used a computerized technique to whittle down odors to their most basic essence. They classified the odors into 10 basic categories; floral, fruity (non-citrus), woody, chemical, minty, sweet, pungent, popcorn, citrus, and decayed. Twenty-four odors used in the final version of Iran-SIT could be classified in eight categories shown in Table 5. None of the odors belonged to chemical and decayed categories. All odors are assumed to stimulate the first cranial nerve (olfactory nerve); however, a few subjects reported that some odors like garlic and mint caused a mild irritation in nose.

The means of identification scores were 20.06 for all subjects, 20.22 for female, and 19.87 for male. In order to examine the effect of gender on olfactory function, unpaired  $t$  test was conducted for identification scores of each subject.  $t$  test demonstrated no significant difference between female and male ( $t(575)=1.15$ ). Furthermore, in order to examine the effect of aging on olfactory function, we conducted one-way factorial analysis of variance (ANOVA) for identification score with the age group as the between-subject factor. ANOVA revealed significant main effect of age group ( $F(12, 564)=58.24$ ). Multiple comparisons by Tukey's method for the significant main effect demonstrated significant differences between some combinations of age groups as

**Table 7** Diagnostic criterion of olfactory disorder using the final version of Iran-SIT

	Anosmia	Severe microsmia	Mild microsmia	Normosmia
Identification score	0–9	10–13	14–18	19–24

shown in Table 6. It is well known that human's chemosensory function declines with aging (Ship and Weiffenbach 1993; Doty et al. 1984a, b; De Jong et al. 1999; Hummel et al. 2003). Murphy et al. (2002) reported that the 24.5 % of people over 53 years of age and 62.5 % of those aged 80–97 years suffered from olfactory impairment. In our study, the decrease of olfactory function was observed over 50 years. Adult aged 20–50 years kept significantly higher olfactory function than children or elderly people. Children, especially those under 10 years of age, markedly obtained the low scores in Iran-SIT due to their insufficient experience.

### Test-Retest Study

We created a bubble chart for the test-retest study in Fig. 2. In order to assess reliability and stability of the final version of Iran-SIT over time, we calculated the Pearson's correlation coefficient ( $r=0.93$ ) and Spearman's rank correlation coefficient ( $\rho=0.89$ ) between test-retest identification scores. Test for non-correlation demonstrated significant correlations for both coefficients ( $p<0.000$ ). In previous studies, Pearson's correlation coefficients between test-retest identification scores was 0.949 for UPSIT with interval of 2 weeks (Doty et al. 1985), 0.918 for UPSIT with interval of 6 months (Doty et al. 1984a, b), 0.71 for cross-cultural version of UPSIT (CC-SIT) (Doty et al. 1996), and 0.73 for Sniffin' Sticks (Kobal et al. 1996). Compared with other olfactory identification tests reported in previous researches, Pearson's correlation coefficient of the final version of Iran-SIT had an acceptable value. We assumed that the final version of Iran-SIT was reliable and stable over time.

### Diagnostic Criterion of Olfactory Disorder Using the Final Version of Iran-SIT

Based on identification scores obtained from adult aged 20–50 years in main study, we determined diagnostic criterion of olfactory disorder using the final version of Iran-SIT (Table 7). Our results showed that 95 % of these subjects correctly identified over 18 odors, so that subjects who gained identification scores from 19 to 24 was considered as normosmia with error at 5 % level. According to the laws of probability in four-alternative forced-choice test using 24 odors, the accumulation probability to identify correctly below 10 odors at chance level is 94.5 %. Hence, subjects who gained identification scores from 0 to 9 were considered as anosmia with error rate at 5.5 % level. Subjects who gained identification scores from 10 to 18 were subdivided into two levels: severe microsmia (the scores from 10 to 13) and mild microsmia (the score from 14 to 18). This heuristic classification has been applied to classify UPSIT data obtained from approximately 4000 subjects (Doty 1995).

### Conclusion

In the present study, we have developed a standardized 24-item smell identification test to assess the olfactory function of Iranian population considering cultural adaptation. Iran-SIT has adequacy to classify adult patients into four levels: ones with the normal olfactory function (normosmia), ones with mildly decreased olfactory function (mild microsmia), ones with severely decreased olfactory function (severe microsmia), and ones with loss of olfactory function (anosmia).

**Acknowledgments** This work was carried out with financial support from the Iran National Science Foundation (INSF). The authors wish to thank the Iranian Legal Medicine Research Center for supporting us to conduct the study on detecting malingering. We are very grateful to Dr. Saloumeh Salarian, Dr. Ebrahim Razmpa, Dr. Amin Amali, Dr. Shahin Bastaninejad, Dr. Masoud Mozafari, and Dr. Maryam Ghaffari for their valuable guidance. Special thanks are owed to all who participated in our study with great patience and kindness.

**Compliance with Ethics Requirements** All procedures followed were in accordance with the ethical standards of the institutional and national research committee and with the Helsinki Declaration of 1975, as revised in 2013. Informed consent was obtained from all individual participants for being included in this study.

**Conflict of Interest** The authors declare that they have no competing interests.

### References

- Asia-Pacific Population Journal, United Nations (2006) A New Direction in Population Policy and Family Planning in the Islamic Republic of Iran. Retrieved 04–14
- Cain WS, Cometto-Muniz JE, De Wijk RA (1992) Techniques in the quantitative study of human olfaction. In: Serby MJ, Chobor KL (eds) Science of olfaction. Springer, New York, pp 279–308
- Cardesin A, Alobid I, Benítez P et al (2006) Barcelona Smell Test—24 (BAST-24): validation and smell characteristics in the healthy Spanish population. *Rhinology* 44:83–89
- Castro JB, Ramanathan A, Chennubhotla CS (2013) Categorical dimensions of human odor descriptor space revealed by non-negative matrix factorization. *PLoS ONE* 8:1
- CĂtanĂ I, Negoia S, Maniu A, Porojan M, Cosgarea M (2012) A Modified version of “SNIFFIN’ STICKS” odor identification test: the Romanian cultural adaptation. *Clujul Med* 85:218–223
- Cho JH, Jeong YS, Lee YJ, Hong SC, Yoon JH, Kim JK (2009) The Korean version of the Sniffin’ stick (KVSS) test and its validity in comparison with the cross-cultural smell identification test (CC-SIT). *Auris Nasus Larynx* 36:280–286
- De Jong N, Mulder I, de Graaf C, van Staveren WA (1999) Impaired sensory functioning in elders: the relation with its potential determinants and nutritional intake. *J Gerontol A Biol Sci Med Sci* 54:324–331
- Deems DA, Doty RL, Settle RG et al (1991) Smell and taste disorders: a study of 750 patients from the University of Pennsylvania Smell and Taste Center. *Arch Otolaryngol Head Neck Surg* 117:519–528
- Doty RL (1995) The smell identification test, administration manual, 3rd edn. Sensonics, Inc., Haddon Heights
- Doty RL (2001) Olfaction. *Annu Rev Psychol* 52:423–452



- Doty RL, Shaman P, Applebaum SL et al (1984a) Smell identification ability: changes with age. *Science* 226:1441–1443
- Doty RL, Shaman P, Dann MS (1984b) Development of the University of Pennsylvania Smell Identification Test: a standardized microcapsulated test of olfactory function. *Physiol Behav* 32:489–502
- Doty RL, Newhouse MG, Azzalina JF (1985) Internal consistency and short term test-retest reliability of the University of Pennsylvania Smell Identification Test. *Chem Senses* 10:297–300
- Doty RL, Marcus A, Lee WW (1996) Development of the 12-item cross-cultural smell identification test (CC-SIT). *Laryngoscope* 106:353–356
- Eibenstein A, Fioretti AB, Lena C, Rosati N, Amabile G, Fusetti M (2005) Modern psychophysical tests to assess olfactory function. *Neurol Sci* 26:147–155
- Frye RE, Schwartz BS, Doty RL (1990) Dose-related effects of cigarette smoking on olfactory function. *JAMA* 263:1233–1236
- Harper R, Bate Smith EC, Land DG (1968) Odour description and odour classification. American Elsevier, New York
- Henkin RI (1994) Drug-induced taste and smell disorders. *Drug Saf* 11: 318–377
- Hummel T, Sekinger B, Wolf S, Pauli E, Kobal G (1997) “Sniffin’ Sticks”: olfactory performance assessed by the combined testing of odor identification, odor discrimination and olfactory threshold. *Chem Senses* 22:39–52
- Hummel T, Futschik T, Frasnelli J, Huttenbrink KB (2003) Effects of olfactory function, age, and gender on trigeminally mediated sensations: a study based on the lateralization of chemosensory stimuli. *Toxicol Lett* 140:273–280
- Ishimaru T, Fujii M (2007) Effects of smoking on odour identification in Japanese subjects. *Rhinology* 45:224–228
- Kobal G, Hummel TH, Sekinger B, Barz S, Roscher S, Wolf S (1996) “Sniffin’ sticks”: screening of olfactory performance. *Rhinology* 34: 222–226
- Landis BN, Hummel T, Lacroix JS (2005) Basic and clinical aspects of olfaction. In *Adv Standards Neurosur* 30:75–77
- Mair RG, Harrison LM (1991) Influence of drugs on smell function. In: Laing DG, Doty RL, Breipohl W (eds) *The human sense of smell*. Springer, Berlin, pp 335–359
- Murphy C (1985) Cognitive and chemosensory influences on age related changes in the ability to identify blended foods. *J Gerontol* 40:47–52
- Murphy C, Schubert CR, Cruickshanks KJ, Klein BE, Klein R, Nondahl DM (2002) Prevalence of olfactory impairment in older adults. *JAMA* 288:2307–2312
- Nordin S, Nyroos M, Maunuksela E, Niskanen T, Tuorila H (2002) Applicability of the Scandinavian odor-identification test: a Finnish-Swedish comparison. *Acta Otolaryngol (Stockh)* 122:294–297
- Oniz A, Erdogan I, Ikiz AO, Evirgen N, Ozgoren M (2013) The modified Sniffin’ sticks test in Turkish population based on odor familiarity survey. *J Neurol Sci* 30:270–280
- Saito S, Ayabe-Kanamura S, Takashima Y et al (2006) Development of a smell identification test using a novel stick-type odor presentation kit. *Chem Senses* 31:379–391
- Ship JA, Weiffenbach JM (1993) Age, gender, medical treatment, and medication effects on smell identification. *J Gerontol Med Sci* 48: 26–32
- Silveira-Moriyama L, Azevedo AM, Ranvaud R, Barbosa ER, Doty RL, Lees AJ (2010) Applying a new version of the Brazilian-Portuguese UPSIT smell test in Brazil. *Arq Neuropsiquiatr* 68:700–705
- Snow JB, Doty RL, Bartoshuk LM (1991) Clinical evaluation of olfactory and gustatory disorders. In: Getchell TV, Doty RL, Bartoshuk LM, Snow JB (eds) *Smell and taste in health and disease*. Raven, New York, pp 463–467
- Thomas-Danguin T, Rouby C, Sicard G et al (2003) Development of the ETOC: a European test of olfactory capabilities. *Rhinology* 41:142–151