

On the Isotropic Constant of Random Polytopes

David Alonso-Gutiérrez · Alexander E. Litvak ·
Nicole Tomczak-Jaegermann

Received: 14 January 2014 / Published online: 27 January 2015
© Mathematica Josephina, Inc. 2015

Abstract Let X_1, \dots, X_N be independent random vectors uniformly distributed on an isotropic convex body $K \subset \mathbb{R}^n$, and let K_N be the symmetric convex hull of X_i 's. We show that with high probability $L_{K_N} \leq C\sqrt{\log(2N/n)}$, where C is an absolute constant. This result closes the gap in known estimates in the range $Cn \leq N \leq n^{1+\delta}$. Furthermore, we extend our estimates to the symmetric convex hulls of vectors $y_1 X_1, \dots, y_N X_N$, where $y = (y_1, \dots, y_N)$ is a vector in \mathbb{R}^N . Finally, we discuss the case of a random vector y .

Keywords Convex bodies · Isotropic measures · Isotropic constant · Random polytopes

Mathematics Subject Classification Primary 52A23 · 46B06 · Secondary 52A22 · 52B11 · 60D05

1 Introduction

In this paper we estimate the isotropic constant of some random polytopes (for the definitions and notation see Sect. 2). It is known (see, e.g., [28]) that among all the

D. Alonso-Gutiérrez
Departament de Matemàtiques, Universitat Jaume I, Campus de Riu Sec, 12071 Castelló de la Plana,
Spain
e-mail: davidalonsogutierrez@gmail.com

A. E. Litvak (✉) · N. Tomczak-Jaegermann
Department of Mathematics and Statistics Sciences, University of Alberta, Edmonton, AB T6G 2G1,
Canada
e-mail: aelitvak@gmail.com

N. Tomczak-Jaegermann
e-mail: nicole.tomczak@ualberta.ca

convex bodies in \mathbb{R}^n the Euclidean ball is the one with the smallest isotropic constant, that is $L_K \geq L_{B_2^n} \geq c$, where c is an absolute positive constant. However, it is still an open problem to determine whether there exists or not an absolute constant C such that for every convex body $K \subset \mathbb{R}^n$ one has $L_K \leq C$. The boundedness of L_K by an absolute constant is equivalent to the long standing hyperplane conjecture [9]. The best general upper bound known up to now is $L_K \leq Cn^{1/4}$ [22]. This estimate slightly improves (by a logarithmic factor) the earlier Bourgain’s upper bound [10].

Since the remarkable result of Gluskin [18] random polytopes are known to provide many examples of convex bodies (and related normed spaces) with a “pathologically bad” behavior of various parameters of a linear and geometric nature (we refer to the survey [27] and references therein; see also recent examples in [19] and [21]). Not surprisingly, they were also a natural candidate for a potential counterexample for the hyperplane conjecture. This was resolved in [23], where it was shown that the convex hull or the symmetric convex hull of independent Gaussian random vectors in \mathbb{R}^n with high probability has the bounded isotropic constant. Some other distributions for vertices were also considered. In all of them the vertices had independent coordinates.

Following the ideas in [23], the problem of estimating the isotropic constant of random polytopes was considered in [4], for independent random vectors distributed uniformly on the sphere S^{n-1} , and in [13], for independent random vectors uniformly distributed on an isotropic unconditional convex body. Also in these cases the isotropic constant of random polytopes generated by these vectors is bounded with high probability. One can check that the same method works for independent random vectors uniformly distributed on a ψ_2 isotropic convex body as well.

In this paper we estimate the isotropic constant of a random polytope in an isotropic convex body (see Sect. 2 for the definitions). It is known (see [6,20] or [5]) that if K_N is a polytope in \mathbb{R}^n with N vertices then

$$L_{K_N} \leq C \min \left\{ \sqrt{\frac{N}{n}}, \log N \right\},$$

where C is an absolute constant.

In [14,15], the authors provided a lower estimate for the volume of a random polytope K_N obtained as the convex hull of $N \leq e^{\sqrt{n}}$ random points, namely

$$|K_N|^{\frac{1}{n}} \geq c \sqrt{\frac{\log \frac{N}{n}}{n}} L_K$$

(see the end of Sect. 2 for more precise formulation and details). On the other hand, the proof of the estimate $L_{K_N} \leq C \log N$ in [5] passes through showing that if X_1, \dots, X_N are the vertices of K_N , then for any affine transformation T we have

$$L_{K_N} \leq \frac{C \max_{1 \leq i \leq N} |T X_i| \log N}{n |T K_N|^{\frac{1}{n}}}.$$

Consequently, taking T to be the identity operator and using the concentration of measure result proved by Paouris [30], we obtain that if K_N is the convex hull or the symmetric convex hull of N independent random vectors (for $n + 1 \leq N \leq e^{\sqrt{n}}$) uniformly distributed on an isotropic convex body, then with high probability

$$L_{K_N} \leq \frac{C \log N}{\sqrt{\log \frac{N}{n}}}. \tag{1.1}$$

Notice that if $N \geq n^{1+\delta}$, $\delta \in (0, 1)$, this estimate does not exceed $(C/\delta)\sqrt{\log \frac{N}{n}}$. However, C/δ tends to infinity as δ tends to 0. On the other hand, if N is proportional to n the isotropic constant of K_N is bounded (by an absolute constant), while the upper bound in (1.1) is not. The following theorem closes the gap between $N \leq cn$ and $N \geq n^{1+\delta}$.

Theorem 1.1 *There exist absolute positive constants c, C such that if $n \leq N$, and X_1, \dots, X_N are independent random vectors uniformly distributed on an isotropic convex body K , and K_N is their symmetric convex hull, then*

$$\mathbb{P}\left(\left\{L_{K_N} \leq C\sqrt{\log \frac{2N}{n}}\right\}\right) \geq 1 - \exp(-c\sqrt{n}).$$

Furthermore, we study a natural more general family of perturbations of random polytopes. Namely, for any $X_1, \dots, X_N \in \mathbb{R}^n$ and $y \in \mathbb{R}^N$, we define

$$K_{N,y} = \text{conv}\left\{\pm y_1 X_1, \dots, \pm y_N X_N\right\}.$$

As it turns out Theorem 1.1 is valid for this family as well. To describe this result we need more notation.

For a vector $y = (y_1, \dots, y_N) \in \mathbb{R}^N$, let $\{y_1^*, \dots, y_N^*\}$ be the sequence of decreasing rearrangement of $\{|y_i|\}_i$. For $1 \leq k \leq N$ and $n \leq m \leq N$ denote

$$\|y\|_{k,2} := \left(\sum_{i=1}^k y_i^{*2}\right)^{1/2} \quad \text{and} \quad \alpha_{y,m} := \left(\prod_{i=m-n+1}^m y_i^*\right)^{\frac{1}{n}} \geq y_m^*.$$

Our main result is the following theorem.

Theorem 1.2 *There exist absolute positive constants c and C such that the following holds. Let $n \leq N$ and let X_1, \dots, X_N be independent random vectors uniformly distributed on an isotropic convex body K . Then for every $y \in \mathbb{R}^N$ the event*

$$L_{K_{N,y}} \leq C \frac{\|y\|_{n,2}}{\sqrt{n}} \inf \left\{ \alpha_{y,m}^{-1} \frac{\log(2N/n)}{\sqrt{\log(2m/n)}} \mid n \leq m \leq N \right\}$$

occurs with probability greater than $1 - e^{-c\sqrt{n}}$. Moreover, the event

$$\sup_{\|y\|_{n,2} \leq 1} L_{K_N,y} \leq C \alpha_{y,m_0}^{-1} \sqrt{\frac{\log(2N/n)}{n}},$$

where $m_0 = N(1 - c/\log N)$, occurs with probability greater than $1 - e^{-c\sqrt{n}}$.

Remark 1 Note that if $n \leq N \leq Cn$ then $L_K \leq C$ for any symmetric polytope K generated by N vectors [6].

Remark 2 Clearly Theorem 1.2 applied to the vector $y = (1, \dots, 1)$ implies Theorem 1.1.

Finally, we apply Theorem 1.2 to the case when the vector describing the perturbation is also random. Such a setting has been recently considered in [7]. In Theorem 4.1 we show that for the Gaussian vector $G = (g_1, \dots, g_N)$ in \mathbb{R}^N with high probability we have

$$L_{K_N,G} \leq C \sqrt{\log \frac{2N}{n}}.$$

2 Preliminaries

In this paper the letters c, C, c_1, C_1, \dots will always denote absolute positive constants, whose values may change from line to line. Given two functions f and g we say that they are equivalent and write $f \approx g$ if $c_1 f \leq g \leq c_2 f$.

By $|\cdot|$ and $\langle \cdot, \cdot \rangle$ we denote the canonical Euclidean norm and the canonical inner product on \mathbb{R}^n . The (unit) Euclidean ball and sphere are denoted by B_2^n and S^{n-1} . Let K be a symmetric convex body in \mathbb{R}^n and let $\|\cdot\|_K$ be its associated norm

$$\|x\|_K = \inf \left\{ \lambda > 0 : x \in \lambda K \right\}.$$

The support function of K is $h_K(y) = \max_{x \in K} \langle x, y \rangle$ and it is the norm associated with the polar body of K ,

$$K^\circ = \left\{ y \in \mathbb{R}^n : \langle x, y \rangle \leq 1 \ \forall x \in K \right\}.$$

Given convex body K we denote by $|K|$ its volume. We also denote by $|E|$ the cardinality of a finite set E . For $E \subset \{1, \dots, N\}$ the coordinate projection on \mathbb{R}^E is denoted by P_E .

We say that a convex body $K \subseteq \mathbb{R}^n$ is isotropic if it has volume $|K| = 1$, its center of mass is at 0 (i.e., $\int_K x dx = 0$) and for every $\theta \in S^{n-1}$ one has

$$\int_K \langle x, \theta \rangle^2 dx = L_K^2,$$

where L_K is a constant independent of θ . L_K is called the isotropic constant of K .

It is known that every convex body has a unique (up to an orthogonal transformation) affine image that is isotropic. This allows to define the isotropic constant of any convex body as the isotropic constant of its isotropic image. It is also known (see, e.g., [28]) that

$$nL_K^2 = \inf \left\{ \frac{1}{|TK|^{1+\frac{2}{n}}} \int_{a+TK} |x|^2 dx : T \in GL(n), a \in \mathbb{R}^n \right\}. \tag{2.1}$$

We need two more results on the distribution of Euclidean norms of random vectors and their sums. Let $X_i, i \leq N$, be independent random vectors uniformly distributed in an isotropic convex body $K \subset \mathbb{R}^n$. Let A be a random $n \times N$ matrix, whose columns are the X_i 's. For $m \leq N$ denote

$$A_m = \sup \left\{ |Ay| \mid y \in B_2^N, |\text{supp } y| \leq m \right\}$$

(supp denotes the support of y). Theorem 3.13 in [1] (note the different normalization) implies the following estimate.

Theorem 2.1 *There is an absolute positive constant C such that for every $\gamma \geq 1$ and every $m \leq N$*

$$\mathbb{P} \left(\left\{ A_m \geq CL_K \gamma \sqrt{m} \log \frac{2N}{m} + 6 \max_i |X_i| \right\} \right) \leq \exp \left(-\gamma \sqrt{m} \log \frac{2N}{m} \right).$$

The following theorem is a combination of Paouris's theorem ([30], see also [3] for a short proof) with the union bound (cf. Lemma 3.1 in [1]).

Theorem 2.2 *There exists an absolute positive constant C such that for any $N \leq \exp(\sqrt{n})$ and for every $\lambda \geq 1$ one has*

$$\mathbb{P} \left(\left\{ \max_{i \leq N} |X_i| \geq C\lambda\sqrt{n} L_K \right\} \right) \leq \exp(-\lambda\sqrt{n}).$$

Finally we need the estimate on the volume of the random polytope

$$K_N = \text{conv}\{\pm X_1, \dots, \pm X_N\},$$

where $X_i, i \leq N$, are independent random vectors uniformly distributed in an isotropic convex body $K \subset \mathbb{R}^n$. The estimates of the following theorem were observed in [14] (see Fact 3.2, the remarks following it, and Fact 3.3 there; see also [32] and Chapter 11 of [11], where the assumption $N \geq Cn$ was reduced to $N \geq n$).

Theorem 2.3 *There are absolute positive constants c_1, c_2 such that for $n \leq N \leq e^{\sqrt{n}}$,*

$$\mathbb{P} \left(\left\{ |K_N|^{1/n} \geq c_1 \sqrt{\frac{\log(N/n)}{n}} L_K \right\} \right) \geq 1 - \exp(-c_2\sqrt{N}).$$

In fact this theorem is a combination of three results. The first says that K_N contains the centroid body. Recall that for $p \geq 1$ the p -centroid body $Z_p(K)$ was introduced in [26] (with a different normalization) as the convex body, whose support function is

$$h_{Z_p(K)}(\theta) = \left(\int_K |\langle x, \theta \rangle|^p dx \right)^{\frac{1}{p}}. \tag{2.2}$$

In [15] (Theorem 1.1) the authors proved that for every parameters $\beta \in (0, 1/2)$ and $\gamma > 1$ one has the inclusion $K_N \supset c_1 Z_p(K)$ for $p = c_2 \beta \log(2N/n)$ and $N \geq c_3 \gamma n$ with the probability greater than

$$1 - \exp(-c_4 N^{1-\beta} n^\beta) - \mathbb{P} \left(\left\{ \|A\| > \gamma L_K \sqrt{N} \right\} \right),$$

where A is the random matrix whose columns are X_1, \dots, X_N . The probability that norm of A (note $\|A\| = A_N$) is large was estimated in [1] (combine Theorems 2.1 and 2.2 above). Finally, from results of [24] and [30] the bound

$$|Z_p(K)|^{1/n} \approx \sqrt{p/n} L_K \tag{2.3}$$

follows provided that $p \leq \sqrt{n}$ (it improves the bound provided in [25]).

In the Sect. 4 we will use the following standard estimate. For the sake of completeness we provide a proof (cf. Example 10 in [16]).

Lemma 2.4 *There exists an absolute positive constant C such that for $Cm \leq N$ and independent standard Gaussian random variables g_1, \dots, g_N one has*

$$\mathbb{P} \left(\left\{ g_m^* \leq \sqrt{\log(eN/m)} \right\} \right) \leq \exp \left(- \frac{\sqrt{mN}}{10 \sqrt{\log \left(\frac{eN}{m} \right)}} \right)$$

Proof Denote $\alpha = \sqrt{\log(eN/m)}$. Note, $g_m^* \leq \alpha$ in particular means that there exists $\sigma \subset \{1, \dots, N\}$ of cardinality $\ell := N - m$ such that $|g_i| \leq \alpha$ for every $i \in \sigma$. Therefore,

$$p := \mathbb{P} \left(\left\{ g_m^* \leq \alpha \right\} \right) \leq \binom{N}{\ell} (\mathbb{P}(\{|g_1| \leq \alpha\}))^\ell.$$

Using that

$$\mathbb{P}(\{|g_1| \leq \alpha\}) = 1 - \sqrt{\frac{2}{\pi}} \int_\alpha^\infty e^{-x^2/2} dx \leq 1 - \frac{e^{-\alpha^2/2}}{\sqrt{2\pi}\alpha} \leq \exp \left(- \frac{e^{-\alpha^2/2}}{\sqrt{2\pi}\alpha} \right),$$

$\alpha = \sqrt{\log(eN/m)}$ and $\binom{N}{\ell} = \binom{N}{m} \leq (eN/m)^m$ we obtain

$$p \leq \exp \left(m \log(eN/m) - \ell \frac{\sqrt{\frac{m}{eN}}}{\sqrt{2\pi \log(\frac{eN}{m})}} \right).$$

As $\ell = N - m$, this implies the result. □

3 Proofs

In this section we prove Theorem 1.2. The proof consists of two propositions.

Proposition 3.1 *Let $n \leq N \leq e\sqrt{n}$ and X_1, \dots, X_N are independent random vectors distributed uniformly on an isotropic convex body K . Then the event*

$$\sup \left\{ \frac{1}{|K_{N,y}|} \int_{K_{N,y}} |x|^2 dx \mid \|y\|_{n,2} \leq 1, y_n^* > 0 \right\} \leq C \frac{1}{n} L_K^2 \log^2 \frac{2N}{n}$$

occurs with probability greater than $1 - \exp(-\sqrt{n} \log(2N/n))$, where C is an absolute constant.

To prove this proposition we need the following lemma.

Lemma 3.2 *Let $1 \leq n \leq N$ be integers and $P = \text{conv}\{P_1, \dots, P_N\}$ be a non-degenerated symmetric polytope in \mathbb{R}^n . Then*

$$\frac{1}{|P|} \int_P |x|^2 dx \leq \frac{1}{(n+1)(n+2)} \sup_E \left(\sum_{i \in E} |P_i|^2 + \left| \sum_{i \in E} P_i \right|^2 \right),$$

where the supremum is taken over all subsets $E \subset \{1, \dots, N\}$ of cardinality n .

Proof We can decompose P as a disjoint union of simplices (up to sets of measure 0), say $P = \cup_{i=1}^{\ell} C_i$, where each C_i is of the form $\text{conv}\{0, P_{i_1}, \dots, P_{i_n}\}$ for some choice of P_{i_j} 's. For every such C_i , denote $F_i := \text{conv}\{P_{i_1}, \dots, P_{i_n}\}$. Then for any integrable function f we have

$$\int_{C_i} f(x) dx = \int_{F_i} \int_0^1 r^{n-1} f(ry) |\langle y, \nu(y) \rangle| dy dr = d(0, F_i) \int_{F_i} \int_0^1 r^{n-1} f(ry) dy dr,$$

where $\nu(y)$ is the outer normal vector to P at the point y and $d(0, F_i)$ is the distance from the origin to the affine subspace spanned by F_i . Thus, as in [23], for every $i \leq \ell$ one has

$$|C_i| = n^{-1} |F_i| d(0, F_i) \quad \text{and} \quad \int_{C_i} |x|^2 = \frac{d(0, F_i)}{n+2} \int_{F_i} |y|^2 dy.$$

In particular,

$$|P| = \sum_{i=1}^{\ell} |C_i| = \frac{1}{n} \sum_{i=1}^{\ell} |F_i| d(0, F_i).$$

Therefore,

$$\begin{aligned} \frac{1}{|P|} \int_P |x|^2 dx &= \frac{1}{|P|} \sum_{i=1}^{\ell} \frac{d(0, F_i)}{n+2} \int_{F_i} |y|^2 dy \\ &\leq \frac{1}{|P|} \sum_{i=1}^{\ell} \frac{d(0, F_i) |F_i|}{n+2} \sup_{1 \leq i \leq \ell} \frac{1}{|F_i|} \int_{F_i} |y|^2 dy \\ &\leq \frac{n}{n+2} \sup_F \frac{1}{|F|} \int_F |y|^2 dy, \end{aligned}$$

where the supremum is taken over all $F = \text{conv}\{P_{i_1}, \dots, P_{i_n}\}$. Note that any such F can be presented as $F = T \Delta^{n-1}$, where $\Delta^{n-1} = \text{conv}\{e_1, \dots, e_n\}$ denotes the regular $n - 1$ dimensional simplex in \mathbb{R}^n and T is the matrix whose columns are the vectors P_{i_j} . Since

$$\frac{1}{|\Delta^{n-1}|} \int_{\Delta^{n-1}} y_i y_j dy = \frac{1 + \delta_{ij}}{n(n+1)},$$

where δ_{ij} is the Kronecker delta, for every $F = \text{conv}\{P_{i_1}, \dots, P_{i_n}\}$ we obtain

$$\frac{1}{|F|} \int_F |y|^2 dy = \frac{1}{n(n+1)} \left(\sum_{j=1}^n |P_{i_j}|^2 + \left| \sum_{j=1}^n P_{i_j} \right|^2 \right).$$

This implies the desired estimate. □

Proof of Proposition 3.1. Note that if $y_n^* > 0$ then the cardinality of the support of y is at least n , so $K_{N,y}$ is not degenerated with probability one. Therefore, with probability one $K_{N,y}$ is non-degenerated for any countable dense set in $B_0 := \{y \in \mathbb{R}^N \mid \|y\|_{n,2} \leq 1, y_n^* > 0\}$. Clearly, the supremum under question is the same over $y \in B_0$ and over such a dense set.

Now, by Lemma 3.2 we have that $\sup_{y \in B_0} |K_{N,y}|^{-1} \int_{K_{N,y}} |x|^2 dx$ is bounded from above by

$$\frac{1}{(n+1)(n+2)} \sup_{y \in B_0} \sup_{|E|=n} \left(\sum_{i \in E} |y_i X_i|^2 + \left| \sum_{i \in E} y_i X_i \right|^2 \right)$$

(formally, we should additionally take supremum over $\varepsilon_i = \pm 1$ and to have $y_i \varepsilon_i X_i$ in the formula under suprema, but, since B_0 is unconditional, the supremum over ε_i 's can be omitted).

Note that

$$\sum_{i \in E} |y_i X_i|^2 \leq \|y\|_{n,2}^2 \max_{i \leq N} |X_i|^2$$

and

$$\left| \sum_{i \in E} y_i X_i \right| = |AP_E y| \leq A_n \|y\|_{n,2},$$

where A is the matrix whose columns are X_1, \dots, X_N . Therefore, applying Theorems 2.1 and 2.2 (with $m = n$ and $\lambda = 2 \log(2N/n)$) we obtain that

$$\sup_{y \in B_0} \sup_{|E|=n} \left(\sum_{i \in E} |y_i X_i|^2 + \left| \sum_{i \in E} y_i X_i \right|^2 \right) \leq C n L_K^2 \log^2 \frac{2N}{n}$$

with probability greater than $1 - \exp(-\sqrt{n} \log(2N/n))$. □

Proposition 3.3 *There exist absolute positive constants c_1, c_2 such that if $n \leq N \leq e\sqrt{n}$ and X_1, \dots, X_N are independent random vectors distributed uniformly on an isotropic convex body K , then for every $y \in \mathbb{R}^N$,*

$$\mathbb{P} \left(\left\{ \forall m \geq n : |K_{N,y}|^{\frac{1}{n}} \geq c_1 \alpha_{y,m} L_K \sqrt{\frac{\log(2m/n)}{n}} \right\} \right) \geq 1 - \exp(-c_2 \sqrt{n}).$$

Moreover, the event

$$\forall m \geq N - \frac{c_1 N}{\log N} \quad \forall y \in \mathbb{R}^N \quad |K_{N,y}|^{\frac{1}{n}} \geq c_1 \alpha_{y,m} L_K \sqrt{\frac{\log(2m/n)}{n}}$$

occurs with probability greater than $1 - \exp(-c_2 \sqrt{N})$.

The probability estimates in Proposition 3.3 are based on an estimate of corresponding probability for a fixed y and the union bound. We start with the following lemma.

Lemma 3.4 *There exist absolute positive constants c_1, c_2 such that the following holds. Let $n \leq m \leq N \leq e\sqrt{n}$ and X_1, \dots, X_N are independent random vectors distributed uniformly on an isotropic convex body K . Then for every $y \in \mathbb{R}^N$ with $y_m^* > 0$ there exists $v = v(y) \in \mathbb{R}^N$ having 0/1 coordinates with exactly m ones such that*

$$|K_{N,y}| \geq \alpha_{y,m}^n |K_{N,v}|$$

and

$$\mathbb{P} \left(\left\{ |K_{N,v}|^{\frac{1}{n}} \geq c_1 L_K \sqrt{\frac{\log(2m/n)}{n}} \right\} \right) \geq 1 - \exp(-c_2 \sqrt{m}).$$

Proof Fix $y \in \mathbb{R}^N$ with $y_m^* > 0$ (i.e., $|\text{supp } y| \geq m$). Let i_1, \dots, i_m be the indices such that $y_{i_j} = y_j^*$ and let $v = v(y) \in \mathbb{R}^N$ be the vector with $v_k = 1$ if $k = i_j$ and 0 otherwise. Decompose the polytope $K_{N,v}$ into a disjoint union of simplices (up to a set of zero measure)

$$K_{N,v} = \bigcup_{k=1}^{\ell} C_k,$$

where $C_k = \text{conv}\{0, \varepsilon_{k_1} X_{k_1}, \dots, \varepsilon_{k_n} X_{k_n}\}$ for some $\varepsilon_{k_j} = \pm 1$ and some vectors X_{k_j} , given by the simplicial decomposition of the facets of $K_{N,v}$. Denote

$$C_{k,y} = \text{conv}\{0, \varepsilon_{k_1} |y_{k_1}| X_{k_1}, \dots, \varepsilon_{k_n} |y_{k_n}| X_{k_n}\} \subset K_{N,y}.$$

Clearly, $C_{k,y}$'s are also disjoint up to a set of zero measure and

$$\begin{aligned} |C_k| &= |\det(\varepsilon_{k_1} X_{k_1}, \dots, \varepsilon_{k_n} X_{k_n})| |\text{conv}\{0, e_1, \dots, e_n\}| \\ &\leq \frac{1}{\alpha_{y,m}^n} |\det(\varepsilon_{k_1} |y_{k_1}| X_{k_1}, \dots, \varepsilon_{k_n} |y_{k_n}| X_{k_n})| |\text{conv}\{0, e_1, \dots, e_n\}|. \end{aligned}$$

This implies

$$|K_{N,v}| = \sum_{k=1}^{\ell} |C_k| \leq \alpha_{y,m}^{-n} \sum_{k=1}^{\ell} |C_{k,y}| \leq \alpha_{y,m}^{-n} |K_{N,y}|.$$

This proves the first estimate. The second one follows by Theorem 2.3, since $K_{N,v}$ is a symmetric random polytope in an isotropic convex body generated by $m \geq n$ random points. □

Proof of Proposition 3.3. Without loss of generality we only consider y 's satisfying $y_n^* > 0$ (otherwise estimates are trivial).

The first estimate follows from Lemma 3.4 and the union bound, since

$$\sum_{m \geq n} e^{-c_2 \sqrt{m}} \leq e^{-c_0 \sqrt{n}} \tag{3.1}$$

for an absolute positive constant c_0 .

To prove the second bound note that the set $\{v(y)\}_{y \in \mathbb{R}^N}$ ($v(y)$ is from Lemma 3.4) has cardinality $\binom{N}{m}$ and that denoting $k = N - m$

$$\binom{N}{m} \exp(-c_2\sqrt{m}) \leq \exp(-c_2\sqrt{m} + k \log(eN/k)) \leq \exp(-c_2\sqrt{m}/2),$$

provided that $k \leq c\sqrt{N}/\log N$. Lemma 3.4 and the union bound imply

$$\mathbb{P}\left(\left\{\forall y \in \mathbb{R}^N : |K_{N,v}|^{\frac{1}{n}} \geq c_1 \alpha_{y,m} L_K \sqrt{\frac{\log(2m/n)}{n}}\right\}\right) \geq 1 - \exp(-c_2\sqrt{m}/2).$$

The result follows by the union bound and (3.1). □

Proof of Theorem 1.2. For $n \leq N \leq e\sqrt{N}$ Propositions 3.1 and 3.3 imply the result, since, by (2.1),

$$nL_{K_{N,y}}^2 \leq \frac{1}{|K_{N,y}|^{1+\frac{2}{n}}} \int_{K_{N,y}} |x|^2 dx.$$

For $N \geq e\sqrt{n}$ the theorem follows from the general estimate $L_K \leq Cn^{1/4}$ for any n -dimensional convex body [22]. □

4 Random Perturbations of Random Polytopes

In this section $G = (g_1, \dots, g_N)$ denotes a standard Gaussian random vector in \mathbb{R}^N , independent of any other random variables. In the following theorem, which is a consequence of Theorem 1.2 and Lemma 2.4, we estimate $L_{K_{N,G}}$.

Theorem 4.1 *Let $n \leq N$. Let X_1, \dots, X_N be independent copies of a random vector uniformly distributed on an isotropic convex body. Then*

$$\mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ L_{K_{N,G}} \leq c_1 \sqrt{\log \frac{2N}{n}} \right\} \right) \geq 1 - \exp(-c_2\sqrt{n}),$$

where c_1 and c_2 are absolute positive constants.

Proof Without loss of generality we assume that $N \geq Cn$ for a sufficiently large absolute constant (see Remark 1 following Theorem 1.2). It is well-known (and can be directly calculated) that for the Gaussian vector $G = (g_1, \dots, g_N)$ one has

$$\mathbb{E} \|G\|_{n,2} \approx \sqrt{n \log \frac{N}{n}}.$$

Using concentration (see, e.g., Theorem 1.5 in [31]), we observe that for some absolute constant $C_1 > 0$,

$$\mathbb{P}_G \left(\|G\|_{n,2} \geq C_1 \sqrt{n \log \frac{N}{n}} \right) \leq \exp(-n \log(N/n)).$$

By Lemma 2.4 with $m = \lceil \sqrt{nN} \rceil$ we have

$$\mathbb{P} \left(\left\{ g_m^* \leq \sqrt{\log(\sqrt{N/n})} \right\} \right) \leq \exp \left(- \frac{N^{3/4} n^{1/4}}{10 \sqrt{\log \left(e \sqrt{\frac{N}{n}} \right)}} \right).$$

Since $\alpha_{G,m} \geq g_m^*$, Theorem 1.2 with $m = \lceil \sqrt{nN} \rceil$ in the infimum implies

$$L_{K_{N,G}} \leq C_2 \frac{\|G\|_{n,2}}{\sqrt{n}} \alpha_{G,m}^{-1} \frac{\log(2N/n)}{\sqrt{\log(2m/n)}} \leq C_3 \sqrt{\log \frac{N}{n}}.$$

with probability at least $1 - \exp(-c\sqrt{n}) - \exp(-n) - \exp\left(-c' \frac{N^{3/4} n^{1/4}}{\sqrt{\log \frac{eN}{n}}}\right)$. This implies the desired result. □

5 Concluding Remarks

In this section we show that under additional (strong) assumption that random vectors are ψ_2 vectors, the polytope $K_{N,G}$ contains Z_p for an appropriate p (recall that G is a standard Gaussian random vector in \mathbb{R}^N and the Z_p body was defined in (2.2)).

We first recall the definition of ψ_α norm. For a real random variable z and $\alpha \in [1, 2]$ we define the ψ_α -norm by

$$\|z\|_{\psi_\alpha} = \inf \{ C > 0 \mid \mathbb{E} \exp(|Y|/C)^\alpha \leq 2 \}.$$

It is well known (see, e.g., [12]) that the condition $\|z\|_{\psi_\alpha} \leq c_1$ is equivalent to the condition

$$\forall p > 1 : (\mathbb{E}|z|^p)^{1/p} \leq c_2 p^{1/\alpha} \mathbb{E}|z|. \tag{5.1}$$

Let X be a centered random vector in \mathbb{R}^n and $\alpha > 0$. We say that X is ψ_α or a ψ_α vector, if

$$\|X\|_{\psi_\alpha} := \sup_{y \in S^{n-1}} \|\langle X, y \rangle\|_{\psi_\alpha} < \infty. \tag{5.2}$$

We also denote

$$\gamma_p := (\mathbb{E}|g_1|^p)^{1/p} \approx \sqrt{p}.$$

Proposition 5.1 *There are absolute positive constants C, c_1 and c_2 such that the following holds. Let $\beta > 2$ and $N \geq C(\log \beta)^2 n \log(3n)/(\log \log(3n))$. Let X_1, \dots, X_N be independent copies of a random vector uniformly distributed on an isotropic convex body $K \subset \mathbb{R}^n$. Assume that $\|X_1\|_{\psi_2} \leq \beta L_K \sqrt{\log \frac{N}{n}}$. Then*

$$\mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ K_{N,G} \supseteq c_1 \sqrt{\log \frac{N}{n}} Z_{\log(N/n)}(K) \right\} \right) \geq 1 - \alpha,$$

where $\alpha = \exp(-c_2 \sqrt{N})$ for $N \geq (n/\log(3n))^2$ and $\alpha = (N/n)^{-c_2 N/n}$ otherwise.

Remark 1 We would like to note that for $N \geq n^2$ this Proposition (with slightly worse probability) was proved in [7] (see Lemma 4 there) without any assumptions on ψ_2 -norm.

Remark 2 Proposition 5.1 together with Proposition 3.1 and volume estimate (2.3) can be used to obtain the estimate of Theorem 4.1 (under additional ψ_2 assumption).

Our proof is very similar to the one given in [7]. We provide it for the sake of completeness. The main new ingredient is the following lemma, which is needed to estimate the norm of matrix A in the proof of Proposition 5.1.

Lemma 5.2 *Let g be a standard Gaussian random variable. Let X be uniformly distributed on an isotropic convex body $K \subset \mathbb{R}^n$. Assume that X is a ψ_2 random vector and that $\|X\|_{\psi_2} = \psi L_K$. Then for every $p \geq 1$,*

$$(\mathbb{E}|gX|^p)^{1/p} \leq CL_K (\sqrt{pn} + p\psi),$$

where C is an absolute positive constant.

Proof We use the following form of Paouris’s theorem, which may be deduced directly from Paouris’s work [30] (as formulated here it first appeared as Theorem 2 in [2], a short proof was given in [3])

$$(\mathbb{E}|X|^p)^{1/p} \leq C \left(\mathbb{E}|X| + \sup_{y \in S^{n-1}} (\mathbb{E}|\langle X, y \rangle|^p)^{1/p} \right).$$

Thus, using assumptions of X , we obtain

$$(\mathbb{E}|gX|^p)^{1/p} \leq \gamma_p C (L_K \sqrt{n} + \sqrt{p} L_K \psi),$$

which implies the result. □

Proof of Proposition 5.1. First note that $g_i X_i, i \leq N$, have ψ_1 norm bounded by $c_1 \beta L_K \sqrt{\log(N/n)}$. Indeed, for any $p \geq 1$ and $\theta \in S^{n-1}$, one has

$$(\mathbb{E}|\langle g_i X_i, \theta \rangle|^p)^{1/p} = \gamma_p (\mathbb{E}|\langle X_i, \theta \rangle|^p)^{1/p} \leq c_2 p \beta L_K \sqrt{\log \frac{N}{n}} \mathbb{E}|\langle X_i, \theta \rangle|.$$

Denote by A the $n \times N$ random matrix whose columns are the vectors $g_i X_i$. By Theorem 3.13 in [1] (cf. Theorem 2.1),

$$\mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \|A\| \geq c_3 \beta L_K \sqrt{N \log \frac{N}{n}} + 6 \max_{i \leq N} |g_i X_i| \right\} \right) \leq \exp(-2\sqrt{N}).$$

Now we estimate $\max_{i \leq N} |g_i X_i|$. If $N \geq (n/\ln(3n))^2$ we choose $p = 4\sqrt{N}$, otherwise $p = 4(N/n) \ln(N/n)$. In both cases,

$$\max \left\{ \sqrt{pn}, p\sqrt{\ln(N/n)} \right\} \leq 4\sqrt{N \ln(N/n)}.$$

Then by Lemma 5.2 and the Chebyshev inequality we have for every $i \leq N$,

$$\mathbb{P} \left(\left\{ |g_i X_i| \geq 8e C \beta L_K \sqrt{N \ln(N/n)} \right\} \right) \leq \frac{\mathbb{E}|g_i X_i|^p}{(8e C \beta L_K \sqrt{N \ln(N/n)})^p} \leq e^{-p}.$$

Union bound implies,

$$\mathbb{P} \left(\left\{ \max_{i \leq N} |g_i X_i| \geq 8e C \beta L_K \sqrt{N \ln(N/n)} \right\} \right) \leq N e^{-p} \leq e^{-p/2}$$

due to conditions on N and n . This implies

$$\mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \|A\| \geq c_5 \beta L_K \sqrt{N \log \frac{N}{n}} \right\} \right) \leq \alpha_0, \tag{5.3}$$

where $\alpha_0 = e^{-\sqrt{N}}$ for $N \geq (n/\ln(3n))^2$ and $\alpha_0 = \exp(-(N/n) \ln(N/n))$ otherwise.

On the other hand, for every $\sigma \subseteq \{1, \dots, N\}, q \geq 1$ and $\theta \in S^{n-1}$, by the Paley–Zygmund inequality,

$$\begin{aligned} & \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \max_{i \in \sigma} |\langle g_i X_i, \theta \rangle| \leq \frac{1}{2} (\mathbb{E}|g_1|^q)^{\frac{1}{q}} (\mathbb{E}|\langle X_1, \theta \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ &= \prod_{i \in \sigma} \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ |\langle g_i X_i, \theta \rangle| \leq \frac{1}{2} (\mathbb{E}|g_1|^q)^{\frac{1}{q}} (\mathbb{E}|\langle X_1, \theta \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ &\leq \left(1 - \left(1 - \left(\frac{1}{2} \right)^q \right)^2 \frac{(\mathbb{E}|g_1|^q \mathbb{E}|\langle X_1, \theta \rangle|^q)^2}{\mathbb{E}|g_1|^{2q} \mathbb{E}|\langle X_1, \theta \rangle|^{2q}} \right)^{|\sigma|}. \end{aligned}$$

Since $\gamma_p \approx \sqrt{p}$, and from Borell’s lemma ([8], see also Appendix III in [29]),

$$\mathbb{E}|\langle X_1, \theta \rangle|^{2q} \leq c_6^q (\mathbb{E}|\langle X_1, \theta \rangle|^q)^2,$$

the quantity above is bounded by

$$\left(1 - \frac{1}{4C^q}\right)^{|\sigma|} \leq \exp\left(-\frac{|\sigma|}{4C^q}\right).$$

Set $m = \lceil N/n \rceil$. Let $\sigma_1, \dots, \sigma_n$ be a partition of $\{1, \dots, N\}$ with $m \leq |\sigma_i|$ for every i and $\|\cdot\|_0$ be the norm

$$\|u\|_0 = \frac{1}{n} \sum_{i=1}^n \max_{j \in \sigma_i} |u_j|.$$

Note that $\|\cdot\|_0 \leq n^{-1/2}|\cdot|$. Since for all $1 \leq i \leq n$ and every $z \in \mathbb{R}^n$

$$h_{K_{N,G}}(z) = \max_{1 \leq j \leq N} |\langle g_j X_j, z \rangle| \geq \max_{j \in \sigma_i} |\langle g_j X_j, z \rangle|,$$

then for every $z \in \mathbb{R}^n$

$$h_{K_{N,G}}(z) \geq \|A^t z\|_0,$$

where A^t is the transpose of A .

Clearly, if $z \in \mathbb{R}^n$ verifies $\|A^t z\|_0 \leq \frac{1}{4}\gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}}$, then there exists a set $I \subseteq \{1, \dots, n\}$ with $|I| \geq \frac{n}{2}$ such that

$$\max_{j \in \sigma_i} |\langle g_j X_j, z \rangle| \leq \frac{1}{2}\gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}}$$

for every $i \in I$. Thus, for every $z \in \mathbb{R}^n$,

$$\begin{aligned} & \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \|A^t z\|_0 \leq \frac{1}{4}\gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ & \leq \sum_{|I|=\lceil \frac{n}{2} \rceil} \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \forall i \in I : \max_{j \in \sigma_i} |\langle g_j X_j, z \rangle| \leq \frac{1}{2}\gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ & \leq \sum_{|I|=\lceil \frac{n}{2} \rceil} \prod_{i \in I} \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \max_{j \in \sigma_i} |\langle g_j X_j, z \rangle| \leq \frac{1}{2}\gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ & \leq \sum_{|I|=\lceil \frac{n}{2} \rceil} \prod_{i \in I} \exp\left(-\frac{|\sigma_i|}{4C^q}\right) \leq 2^n \exp\left(-\frac{nm}{4C^q}\right) \leq 2^n \exp\left(-\frac{N}{8C^q}\right) \end{aligned}$$

$$\leq \exp\left(-\frac{\sqrt{Nn}}{16}\right)$$

provided that $q := (1/2) \log(N/n)$ and $N > 125n$.

Now, let

$$S = \left\{ z \in \mathbb{R}^n \mid \frac{1}{2} \gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} = 1 \right\}$$

and let $U \subset S$ be a δ -net (in metric defined by S) with cardinality $|U| \leq (\frac{3}{\delta})^n$, i.e., for every $z \in S$ there is $u \in U$ such that $\frac{1}{2} \gamma_q (\mathbb{E}|\langle X_1, z - u \rangle|^q)^{\frac{1}{q}} \leq \delta$. Then

$$\mathbb{P}\left(\left\{\exists u \in U : \|A^t u\|_0 \leq \frac{1}{2}\right\}\right) \leq \exp\left(n \log \frac{3}{\delta} - \sqrt{Nn}/16\right).$$

By isotropicity we have that $(\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \geq L_k |z|$ (because we have chosen $q = (1/2) \log(N/n) > 2$). Thus, assuming $\|A^t\| \leq c_5 \beta L_K \sqrt{N \log \frac{N}{n}}$, we have

$$\begin{aligned} \|A^t z\|_0 &\leq \frac{1}{\sqrt{n}} |A^t z| \leq c_5 \beta L_K \sqrt{\frac{N}{n}} \sqrt{\log \frac{N}{n}} |z| \leq c_5 \beta \sqrt{\frac{N}{n}} \sqrt{2q} (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \\ &\leq c_6 \beta \gamma_q \sqrt{\frac{N}{n}} (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}}. \end{aligned}$$

Therefore, if $u \in U$ approximates $z \in S$, that is if $\frac{1}{2} \gamma_q (\mathbb{E}|\langle X_1, z - u \rangle|^q)^{\frac{1}{q}} \leq \delta$, then u also satisfies

$$\|A^t u\|_0 \leq \|A^t z\|_0 + c_7 \beta \sqrt{\frac{N}{n}} \delta.$$

Choosing $\delta = \sqrt{n}/(4\beta c_7 \sqrt{N})$ and denoting the event

$$\Omega_0 := \left\{ \|A\| \leq c_5 \beta L_K \sqrt{N \log \frac{N}{n}} \right\}$$

we obtain

$$\begin{aligned} &\mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \omega \in \Omega_0 \mid \exists z \in \mathbb{R}^n : \|A^t z\|_0 \leq \frac{1}{8} \gamma_q (\mathbb{E}|\langle X_1, z \rangle|^q)^{\frac{1}{q}} \right\} \right) \\ &= \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \omega \in \Omega_0 \mid \exists z \in S : \|A^t z\|_0 \leq \frac{1}{4} \right\} \right) \\ &\leq \mathbb{P}_{G, X_1, \dots, X_N} \left(\left\{ \omega \in \Omega_0 \mid \exists u \in U : \|A^t u\|_0 \leq \frac{1}{2} \right\} \right) \end{aligned}$$

$$\leq \exp\left(n \log \frac{12c_7 \beta \sqrt{N}}{\sqrt{n}} - \sqrt{Nn}/16\right) \leq \exp\left(-\sqrt{Nn}/20\right),$$

provided $N \geq C(\log \beta)^2 n$ for a big enough absolute constant C . Since $h_{K_{N,G}}(z) = \|A^t z\|_\infty \geq \|A^t z\|_0$, this together with (5.3) and the definition (2.2), implies that with probability at least $1 - \alpha_0 - \exp\left(-\sqrt{Nn}/20\right)$,

$$K_{N,G} \supseteq \frac{1}{8} \gamma_q Z_q(K) \supseteq c \sqrt{\log \frac{N}{n}} Z_{\log(N/n)}(K).$$

□

Acknowledgments Part of this work was done while the first named author was a postdoctoral fellow at the Department of Mathematical and Statistical Sciences at University of Alberta, and continued while he was a “Juan de la Cierva” postdoctoral researcher at the Mathematics Department in Universidad de Murcia. He would like to thank both departments for providing support, good environment and working conditions. We would also like to thank the anonymous referee for careful reading of the manuscript and for informing us that Theorem 1.1 has subsequently been proved also in [17] using essentially the same approach. Finally we are very thankful to Rafał Latała for showing us the argument in Lemma 5.2. David Alonso-Gutiérrez is Partially supported by MICINN project MTM2010-16679, MICINN-FEDER project MTM2009-10418, “Programa de Ayudas a Grupos de Excelencia de la Región de Murcia”, Fundación Séneca, 04540/GERM/06 and Institut Universitari de Matemàtiques i Aplicacions de Castelló Alexander E. Litvak Research partially supported by the E.W.R. Steacie Memorial Fellowship Nicole Tomczak-Jaegermann holds the Canada Research Chair in Geometric Analysis.

References

1. Adamczak, R., Litvak, A.E., Pajor, A., Tomczak-Jaegermann, N.: Quantitative estimates of the convergence of the empirical covariance matrix in log-concave ensembles. *J. Am. Math. Soc.* **23**(2), 535–561 (2010)
2. Adamczak, R., Latała, R., Litvak, A.E., Pajor, A., Tomczak-Jaegermann, N.: Geometry of log-concave Ensembles of random matrices and approximate reconstruction. *C.R. Math. Acad. Sci. Paris* **349**, 783–786 (2011)
3. Adamczak, R., Latała, R., Litvak, A.E., Oleszkiewicz, K., Pajor, A., Tomczak-Jaegermann, N.: A short proof of Paouris’s inequality. *Can. Math. Bull.* **57**, 3–8 (2014)
4. Alonso-Gutiérrez, D.: On the isotropy constant of random convex sets. *Proc. AMS* **136**(9), 3293–3300 (2008)
5. Alonso-Gutiérrez, D.: A remark on the isotropy constant of polytopes. *Proc. AMS* **139**(7), 2565–2569 (2011)
6. Alonso-Gutiérrez, D., Bastero, J., Bernués, J., Wolff, P.: On the isotropy constant of projections of polytopes. *J. Funct. Anal.* **258**(5), 1452–1465 (2010)
7. Alonso-Gutiérrez, D., Prochno, J.: Mean width of random perturbations of random polytopes. preprint
8. Borell, C.: Convex measures on locally convex spaces. *Ark. Mat.* **12**, 239–252 (1974)
9. Bourgain, J.: On high-dimensional maximal functions associated with convex bodies. *Am. J. Math.* **108**(6), 1467–1476 (1986)
10. Bourgain, J.: On the distribution of polynomials on high-dimensional convex sets. *GAF*A (1989–90), Lecture Notes in Math., vol. 1469, pp. 127–137. Springer, Berlin (1991)
11. Brazitikos, S., Giannopoulos, A., Valettas, P., Vritsiou, B.: *Geometry of Isotropic Convex Bodies*. Mathematical Surveys and Monographs, vol. 196. American Mathematical Society, Providence (2014)
12. Chafaï, D., Guédon, O., Lecué, G., Pajor, A.: Interactions between compressed sensing random matrices and high dimensional geometry, *Panoramas et Synthèses [Panoramas and Syntheses]*, 37. Soc. Math. de France, Paris (2012)

13. Dafnis, N., Giannopoulos, A., Guédon, O.: On the isotropic constant of random polytopes. *Adv. Geom.* **10**, 311–321 (2010)
14. Dafnis, N., Giannopoulos, A., Tsolomitis, A.: Quermass integrals and asymptotic shape of random polytopes in an isotropic convex body. *Mich. Math. J.* **62**, 59–79 (2013)
15. Dafnis, N., Giannopoulos, A., Tsolomitis, A.: Asymptotic shape of a random polytope in a convex body. *J. Funct. Anal.* **257**, 2820–2839 (2009)
16. Gordon, Y., Litvak, A.E., Schuett, C., Werner, E.: On the minimum of several random variables. *Proc. AMS* **134**, 3665–3675 (2006)
17. Giannopoulos, A., Hioni, L., Tsolomitis, A.: Asymptotic shape of the convex hull of isotropic log-concave random vectors. preprint
18. Gluskin, E.D.: The diameter of Minkowski compactum roughly equals to n . *Funktional. Anal. i Prilozhen.* **15**(1), 72–73 (1981). English Trans: *Funct. Anal. Appl.* **15**(1), 57–58 (1981)
19. Gluskin E.D., Litvak A.E.: A remark on vertex index of the convex bodies, *GAFa*. Lecture Notes in Math., vol. 2050, pp. 255–265. Springer, Berlin (2012)
20. Junge, M.: Hyperplane conjecture for quotient spaces of L_p . *Forum Math.* **6**, 617–635 (1994)
21. Litvak, A.E., Rudelson, M., Tomczak-Jaegermann, N.: On approximation by projections of polytopes with few facets. *Israel J. Math.* **203**, 141–160 (2014)
22. Klartag, B.: On convex perturbations with a bounded isotropic constant. *Geom. Funct. Anal. (GAFa)* **16**(6), 1274–1290 (2006)
23. Klartag, B., Kozma, G.: On the hyperplane conjecture for random convex sets. *Israel J. Math.* **170**(1), 1–33 (2009)
24. Klartag, B., Milman, E.: Centroid bodies and the logarithmic Laplace transform—a unified approach. *J. Funct. Anal.* **262**(1), 10–34 (2012)
25. Lutwak, E., Yang, D., Zhang, G.: L_p affine isoperimetric inequalities. *J. Diff. Geom.* **56**(1), 111–132 (2000)
26. Lutwak, E., Zhang, G.: Blaschke–Santaló inequalities. *J. Diff. Geom.* **47**(1), 1–16 (1997)
27. Mankiewicz, P., Tomczak-Jaegermann, N.: Quotients of finite-dimensional Banach spaces; random phenomena. In: Johnson, W.B., Lindenstrauss, J. (eds.) *Handbook in the Geometry of Banach Spaces*, vol 2, pp. 1201–1246. Elsevier, Amsterdam (2001)
28. Milman, V.D., Pajor, A.: Isotropic positions and inertia ellipsoids and zonoids of the unit ball of a normed n -dimensional space. *GAFa Seminar 87–89*, Springer. Lecture Notes in Math., vol. 1376, pp. 64–104. Springer, New York (1989)
29. Milman, V.D., Schechtman, G.: Asymptotic theory of finite-dimensional normed spaces. *Lecture Notes in Math.*, vol. 1200. Springer, Berlin-New York (1985)
30. Paouris, G.: Concentration of mass on convex bodies. *Geom. Funct. Anal.* **16**, 1021–1049 (2006)
31. Pisier, G.: Probabilistic methods in the geometry of Banach spaces, probability and analysis (Varenna, 1985). *Lecture Notes in Math.*, vol. 1206, pp. 167–241. Springer, Berlin (1986)
32. Pivovarov, P.: On determinants and the volume of random polytopes in isotropic convex bodies. *Geom. Dedicata* **149**, 45–58 (2010)