

Nutrients in the Western Wadden Sea: Freshwater Input Versus Internal Recycling

Catarina Leote¹ · Lieke L. Mulder² · Catharina J. M. Philippart¹ · Eric H. G. Epping¹

Received: 27 November 2014 / Revised: 20 April 2015 / Accepted: 22 April 2015 / Published online: 6 May 2015
© Coastal and Estuarine Research Federation 2015

Abstract At present, phosphorus (P) is seen as the main limiting nutrient for phytoplankton growth in the western Wadden Sea. Six cruises were performed for water sampling at selected stations covering a full tidal cycle for later determination of dissolved and particulate nutrient concentrations. The major P sources were identified on a seasonal basis, by comparing the contribution of freshwater discharge and sediment release, calculated in a previous study, to the concentrations in the water column. A close relationship was found between the pelagic concentrations of dissolved inorganic nutrients and chlorophyll *a*, with a concomitant decrease in nutrients and increase in chlorophyll. This was observed in early spring and was followed by a later increase in the nutrient concentrations in spring–summer. The low concentrations found for the freshwater and seawater end-members for this period ruled out their importance as nutrient sources, suggesting that this increase resulted mainly from internal recycling in the Wadden Sea. Even though P limitation was observed during most of the year, a potential seasonal change in the limiting nutrient, from P to silica, was observed. The comparison between P supply to the Wadden Sea by freshwater discharge and sediment release showed a much higher contribution of the latter, especially in April–November. To our knowledge, this is the first study clearly presenting internal recycling as

the main nutrient source to the western Wadden Sea in spring–autumn, instead of freshwater discharge or the North Sea.

Keywords Nutrients · Mineralization · Wadden Sea · Primary production · Sorption · Freshwater discharge

Introduction

The Wadden Sea is a shallow coastal sea, separated from the North Sea by a chain of barrier islands, stretching for 600 km from the Netherlands to Denmark. It represents a complex system of intertidal flats and gullies creating a highly diverse ecosystem of high biological productivity (Beukema et al. 2002). The primary supply of nutrients to its westernmost area, the Marsdiep basin, occurs via direct freshwater discharge from the adjacent Lake IJssel and indirectly via the coastal bound residual flow of Rhine and Scheldt river waters admixed with North Sea water. The atmospheric contribution to the overall nutrient input through wet and dry deposition is considered secondary (van Raaphorst et al. 2000). Nutrients are supplied in dissolved and particulate organic and inorganic forms, where the form dominating the input depends on the element and stage of the seasonal productivity cycle. Especially, the input of particulate organic matter from the North Sea has since long received considerable attention as to explain the high productivity and heterotrophic status of this basin (Postma 1954; van Beusekom and de Jonge 2002; van Beusekom 2005).

Similar to many European coastal waters, the western Wadden Sea experienced a major increase in the nutrient loading from freshwater sources. From the 1950s until the 1980s, the loading of nitrogen (N) and phosphorus (P) increased by a factor of 12 and 10, respectively, due to anthropogenic activities (van Raaphorst and de Jonge 2004). The increase in

Communicated by James L. Pinckney

✉ Catarina Leote
catleote@gmail.com

¹ Royal Netherlands Institute for Sea Research (NIOZ)—Division Texel, PO box 59, 1790 AB, Texel Den Burg, The Netherlands

² Royal Netherlands Institute for Sea Research (NIOZ)—Division Yerseke, PO box 140, 4400 AC Yerseke, The Netherlands

nutrient supply and presumably availability was paralleled by a rise in primary production from an estimated $20 \text{ g C m}^{-2} \text{ year}^{-1}$ prior to the 1950s up to $520 \text{ g C m}^{-2} \text{ year}^{-1}$ in 1986, supporting the view of a strict and causal relationship between freshwater discharge and primary production (de Jonge 1990). Restrictive policies in nutrient riverine discharge to the coastal zone resulted in a reduction of the loadings to values close to the pre-1950s (de Jonge 1997; Colijn and van Beusekom 2005), followed by a proportional response in the nutrient concentrations in the Wadden Sea (de Vries et al. 1998; Kuipers and van Noort 2008). This was particularly effective for P resulting in dissolved N/P ratios up to 100, suggesting severe P limitation on the primary production (van Raaphorst and de Jonge 2004; Philippart et al. 2007; Ly et al. 2014). Despite the decrease in nutrient loadings, high levels of biomass primary productivity were maintained, and only since the 2000s, a significant decrease was observed in biomass during the autumn blooms and in production, to present-day values of $\sim 200 \text{ g C m}^{-2} \text{ year}^{-1}$ (de Jonge 1990; Cadée and Hegeman 1993; de Jonge et al. 1996; van Beusekom 2005; Kuipers and van Noort 2008). The lag in productivity with decreased nutrient loading is generally attributed to the temporal storage of nutrients, notably P, in the sediment and controlled exchange with the overlying water, dampening the decrease in freshwater loading.

The goal of this study was to understand the interaction between phytoplankton growth and nutrient availability in the water column and to identify the main seasonal nutrient, especially P, sources to primary producers in the modern western Wadden Sea. Assumed as the main nutrient sources to the study area, the contributions of freshwater discharge and sediment release to the pelagic P concentrations were compared on a seasonal scale. In addition to dissolved nutrients, the concentration and composition of suspended particulate matter and chlorophyll *a* content were considered. This may provide information not only about the growth cycle of primary producers but also on the potential availability of nutrients following the degradation of the particulate material.

Methods

Study Site

The Marsdiep basin, the westernmost tidal basin of the Wadden Sea, covers an area of about 700 km^2 , $\sim 580 \text{ km}^2$ of which are tidal channels (Dastgheib 2007) (Fig. 1). The basin, with an average depth of 3.3 m, is subject to a mesotidal regime, with a mean tidal range of 1.37 m and a tidal frequency of 1.92 tides day^{-1} . The tidal residual transport ranges $1\text{--}3 \times 10^3 \text{ m}^3 \text{ s}^{-1}$ (Ridderinkhof et al. 2002; Nauw et al. 2014). Because of freshwater discharge from Lake IJssel and other minor sources, the area behaves

like an estuary, covering a salinity range from 0.7 (near the sluices) to 30 (near the tidal inlets), with a flushing time of the discharged water of 5.5 days (van Aken 2008; Nauw et al. 2014; this study). On average, the water in this basin is composed of $\sim 85 \%$ North Sea seawater, with a 15 % admixture of Lake IJssel water (de Jonge et al. 1996), and the bulk of the sediment and particulate matter supply mainly originates from the North Sea (Oost and de Boer 1994). The suspended particulate matter concentrations are relatively high ranging $10\text{--}30 \text{ mg l}^{-1}$ (Nauw et al. 2014; Philippart et al. 2013), resulting in light penetration depths of 0.1–0.5 m (Hommerson et al. 2009; Loebel et al. 2009). Therefore, benthic photosynthesis is unlikely in the subtidal domain. The Marsdiep basin is largely composed of sandy sediments with a variable silt (particle size $< 63 \mu\text{m}$) content (1–10 %). Finer-grained sediments, with silt contents up to 72 %, represent only 2 % of the basin and are found adjacent to the coast (Leote et al. 2014). Most sediment P is in a non-easily exchangeable form, with the bioavailable fraction corresponding to $\sim 1/3$ of the total sediment P pool (Leote et al. 2014).

Sampling Scheme

Sixteen stations were selected to cover the Marsdiep tidal basin and the freshwater (Lake IJssel) and seawater (North Sea) end-members (Fig. 1). Six stations (2, 5, 6, 11, 14, and 17) were seasonally sampled during a full tidal cycle ($\sim 13 \text{ h}$) whereas station 16, located in Lake IJssel, was sampled only once per cruise. Sampling campaigns were carried out on 20–24 April and 2–7 November 2009 and on 15–17 February, 22–26 March, 3–7 May, and 6–10 September 2010 for stations 2, 5, and 11. Because of logistic and weather constraints, stations 6 and 17 were not sampled in March 2010 and April 2009, respectively. Station 14 was only sampled in April 2009 and March 2010, while station 16 was sampled in April 2009 and in March, May, and September 2010. All stations were sampled within the same week. For each station, water samples were collected every hour with a Niskin bottle from the side of the ship at three depths: surface, mid column, and bottom. Each sampling event was preceded by conductivity, temperature, and depth (CTD) profiling to retrieve information on the tidal stage locally.

Dissolved Nutrients

Water samples were collected with a syringe and filtered ($0.2\text{-}\mu\text{m}$ pore diameter Acrodisc filter, Supor® membrane) for later analysis of NO_3 , NO_2 , NH_4 , total dissolved nitrogen (TDN), Si, PO_4 , and total dissolved phosphorus (TDP). Samples were stored in pony vials at $-20 \text{ }^\circ\text{C}$ for NO_3 , NO_2 , NH_4 , TDN, and TDP and at $4 \text{ }^\circ\text{C}$ for PO_4 and Si.




Fig. 1 Map of the western Wadden Sea with the sampling locations in the Wadden Sea (1–13, 17), North Sea (14, 15), and Lake IJssel (16). The enclosed stations were seasonally sampled

Particulate Matter

To determine the total suspended matter (TSM), particulate total C, N, and P, organic C, and chlorophyll *a* contents, 100–300 ml of water were collected every 2 h and filtered using pre-ashed (500 °C for 4 h) Whatman GF/F filters. Filters for the analysis of TSM, total C, total N, organic C, and chlorophyll *a* content were stored at –20 °C until analysis, while the extraction for total particulate P started immediately after filtration. The chlorophyll *a* contents were only determined in 2010, while the TSM and particulate C, N, and P were sampled from November 2009 on.

Analytical Procedures

All analyses on dissolved nutrients were performed on a TRAACS 800 segmented continuous flow analyser. NO_x ($\text{NO}_3 + \text{NO}_2$) and NH_4 were analysed according to Grasshoff et al. (1999) and Helder and de Vries (1979) with detection limits of 0.04 and 0.07 μM and precisions of 0.02 and 0.03 μM , respectively. PO_4 and Si were analysed according to the methods of Murphy and Riley (1962) and Strickland and Parsons (1968), with detection limits of 0.007 and

0.03 μM and a precision of 0.002 and 0.016 μM , respectively. The analyses of TDN and TDP followed the NIOZ Nutrient Lab in-house method based on Schreurs and Nijssse (2000) and Nijssse and Spronk (2000), with a detection limit and precision of 0.03 μM . All analyses had an accuracy within the 95 % confidence interval assigned for the reference materials used, as confirmed in the intercomparison rounds of QUASIMEME and MRI Japan (Aoyama et al. 2010). The concentration of dissolved organic N (DON) was determined as the difference between TDN and $\text{NO}_3 + \text{NO}_2 + \text{NH}_4$, while the dissolved organic P (DOP) was calculated as $\text{TDP} - \text{PO}_4$. The TSM concentration was determined by subtracting the filter weight from the weight of the freeze-dried filter with sample. The contents of total particulate C and N were measured on the filters using a Thermo-Interscience Flash EA1112 series elemental analyser according to Verardo et al. (1990), with detection limits of 2.5 and 4.8 wt%, a precision of 9.7 and 2.7 %, and an accuracy of 0.4 and 1.0 % for N and C, respectively. The organic C was determined following the same procedure after removal of the carbonates with 2 M HCl. The total particulate P content was determined according to the persulfate oxidation method from Valderrama (1981), improved to a digestion time of 90 min by G. Kramer and K.

Bakker (NIOZ Nutrient Lab). Chlorophyll *a* was analysed in a high-performance liquid chromatographer (HPLC) after extraction in a 90 % acetone medium (Jeffrey et al. 1997).

Data Analyses

Statistical Significance

The significance of the seasonal differences in the concentrations of dissolved and particulate nutrients, TSM, and chlorophyll *a* was assessed by Student's *t* test in Excel (Microsoft ©), assuming a two-tailed distribution and an unequal variance.

Mixing Diagrams

In estuarine systems, a plot of the nutrient concentration versus a conservative tracer such as salinity may provide information concerning the fate and origin of nutrients between the freshwater and seawater end-members. For the Marsdiep tidal basin, such mixing diagrams were compiled for each cruise with the concentrations of dissolved nutrients (NO_3 , NH_4 , PO_4 , and Si) plotted against salinity. A negative relationship between salinity and dissolved nutrient concentration implied that the freshwater end-member was the main source of the nutrient for this tidal basin, while a positive relationship indicated that the oceanic end-member was the main source. The occurrence of a concave relation or the presence of lower concentrations at intermediate salinities indicated a loss of that nutrient within the tidal basin (e.g. denitrification or sedimentation). Oppositely, a convex relationship or higher concentrations at intermediate salinities indicated production or additional sources within the tidal basin (e.g. mineralization or local input).

Quantification of the P Sources

A comparison was made between the relative importance to the pelagic P concentrations of freshwater discharge and release from the sediments. To do so, the monthly freshwater P discharge from the Lake IJssel sluices and the harbour of Den Helder was estimated and compared with the monthly sediment P release for the Marsdiep basin according to Leote and Epping (2015). The discharge from Lake IJssel was calculated by multiplying the summed sluice debits in Den Oever and Kornwerderzand (Fig. 1) by the PO_4 concentrations from Lake IJssel (station Vrouwezand) using data from the Rijkswaterstaat Waterbase (live.waterbase.nl). The database provided one daily debit value (in $\text{m}^3 \text{s}^{-1}$), which was assumed to be valid for 24 h. The debit in Den Helder was assumed to be 1/20 of the total debit from Lake IJssel based on van Raaphorst and van der Veer (1990). The database provided one monthly PO_4 concentration value for station Vrouwezand, which was assumed valid for the period between

samplings. The PO_4 concentration of the freshwater discharging in Den Helder was assumed to correspond to the concentrations from the 1950s (Postma 1954), which agree well with the present concentrations from Lake IJssel. The freshwater discharge was calculated between January 2009 and December 2010, so that the monthly discharge corresponded to an average value for the 2 years. Values for sediment nutrient release were taken from Leote and Epping (2015) who estimated the PO_4 release for the entire Marsdiep basin based on a relationship found between the silt content and the sediment-water exchange for the same stations presented in this study. The sediment-water exchange was estimated based on porewater profiles of nutrient concentration using Fick's law and correcting for the effect of macrofauna activity. No benthic microphotosynthetic communities were detected since oxygen microprofiles under different light intensities did not change with increased light availability (data not shown).

Results

Dissolved Nutrients


Measured Values

The concentrations of the major dissolved nutrients (NO_3 , NH_4 , PO_4 , Si, DON, and DOP) were, in general, similar for all depths, with the exception of station 5, close to the freshwater sluices of Den Oever (Fig. 1), indicating a well-mixed water column with stratification only near the sluices or following a major freshwater discharge. The average NO_3 concentrations ranged 2–192 μM , while NH_4 had a much smaller variation (0.2–17 μM) (Fig. 2). PO_4 varied between 0 and 1 μM , while Si reached a maximum of 76 μM and a minimum of 0 μM . The DOP concentrations ranged 0–0.5 μM , while DON ranged 6–41 μM .

Seasonal Pattern

A clear seasonal pattern was observed (Fig. 2). For Si, PO_4 , and NH_4 , a dramatic decrease in concentrations was observed in February–April, when nutrient depletion was observed for all stations except 5 and 16 (located close to and in Lake IJssel, respectively), followed by an increase in May, especially for NH_4 and PO_4 . A second drop in the concentrations was observed in September for PO_4 and NH_4 and in November for Si. This similar seasonal trend may point to common sources and sinks for these three nutrients. Oppositely, NO_3 concentrations decreased throughout the year for all Wadden Sea stations, suggesting different sources and sinks. The DOP and DON concentrations increased in February–April,

Fig. 2 Seasonal variation of the concentrations in the water column of inorganic (NO_3 , Si, PO_4 , and NH_4) and organic (DOP and DON) dissolved nutrients, including measurements from 2009 to 2010. The average values for all depths and for a full tidal cycle are presented with standard deviations



followed by a decrease until November. The DON showed more spatial variation than the DOP, with higher concentrations for stations 5 and 16, clearly associated with freshwater. The seasonal trends of inorganic and organic nutrients were opposite, with higher concentrations of organic nutrients (DOP, DON) in parallel with a decrease in the inorganic nutrients (PO_4 , NH_4). The concentration differences between months were always significant ($p < 0.01$), with the exception of April–May for NO_3 ($p = 0.53$), March–April for Si ($p = 0.03$), and March–April for NH_4 ($p = 0.62$).

Mixing Diagrams

Regarding NO_3 , Lake IJssel was clearly a source for the Marsdiep basin in February–May, with a near-conservative behaviour in February and May and net removal in March and probably in April (Fig. 3). In

September–November, the concentrations in Lake IJssel and the Wadden Sea decreased substantially to close to zero and the North Sea became a minor source of NO_3 . The NH_4 concentration was always lower in the freshwater end-member. Therefore, the elevated concentrations along the salinity gradient likely resulted from production/regeneration within the Wadden Sea, most prominent in May, but also evident in February, September, and November. The PO_4 concentration was also always lower in Lake IJssel, except in May. In February, an approximately conservative behaviour was observed, with the North Sea as the source. In March–April, the concentrations were very low, while in May, the concentrations for the intermediate salinities were clearly above the predicted, indicating PO_4 production, while in September–November, they were below, suggesting consumption or eventual adsorption to Fe oxyhydroxides from surface sediments. Si concentration




Fig. 3 Mixing diagrams for the major inorganic nutrients (NO_3 , NH_4 , PO_4 , and Si) plotted separately for each cruise. Surface values for all sampled stations are presented: station 2 (closed triangles), station 5 (open circles), station 6 (dark grey hexagons), station 11 (open triangles), station 14 (light grey rhombuses), station 16 (closed circles), and station 17 (open squares). The other stations sampled in April 2009

(1, 3, 4, 7–9, 12, 13, 15) are represented by closed circles. The line represents the predicted concentrations assuming mixing between the end-members as the only process affecting the concentration distribution. The end-member concentrations (North Sea and Lake IJssel) were taken from the Rijkswaterstaat Waterbase (live.waterbase.nl)

was mostly higher in the freshwater end-member. In February, Si showed a conservative behaviour, while in March, the concentrations decreased, with consumption at intermediate salinities. In April–May, all concentrations were close to zero, with a slight increase at the intermediate salinities in May. In September, the concentrations were below the predicted with Lake IJssel as the source, while in November, the concentrations levelled off between the freshwater and seawater end-members.

Particulate Matter

Measured Values

The concentration of TSM varied between 15 and 120 mg l^{-1} (Fig. 4). The total and organic C contents ranged 6–25 and 1–21 wt%, respectively, with the highest values for Lake IJssel in March–September. The total N content was much lower, 0.5–3 wt%, while the total particulate P ranged 14–69 $\mu\text{mol g}^{-1}$, being higher in Lake IJssel in March and September, respectively. The organic C/total P ratio provides additional information concerning the composition of particulate P, i.e. biogenic- or terrigenous/mineral-associated. During our sampling period, it ranged 38–841, mostly above the Redfield

ratio (C:P=106:1), being higher in February in Den Oever and in March and May in Lake IJssel.

Seasonal Pattern

Like the dissolved nutrients, the TSM, total C, N, P, organic C, and organic P/total C ratio showed a clear seasonal variation (Fig. 4). The TSM concentration and standard deviation were lower in February–March, increasing in May–September and decreasing in November. The contents of total and organic C, total N, and especially the particulate P showed a comparable seasonal pattern. The organic C/total P ratio showed an opposite seasonal profile, with higher values in February–March, decreasing in May–September and slightly increasing in November. The differences in concentrations between the months were always significant ($p < 0.01$), with the exception of February–March and May–September for both TSM and total P ($p = 0.95$ and 0.92 for TSM; $p = 0.07$ and 0.08 for total P, respectively).

Chlorophyll *a*

The average chlorophyll *a* concentrations varied between $\sim 4 \mu\text{g l}^{-1}$ in February and $84 \mu\text{g l}^{-1}$ in




Fig. 4 Seasonal variation of the concentration of total suspended matter (TSM), total C, organic C, total N, total P, and organic C/total P ratio in the water column for all seasonally sampled stations. Values presented

correspond to an average for all depths and for a full tidal cycle with the respective standard deviations

September (Fig. 5). The concentrations and respective standard deviations were lower in February and May and higher in March and September, especially for the surface samples, with lower values for the mid-column and bottom depths (data not shown). The spatial variation was also higher in March and September. The seasonal differences observed were statistically significant ($p < 0.01$).




Fig. 5 Seasonal variation in the concentrations of chlorophyll *a* in the water column for all the seasonally sampled stations. The plot includes the average values for all depths and for a full tidal cycle with the respective standard deviations

Tidal Variation

In addition to the seasonal pattern, a tidal pattern was observed for the concentrations of particulate matter in the water column. Taking station 17, located in the Balgzand tidal flat, as an example, in February–March, the TSM concentration was very low and similar for all depths, with no significant variation during the tidal cycle (Fig. 6). However, in May–September, the concentrations increased approximately after maximum tidal current speed (i.e. between high and low tide), especially at the bottom, suggesting resuspension of deposited material. A similar pattern was observed for chlorophyll *a*, indicating that a part of the resuspended material corresponded to fresh organic matter. The plots of the concentration of chlorophyll *a* versus the TSM were highly correlated, with chlorophyll *a* representing a larger fraction of the TSM in March–September.

Discussion

Nutrient Concentrations Linked to Primary Production

Throughout the year, a clear relationship between the concentrations of chlorophyll *a* and the major inorganic and organic




Fig. 6 Total suspended matter (TSM) and chlorophyll *a* concentrations plotted for a full tidal cycle and chlorophyll *a* versus TSM concentrations. Values presented are for station 17 for surface (closed circles), mid (open

circles), and bottom (closed triangles) depth. The line in the first two rows indicates the tide height

nutrients was observed in the Wadden Sea and Lake IJssel (Cadée and Hegeman 1993; Loebel et al. 2007; van Beusekom et al. 2009). This is based on the assumption that the concentrations of dissolved and particulate nutrients and chlorophyll *a* did not differ greatly between 2009 and 2010 since we use a combination of data from both years. Even though this is not ideal, studies on the seasonal variation of nutrients, turbidity, salinity, and chlorophyll *a* suggest this is a fair assumption (Cadée and Hegeman 2002; van Beusekom and de Jonge 2002; Colijn and Cadée 2003; Kuipers and van Noort 2008; van Aken 2008; Philippart et al. 2000; 2010; 2013). In February, the concentrations of inorganic nutrients were within the highest measured (Fig. 2). Lake IJssel was clearly the source of NO_3 and Si, according to the mixing diagrams, with the concentrations at intermediate salinities resulting from mixing between the two end-members, i.e. conservative behaviour (Fig. 3). However, this did not seem to be the case for PO_4 and NH_4 , which showed very low concentrations in the freshwater end-member and no conservative behaviour. The higher concentrations of chlorophyll *a* for

station 5 (close to a freshwater sluice) suggest the occurrence of an earlier phytoplankton bloom in Lake IJssel consuming PO_4 and NH_4 , in agreement with previous studies (Cadée 1986; de Vries et al. 1998). Later in March and April, the increase in the chlorophyll *a* concentrations (Fig. 5) was coupled with depletion in the water column of NH_4 , PO_4 , and Si, the most important nutrients for phytoplankton growth (Crouzet et al. 1999) (Fig. 2), suggesting a first phytoplankton bloom in the western Wadden Sea between February and March. Consumption overcame production/supply, and the nutrient concentrations in the freshwater and seawater end-members and in the Wadden Sea levelled off. The higher contents of organic C and total N in the particulate matter (Fig. 4), indicating that a large fraction of the suspended material was organic, corroborate this idea. Moreover, a first phytoplankton bloom in early March is in line with previous studies (Cadée 1986; Cadée and Hegeman 1993; Philippart et al. 2000; Kuipers and van Noort 2008; Philippart et al. 2010). After the initial bloom, nutrient depletion likely limited the growth of primary producers, as apparently seen in May,

with a decrease in the concentrations of chlorophyll *a* and particulate organic C and total N (Figs. 4 and 5). In May, the increase in the concentrations of PO₄, NH₄, and Si indicates that the supply of nutrients, probably from mineralization of organic material, exceeded the consumption by primary producers (Figs. 2 and 3). The growth of primary producers was no longer limited by the availability of nutrients, allowing for a second bloom in September, evidenced by the concomitant increase in the concentrations of chlorophyll *a*, total C, total N, and organic C concentrations and decrease in the concentrations of NH₄, PO₄, and Si (Figs. 2, 4, and 5). Nutrients were apparently available in the water column throughout summer, indicating that, in spite of a relatively high primary production during that period, the supply of nutrients compensated for the biological uptake. Even though part of the measured chlorophyll *a* might correspond to resuspended benthic microalgae, most is likely of pelagic origin since the high turbidity limits benthic primary production to intertidal areas, which cover only 8 % of the Marsdiep basin (Compton et al. 2013). This suggests mineralization as an important responsible for nutrient replenishment in the water column since the high temperatures and availability of labile organic material promote a higher degradation rate during summer.

Internal Recycling and Sediment Release as a Source of Nutrients to the Water Column

The mixing diagrams (Fig. 3) support the idea of mineralization in the Wadden Sea as an important source of NH₄, PO₄, and Si to the water column, especially in spring and summer. As mentioned above, even though freshwater appeared to be the source of NO₃ and Si to the western Wadden Sea in winter, as also observed for the East Frisian Wadden Sea (Grunwald et al. 2010), the same did not apply to PO₄ and NH₄ likely due to an earlier phytoplankton bloom in Lake IJssel, as mentioned above. In addition, retention of NH₄ and PO₄ by sorption in the sediments may be more efficient in freshwater than in seawater, due to the reduced presence of competing ions (Caraco et al. 1989; Hawke et al. 1989). Instead, the winter availability of NH₄ and PO₄ in the Wadden Sea seemed to result from internal “production”, whether by slow degradation of refractory organic matter and/or desorption of nutrients stored over winter in the sediments (van Raaphorst et al. 1988; van Raaphorst and Kloosterhuis 1994; de Vries et al. 1998; Leote et al. 2013). The first phytoplankton bloom in the Marsdiep in March resulted in a high demand for NH₄, PO₄, and Si in both end-members and the Wadden Sea, with the concentrations of dissolved inorganic nutrients dropping to extremely low values after conversion to particulate organic material. However, the degradation of the dead phytoplankton cells can replenish the water column with dissolved nutrients (van Beusekom and de Jonge 2012), as evidenced by the increase in DON and DOP concentrations in April (Fig. 2).

This indicates an initial degradation of the organic material, with nutrients becoming available to heterotrophs or autotrophs after external oxidation or microbial degradation (Nausch and Nausch 2006; 2007). This drop in PO₄ and Si concentrations in spring, followed by an increase towards summer has been observed for other basins of the Wadden Sea, namely the East Frisian Wadden Sea, close to Spiekeroog Island (Grunwald et al. 2010; Beck and Brumsack 2012). For the List Tidal Basin, northern Wadden Sea, Loebl et al. (2007) and van Beusekom et al. (2009) have also reported an increase in the PO₄ concentrations in early summer, after the first bloom; however, the Si and NH₄ concentrations remained relatively low until autumn. Similarly, higher DOP concentrations during spring and summer, an opposite behaviour between DOP and PO₄, and a positive relationship between DOP and chlorophyll *a* have also been reported for the Dutch Wadden Sea and Sylt-Rømø area (van Beusekom and de Jonge 2012). In May, comparing the increase in the NH₄, PO₄, and Si concentrations in the water column with the mixing diagrams (Fig. 3), it becomes clear that neither Lake IJssel nor the North Sea were the main nutrient sources. Instead, this increase was only observed at intermediate salinities, suggesting that mineralization within the Wadden Sea, enhanced by the increased availability of fresh organic matter and higher temperatures, was the main nutrient source (van Beusekom et al. 2009). Silicate showed a less marked increase in the concentrations compared to NH₄ and PO₄ since the dissolution of biogenic Si requires an initial degradation of the organic coating of diatom frustules (Van Cappellen and Qiu 1997) and can be delayed by aggregation of the diatom cells (Loucaides 2009), unlike NH₄ and PO₄. The rapid availability of fresh organic material and consequent reduced oxygen availability in the sediments, as presented in Leote and Epping (2015) for this period, likely resulted in an increased demand for electron acceptors to be used in the oxidation of organic matter. Since NO₃ is the next most favourable oxidizer after oxygen, its consumption is expected to increase in this period, mainly via denitrification. Annual sediment denitrification rates of 110 mmol N m⁻² have been reported for the western Wadden Sea (Kieskamp et al. 1991), while a total N loss within the basin of 4 mmol N m⁻² days⁻¹ was reported by Philippart et al. (2000). This explains why NO₃ concentrations were independent of chlorophyll *a*, continuously decreasing from February until September, i.e. from the lowest to the highest temperature, with no increase in May–September, since consumption by denitrification prevented its accumulation in the water column. Other Wadden Sea regions, such as the East Frisian Wadden Sea also showed a later increase in NO₃ concentrations when compared to the other inorganic nutrients (Grunwald et al. 2010; Beck and Brumsack 2012).

The conclusions above match well with the previous findings of Leote and Epping (2015), who found higher oxygen

consumption in late spring. The same authors also found increasing sediment-water exchange rates for dissolved inorganic carbon between March and September. Since dissolved inorganic carbon is an end product of mineralization, its higher release rates further confirm a higher mineralization during this period, at least in the sediment, with release rates ranging between 0 (in winter) and $52 \text{ mmol m}^{-2} \text{ days}^{-1}$ (in late summer).

The shallowness of the western Wadden Sea suggests a major role of the sediments in nutrient regeneration with 70 % efficiency in the recycling of organic P compounds reported by van Raaphorst et al. (1988). In addition, resuspension, due to tidal currents and wind-driven waves, may enhance the release of nutrients, by disruption of the sediment-water interface and desorption from sediment particles due to changing equilibrium conditions (Leote et al. 2013).

Resuspension as an Enhancer of Mineralization

The TSM concentrations measured for the western Wadden Sea were slightly higher than those reported by van Beusekom (2005) and Reynhout (2002) in February, March, and November, when no major differences were observed during a tidal cycle (Figs. 4 and 6). However, in May and September, the concentrations were much higher, increasing to 205 mg l^{-1} after the maximum tidal current speeds and especially for the bottom samples (Fig. 6), suggesting resuspension of the deposited material (Postma 1961). A certain time lag was observed between the expected maximum current speed and the maximum TSM concentrations, similarly to Lunau et al. (2006). These increased TSM concentrations at higher current speeds have already been observed in the Wadden Sea and specifically in the Marsdiep basin (Hommerson et al. 2009; Nauw et al. 2014). The concentrations of chlorophyll *a* followed a similar pattern (Fig. 5), indicating that part of the resuspended material was fresh organic material, in agreement with the high organic C/total P ratio found throughout the year (Fig. 4). Even though bottom currents are potentially higher in winter when wind-induced mixing is stronger (Reynhout 2002), the higher concentrations of suspended material were observed in spring and in summer, as observed by Hommerson et al. (2009). This suggests that the higher TSM concentrations did not result from increased current speeds but might be attributed to the increased availability of easily resuspendable material, i.e. an observed fluffy layer of deposited fresh organic material instead of a well-defined interface of clay/sand particles (Pempkowiak et al. 2005). The resuspension of this organic rich material (Fig. 4) may have important implications in the regeneration of nutrients by increasing mineralization in the water column, due to exposure of a larger surface area of the organic matter to a well-oxygenated environment (Ståhlberg et al. 2006).

Nutrient Availability and Potential Relation with the Phytoplankton Community

Irradiance and nutrient (N, P, and Si) availability are the main controls of phytoplankton growth in the Wadden Sea, even though debate remains concerning the limiting factor (de Jonge et al. 1996; Philippart and Cadée 2000; Colijn and Cadée 2003; Colijn and van Beusekom 2005). Recent findings for the Marsdiep basin on the absence of trends in light conditions during the last 40 years (Philippart et al. 2013) and the occurrence of P limitation during spring blooms using various techniques (Ly et al. 2014) strongly suggest the structuring role of P on phytoplankton growth in this area for the recent years. Due to different nutritional needs, the proportional availability of nutrients will determine the species composition of the phytoplankton community. Based on Redfield stoichiometry, PO_4 was apparently the limiting nutrient during most of the year (Fig. 7) as found by Philippart et al. (2007) and Kuipers and van Noort (2008). A P-Si-N limitation was observed in the end of winter allowing the growth of diatoms (Ly et al. 2014), as evidenced by the depletion of Si observed in March. However, in spring, Si became the limiting nutrient, with concentrations often below $2 \text{ }\mu\text{M}$, a threshold value for dinoflagellate growth to take over diatoms (Egge and Aksnes 1992). The growth of flagellates and *Phaeocystis* spp. was then favoured since they do not require Si, as shown by Ly et al. (2014) for the Marsdiep basin in 2010 and for the List Tidal Basin (Loebl et al. 2007). In addition, *Phaeocystis* spp. can use DOP as a source of P (Schoemann et al. 2005; Verity et al. 2007), giving it an advantage in April, when NH_4 but no PO_4 was available (Fig. 2). In September, Si seemed to no longer be the limiting nutrient (P-Si-N or P-N-Si limitation) and the drop in Si concentrations in November suggested a secondary diatom bloom, consistent with the findings of Cadée (1986).

Sources of P to the Western Wadden Sea

Since P was identified as the limiting nutrient during most of the year, in agreement with recent literature (Philippart et al. 2007; Kuipers and van Noort 2008; Ly et al. 2014), we will only compare its different sources to the western Wadden Sea. The main source of PO_4 to the western Wadden Sea remains unclear, with the North Sea, Lake IJssel, and internal recycling as the main candidates (de Jonge 1990; van Raaphorst and van der Veer 1990; de Jonge et al. 1996; de Jonge 1997; Philippart et al. 2000; van Raaphorst and de Jonge 2004). Postma (1954) suggested that the high PO_4 concentrations in spring–summer in the study area result from the mineralization of particulate organic matter from the North Sea. In agreement, our results confirm the importance of mineralization within the Wadden Sea, with the concentrations of NH_4 , PO_4 , and Si increasing at intermediate salinities in May, while the lower concentrations




Fig. 7 Diagrams based on the Redfield-Brzezinski ratio showing the major nutrient (N, P, Si) limitation (Rocha et al. 2002) throughout the year and plotted separately for each seasonally sampled station. Plotting the DIN/DIP against the Si/DIN molar ratios in the water column and

comparing them with the Redfield-Brzezinski ratio allows the definition of “zones” of sequential nutrient limitation. The diagram below indicates those “zones” where, for example, in the lower left quadrant, Si is the main limiting nutrient, followed by N and P

for Lake IJssel and the North Sea (Fig. 3) make them improbable major sources of inorganic P. Besides, the discharge of freshwater was low during this period with usually <25 % of freshwater in the water column composition. A comparison between PO_4 release from the sediments and freshwater discharge from Den Helder and the Afsluitdijk sluices, which correspond to approximately 60 % of the discharged water into the western Wadden Sea (Ridderinkhof 1988; van Raaphorst and van der Veer 1990), showed a much higher contribution of sediment release to the water column loading, throughout the year and especially for May–November (Fig. 8). A similar low contribution of freshwater in terms of nutrient supply has also been found for the East Frisian Wadden Sea (Grunwald et al. 2010). In addition, the contribution of mineralization within the Wadden Sea is likely underestimated since it is based on a conservative estimate of sediment release using porewater profiles of nutrient concentrations (Leote and Epping 2015) and does not include




Fig. 8 Comparison between the contribution of freshwater (Rijkswaterstaat Waterbase (live.waterbase.nl), assuming discharge from Den Oever, Kornwerderzand, and Den Helder) and sediment release (Leote and Epping 2015) in terms of PO_4 loading to the western Wadden Sea

mineralization or desorption in the water column, potentially enhanced by resuspension. No bias in the fluxes due to microphytobenthos activity (Sundbäck et al. 1991) was expected, given the reduced light penetration depth for the Marsdiep basin. In addition, flux overestimation due to advection-driven fluxes is also not likely considering the relatively poor sorting of our stations, in spite of their high sand content, meaning that the sediment was well-compacted (Leote and Epping 2015). However, the contribution from submarine groundwater discharge is unknown for this area but has been reported for other regions of the Wadden Sea (e.g. East Frisian Wadden Sea) as having an impact on some nutrient concentrations in the water column (Riedel et al. 2010; Moore et al. 2011). Even though we have no data to assess its contribution, we expect it to be low when compared to the East Frisian Wadden Sea, considering the smaller intertidal area of the Marsdiep basin and the low hydraulic gradient generated by the land adjacent to the Marsdiep basin (Moore et al. 2011; Compton et al. 2013). Concluding, in spite of the past large contribution of freshwater discharge in terms of PO_4 loading (van Raaphorst and van der Veer 1990; van Raaphorst and de Jonge 2004), at present, internal recycling of organic material, likely of both autochthonous and allochthonous origin, seems to be the main source of PO_4 to the western Wadden Sea, largely fuelling primary production during spring and summer.

Conclusions

In this study, we presented updated dissolved and particulate nutrient information for the Marsdiep basin collected over full tidal cycles. Regarding seasonality, the concentrations of dissolved nutrients in the western Wadden Sea were closely linked to the growth of primary producers with a negative relationship between the concentrations of chlorophyll *a* and the major inorganic nutrients. Phosphorus limitation, according to the Redfield ratio, was observed during most of the year, even though Si was also limiting in late spring. Recycling of fresh organic material resulting from the first phytoplankton bloom in spring seemed to be the major source of inorganic nutrients in late spring–summer since the low concentrations measured for the North Sea and Lake IJssel made them unlikely sources for that period. A comparison between PO_4 loading to the Wadden Sea by freshwater discharge and sediment release revealed a much higher contribution of sediment release. In addition, sediment resuspension was observed in spring–summer, potentially further favouring sediment nutrient release. This is in contrast with most previous work, where freshwater discharge was seen as the main source of nutrients to the Marsdiep basin.

Acknowledgments This research was funded by the Portuguese Foundation for Science and Technology (FCT PhD grant SFRH/BD/38856/2007) to CL, the P REDUCE project (NWO grant 839.08.342) to LLM, and the IN PLACE project (NWO grant 839.08.210). The authors would like to gratefully acknowledge the crews from the RV *Navicula*, RV *Stern*, and the IN PLACE team for their support. A thank you note is also due to Jan van Ooijen, Karel Bakker, Sharyn Crayford, and Evaline van Weerlee for all nutrient analyses performed by the NIOZ Nutrient Lab. We also thank Bert Brinkman and two anonymous reviewers for their constructive comments and discussion.

References

- Aoyama M, Anstey C, Barwell-Clarke J, Baurand F, Becker S, et al. .2010. *2008 Inter-laboratory comparison study of a reference material for nutrients in seawater*. Technical Reports of the Meteorological Research Institute No 60, Japan. DOI: 10.11483/mritechrepo.60.
- Beck, M., and H.-J. Brumsack. 2012. Biogeochemical cycles in sediment and water column of the Wadden Sea: The example Spiekeroog Island in a regional context. *Ocean and Coastal Management* 68: 102–113.
- Beukema, J.J., G.C. Cadée, and R. Dekker. 2002. Zoobenthic biomass limited by phytoplankton abundance: Evidence from parallel changes in two long-term data series in the Wadden Sea. *Journal of Sea Research* 48: 111–125.
- Cadée, G.C. 1986. Recurrent and changing seasonal patterns in phytoplankton of the westernmost inlet of the Dutch Wadden Sea from 1969 to 1985. *Marine Biology* 93: 281–289.
- Cadée, G.C., and J. Hegeman. 1993. Persisting high levels of primary production at declining phosphate concentrations in the Dutch Coastal Area (Marsdiep). *Netherlands Journal of Sea Research* 31(2): 147–152.
- Cadée, G.C., and J. Hegeman. 2002. Phytoplankton in the Marsdiep at the end of the 20th century 30 years monitoring biomass, primary production, and Phaeocystis blooms. *Journal of Sea Research* 48: 97–110.
- Caraco, N.F., J.J. Cole, and G.E. Likens. 1989. Evidence for sulphate controlled phosphorus release from sediments of aquatic ecosystems. *Nature* 341: 316–317.
- Colijn, F., and G.C. Cadée. 2003. Is phytoplankton growth in the Wadden Sea light or nitrogen limited? *Journal of Sea Research* 49: 83–93.
- Colijn, F., and J.E.E. van Beusekom. 2005. Effects of eutrophication on phytoplankton productivity and growth in the Wadden Sea. In *The intertidal ecosystem: The value of Ireland's shores*, ed. J. Wilson, 56–68. Dublin: Royal Irish Academy.
- Compton, T.J., S. Holthuijsen, A. Koolhaas, A. Dekinga, J. ten Horn, J. Smith, Y. Galama, et al. 2013. Distinctly variable mudscapes: Distribution gradients of intertidal macrofauna across the Dutch Wadden Sea. *Journal of Sea Research* 82: 103–116. doi:10.1016/j.seares.2013.02.002.
- Crouzet P, Nixon S, Rees Y, Parr W, Laffon L, Bogestrand J, Kristensen P, et al. .1999. *Nutrients in European Ecosystems*. European Environment Agency.
- Dastgheib A .2007. *Long-term morphological modeling of Marsdiep Basin in the Dutch Wadden Sea*, The Netherlands. MSc thesis. Delft University, UNESCO-IHE
- de Jonge, V.N. 1997. High remaining productivity in the Dutch Western Wadden Sea despite decreasing nutrient inputs from riverine sources. *Marine Pollution Bulletin* 34(6): 427–436.
- de Jonge, V.N. 1990. Response of the Dutch Wadden Sea ecosystem to phosphorus discharges from the River Rhine. *Hydrobiologia* 195: 49–62.

- de Jonge, V.N., J.F. Bakker, and M. van Stralen. 1996. Recent changes in the contributions of river Rhine and North Sea to the eutrophication of the Western Dutch Wadden Sea. *Netherlands Journal of Aquatic Ecology* 30(1): 26–39.
- de Vries, I., R.N.M. Duin, J.C.H. Peeters, F.J. Los, M. Bokhorst, and R.W.P.M. Laane. 1998. Patterns and trends in nutrients and phytoplankton in Dutch Coastal Waters: Comparison of time-series analysis, ecological model simulation, and mesocosm experiments. *ICES Journal of Marine Science: Journal Du Conseil* 55(4): 620–634. doi:10.1006/jmsc.1998.0399.
- Egge, J.K., and J.D.L. Aksnes. 1992. Silicate as a regulating nutrient in phytoplankton competition. *Marine Ecology Progress Series* 833: 281–289.
- Grasshoff, K., M. Ehrhardt, and K. Kremling. 1999. *Methods of seawater analysis*, 3rd ed. Weinheim: Wiley-VCH.
- Grunwald, M., O. Dellwig, C. Kohlmeier, N. Kowalski, M. Beck, T.H. Badewien, S. Kotzur, G. Liebezeit, and H.-J. Brumsack. 2010. Nutrient dynamics in a back barrier tidal basin of the Southern North Sea: Time-series, model simulations, and budget estimates. *Journal of Sea Research* 64: 199–212.
- Hawke, D., P.D. Carpenter, and K.A. Hunter. 1989. Competitive adsorption of phosphate on goethite in marine electrolytes. *Environmental Science & Technology* 23: 187–191.
- Helder, W., and R.T.P. de Vries. 1979. An automatic phenol-hypochlorite method for the determination of ammonia in sea and brackish waters. *Netherlands Journal of Sea Research* 13(1): 154–160.
- Hommerson, A., S. Peter, M.R. Wernand, and J. de Boer. 2009. Spatial and temporal variability in bio-optical properties of the Wadden Sea. *Estuarine, Coastal and Shelf Science* 83: 360–370.
- Jeffrey, S.W., R.F.C. Mantoura, and S.W. Wright (eds.). 1997. *Phytoplankton pigments in oceanography: Guidelines to modern methods*, 661. Paris: Unesco Publishing.
- Kieskamp, W.M., L. Lohse, E. Epping, and W. Helder. 1991. Seasonal variation in denitrification rates and nitrous oxide fluxes in intertidal sediments of the Western Wadden Sea. *Marine Ecology Progress Series* 72: 145–151.
- Kuipers, B.R., and G.J. van Noort. 2008. Towards a natural Wadden Sea? *Journal of Sea Research* 60(1–2): 44–53.
- Leote, C., E.H.G. Epping, and P. Van Cappellen. 2013. Phosphate sorption from seawater solutions: Particle concentration effect. *Marine Chemistry* 148: 52–62.
- Leote, C., L. Mulder, and E. Epping. 2014. A budget of bioavailable inorganic phosphorus in the sediment for the Western Wadden Sea. *Journal of Sea Research* 87: 79–90.
- Leote, C., and E. Epping. 2015. Sediment-water exchange of nutrients in the Marsdiep Basin, Western Wadden Sea: Phosphorus limitation induced by a controlled release? *Continental Shelf Research* 92: 44–58.
- Loebl, M., F. Colijn, J.E.E. van Beusekom, J.G. Baretta-Bekker, C. Lancelot, C.J.M. Philippart, V. Rousseau, and K.H. Wiltshire. 2009. Recent patterns in potential phytoplankton limitation along the Northwest European continental coast. *Journal of Sea Research* 61: 34–43.
- Loebl, M., T. Dolch, and J.E.E. van Beusekom. 2007. Annual dynamics of pelagic primary production and respiration in a shallow coastal basin. *Journal of Sea Research* 58: 269–282.
- Loucaides S. 2009. *Dissolution of biogenic silica: Roles of pH, salinity, pressure, electrical charging and reverse weathering*. PhD Thesis. Utrecht University.
- Lunau, M., A. Lemke, O. Dellwig, and M. Simon. 2006. Physical and biogeochemical controls of microaggregate dynamics in a tidally affected coastal ecosystem. *Limnology and Oceanography* 51(2): 847–859.
- Ly, J., C.J.M. Philippart, and J.C. Kromkamp. 2014. Phosphorus limitation during a phytoplankton spring bloom in the Western Dutch Wadden Sea. *Journal of Sea Research* 88: 109–120.
- Moore, W.S., M. Beck, T. Riedel, M. Rutgers van der Loeff, O. Dellwig, T.J. Shaw, B. Schnetger, and H.-J. Brumsack. 2011. Radium-based pore water fluxes of silica, alkalinity, manganese, DOC, and uranium: A decade of studies in the German Wadden Sea. *Geochimica et Cosmochimica Acta* 75: 6535–6555.
- Murphy, J., and J.P. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Analytica Chimica Acta* 27: 31–36.
- Nausch, M., and G. Nausch. 2006. Bioavailability of dissolved organic phosphorus in the Baltic Sea. *Marine Ecology Progress Series* 321: 9–17.
- Nausch, M., and G. Nausch. 2007. Bioavailable dissolved organic phosphorus and phosphorus use by heterotrophic bacteria. *Aquatic Biology* 1: 151–160. doi:10.3354/ab00012.
- Nauw, J.J., L.M. Merckelbach, H. Ridderinkhof, and H.M. van Aken. 2014. Long-term ferry-based observations of the suspended sediment fluxes through the Marsdiep inlet using acoustic doppler current profilers. *Journal of Sea Research* 87: 17–29.
- Nijssen A, Spronk G. 2000. *De Bepaling van het Gehalte van Fosfor van in water Opgeloste Fosforverbindingen via U.V.-Destructie-continuous flow: Skalar Systeem*. RIKZ rapport nr: A100, rapport version 4.3
- Oost, A.P., and P.L. de Boer. 1994. Sedimentology and development of barrier islands, ebb-tidal deltas, inlets and backbarrier areas of the Dutch Wadden Sea. *Senckenbergiana Maritima* 24(1/6): 65–115.
- Pempkowiak, J., J. Beldowski, K. Pazdro, A. Staniszewski, A. Zaborska, T. Leipe, and K. Emeis. 2005. Factors influencing fluffy layer suspended matter (FLSM) properties in the Odra River-Pomeranian Bay-Arkona Deep System (Baltic Sea) as derived by principal components analysis (PCA), and cluster analysis (CA). *Hydrology and Earth System Sciences* 9: 67–80.
- Philippart, C.J.M., M.S. Salama, J.C. Kromkamp, H.J. van der Woerd, A.F. Zuur, and G.C. Cadée. 2013. Four decades of variability in turbidity in the Western Wadden Sea as derived from corrected Secchi disk readings. *Journal of Sea Research* 82: 67–79.
- Philippart, C.J.M., J.M. van Iperen, G.C. Cadée, and A.F. Zuur. 2010. Long-term field observations on seasonality in chlorophyll-a concentrations in a shallow coastal marine ecosystem, the Wadden Sea. *Estuaries and Coasts* 33: 286–294.
- Philippart, C.J.M., J.J. Beukema, G.C. Cadée, R. Dekker, P.W. Goedhart, J.M. van Iperen, M.F. Leopold, and P.M.J. Herman. 2007. Impacts of nutrient reduction on coastal communities. *Ecosystems*. doi:10.1007/s10021-006-9006-7.
- Philippart, C.J.M., G.C. Cadée, W. van Raaphorst, and R. Riegman. 2000. Long-term phytoplankton-nutrient interactions in a shallow coastal area: Algal community structure, nutrient budgets and denitrification potential. *Limnology and Oceanography* 45(1): 131–144.
- Philippart, C.J.M., and G.C. Cadée. 2000. Was total primary production in the Western Wadden Sea stimulated by nitrogen loading? *Helgoland Marine Research* 54: 55–62.
- Postma H. 1954. *Hydrography of the Dutch Wadden Sea*. PhD Thesis, University of Groningen.
- Postma, H. 1961. Transport and accumulation of suspended matter in the Dutch Wadden Sea. *Netherlands Journal of Sea Research* 1: 148–190.
- Reynhout EJM 2002. *The influence of the tide on the concentration of suspended matter in the Marsdiep*. MSc thesis, Amsterdam, The Netherlands: University of Amsterdam
- Ridderinkhof H, van Haren H, Eijgenraam F, Hillebrand T. 2002. Ferry observation on temperature, salinity and currents in the Marsdiep Inlet between the North Sea and Wadden Sea. In: *Operational Oceanography. Implementation at the European and Regional Scales*. Elsevier Oceanography Series 66: 139–148, Proceedings of the second international conference on EUROGOOS

- Ridderinkhof, H. 1988. Tidal and residual flows in the Western Dutch Wadden Sea I: Numerical model results. *Netherlands Journal of Sea Research* 22(1): 1–21.
- Riedel, T., K. Lettmann, M. Beck, and H.-J. Brumsack. 2010. Tidal variations in groundwater storage and associated discharge from an intertidal coastal aquifer. *Journal of Geophysical Research* 115: C04013.
- Rocha, C., H. Galvão, and A. Barbosa. 2002. Role of transient silicon limitation in the development of cyanobacteria blooms in the Guadiana Estuary, South-Western Iberia. *Marine Ecology Progress Series* 228: 35–45.
- Schreurs W, Nijssen A .2000. *De Bepaling van het Gehalte aan Stikstof van in Water Opgeloste Stikstofverbindingen via U.V.-Destructie-continue flow: Skalar-Systeem*. A. RIKZ rapport nr.: A089 version 4.2
- Schoemann, V., S. Becquevort, J. Stefels, V. Rousseau, and C. Lancelot. 2005. Phaeocystis blooms in the global ocean and their controlling mechanisms: A review. *Journal of Sea Research* 53: 43–66.
- Ståhlberg, C., D. Bastviken, B.H. Svensson, and L. Rahm. 2006. Mineralisation of organic matter in coastal sediments at different frequency and duration of resuspension. *Estuarine, Coastal and Shelf Science* 70: 317–325.
- Strickland, J.D.H., and T.R. Parsons. 1968. *A practical book of seawater analysis*. *Bulletin 167*. Canada: Fisheries Research Board of Canada.
- Sundbäck, K., V. Enoksson, W. Granéli, and K. Pettersson. 1991. Influence of Sublittoral microphytobenthos on the oxygen and nutrient flux between sediment and water: A laboratory continuous-flow study. *Marine Ecology Progress Series* 74: 263–279.
- Valderrama, J.C. 1981. The simultaneous analysis of total nitrogen and total phosphorus in natural waters. *Marine Chemistry* 10: 109–122.
- van Aken, H.M. 2008. Variability of the salinity in the Western Wadden Sea on tidal to centennial time scales. *Journal of Sea Research* 59(3): 121–132. doi:10.1016/j.seares.2007.11.001.
- van Beusekom, J.E.E., and V.N. de Jonge. 2012. Dissolved organic phosphorus: An indicator of organic matter turnover? *Estuarine, Coastal and Shelf Science* 108: 29–36.
- van Beusekom, J.E.E. 2005. A historic perspective on Wadden Sea eutrophication. *Helgoland Marine Research* 59: 45–54.
- van Beusekom, J.E.E., and V.N. de Jonge. 2002. Long term changes in Wadden Sea nutrient cycles: Importance of organic matter from the North Sea. *Hydrobiologia* 475(476): 185–194.
- van Beusekom, J.E.E., M. Loebl, and P. Martens. 2009. Distant riverine nutrient supply and local temperature drive the long-term phytoplankton development in a temperate Coastal Basin. *Journal of Sea Research* 61: 26–33.
- Van Cappellen P, Qiu L .1997. *Biogenic silica dissolution in sediments of the Southern Ocean*. II. Kinetics. *Deep-Sea Research II* 44 (5)
- van Raaphorst, W., and V.N. de Jonge. 2004. Reconstruction of the total N and P inputs from the IJsselmeer into the Western Wadden Sea between 1935 and 1998. *Journal of Sea Research* 51: 109–131.
- van Raaphorst W, de Jonge VN, Dijkhuizen D, Frederiks B .2000. *Natural background concentrations of phosphorus and nitrogen in the Dutch Wadden Sea*. Directoraat-Generaal Rijkswaterstaat.
- van Raaphorst, W., and H.T. Kloosterhuis. 1994. Phosphate sorption in superficial intertidal sediments. *Marine Chemistry* 48: 1–16.
- van Raaphorst, W., P. Ruardij, and A.G. Brinkman. 1988. The assessment of benthic phosphorus regeneration in an estuarine ecosystem model. *Netherlands Journal of Sea Research* 22(1): 23–36.
- van Raaphorst, W., and H.W. van der Veer. 1990. The phosphorus budget of the Marsdiep Tidal Basin (Dutch Wadden Sea) in the period 1950–1985: Importance of the exchange with the North Sea. *Hydrobiologia* 195: 21–38.
- Verardo, D.J., P.N. Froelich, and A. McIntyre. 1990. Determination of organic carbon and nitrogen in marine sediments using the Carlo Erba NA-1500 N analyzer. *Deep-Sea Research* 37(1): 157–165.
- Verity, P.G., C.P. Brussaard, J.C. Nejtgaard, M.A. van Leeuwe, C. Lancelot, and L.K. Medlin. 2007. Current understanding of Phaeocystis ecology and biogeochemistry, and perspectives for future research. *Biogeochemistry* 83: 311–330.