

Regulation of mitophagy in ischemic brain injury

Yang Yuan, Xiangnan Zhang, Yanrong Zheng, Zhong Chen

Department of Pharmacology, Key Laboratory of Medical Neurobiology (Ministry of Health of China), College of Pharmaceutical Sciences, Zhejiang University, Hangzhou 310058, China

Corresponding author: Zhong Chen. E-mail: chenzhong@zju.edu.cn

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2015

The selective degradation of damaged or excessive mitochondria by autophagy is termed mitophagy. Mitophagy is crucial for mitochondrial quality control and has been implicated in several neurodegenerative disorders as well as in ischemic brain injury. Emerging evidence suggested that the role of mitophagy in cerebral ischemia may depend on different pathological processes. In particular, a neuroprotective role of mitophagy has been proposed, and the regulation of mitophagy seems to be important in cell survival. For these reasons, extensive investigations aimed to profile the mitophagy process and its underlying molecular mechanisms have been executed in recent years. In this review, we summarize the current knowledge regarding the mitophagy process and its role in cerebral ischemia, and focus on the pathological events and molecules that regulate mitophagy in ischemic brain injury.

Keywords: mitophagy; ischemic brain injury; mitochondria

Introduction

One of the first autophagy investigations was performed by Keith R. Porter, who in 1962 discovered an increased number of vacuoles in rat liver cells treated with glucagon^[1]. Autophagy is a pivotal intracellular process for the bulk degradation and recycling of unnecessary or dysfunctional proteins and organelles by lysosomes^[2]. There are three forms: microautophagy, chaperone-mediated autophagy, and macroautophagy^[3]. Macroautophagy is the most prevalent form and here is referred to as autophagy. It is critical for maintaining cell functions and deciding cell fate, and its intracellular processes have been elucidated in recent decades^[3, 4]. Generally, double-membrane vesicles termed autophagosomes sequester target intracellular cargoes and then fuse with endosomes or directly with lysosomes for degradation. Autophagy is canonically activated under starvation^[5], hypoxia^[6] and intracellular stress^[7, 8]. The mammalian target of rapamycin (mTOR) kinase is a critical regulator of autophagy induction: activation of mTOR by AMPK-p53 signaling promotes it^[9], while inhibition of mTOR *via* PI3K-Akt signaling is

suppressive^[10]. The current knowledge of cell signaling on the regulation of autophagy has been summarized in several comprehensive review articles^[11–13].

Autophagy is widely implicated in central nervous system (CNS) disorders^[2]. A well-supported theory is that dysfunctional autophagy leads to aberrant accumulations of toxic protein aggregates in specific sites within the CNS and thus causes neurodegenerative diseases^[14]. Given that autophagy can promptly respond to insufficient nutrients and energy supply, it is not surprising that it is extensively observed in ischemic brain tissues^[6, 15, 16], where it was primarily postulated to promote cell death^[17], whereas lines of evidence indicated that it is required for neuronal survival during development and neurodegeneration, as well as in several models of cerebral ischemia^[15, 18, 19]. Unlike apoptosis and necrosis, which certainly contribute to ischemic brain injury, autophagy differs in that it may serve as a potential therapeutic target against ischemia^[20–22]. Therefore, although they remain controversial, these emerging data suggest that autophagy may play a vital role in neuroprotection against ischemic brain injury, but the underlying mechanisms await clarification.

Selective elimination of mitochondria by autophagy, termed mitophagy, may be a responsible mechanism. Mitophagy is highly evolutionarily conserved; many of its essential proteins in yeast have homologs in mammals^[23]. As the powerhouse of cells, mitochondria have a much shorter average lifespan of 10–25 days, compared with neurons, which are postmitotic cells with high energy demands. As a result, mitochondrial turnover by mitophagy is extremely important for neuronal survival. Dysfunctional mitophagy has been implicated in several human CNS diseases, including Alzheimer's disease^[24] and Parkinson's disease^[25]. Intriguingly, it has also been reported in ischemic brain injury by several independent groups^[15, 26–28]. By clearing dysfunctional mitochondria and promoting mitochondrial turnover, cell death resulting from ischemic insult is attenuated^[15]. Thus, mitophagy seems to be involved in ischemic brain injury, but questions remain. In this review, we summarize the current progresses on the regulation of mitophagy and its specific role in ischemic brain injury.

Mitophagy in Ischemic Brain Injury

Stroke is the third leading cause of death and disability worldwide. Cerebral ischemia makes up 65%–80% of total stroke events. Ischemic brain injury is generally

caused by the sudden block of blood supply, leading to decreased oxygen and glucose supply to the brain tissue. The pathological mechanisms include, but are not limited to, neuro-excitotoxicity, brain inflammation, microglial activation, and endothelial injury^[29–32]. In general, apoptosis is primarily found in striatal and cortical neurons after ischemic reperfusion, followed by necrotic neuronal death in the ischemic core area^[33]. Autophagy, however, is widely involved in ischemic brain injury^[15–18], and the autophagosome marker LC3-II is strongly elevated at the onset of ischemia^[15], indicating the robust activation of autophagy, which might represent a cell defense mechanism against ischemic insult. Further, within 6 h after reperfusion, electron microscopy has shown damaged mitochondria surrounded by autophagosomes in the ischemic penumbra^[15], and the numbers of mitochondria assessed by the constitutively-expressed mitochondrial proteins TOMM20 and COX411 are significantly reduced^[15], suggesting that mitochondria are degraded by autophagy. In primary cultured neurons with oxygen and glucose deprivation (OGD) treatment, mitophagy is observed within 1 h, revealed by the co-localization of Mito-DsRed-labeled mitochondria and GFP-LC3-labeled autophagosomes (Fig. 1). The data indicated that mitophagy may reach a maximum within 3 h of OGD. Mitophagy is mainly evident




Fig. 1. Time-lapse imaging of mitophagy. Primary cultured mouse cortical neurons expressing Mito-DsRed and GFP-LC3 were exposed to oxygen and glucose deprivation for 2 h, and imaged at 1 frame/30 min. Mitophagy was determined by the co-localization of Mito-DsRed-labeled mitochondria and GFP-LC3-labeled autophagosomes. The boxed regions were amplified 4 times in the bottom right boxes. Scale bar, 10 μ m.

in neurons undergoing brain ischemia^[15, 19, 34], although astrocytes have also been documented^[35, 36].

The contributions of autophagy and/or mitophagy to ischemic brain injury have been controversial for years^[6, 16]. In long-term ischemia, for example, permanent middle cerebral artery occlusion (MCAO) models, the excessive activation of autophagy has been proposed to be associated with increased injury by inducing autophagic cell death, and inhibition of autophagy by 3-MA is neuroprotective^[18]. Similarly, excessive induction of mitophagy leads to apoptosis in a neonatal ischemia/hypoxia model^[37], in which the restoration of circulation is deficient due to intravascular coagulation. However, in the reperfusion phase after ischemia, mitophagy plays beneficial roles^[15]. Inhibition of autophagy either pharmacologically or genetically rescues the ischemic brain^[15] and conversely, injection of rapamycin, an autophagy inducer, has the opposite effects in rats with transient MCAO^[28]. More interestingly, inhibition of autophagy immediately after reperfusion aggravates ischemic brain injury by reducing mitochondria-based apoptosis^[15], while delayed inhibition of autophagy by 3-MA after reperfusion (3 h later) does not significantly increase brain damage (unpublished data), indicating that the protective action of autophagy occurs only at the initial step. In another study, rapamycin treatment was shown to reduce necrotic cell death in neonatal hypoxia-ischemia-induced brain injury^[38]. Increasing evidence implies that autophagy may switch its contributions to ischemic stroke along with the pathological stages, but the reason is not clear. Energy restoration upon reperfusion^[39] is perhaps the most plausible since autophagy/mitophagy is an energy-consuming process. But how the energy state may exquisitely regulate autophagy is far from clear. Alternation of mitochondrial dynamics during ischemia and reperfusion is an intriguing speculation^[26, 40]. Upon reperfusion, mitochondria are fragmented by Drp1, a mitofission protein^[26]. It is hypothesized that mitofission allows mitochondria to be engulfed by autophagosomes much more easily. The distinct mechanisms of permanent and transient ischemia may be another factor. In the reperfusion phase after ischemia, apoptosis is prone to be induced otherwise the prolonged ischemia normally results in necrosis^[15, 41-44]. This may provide a chance for mitophagy to confer neuroprotection by clearing the damaged

mitochondria. Subsequently, cytochrome *c* released from mitochondria is reduced and thus mitochondria-dependent apoptosis can be inhibited^[15]. Although remaining enigmatic, the simplistic tagging of autophagy as 'bad' or 'good' in ischemic brains deserves to be reconsidered.

Pathological Events in Mitophagy Activation in Ischemic Brain

Although lines of evidence have implicated mitophagy in ischemic brains, the specific events initiating it and their integrated mechanisms remain largely unsolved. Recent work on endoplasmic reticulum (ER) stress, oxidative stress, and excitotoxicity may offer a solution to these questions.

Endoplasmic Reticulum Stress

ER stress has been reported to be widely involved in ischemic injury^[45, 46]. However, consensus has not been reached on its role in this process. ER stress leads to apoptosis^[47] and inhibition of the ER-associated pro-apoptotic factor CHOP confers remarkable neuroprotection in the mouse bilateral common carotid arteries occlusion model^[48]. Consistently, several reagents that relieve ER stress protect against brain ischemia-reperfusion injury^[49, 50], indicating that ER stress may be responsible for it. However, ER stress also triggers the unfolded protein response to restore ER functions by activating the ER transmembrane receptors PERK, IRE1, and ATF6^[51]. It is well-accepted that increased ER stress leads to autophagy, and correspondingly autophagy helps to suppress ER stress^[7, 52]. In the context of ischemic myocardial injury, treatment with ER stressors before operation induces autophagy^[53]. Furthermore, mild induction of ER stress selectively reinforces ischemia-reperfusion-induced mitophagy *via* the PERK-EIF2S1-ATF4-Parkin signaling pathway^[54]. Notably, the autophagic machinery does not accordingly increase with regard to the remarkable activation of mitophagy, as revealed by the numbers of autophagic vacuoles on mitochondria^[54]. Since the stimulation of ER stress has been reported to result in mitochondrial dysfunction and thus impaired mitophagy in lung epithelial cells^[55], it is of interest to explore why moderate ER stress reinforces mitophagy. The ER-mitochondria contact site (MAM) has recently been reported to be the location of autophagosome generation^[56]. It is likely that mitochondria have priority to

be targeted by autophagosomes generated in the MAM, and appropriate stimulation of ER stress may accelerate the mitophagy originated by the ER. The mechanisms underlying ER stress-induced mitophagy, however, are not fully understood. Further, the extent to which ER stress favors mitophagy, and how to target mitophagy for therapy through ER stress activation, need to be further addressed.

Reactive Oxygen Species

Reactive oxygen species (ROS) are byproducts of oxygen metabolism, extensively and promptly generated in ischemic tissues after reperfusion^[57-60]. ROS accumulation results in oxidative damage, opening of the mitochondrial permeability transition pore, and mitochondria-dependent cell death. Mitochondria produce most of the cellular ROS via the mitochondrial electron-transport chain^[61], and play an important role in the pathogenesis of ischemic brain injury. ROS initiate autophagosome formation^[62], and excessive ROS formation triggers bulk autophagy^[63], while conversely, autophagy helps to reduce ROS levels by removing damaged organelles and abnormal proteins^[64]. Interestingly, moderate ROS levels specifically induce mitophagy but not general autophagy to protect against cell death in a Drp1-dependent manner^[63], indicating the ability of ROS to stimulate mitophagy. In support of this, in another study, increased ROS levels were shown to lower the mitochondrial membrane potential and activate Parkin-dependent mitophagy. Conversely, overexpression of superoxide dismutase-2, a mitochondrial antioxidant protein, blocks mitophagy by clearing photosensitizer-induced mitochondrial ROS in HeLa cells^[62]. However, the association of ROS extension with the selective activation of mitophagy is still not clear.

In the context of ischemia, like ER stress, ROS may also be a double-edged sword with regard to cell survival (Fig. 2). On one hand, antioxidants that relieve ROS have been reported to help minimize ischemic injury^[65-67]; on the other hand, ROS-induced mitophagy protects against ischemic myocardial injury. In p53-deficient mice, a p53-TIGAR-mediated decrease in the ROS signal reduces Bnip3-dependent mitophagy and enlarges the infarcted myocardium, while these are reversed by injection of the antioxidant N-acetylcysteine^[68]. Therefore, it is likely that ROS are also responsible for brain ischemia-reperfusion-induced mitophagy, and serve as a potential target for clinical therapy.

Excitotoxicity

Excitotoxicity is evoked by excessive activation of neurotransmitters such as glutamate, which is the main neurotransmitter in the CNS. Excitatory neurotransmitters are widely involved in neurodegenerative diseases like Alzheimer's disease^[24], Parkinson's disease^[69], and stroke^[59, 60, 70]. Excitotoxicity is induced by high levels of glutamate caused by over-activation of N-methyl-D-aspartate receptors (NMDARs) and α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors in ischemic brain injury. NMDA antagonists such as memantine are neuroprotective both *in vivo* and *in vitro* in brain ischemia models^[70, 71]. Excitotoxicity has been postulated to be associated with autophagy/mitophagy in mammals. In rats treated with the NMDAR agonist quinolinic acid, excitotoxicity is induced and it aggravates the neuronal injury by p53-activated autophagy^[9]. A more recent study indicated that glutamate exposure results in the translocation of Parkin to mitochondria in a NMDAR-dependent manner, but is not sufficient to activate mitophagy. However, when co-treated with the antioxidant N-acetylcysteine, mitophagy is promoted^[72], indicating that excitotoxicity can induce autophagy, but is insufficient to activate it alone. Therefore, excitotoxicity may not always be deleterious, since it facilitates the translocation of Parkin to mitochondria, which favors mitophagy, especially in the complex pathological process of ischemic brain injury.

With the progress made so far, multiple pathways could be involved in mitophagy in ischemic brain injury (Fig. 2). And more pathological events might be involved in mitophagy activation in the ischemic brain that need to be explored. How mitophagy occurs and how to promote it require further studies.

PINK1- and Parkin-Mediated Mitophagy

PINK1 (PTEN-induced kinase 1)- and Parkin-mediated mitophagy is perhaps the most extensively-studied mechanism underlying mitophagy. Both having loss-of-function mutations in familial Parkinson's disease, PINK1 and Parkin physically interact and work together in the same pathway in *Drosophila*^[73]. In dysfunctional mitochondria, due to depolarization of the mitochondrial membrane, PINK1 is concentrated in the outer mitochondrial membrane, and then Parkin, an E3 ubiquitin




Fig. 2. Regulation of mitophagy in ischemic brain injury. Neuronal mitochondria are damaged by ER stress, the activation of oxidative stress, and/or excitotoxicity upon brain ischemia. Subsequently, the release of mitochondrial proteins such as cytochrome *c* ultimately leads to apoptotic cell death. However, on the other hand, mild ER stress, ROS generation, and excitotoxicity trigger the degradation of damaged mitochondria by mitophagy and therefore protect against ischemic brain injury. PINK1/Parkin, Bnip3, Nix, and Fundc1 are critical molecules responsible for mitophagy.

ligase, is phosphorylated by PINK1, and is further recruited to the damaged mitochondria^[74] (for review, see ^[75]).

Lines of evidence have suggested that PINK1-Parkin are involved in ischemic brain injury. Parkin is markedly down-regulated after reperfusion for 3 h in a mouse model of transient MCAO^[39]. This is temporally consistent with the peak of autophagic flux^[15], implying that Parkin-labeled mitochondria can be eliminated by autophagy. Besides, ubiquitinated proteins are also present in the same model^[76], regardless of whether Parkin is responsible for the ubiquitination or not, and this is now under discussion. Parkin is known to ubiquitinate several important substrates in cardiac injury^[77], indicating that it may participate in ischemic brain injury by activating mitophagy. In ischemic preconditioning (IPC) of the rat heart, Parkin is

translocated to mitochondria, and its ablation abolishes the cardioprotective effects of IPC^[27], suggesting that Parkin-mediated mitophagy is required in this process. In support, we also found translocation of Parkin to mitochondria in primary cultured neurons exposed to OGD-reperfusion, and knockdown of Parkin impairs OGD-reperfusion-induced mitophagy^[54]. Interestingly, the translocation of Parkin to mitochondria is accelerated by moderate activation of ER stress, ROS, and excitotoxicity as noted above, indicating that multiple stimulations may trigger Parkin-mediated mitophagy, which, from another perspective, ensures the integrity of execution of Parkin-mediated mitophagy in ischemic brain injury. Above all, Parkin-dependent mitophagy could be a target for ischemic brain injury therapy.

Bnip3- and Nix/Bnip3L-Mediated Mitophagy

Both Bnip3 and Nix (Bnip3L) are BH3-only proteins implicated in apoptosis and autophagy^[78-80]. Bnip3 has been reported to participate in hypoxia-induced mitophagy in rat heart. Nix was first found to be essential for the induction of mitophagy during the maturation of red blood cells^[79]. Both Bnip3 and Nix are primarily localized on the outer mitochondrial membrane, and share a similar mechanism to induce mitophagy (for reviews, see^[80, 81]). The difference between the two proteins during the induction of mitophagy is of interest. Nix up-regulation cannot compensate for the loss of Bnip3-induced mitophagy^[37], indicating clear differences between Bnip3- and Nix-dependent mitophagy. Bnip3 exclusively activates excessive mitophagy by interacting with LC3 in a mouse model of neonatal ischemia/hypoxia, and loss of Bnip3 significantly decreases mitophagy and reduces neuronal apoptosis. However, Nix has been proposed to function only under physiological conditions^[37]. Nevertheless, a recent investigation found that Nix overexpression promotes carbonylcyanide-3-chlorophenylhydrazone (CCCP)-induced mitophagy in human embryonic kidney cells^[82], suggesting that Nix also participates in stress-induced mitophagy. A possible hypothesis is that Nix may act as a substrate of Parkin, through which it senses damage to mitochondria^[82]. Nevertheless, whether Nix-mediated mitophagy is involved in ischemic brain injury needs further investigation.

FUNDC1-Mediated Mitophagy

FUNDC1, which is located on the mitochondrial outer membrane, contains three transmembrane domains and has been newly identified as the mitophagy receptor in hypoxia^[83]. FUNDC1 directly binds to LC3 *via* its conserved LIR motif. Both knockdown of FUNDC1 and mutation of the LIR motif inhibit mitophagy^[83]. This binding is strengthened under hypoxic conditions, and is much stronger than that of Nix, suggesting that FUNDC1 has important implications for ischemia. Further, FUNDC1 can be phosphorylated by ULK1 at serine 17, which is critical for mitophagy induction under hypoxia. In contrast, FUNDC1 promotes the recruitment of ULK1 to damaged mitochondria^[84]. Nonetheless, the role of FUNDC1 in brain ischemia has not been discussed.

Interestingly, increasing numbers of studies have indicated that these mitophagy pathways work cooperatively in pathological processes to ensure the effectiveness of mitophagy, especially in mammals. For example, Nix initiates autophagy and promotes the CCCP-induced translocation of Parkin^[85]. In addition, Nix has very recently been reported to be the substrate of Parkin in PD^[82], suggesting that Nix functions downstream of Parkin. Besides, mitophagy is nevertheless induced in HeLa cells that lack the Parkin protein^[82]. Finally, Bnip3, Nix, and FUNDC1 have all been reported to be involved in hypoxia-induced mitophagy, suggesting that these proteins work together in specific mitochondrial diseases. However, how these pathways cooperate and which occupies a dominant role in ischemic brain injury need to be addressed.

Proteins That Potentially Regulate Mitophagy in Ischemic Brain Injury

Many mitophagy-related proteins have been implicated in ischemic brain injury (Table 1). However, whether they all participate in the process by regulating brain injury-induced mitophagy remains unclear. We summarize the features of several frequently-reported proteins below.

Beclin1

Beclin1 is an autophagy regulator and is implicated in mitophagy^[86]. In myocardial ischemia, autophagy is activated by an AMPK-dependent mechanism, but in the reperfusion phase, Beclin1 is required^[87]. Knockdown of Beclin1 by RNA interference protects against cerebral ischemic injury in rats by activating autophagy^[88]. In addition, Beclin1 interacts with PINK1^[86], and turns on autophagic flux. Nevertheless, mitophagy can also be triggered in a Beclin1-independent pathway with the parkinsonian neurotoxin MPP⁺ treatment of neurons^[89], suggesting that Beclin1 may not be as important in ischemic brain injury-induced mitophagy as in myocardial ischemia.

VDAC1

VDAC1 (voltage-dependent anion channel 1) is a poly-ubiquitination substrate of Parkin^[90]. It is located on mitochondrial outer membrane, and has recently been reported to promote mitophagy and protect neurons in subarachnoid hemorrhage^[34]. While VDAC1 is required for

Table 1. Mitophagy-related proteins involved in ischemic brain injury

Protein	Location	Function	Ref
SQSTM1	Cytosol	Autophagy receptor, ubiquitination substrate of Parkin	[98, 99]
ALOX15	Cytosol	Key enzyme in mitophagy induction in reticulocytes	[100]
CDC37	Cytosol	Co-chaperone of HSP90	[101]
P38	Cytosol	Phosphorylates LC3	[102, 103]
HSP90	Cytosol	Regulates ULK1- and ATG13-mediated mitophagy	[101]
TNF α	Cytosol	Induces mitophagy in mouse macrophages	[104]
RIPK2	Cytosol	Regulates mitophagy by phosphorylating ULK1	[105]
PINK1	Mitochondria or Cytosol	Involved in mitophagy by promoting translocation of Parkin	[75, 106]
Parkin	Mitochondria or Cytosol	E3 Ub ligase, ubiquitylates OMM proteins to promote mitophagy	[54, 75]
FUNDC1	Mitochondria	Receptor for hypoxia-induced mitophagy	[83]
VDAC1	Mitochondria	Ubiquitination substrate of Parkin	[34]
Drp1	Mitochondria	Required in mitochondrial division	[97]
Mitofusin1	Mitochondria	Mediates mitochondrial fusion	[107]
Mitofusin2	Mitochondria	Mediates mitochondrial fusion	[107]
PARL	Mitochondria	Prevents release of mitochondrial cytochrome C	[108, 109]
Beclin1	Mitochondria	Induces autophagy	[86, 88]
Bcl-2	Mitochondria	Regulates PINK1-Parkin-mediated mitophagy	[110]
Bcl-2L1	Mitochondria	Suppresses FUNDC1-mediated mitophagy	[111]
TRAF2	Mitochondria	E3 ubiquitin ligase	[112]
Nix	Mitochondria and ER	Mitophagy receptor, interacts with LC3 and GABARAP	[85, 113]
Bnip3	Mitochondria and ER	Mitophagy receptor, interacts with LC3	[78, 80]
HDAC6	Nucleus	Regulates Parkin-induced mitophagy	[114]
HMGB1	Nucleus	Regulates mitochondrial function and morphology	[115]
HSP27	Nucleus	Required for mitochondrial quality control	[116]

GABARAP, GABA receptor-associated protein.

Parkin-dependent mitophagy^[90], whether it is dispensable for the induction of mitophagy is still under debate, since it cannot fully account for the mitochondrial K63-linked ubiquitin immunoreactivity after mitochondria depolarization^[91].

Drp1

Mitochondrial dynamics and mitophagy are closely related (see reviews^[92, 93]). Mitochondrial fission can produce an impaired unit that undergoes autophagic elimination^[94]. Dynamin-related protein 1 (Drp1) is required for mitochondrial fission, and it controls the integrity of mitochondrial structure^[95]. By interacting with Parkin, Drp1 is ubiquitylated and degraded by proteasomes, leading to mitofusion that blocks mitochondria from removal by

autophagy^[96]. Parkin and Drp1 may work synergistically, and interestingly when Drp1 is absent, Parkin becomes much more critical^[97], suggesting that Parkin may help to initiate a compensatory pathway. Further, selective inhibition of Drp1 by mdivi-1 prevents mitochondrial division and mitophagy in brain ischemia-reperfusion, as well as exacerbating brain infarct volume^[15].

Clinical Advances by Targeting Mitophagy

Molecular targets of autophagy have been identified for the discovery of inhibitors or enhancers (see review^[117]), and promising patents are emerging^[117]. For example, by targeting a functional region of Beclin1, Tat-Beclin1 peptide

interacts with GABP-1, a negative regulator of autophagy, to induce protective autophagy^[118]. Because of the crucial role of mitophagy in promoting cell survival, mitophagy has been considered as a clinical target for conquering diverse diseases^[64, 119]. However, the specific selectivity of mitophagy, i.e., that specifically targets damaged mitochondria with autophagosomes, makes it much harder to take advantage of mitophagy for clinical applications. However, several mitophagy-related proteins, such as Parkin and Beclin1, have been proposed to be beneficial targets for ischemic brain injury treatment^[54, 88]. Besides, the pathologic processes of ischemic brain injury have become increasingly clearer, and drugs that target mitophagy could be possible in the near future.

Conclusions and Future Directions

Looking deep into mitophagy helps to better understand the pathology of ischemic brain injury. The molecular regulation of mitophagy has achieved great progress, but many questions remain to be solved, for example, the role(s) of mitophagy in different models of brain ischemia, how mitophagy is initiated after ischemia-reperfusion, the detailed underlying mechanisms, and how to take advantage of mitophagy for clinical therapy. These questions still need further investigations in the future.

ACKNOWLEDGEMENTS

This review was supported by the National Natural Science Foundation of China (81373393 and 81273506), the Zhejiang Provincial Natural Science Foundation (LR15H310001), the Program for Zhejiang Provincial Leading Team of S&T Innovation Team (2011R50014), and the Fundamental Research Funds for the Central Universities, China (2015FZA7018). We are grateful to the Imaging Facilities, Zhejiang University School of Medicine, for help with confocal microscopy.

Received date: 2015-04-30; Accepted date: 2015-07-09

REFERENCES

- [1] Ashford TP, Porter KR. Cytoplasmic components in hepatic cell lysosomes. *J Cell Biol* 1962, 12: 198–202.
- [2] Rajawat YS, Bossis I. Autophagy in aging and in neurodegenerative disorders. *Hormones (Athens)* 2008, 7: 46–61.
- [3] Mizushima N. Autophagy: process and function. *Genes Dev* 2007, 21: 2861–2873.
- [4] Bamber BA, Rowland AM. Shaping cellular form and function by autophagy. *Autophagy* 2006, 2: 247–249.
- [5] Kuma A, Hatano M, Matsui M, Yamamoto A, Nakaya H, Yoshimori T, *et al.* The role of autophagy during the early neonatal starvation period. *Nature* 2004, 432: 1032–1036.
- [6] Balduini W, Carloni S, Buonocore G. Autophagy in hypoxia-ischemia induced brain injury: evidence and speculations. *Autophagy* 2009, 5: 221–223.
- [7] Yorimitsu T, Nair U, Yang Z, Klionsky DJ. Endoplasmic reticulum stress triggers autophagy. *J Biol Chem* 2006, 281: 30299–30304.
- [8] Ding Z, Liu S, Wang X, Dai Y, Khaidakov M, Romeo F, *et al.* LOX-1, oxidant stress, mtDNA damage, autophagy, and immune response in atherosclerosis. *Can J Physiol Pharmacol* 2014, 92: 524–530.
- [9] Wang Y, Dong XX, Cao Y, Liang ZQ, Han R, Wu JC, *et al.* p53 induction contributes to excitotoxic neuronal death in rat striatum through apoptotic and autophagic mechanisms. *Eur J Neurosci* 2009, 30: 2258–2270.
- [10] Heras-Sandoval D, Perez-Rojas JM, Hernandez-Damian J, Pedraza-Chaverri J. The role of PI3K/AKT/mTOR pathway in the modulation of autophagy and the clearance of protein aggregates in neurodegeneration. *Cell Signal* 2014, 26: 2694–2701.
- [11] Kim YC, Guan KL. mTOR: a pharmacologic target for autophagy regulation. *J Clin Invest* 2015, 125: 25–32.
- [12] Hale AN, Ledbetter DJ, Gawriluk TR, Rucker EB, 3rd. Autophagy: regulation and role in development. *Autophagy* 2013, 9: 951–972.
- [13] Russell RC, Yuan HX, Guan KL. Autophagy regulation by nutrient signaling. *Cell Res* 2014, 24: 42–57.
- [14] Ross CA, Poirier MA. Protein aggregation and neurodegenerative disease. *Nat Med* 2004, 10 Suppl: S10–17.
- [15] Zhang X, Yan H, Yuan Y, Gao J, Shen Z, Cheng Y, *et al.* Cerebral ischemia-reperfusion-induced autophagy protects against neuronal injury by mitochondrial clearance. *Autophagy* 2013, 9: 1321–1333.
- [16] Xu F, Gu JH, Qin ZH. Neuronal autophagy in cerebral ischemia. *Neurosci Bull* 2012, 28: 658–666.
- [17] Wen YD, Sheng R, Zhang LS, Han R, Zhang X, Zhang XD, *et al.* Neuronal injury in rat model of permanent focal cerebral ischemia is associated with activation of autophagic and lysosomal pathways. *Autophagy* 2008, 4: 762–769.
- [18] Sheng R, Zhang LS, Han R, Liu XQ, Gao B, Qin ZH. Autophagy activation is associated with neuroprotection in a rat model of focal cerebral ischemic preconditioning. *Autophagy* 2010, 6: 482–494.
- [19] Yan H, Zhang X, Hu W, Ma J, Hou W, Zhang X, *et al.* Histamine H3 receptors aggravate cerebral ischaemic injury

- by histamine-independent mechanisms. *Nat Commun* 2014, 5: 3334.
- [20] Gabryel B, Kost A, Kasprowska D. Neuronal autophagy in cerebral ischemia--a potential target for neuroprotective strategies? *Pharmacol Rep* 2012, 64: 1–15.
- [21] Puyal J, Vaslin A, Mottier V, Clarke PG. Postischemic treatment of neonatal cerebral ischemia should target autophagy. *Ann Neurol* 2009, 66: 378–389.
- [22] Wei K, Wang P, Miao CY. A double-edged sword with therapeutic potential: an updated role of autophagy in ischemic cerebral injury. *CNS Neurosci Ther* 2012, 18: 879–886.
- [23] Liu L, Sakakibara K, Chen Q, Okamoto K. Receptor-mediated mitophagy in yeast and mammalian systems. *Cell Res* 2014, 24: 787–795.
- [24] Santos RX, Correia SC, Wang X, Perry G, Smith MA, Moreira PI, *et al.* A synergistic dysfunction of mitochondrial fission/fusion dynamics and mitophagy in Alzheimer's disease. *J Alzheimers Dis* 2010, 20 Suppl 2: S401–412.
- [25] Vives-Bauza C, Przedborski S. Mitophagy: the latest problem for Parkinson's disease. *Trends Mol Med* 2011, 17: 158–165.
- [26] Zuo W, Zhang S, Xia CY, Guo XF, He WB, Chen NH. Mitochondria autophagy is induced after hypoxic/ischemic stress in a Drp1 dependent manner: the role of inhibition of Drp1 in ischemic brain damage. *Neuropharmacology* 2014, 86: 103–115.
- [27] Huang C, Andres AM, Ratliff EP, Hernandez G, Lee P, Gottlieb RA. Preconditioning involves selective mitophagy mediated by Parkin and p62/SQSTM1. *PLoS One* 2011, 6: e20975.
- [28] Li Q, Zhang T, Wang J, Zhang Z, Zhai Y, Yang GY, *et al.* Rapamycin attenuates mitochondrial dysfunction via activation of mitophagy in experimental ischemic stroke. *Biochem Biophys Res Commun* 2014, 444: 182–188.
- [29] Kariman K. Mechanism of cell damage in brain ischemia: a hypothesis. *Life Sci* 1985, 37: 71–73.
- [30] Umemura A, Nagai H, Mabe H. Biochemistry of brain ischemia--mechanism of delayed neuronal death. *Nihon Rinsho* 1993, 51 Suppl: 405–412.
- [31] Hu W, Xu L, Pan J, Zheng X, Chen Z. Effect of cerebral ischemia on brain mast cells in rats. *Brain Res* 2004, 1019: 275–280.
- [32] Guo ZH, Li F, Wang WZ. The mechanisms of brain ischemic insult and potential protective interventions. *Neurosci Bull* 2009, 25: 139–152.
- [33] Raichle ME. The pathophysiology of brain ischemia. *Ann Neurol* 1983, 13: 2–10.
- [34] Li J, Lu J, Mi Y, Shi Z, Chen C, Riley J, *et al.* Voltage-dependent anion channels (VDACs) promote mitophagy to protect neuron from death in an early brain injury following a subarachnoid hemorrhage in rats. *Brain Res* 2014, 1573: 74–83.
- [35] Motori E, Puyal J, Toni N, Ghanem A, Angeloni C, Malaguti M, *et al.* Inflammation-induced alteration of astrocyte mitochondrial dynamics requires autophagy for mitochondrial network maintenance. *Cell Metab* 2013, 18: 844–859.
- [36] Qin AP, Liu CF, Qin YY, Hong LZ, Xu M, Yang L, *et al.* Autophagy was activated in injured astrocytes and mildly decreased cell survival following glucose and oxygen deprivation and focal cerebral ischemia. *Autophagy* 2010, 6: 738–753.
- [37] Shi RY, Zhu SH, Li V, Gibson SB, Xu XS, Kong JM. BNIP3 interacting with LC3 triggers excessive mitophagy in delayed neuronal death in stroke. *CNS Neurosci Ther* 2014, 20: 1045–1055.
- [38] Carloni S, Buonocore G, Balduini W. Protective role of autophagy in neonatal hypoxia-ischemia induced brain injury. *Neurobiol Dis* 2008, 32: 329–339.
- [39] Mengesdorf T, Jensen PH, Mies G, Aufenberg C, Paschen W. Down-regulation of parkin protein in transient focal cerebral ischemia: A link between stroke and degenerative disease? *Proc Natl Acad Sci U S A* 2002, 99: 15042–15047.
- [40] Campello S, Strappazon F, Cecconi F. Mitochondrial dismissal in mammals, from protein degradation to mitophagy. *Biochim Biophys Acta* 2014, 1837: 451–460.
- [41] Miyamoto O, Auer RN. Hypoxia, hyperoxia, ischemia, and brain necrosis. *Neurology* 2000, 54: 362–371.
- [42] Li H, Liu X, Zhu Y, Liu Y, Wang Y. Magnolol derivative 002C-3 protects brain against ischemia-reperfusion injury via inhibiting apoptosis and autophagy. *Neurosci Lett* 2015, 588: 178–183.
- [43] Liu X, Wang M, Chen H, Guo Y, Ma F, Shi F, *et al.* Hypothermia protects the brain from transient global ischemia/reperfusion by attenuating endoplasmic reticulum response-induced apoptosis through CHOP. *PLoS One* 2013, 8: e53431.
- [44] Chen CH, Jiang Z, Yan JH, Yang L, Wang K, Chen YY, *et al.* The involvement of programmed cell death 5 (PDCD5) in the regulation of apoptosis in cerebral ischemia/reperfusion injury. *CNS Neurosci Ther* 2013, 19: 566–576.
- [45] Nakka VP, Gusain A, Raghur R. Endoplasmic reticulum stress plays critical role in brain damage after cerebral ischemia/reperfusion in rats. *Neurotox Res* 2010, 17: 189–202.
- [46] Sanderson TH, Gallaway M, Kumar R. Unfolding the unfolded protein response: unique insights into brain ischemia. *Int J Mol Sci* 2015, 16: 7133–7142.
- [47] Szegezdi E, Duffy A, O'Mahoney ME, Logue SE, Mylotte LA, O'Brien T, *et al.* ER stress contributes to ischemia-induced cardiomyocyte apoptosis. *Biochem Biophys Res Commun*

- 2006, 349: 1406–1411.
- [48] Tajiri S, Oyadomari S, Yano S, Morioka M, Gotoh T, Hamada JI, *et al.* Ischemia-induced neuronal cell death is mediated by the endoplasmic reticulum stress pathway involving CHOP. *Cell Death Differ* 2004, 11: 403–415.
- [49] Sokka AL, Putkonen N, Mudo G, Pryazhnikov E, Reijonen S, Khiroug L, *et al.* Endoplasmic reticulum stress inhibition protects against excitotoxic neuronal injury in the rat brain. *J Neurosci* 2007, 27: 901–908.
- [50] Begum G, Kintner D, Liu Y, Cramer SW, Sun D. DHA inhibits ER Ca²⁺ release and ER stress in astrocytes following in vitro ischemia. *J Neurochem* 2012, 120: 622–630.
- [51] Roussel BD, Kruppa AJ, Miranda E, Crowther DC, Lomas DA, Marciniak SJ. Endoplasmic reticulum dysfunction in neurological disease. *Lancet Neurol* 2013, 12: 105–118.
- [52] Sheng R, Liu XQ, Zhang LS, Gao B, Han R, Wu YQ, *et al.* Autophagy regulates endoplasmic reticulum stress in ischemic preconditioning. *Autophagy* 2012, 8: 310–325.
- [53] Petrovski G, Das S, Juhasz B, Kertesz A, Tosaki A, Das DK. Cardioprotection by endoplasmic reticulum stress-induced autophagy. *Antioxid Redox Signal* 2011, 14: 2191–2200.
- [54] Zhang X, Yuan Y, Jiang L, Zhang J, Gao J, Shen Z, *et al.* Endoplasmic reticulum stress induced by tunicamycin and thapsigargin protects against transient ischemic brain injury: Involvement of PARK2-dependent mitophagy. *Autophagy* 2014, 10: 1801–1813.
- [55] Bueno M, Lai YC, Romero Y, Brands J, St Croix CM, Kamga C, *et al.* PINK1 deficiency impairs mitochondrial homeostasis and promotes lung fibrosis. *J Clin Invest* 2015, 125: 521–538.
- [56] Hamasaki M, Furuta N, Matsuda A, Nezu A, Yamamoto A, Fujita N, *et al.* Autophagosomes form at ER-mitochondria contact sites. *Nature* 2013, 495: 389–393.
- [57] Takahashi S. Astroglial protective mechanisms against ROS under brain ischemia. *Rinsho Shinkeigaku* 2011, 51: 1032–1035.
- [58] Sun M, Li M, Huang Q, Han F, Gu JH, Xie J, *et al.* Ischemia/reperfusion-induced upregulation of TIGAR in brain is mediated by SP1 and modulated by ROS and hormones involved in glucose metabolism. *Neurochem Int* 2015, 80: 99–109.
- [59] Kalogeris T, Bao Y, Korhuis RJ. Mitochondrial reactive oxygen species: a double edged sword in ischemia/reperfusion vs preconditioning. *Redox Biol* 2014, 2: 702–714.
- [60] Shen Y, He P, Fan YY, Zhang JX, Yan HJ, Hu WW, *et al.* Carnosine protects against permanent cerebral ischemia in histidine decarboxylase knockout mice by reducing glutamate excitotoxicity. *Free Radic Biol Med* 2010, 48: 727–735.
- [61] Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J* 2009, 417: 1–13.
- [62] Wang Y, Nartiss Y, Steipe B, McQuibban GA, Kim PK. ROS-induced mitochondrial depolarization initiates PARK2/PARKIN-dependent mitochondrial degradation by autophagy. *Autophagy* 2012, 8: 1462–1476.
- [63] Frank M, Duvezin-Caubet S, Koob S, Occhipinti A, Jagasia R, Petcherski A, *et al.* Mitophagy is triggered by mild oxidative stress in a mitochondrial fission dependent manner. *Biochim Biophys Acta* 2012, 1823: 2297–2310.
- [64] Scherz-Shouval R, Elazar Z. Regulation of autophagy by ROS: physiology and pathology. *Trends Biochem Sci* 2011, 36: 30–38.
- [65] Yoshida S, Abe K, Busto R, Watson BD, Kogure K, Ginsberg MD. Influence of transient ischemia on lipid-soluble antioxidants, free fatty acids and energy metabolites in rat brain. *Brain Res* 1982, 245: 307–316.
- [66] Sheng H, Enghild JJ, Bowler R, Patel M, Batinic-Haberle I, Calvi CL, *et al.* Effects of metalloporphyrin catalytic antioxidants in experimental brain ischemia. *Free Radic Biol Med* 2002, 33: 947–961.
- [67] Shen WH, Zhang CY, Zhang GY. Antioxidants attenuate reperfusion injury after global brain ischemia through inhibiting nuclear factor-kappa B activity in rats. *Acta Pharmacol Sin* 2003, 24: 1125–1130.
- [68] Hoshino A, Matoba S, Iwai-Kanai E, Nakamura H, Kimata M, Nakaoka M, *et al.* p53-TIGAR axis attenuates mitophagy to exacerbate cardiac damage after ischemia. *J Mol Cell Cardiol* 2012, 52: 175–184.
- [69] Xu XM, Moller SG. ROS removal by DJ-1: Arabidopsis as a new model to understand Parkinson's Disease. *Plant Signal Behav* 2010, 5: 1034–1036.
- [70] Shen Y, Hu WW, Fan YY, Dai HB, Fu QL, Wei EQ, *et al.* Carnosine protects against NMDA-induced neurotoxicity in differentiated rat PC12 cells through carnosine-histidine-histamine pathway and H(1)/H(3) receptors. *Biochem Pharmacol* 2007, 73: 709–717.
- [71] Molinuevo JL, Llado A, Rami L. Memantine: targeting glutamate excitotoxicity in Alzheimer's disease and other dementias. *Am J Alzheimers Dis Other Demen* 2005, 20: 77–85.
- [72] Van Laar VS, Roy N, Liu A, Rajprohat S, Arnold B, Dukes AA, *et al.* Glutamate excitotoxicity in neurons triggers mitochondrial and endoplasmic reticulum accumulation of Parkin, and, in the presence of N-acetyl cysteine, mitophagy. *Neurobiol Dis* 2015, 74: 180–193.
- [73] Clark IE, Dodson MW, Jiang C, Cao JH, Huh JR, Seol JH, *et al.* *Drosophila pink1* is required for mitochondrial function and interacts genetically with parkin. *Nature* 2006, 441: 1162–1166.
- [74] Narendra DP, Jin SM, Tanaka A, Suen DF, Gautier CA, Shen J, *et al.* PINK1 is selectively stabilized on impaired mitochondria to activate Parkin. *PLoS Biol* 2010, 8: e1000298.

- [75] Eiyama A, Okamoto K. PINK1/Parkin-mediated mitophagy in mammalian cells. *Curr Opin Cell Biol* 2015, 33: 95–101.
- [76] Iwabuchi M, Sheng H, Thompson JW, Wang L, Dubois LG, Gooden D, *et al.* Characterization of the ubiquitin-modified proteome regulated by transient forebrain ischemia. *J Cereb Blood Flow Metab* 2014, 34: 425–432.
- [77] Kubli DA, Zhang X, Lee Y, Hanna RA, Quinsay MN, Nguyen CK, *et al.* Parkin protein deficiency exacerbates cardiac injury and reduces survival following myocardial infarction. *J Biol Chem* 2013, 288: 915–926.
- [78] Thomas RL, Kubli DA, Gustafsson AB. Bnip3-mediated defects in oxidative phosphorylation promote mitophagy. *Autophagy* 2011, 7: 775–777.
- [79] Sandoval H, Thiagarajan P, Dasgupta SK, Schumacher A, Prchal JT, Chen M, *et al.* Essential role for Nix in autophagic maturation of erythroid cells. *Nature* 2008, 454: 232–235.
- [80] Zhang J, Ney PA. Role of BNIP3 and NIX in cell death, autophagy, and mitophagy. *Cell Death Differ* 2009, 16: 939–946.
- [81] Ney PA. Mitochondrial autophagy: Origins, significance, and role of BNIP3 and NIX. *Biochim Biophys Acta* 2015.
- [82] Gao F, Chen D, Si J, Hu Q, Qin Z, Fang M, *et al.* The mitochondrial protein BNIP3L is the substrate of PARK2 and mediates mitophagy in PINK1/PARK2 pathway. *Hum Mol Genet* 2015, 24: 2528–2538.
- [83] Liu L, Feng D, Chen G, Chen M, Zheng Q, Song P, *et al.* Mitochondrial outer-membrane protein FUNDC1 mediates hypoxia-induced mitophagy in mammalian cells. *Nat Cell Biol* 2012, 14: 177–185.
- [84] Wu W, Tian W, Hu Z, Chen G, Huang L, Li W, *et al.* ULK1 translocates to mitochondria and phosphorylates FUNDC1 to regulate mitophagy. *EMBO Rep* 2014, 15: 566–575.
- [85] Ding WX, Ni HM, Li M, Liao Y, Chen X, Stolz DB, *et al.* Nix is critical to two distinct phases of mitophagy, reactive oxygen species-mediated autophagy induction and Parkin-ubiquitin-p62-mediated mitochondrial priming. *J Biol Chem* 2010, 285: 27879–27890.
- [86] Michiorri S, Gelmetti V, Giarda E, Lombardi F, Romano F, Marongiu R, *et al.* The Parkinson-associated protein PINK1 interacts with Beclin1 and promotes autophagy. *Cell Death Differ* 2010, 17: 962–974.
- [87] Matsui Y, Takagi H, Qu X, Abdellatif M, Sakoda H, Asano T, *et al.* Distinct roles of autophagy in the heart during ischemia and reperfusion: roles of AMP-activated protein kinase and Beclin 1 in mediating autophagy. *Circ Res* 2007, 100: 914–922.
- [88] Zheng YQ, Liu JX, Li XZ, Xu L, Xu YG. RNA interference-mediated downregulation of Beclin1 attenuates cerebral ischemic injury in rats. *Acta Pharmacol Sin* 2009, 30: 919–927.
- [89] Zhu M, Zhou M, Shi Y, Li WW. Effects of echinacoside on MPP(+)-induced mitochondrial fragmentation, mitophagy and cell apoptosis in SH-SY5Y cells. *Zhong Xi Yi Jie He Xue Bao* 2012, 10: 1427–1432.
- [90] Geisler S, Holmstrom KM, Skujat D, Fiesel FC, Rothfuss OC, Kahle PJ, *et al.* PINK1/Parkin-mediated mitophagy is dependent on VDAC1 and p62/SQSTM1. *Nat Cell Biol* 2010, 12: 119–131.
- [91] Narendra D, Kane LA, Hauser DN, Fearnley IM, Youle RJ. p62/SQSTM1 is required for Parkin-induced mitochondrial clustering but not mitophagy; VDAC1 is dispensable for both. *Autophagy* 2010, 6: 1090–1106.
- [92] Chen H, Chan DC. Mitochondrial dynamics—fusion, fission, movement, and mitophagy—in neurodegenerative diseases. *Hum Mol Genet* 2009, 18: R169–176.
- [93] Twig G, Shirihai OS. The interplay between mitochondrial dynamics and mitophagy. *Antioxid Redox Signal* 2011, 14: 1939–1951.
- [94] Mao K, Klionsky DJ. Participation of mitochondrial fission during mitophagy. *Cell Cycle* 2013, 12: 3131–3132.
- [95] Santel A, Frank S. Shaping mitochondria: The complex posttranslational regulation of the mitochondrial fission protein DRP1. *IUBMB Life* 2008, 60: 448–455.
- [96] Poole AC, Thomas RE, Yu S, Vincow ES, Pallanck L. The mitochondrial fusion-promoting factor mitofusin is a substrate of the PINK1/parkin pathway. *PLoS One* 2010, 5: e10054.
- [97] Lutz AK, Exner N, Fett ME, Schlehe JS, Kloos K, Lammermann K, *et al.* Loss of parkin or PINK1 function increases Drp1-dependent mitochondrial fragmentation. *J Biol Chem* 2009, 284: 22938–22951.
- [98] Okatsu K, Saisho K, Shimanuki M, Nakada K, Shitara H, Sou YS, *et al.* p62/SQSTM1 cooperates with Parkin for perinuclear clustering of depolarized mitochondria. *Genes Cells* 2010, 15: 887–900.
- [99] Pankiv S, Clausen TH, Lamark T, Brech A, Bruun JA, Outzen H, *et al.* p62/SQSTM1 binds directly to Atg8/LC3 to facilitate degradation of ubiquitinated protein aggregates by autophagy. *J Biol Chem* 2007, 282: 24131–24145.
- [100] Grullich C, Duvoisin RM, Wiedmann M, van Leyen K. Inhibition of 15-lipoxygenase leads to delayed organelle degradation in the reticulocyte. *FEBS Lett* 2001, 489: 51–54.
- [101] Joo JH, Dorsey FC, Joshi A, Hennessy-Walters KM, Rose KL, McCastlain K, *et al.* Hsp90-Cdc37 chaperone complex regulates Uik1- and Atg13-mediated mitophagy. *Mol Cell* 2011, 43: 572–585.
- [102] Mao K, Wang K, Zhao M, Xu T, Klionsky DJ. Two MAPK-signaling pathways are required for mitophagy in *Saccharomyces cerevisiae*. *J Cell Biol* 2011, 193: 755–767.
- [103] Piao CS, Kim JB, Han PL, Lee JK. Administration of the p38 MAPK inhibitor SB203580 affords brain protection with

- a wide therapeutic window against focal ischemic insult. *J Neurosci Res* 2003, 73: 537–544.
- [104] Pei H, Song X, Peng C, Tan Y, Li Y, Li X, *et al.* TNF-alpha inhibitor protects against myocardial ischemia/reperfusion injury via Notch1-mediated suppression of oxidative/nitrative stress. *Free Radic Biol Med* 2015, 82: 114–121.
- [105] Lupfer C, Thomas PG, Anand PK, Vogel P, Milasta S, Martinez J, *et al.* Receptor interacting protein kinase 2-mediated mitophagy regulates inflammasome activation during virus infection. *Nat Immunol* 2013, 14: 480–488.
- [106] Lazarou M, Jin SM, Kane LA, Youle RJ. Role of PINK1 binding to the TOM complex and alternate intracellular membranes in recruitment and activation of the E3 ligase Parkin. *Dev Cell* 2012, 22: 320–333.
- [107] Gegg ME, Cooper JM, Chau KY, Rojo M, Schapira AH, Taanman JW. Mitofusin 1 and mitofusin 2 are ubiquitinated in a PINK1/parkin-dependent manner upon induction of mitophagy. *Hum Mol Genet* 2010, 19: 4861–4870.
- [108] Yoshioka H, Katsu M, Sakata H, Okami N, Wakai T, Kinouchi H, *et al.* The role of PARL and HtrA2 in striatal neuronal injury after transient global cerebral ischemia. *J Cereb Blood Flow Metab* 2013, 33: 1658–1665.
- [109] Meissner C, Lorenz H, Weihofen A, Selkoe DJ, Lemberg MK. The mitochondrial intramembrane protease PARL cleaves human Pink1 to regulate Pink1 trafficking. *J Neurochem* 2011, 117: 856–867.
- [110] Hollville E, Carroll RG, Cullen SP, Martin SJ. Bcl-2 family proteins participate in mitochondrial quality control by regulating Parkin/PINK1-dependent mitophagy. *Mol Cell* 2014, 55: 451–466.
- [111] Wu H, Xue D, Chen G, Han Z, Huang L, Zhu C, *et al.* The BCL2L1 and PGAM5 axis defines hypoxia-induced receptor-mediated mitophagy. *Autophagy* 2014, 10: 1712–1725.
- [112] Yang KC, Ma X, Liu H, Murphy J, Barger PM, Mann DL, *et al.* Tumor necrosis factor receptor-associated factor 2 mediates mitochondrial autophagy. *Circ Heart Fail* 2015, 8: 175–187.
- [113] Kanki T. Nix, a receptor protein for mitophagy in mammals. *Autophagy* 2010, 6: 433–435.
- [114] Lee JY, Nagano Y, Taylor JP, Lim KL, Yao TP. Disease-causing mutations in parkin impair mitochondrial ubiquitination, aggregation, and HDAC6-dependent mitophagy. *J Cell Biol* 2010, 189: 671–679.
- [115] Muhammad S, Barakat W, Stoyanov S, Murikinati S, Yang H, Tracey KJ, *et al.* The HMGB1 receptor RAGE mediates ischemic brain damage. *J Neurosci* 2008, 28: 12023–12031.
- [116] Stetler RA, Gao Y, Zhang L, Weng Z, Zhang F, Hu X, *et al.* Phosphorylation of HSP27 by protein kinase D is essential for mediating neuroprotection against ischemic neuronal injury. *J Neurosci* 2012, 32: 2667–2682.
- [117] Bischoff P, Josset E, Dumont FJ. Novel pharmacological modulators of autophagy and therapeutic prospects. *Expert Opin Ther Pat* 2012, 22: 1053–1079.
- [118] Shoji-Kawata S, Sumpter R, Leveno M, Campbell GR, Zou Z, Kinch L, *et al.* Identification of a candidate therapeutic autophagy-inducing peptide. *Nature* 2013, 494: 201–206.
- [119] Jimenez RE, Kubli DA, Gustafsson AB. Autophagy and mitophagy in the myocardium: therapeutic potential and concerns. *Br J Pharmacol* 2014, 171: 1907–1916.