

# FACTORS INFLUENCING LONGITUDINAL STAIR CLIMB PERFORMANCE FROM MIDLIFE TO EARLY LATE LIFE: THE STUDY OF WOMEN'S HEALTH ACROSS THE NATION CHICAGO AND MICHIGAN SITES

B.S. LANGE-MAIA<sup>1,2</sup>, C.A. KARVONEN-GUTIERREZ<sup>3</sup>, E.S. STROTMAYER<sup>4</sup>, E.F. AVERY<sup>1,2</sup>,  
B.M. APPELHANS<sup>1</sup>, S.L. FITZPATRICK<sup>5</sup>, I. JANSSEN<sup>1</sup>, S.A. DUGAN<sup>6</sup>, H.M. KRAVITZ<sup>1,7</sup>

1. Department of Preventive Medicine, Rush University Medical Center, Chicago, IL, USA; 2. Center for Community Health Equity, Rush University Medical Center, Chicago, IL, USA; 3. Department of Epidemiology, School of Public Health, University of Michigan, Ann Arbor, MI, USA; 4. Department of Epidemiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA; 5. Kaiser Permanente Center for Health Research, Portland, OR, USA; 6. Department of Physical Medicine & Rehabilitation, Rush University Medical Center, Chicago, IL, USA; 7. Department of Psychiatry, Rush University Medical Center, Chicago, IL, USA. Corresponding author: Brittney Lange-Maia, PhD, MPH, 1700 W. Van Buren St., Suite 470, Chicago, IL 60612, Phone: 312-942-9697; Fax: 312-942-8119; Email: Brittney\_Lange-Maia@Rush.edu

**Abstract:** *Objectives:* To quantify the longitudinal change in stair climb performance (a measure indicative of both physical function and muscle power), determine whether physical activity is related to slower decline in performance, and to identify factors that modify the longitudinal change in performance among women from midlife to late life. *Design:* Longitudinal cohort study with up to 15 study visits. *Setting:* Two sites of the Study of Women's Health Across the Nation. *Participants:* Black (n=411) and white (N=419) women followed from median age 47.0 (44.6-49.6) to 62.0 (55.8-65.3) years. *Interventions:* N/A. *Measurements:* Performance on a stair climb test (ascend/descend 4 steps, 3 cycles) was timed. Physical activity (PA) was assessed using the Kaiser Physical Activity Survey (KPAS; possible range 0-15 points). Sociodemographic and health factors were assessed via self-report. BMI was calculated with measured height and weight. Mixed-effects regression modeled longitudinal change in stair climb performance. *Results:* Average baseline stair climb time was 18.12 seconds (95% CI: 17.83-18.41), with 0.98% (95% CI: 0.84%-1.11%) annual slowing. In fully adjusted models, higher levels of PA were associated with faster stair climb times (2.09% faster per point higher, 95% CI: -2.87%-1.30%), and black women had 5.22% (95% CI: 2.43%-8.01%) slower performance compared to white women. Smoking, financial strain, diabetes, osteoarthritis, fair/poor health, and stroke were associated with 3.36% (95% CI: 0.07%-6.65%), 7.56% (95% CI: 4.75%-10.37%), 8.40% (95% CI: 2.89%-13.92%), 8.46% (95% CI: 5.12%-11.79%), 9.16% (95% CI: 4.72%-13.60%), and 16.94% (95% CI: 5.37%-28.51%) slower performance, respectively. In separate models, higher BMI (per 1-unit), osteoarthritis, fair/poor health, and diabetes, were each associated with 0.06% (95% CI: 0.04%-0.08%), 0.48% (95% CI: 0.12%-0.84%), 0.81% (95% CI: 0.35%-1.28%), and 0.84% (95% CI: 0.22%-1.46%), additional slowing per year over time. *Conclusion:* Significant declines in function were evident as women transitioned from midlife to early late life. Declines were amplified by indicators of poor health, emphasizing the importance of health in midlife for promoting healthy aging.

**Key words:** Physical function, aging, midlife women.

## Introduction

At all ages, women are disproportionately affected by limitations in physical function compared to men. During the midlife, physical function for women is influenced both by processes related to chronologic aging and hormonal changes occurring with ovarian aging (1-4). Though limitations in physical function are most prevalent in late life (ages 65+ years), approximately 20% of women report limitations in physical functioning in midlife, the life stage reflecting ages 40-65 years (5). Though much research has focused on physical function during either mid- or late life, considerably less is known about how physical function changes during the critical transition period moving from midlife to late life.

In a broad sense, risk factors for worse physical function during midlife are largely consistent with risk factors during late life, including body composition and factors related health status (6-9). In addition, considerable racial disparities have been previously identified in physical function at midlife

between black and white women, with black women being more likely to have worse performance-based and self-reported physical function, as well as a higher prevalence of disability (2, 7, 10-12). As in older adults, physical activity is an important protective factor for physical function during midlife (13-15).

Limitations in midlife are evident in both self-report (2, 5, 16) and performance-based measures of physical function (7), which is critical as self-report and performance-based measures are known to capture related, but distinct, concepts (17-20). One commonly assessed aspect of physical function (self-report and performance-based) is stair climbing, as the ability to negotiate stairs is important in maintaining independence and community living. Slower performance-based stair negotiation time has been shown to be predictive of the development of functional limitations in older adults, even among those who report no difficulty with stairs (21). Ascending and descending stairs encompasses multiple facets of physical and muscle function, and is a more complex kinematic task

## LONGITUDINAL STAIR CLIMB IN MIDLIFE WOMEN

than walking, another commonly assessed aspect of physical function. Compared to other activities of daily living and aspects of mobility, the ability to climb stairs is among the first—and fastest—to decline with age (22-24). Thus, this test may be preferable for assessing longitudinal physical function changes from mid to late life, a time period where changes in performance-based function may be subtle.

The purpose of this study was to quantify the longitudinal change in stair climb performance among black and white midlife women transitioning to early late life. We also sought to determine whether physical activity was associated with slower decline in performance over time, if performance differed between black and white women, and to identify other factors that modify the longitudinal change in stair climb performance from midlife to early late life.

### Methods


#### Participants

Participants were from the Michigan (southeast Michigan, Detroit-area) and Chicago, IL sites of the Study of Women's Health Across the Nation (SWAN), a longitudinal, multi-racial/ethnic, community-based cohort study of the menopausal transition. Recruitment and design of the SWAN study have been described in detail elsewhere (25). Briefly, women were eligible for the SWAN study if they were between the ages of 42-52 years at baseline in 1996/97, had an intact uterus and at least one ovary, were premenopausal or early peri-menopausal (i.e., had at least one menstrual period in the past 3 months), and were not pregnant, breastfeeding or lactating. Women at the Chicago, IL site were recruited via a random population-based sampling in a contiguous area on the south side of Chicago (N=868) (26). Of these, 457 were recruited for the parent study (Chicago "core" cohort), with the remaining women being part of a sub-study (Chicago "site-specific" cohort). Both Chicago cohorts underwent identical clinical assessments. The Michigan site recruited from two communities in southeast Michigan using a community census based on electrical utility listings in Ypsilanti and Inkster, MI (N=543). By design, both the Michigan and Chicago sites recruited non-Hispanic black and white women. The Chicago and Michigan site protocols were approved by the Rush University Medical Center or the University of Michigan Health Sciences Institutional Review Boards, respectively. All participants provided written informed consent before enrolling and at each clinic visit.

A stair climbing test was assessed repeatedly at these SWAN sites. Stair climb tests commenced for the Michigan cohort at the main study baseline in 1996/97, the Chicago site-specific cohorts at their baseline in 1997/98, whereas the SWAN Chicago core cohort did not begin stair climb data collection until the 1998/1999 (Follow-up visit 2). Participants were asked to complete the stair climb test at near-annual follow-up visits. The initial visit where the stair climb test occurred in conjunction with a physical activity assessment was considered

the analytic baseline for this analysis. Because physical activity was a particular predictor of interest and was not assessed in Follow-up visit 2, the 1999/2000 visit (Follow-up visit 3) study visit was used as analytic baseline for the Chicago core cohort. Participants were included in analyses if they completed each of the following: 1) the stair climb test at the analytic baseline or the immediate follow-up visit, 2) the stair climb test on at least one additional visit, and 3) the physical activity questionnaire at the analytic baseline. A total of 830 participants were included in longitudinal analyses using data through Visit 15, which occurred in 2015-17 (Figure 1). In total, up to 15 stair climb observations were possible for the Michigan and Chicago site-specific cohorts and up to 12 visits for the Chicago core cohort. Mean visits per participant per site were as follows: Michigan site  $10.6 \pm 4.0$  (range 2-15), Chicago site-specific  $6.5 \pm 3.0$  (range 2-12), Chicago core cohort  $8.4 \pm 3.0$  (range 2-12).

**Figure 1**  
Participants from Michigan and Chicago SWAN Sites



Women who were excluded (N=582) versus those who were included did not differ by age ( $46.9 \pm 3.1$  years vs.  $47.1 \pm 3.2$  years,  $p=0.21$ ). However, excluded women were more likely to be black (57.2% vs. 49.5%,  $p=0.004$ ), from one of the Chicago sites (82.7% vs. 46.7%,  $p<0.001$ ), premenopausal (41.8% vs. 33.6%,  $p=0.03$ ), and report fair or poor health (16.7% vs. 11.6%,  $p=0.01$ ). Other demographic factors, including education, financial difficulty paying for basics, and marital status did not differ by inclusion status.

#### Stair Climb Test

The timed stair climb task required participants to ascend and descend four standard stairs for three consecutive cycles. Participants were permitted to use the hand rail if necessary. Times at the Chicago site were collected in seconds, while times at the Michigan site were collected to the tenth of a

second. For consistency, times from the Michigan site were rounded to the nearest second for analysis.

### **Physical Activity Assessment**

Self-reported physical activity was collected using the Kaiser Physical Activity Survey (KPAS) at analytic baseline (27). This self-administered questionnaire is comprised of 38 items assessing physical activity in the past year, and was originally modified from the Baecke physical activity questionnaire (28). Physical activity from three domains was assessed, including sports/exercise, active living, and household/caregiving. Responses are primarily categorical to assess frequency, duration, and relative intensity, with an open-ended question about sport/exercise participation. Total scores can range from 0 to 15, with higher values indicating greater physical activity participation.

### **Covariates**

All covariates were used from the analytic baseline for each respective group unless otherwise specified. Sociodemographic characteristics including race (black or white), educational attainment, marital status, and financial difficulty were assessed via self-report. Age was calculated using date of birth and date of first included stair climb test. Health factors included menopausal hormone use (lifetime ever use, yes/no), menopausal status (pre-menopausal, peri-menopausal, post-menopausal, or unknown) characterized using bleeding criteria, and smoking status (never/former/current, combined to current vs. former/never for analyses). BMI ( $\text{kg}/\text{m}^2$ ) was calculated from measured height and weight. Self-reported health status was self-reported as excellent, very good, good, fair, or poor (excellent/very good/good vs fair/poor combined for analyses). Severity of bodily pain in the past 4 weeks was assessed using a single-item question from the Short Form Health Survey (SF-36), with responses grouped to create categories of mild/very mild, moderate, or severe/very severe. Doctor-diagnosed history of stroke and osteoarthritis was collected by self-report. Diabetes status was determined via self-reported healthcare provider diagnosis, use of hypoglycemic medications, or fasting glucose  $\geq 126$  mg/dl. Hypertension was defined as measured resting blood pressure  $\geq 140$  mmHg (systolic) or  $\geq 90$  mmHg (diastolic) or use of antihypertensive medications. Depressive symptoms were assessed using the Center for Epidemiological Studies Depression Scale (CES-D; score of  $\geq 16$  indicating clinically significant level of depressive symptoms vs.  $< 16$ ) (29, 30).

### **Statistical Analyses**

Descriptive statistics were compared between black and white women using t-tests or nonparametric alternatives as necessary for continuous factors and chi-squared tests for categorical factors. All statistical analyses were performed using SAS version 9.3 (SAS Institute, Inc., Cary, NC).

Stair climb times were natural-log transformed for analyses

due to right-skewed data distribution. Repeated measures analysis using linear mixed models (SAS Proc Mixed) were performed including random slopes and intercepts and an unstructured covariance matrix. An unadjusted model with years since baseline as the only predictor was used to determine the average predicted stair climb time and the average change per year. Then, four sets of models were built: Model 1 included site, age, race, and time; Model 2 added physical activity; Model 3 added education, financial difficulty, and marital status; and Model 4 added BMI, bodily pain, health status, hormone use, menopausal status, smoking status, osteoarthritis, hypertension, stroke, diabetes, and depressive symptoms. In exploratory analyses, interactions with time were considered individually for factors included in the final model to determine whether the rate of change in stair climb time varied by sociodemographic, health, or health behavioral factors. In each model, though we considered several covariates, only factors reaching statistical significance ( $p < 0.05$ ) were retained. However, age and site were included a priori regardless of significance. Because the different sites had different numbers of total stair climb visits, sensitivity analyses restricting the Michigan and Chicago Site-Specific participants to only visits 1-13 so all sites had a maximum of 13 stair climb assessments were performed.

## **Results**

Overall, 830 women (Figure 1;  $N=411$  black and  $N=419$  white) were included in analyses. Median follow-up time was 15.5 years (interquartile range 8.4-18.3 years), and women were followed from a median age of 47.0 (44.6-49.6) to 62.0 (55.8-65.3) years.

Participant characteristics generally differed by race (Table 1), with white women being more likely to have more than a HS diploma (83.8% vs. 72.5%), more likely to be currently married (73.3% vs. 49.1%), less likely to have financial difficulty paying for basics (31.7% vs. 45.1%), and more likely to have ever used hormones (28.4% vs. 13.7%);  $p < 0.001$  for all comparisons. Regarding health status, black women generally had worse self-reported health ( $p < 0.001$ ), were more likely to be current smokers (25.1% vs. 18.9%,  $p = 0.03$ ), had higher BMI (30.8  $\text{kg}/\text{m}^2$ , IQR [interquartile range]: 26.3-36.8  $\text{kg}/\text{m}^2$  vs. 28.2  $\text{kg}/\text{m}^2$ , IQR: 24.3-34.3  $\text{kg}/\text{m}^2$ ;  $p < 0.001$ ), were more likely to report bodily pain ( $p < 0.001$ ), and were more likely to have CES-D scores  $\geq 16$  (25.5% vs 17.2%;  $p < 0.001$ ). Age, prevalence of diabetes, stroke, and osteoarthritis did not vary by race. Compared to white women, black women had lower mean KPAS scores ( $7.3 \pm 1.8$  vs.  $8.1 \pm 1.8$ ,  $p < 0.001$ ), indicating that they were less physically active.

In unadjusted models, the mean baseline stair climb time was 18.12 seconds (95% CI: 17.8-18.4 seconds), and women slowed on average 0.98% (95% CI: 0.84%-1.11%) per year (results not shown). In adjusted models, smoking, financial strain, fair/poor health, osteoarthritis, stroke and diabetes were associated 3.36%

*LONGITUDINAL STAIR CLIMB IN MIDLIFE WOMEN*

**Table 1**  
 Participant Characteristics<sup>1</sup> at Initial Stair Climb Visit Overall and by Race

<b>Participant Characteristics</b>	<b>All Women N=830</b>	<b>Black Women N=411</b>	<b>White Women N=419</b>	<b>P-Value</b>
Age, years, Median (IQR) <sup>2</sup>	47.0 (44.6, 49.6)	46.7 (44.4, 49.5)	47.1 (44.8, 49.7)	0.28
Site Group				<0.001
Michigan	442 (53.3)	262 (63.7)	180 (43.0)	
Chicago Core	212 (25.5)	104 (25.3)	108 (25.8)	
Chicago Site Specific	176 (21.2)	45 (10.9)	131 (31.3)	
Education ≤High School	617 (78.2)	285 (72.5)	332 (83.8)	<0.001
Currently Married	488 (61.2)	197 (49.1)	291 (73.3)	<0.001
Financial difficulty paying for basics, n (%)	305 (38.5)	180 (45.1)	125 (31.7)	<0.001
Menopausal Status, n (%)				0.21
Postmenopausal	31 (3.9)	19 (4.8)	12 (3.0)	
Perimenopausal	403 (50.8)	211 (52.9)	192 (48.7)	
Premenopausal	333 (42.0)	159 (39.8)	174 (44.2)	
Unknown/Missing	26 (3.3)	10 (2.5)	16 (4.1)	
Hormone Use: Ever, n (%)	167 (21.0)	55 (13.7)	112 (28.4)	<0.001
Self-Rated Health, n (%)				<0.001
Excellent	143 (18.0)	52 (13.0)	91 (23.1)	
Very Good	301 (37.9)	135 (33.8)	166 (42.1)	
Good	258 (32.5)	150 (37.5)	108 (27.4)	
Fair	78 (9.8)	53 (13.3)	25 (6.3)	
Poor	14 (1.8)	10 (2.5)	4 (1.0)	
BMI, median (IQR)	29.7 (25.0, 35.5)	30.8 (26.3, 36.8)	28.2 (24.3, 34.3)	<0.001
Current Smoker, n (%)	181 (21.9)	102 (25.1)	79 (18.9)	0.03
Prevalent Diseases/Conditions, n (%)				
Hypertension	299 (36.2)	185 (45.1)	114 (27.4)	<0.001
Diabetes	50 (6.2)	30 (7.5)	20 (5.0)	0.14
Stroke	10 (1.2)	7 (1.7)	3 (0.7)	0.19
Osteoarthritis	186 (22.5)	97 (23.7)	89 (21.3)	0.41
Bodily Pain				<0.001
None	110 (13.3)	61 (14.8)	49 (11.7)	
Mild/Very Mild	466 (56.1)	201 (48.9)	265 (63.2)	
Moderate	200 (24.1)	108 (26.3)	92 (22.0)	
Severe/Very Severe	54 (6.5)	41 (10.0)	13 (3.1)	
CES-D <sup>3</sup> Score ≥16	177 (21.3)	105 (25.5)	72 (17.2)	<0.001
KPAS <sup>4</sup> Score, mean (SD <sup>5</sup> )	7.7 (1.8)	7.3 (1.8)	8.1 (1.8)	<0.001

1. Percentages are reflective of participants with complete data for the measure. Missing data <5%. 2. Interquartile range. 3. Center for Epidemiological Studies Depression Scale; 4. Kaiser Physical Activity Questionnaire; 5. Standard deviation


**Table 2**  
Longitudinal stair climb performance linear mixed models

Model Parameters	Model 1			Model 2			Model 3 <sup>1</sup>			Model 4 <sup>2</sup>		
	Estimate	95% CI Lower	95% CI Upper	Estimate	95% CI Lower	95% CI Upper	Estimate	95% CI Lower	95% CI Upper	Estimate	95% CI Lower	95% CI Upper
Intercept (Seconds)	17.72	17.23	18.22	23.29	21.74	24.95	23.18	21.56	24.93	18.96	17.71	20.30
Time (Effect per year)	0.92%	0.77%	1.07%	0.92%	0.77%	1.07%	0.97%	0.81%	1.12%	0.95%	0.80%	1.11%
Age <sup>3</sup>	0.81%	0.28%	1.33%	0.75%	0.25%	1.26%	0.82%	0.31%	1.34%	0.77%	0.30%	1.24%
Black Race	9.72%	6.63%	12.81%	7.43%	4.42%	10.44%	6.19%	3.14%	9.24%	5.22%	2.43%	8.01%
Physical Activity				-3.53%	-4.35%	-2.71%	-3.59%	-4.42%	-2.77%	-2.09%	-2.87%	-1.30%
Some college or higher							-3.94%	-7.63%	-0.25%	NS	---	---
Difficulty Paying for Basics							10.41%	7.38%	13.45%	7.56%	4.75%	10.37%
Health Fair or Poor										9.16%	4.72%	13.60%
BMI <sup>3</sup>										0.95%	0.76%	1.15%
Current Smoker										3.36%	0.07%	6.65%
Diabetes										8.40%	2.89%	13.92%
Stroke										16.94%	5.37%	28.51%
Osteoarthritis										8.46%	5.12%	11.79%

1. Model sample size N=781, also considered marital status but removed due to non-significance; 2. Model sample size N=755, also considered bodily pain, menopausal status, hormone use, hypertension, and depressive symptoms, but removed due to non-significance. Education was removed in Model 4 as it was attenuated to non-significance by the addition of other covariates. All models also adjusted for site. 3. Age and BMI each centered at their respective medians. Note: Beta estimates for all parameters (except intercept) were multiplied by 100 to interpret as percentage difference in stair climb time. Negative coefficients indicate lower stair climb time, and thus better performance (i.e. faster), while positive values indicate slower stair climb time.

**Figure 2**

Longitudinal change in stair test completion time; depicting interactions for A) self-reported health status, B) body mass index (BMI), C) osteoarthritis, and D) diabetes



LONGITUDINAL STAIR CLIMB IN MIDLIFE WOMEN

**Table 3**

Separate longitudinal mixed models predicting longitudinal change in stair climb time, significant interactions<sup>1</sup>

	<b>BMI Interaction</b>	<b>Health Status (Fair/Poor Health) Interaction</b>	<b>Diabetes Interaction</b>	<b>Osteoarthritis Interaction</b>
Model Parameters	Estimate (95% CI)	Estimate (95% CI)	Estimate (95% CI)	Estimate (95% CI)
Intercept (Seconds)	18.99 (17.73, 20.33)	19.02 (17.76, 20.37)	18.99 (17.73, 20.33)	19.03 (17.77, 20.37)
Time (Effect per year)	-0.99 (-1.62, -0.36)	0.86 (0.70, 1.02)	0.90 (0.74, 1.06)	0.84 (0.67, 1.02)
Main Effect	0.76 (0.56, 0.96)	6.65 (1.99, 11.32)	5.80 (-0.04, 11.64)	6.96 (3.44, 10.48)
Interaction with Time	0.06 (0.04, 0.08)	0.81 (0.35, 1.28)	0.84 (0.22, 1.46)	0.48 (0.12, 0.84)

1. Adjusted for all factors included in Table 2, Model 4, including site, age, race, physical activity, difficulty paying for basics, health status, BMI, current smoking, diabetes, stroke, and osteoarthritis; Note: Beta estimates for all parameters (except intercept) were multiplied by 100 to interpret as percentage difference in stair climb time. Negative coefficients indicate lower stair climb time, and thus better performance (i.e. faster), while positive values indicate slower stair climb time.

(95% CI: 0.07%-6.65%), 7.56% (95% CI:4.75%-10.37%), 8.46% (95% CI: 5.12%-11.79%), 9.16% (95% CI: 4.72%-13.60%), 16.94% (95% CI: 5.37%-28.51%), and slower stair performance, respectively (Table 2, Model 4). Each unit higher of BMI was associated with 0.95% (95% CI: 0.76%-1.15%) slower stair climb times. We also considered bodily pain, menopausal status, hormone use, hypertension, and depressive symptoms, though these factors were not independently associated with stair climb time ( $p>0.05$  for each) and were not retained in the models.

In separate models for the exploratory analyses testing for interactions among significant factors in Model 4, several factors were related to a difference in performance over time. Significant interactions with time were found such that higher BMI (per 1-unit higher), osteoarthritis, fair/poor health, and diabetes, and were each associated with 0.06% (95% CI:0.04%-0.08%), 0.48% (95% CI:0.12%-0.84%), 0.81% (95% CI:0.4%-1.3%), 0.84% (95% CI:0.22%-1.46%), additional slowing per year over time, respectively, (Table 3; interactions also displayed in Figure 2).

Results remained consistent in sensitivity analyses restricting all sites to only 12 possible assessments (results not shown).

**Discussion**

In this longitudinal study, we found overall significant changes in stair climb time among women transitioning from midlife to early late life, followed for an average of nearly 16 years. This degree of decline in physical function performance is notable, given that it occurs earlier than when physical function is typically considered to decline, suggesting the stair climb test may be a sensitive performance measure for early aging. Further, higher BMI, and certain prevalent conditions—including diabetes, history of stroke, and osteoarthritis—were each associated with accelerated performance deterioration, supporting the importance of promoting health at midlife to improve the likelihood of physical independence at old age.

We found a significant racial difference in stair climb times between black and white women. The 9.72% race group difference in stair climb time from the minimally adjusted

model was ultimately attenuated nearly 50%, and a 5.22% difference between black and white women remained in the final model. Prior work has shown clear racial disparities among mid- and late life women in a variety of aspects of physical function including a higher likelihood of clinically slow gait speed, worse performance-based physical function and self-reported functional limitations, and more disability in black women (2, 7, 10-12, 31-34) Still, in some of these studies racial differences were attenuated to non-significance when controlling for covariates, particularly prevalent health conditions, socioeconomic factors, BMI, pain, and osteoarthritis (2, 7, 33). Additional research addressing the mechanisms underlying this racial disparity is warranted, though our results and others suggest that health inequities may play a role.

Physical activity was related to better stair climb performance at every time point, but it did not impact the rate of decline in stair climb over time. This is consistent with work in older adults showing that physical activity is related to better cardiovascular endurance, but does not influence the rate of endurance decline occurring with age (35, 36). Thus, our results suggest that the level of physical activity in midlife is particularly important to achieve a higher physical function prior to the decline associated with the aging process in early late life.

Though physical activity did not influence the rate of change in performance over time, several other factors did. These factors—including self-rated fair or poor health, higher BMI, osteoarthritis, stroke, and diabetes—could be important for developing targeted secondary prevention efforts in order for midlife women to maintain intact function through late life. Prior work has shown accelerated physical function decline and increased disability risk among adults with diabetes, with this effect being particularly profound in women (37). Obesity at midlife is associated with higher risk of mobility limitations in late life compared to normal weight or overweight in midlife (38). Reducing obesity in midlife is a particularly relevant intervention target in order to reduce diabetes, stroke, and osteoarthritis associated with functional decline.

Declines in stair climb performance were evident over a 15 year period among midlife women transitioning to late

life, indicating this measure shows promise for wider use for assessing performance in midlife populations. A future direction is to establish age-based normative values and determine values associated with poor outcomes.

Though other studies have utilized stair climb tests, large protocol variations exist, making the comparison to prior work difficult (39). In older adults (age 70+), age-normative values for stair ascent and descent times have been determined, as well as values of large meaningful change in stair ascent and descent performance, with a one year change of 1.15 seconds for ascent and 1.25 seconds for descent indicating large meaningful changes (21, 40). However, these studies utilized a substantially different protocol and are not directly comparable. A further variation is in using stair climb tests for assessing lower-extremity power based on ascend time (41, 42). Stair climb power is strongly related to mobility and lower-extremity performance in older adults (41). Power could not be calculated; split times were recently completed as part of the updated protocol in SWAN at all sites and will be a future direction for analyses. While ascending and descending stairs is a common task for community living, the protocol for this particular stair climb test does not necessarily mimic stair climbing done in everyday life. Had we used a self-report assessment of stair climbing ability, we may have seen different longitudinal patterns.

Strengths of this study include the longitudinal design with a median follow-up of 15.5 years. This substantial follow-up allowed us to observe women's physical performance as they transitioned from midlife to early late life. This work builds upon prior work in SWAN (limited to only the Michigan site and only up to study Visit 12 in 2008) utilizing the stair climb test that focused on specific risk factors—peripheral neuropathy (6) and depressive symptoms (8)—by showing the role of physical activity in longitudinal stair climb performance, and also demonstrating how longitudinal performance is modified by factors related to poor health. In addition, the participants in this study were from two sites of a well-characterized cohort of initially midlife women, both black and white, which allowed us to examine a host of potential covariates to assess confounding. Importantly, the stair climb test is likely more sensitive to early changes in physical function compared to traditional tests in older populations such as gait speed or chair rise tests as stair climbing ability is one of the first aspects of physical function to decline with age (22-24).

Limitations of this work must be considered when interpreting these results. First, the difference by site in timelines of when the stair climb test was introduced made data alignment complicated, and made for different follow-up periods by site. We performed sensitivity analyses restricting sites to the same possible number of follow-up visits, and adjusted all models for age and site to mitigate this limitation. In addition, women had to have at least two stair climb times in order to be included, with one of those being either the first or second visit where the test was offered. Potentially, women

with the greatest decreases in function may have been less likely to return for clinic visits, excluding them from these analyses—leading to an underestimation of the change over time. In addition, though the stair climb protocol was the same at both the Chicago and Michigan sites, slight variations existed in the data collection forms. At the Chicago site stair climb times were recorded to the nearest second, while Michigan recorded times to the nearest tenth of a second. Additionally, use of the hand rail was not collected consistently by site, and therefore we could not adjust for hand rail use. Finally, racial/ethnic disparities in health are not limited to differences only between black and white women, though we were limited to only these comparisons due to recruitment at these sites, and SWAN by design only includes women and not men.

In conclusion, this study demonstrated worsening performance on a stair climb task in a cohort of midlife black and white women transitioning from midlife to late life. Women with higher levels of physical activity in midlife had better stair climb performance at midlife through early old age compared to women who were less physically active. Additionally, black women had overall slower stair climb times compared to white women, but there was no difference in their rate of change over time. Specific health-related factors (BMI, self-rated health, prior stroke, and diabetes) were associated with more rapid stair climb performance decline, and secondary prevention interventions should potentially target midlife women who already have chronic conditions, obesity, or report ill health in order to delay and prevent late life disability. Because midlife is a critical period when limitations in physical function begin, a greater focus on changes in physical function in this age group as well as sensitive methods to measure small, though significant, early changes is needed in order to identify women in need of early interventions to prevent significant disability in late life.

*Acknowledgments:* The Study of Women's Health Across the Nation (SWAN) has grant support from the National Institutes of Health (NIH), DHHS, through the National Institute on Aging (NIA), the National Institute of Nursing Research (NINR) and the NIH Office of Research on Women's Health (ORWH) (Grants U01NR004061; U01AG012505, U01AG012535, U01AG012531, U01AG012539, U01AG012546, U01AG012553, U01AG012554, U01AG012495). The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the NIA, NINR, ORWH or the NIH. Clinical Centers: University of Michigan, Ann Arbor – Siobán Harlow, PI 2011 – present; MaryFran Sowers, PI 1994-2011; Massachusetts General Hospital, Boston, MA – Joel Finkelstein, PI 1999 – present; Robert Neer, PI 1994 – 1999; Rush University, Rush University Medical Center, Chicago, IL – Howard Kravitz, PI 2009 – present; Lynda Powell, PI 1994 – 2009; University of California, Davis/Kaiser – Ellen Gold, PI; University of California, Los Angeles – Gail Greendale, PI; Albert Einstein College of Medicine, Bronx, NY – Carol Derby, PI 2011 – present; Rachel Wildman, PI 2010 – 2011; Nanette Santoro, PI 2004 – 2010; University of Medicine and Dentistry – New Jersey Medical School, Newark – Gerson Weiss, PI 1994 – 2004; and the University of Pittsburgh, Pittsburgh, PA – Karen Matthews, PI. NIH Program Office: National Institute on Aging, Bethesda, MD – Chhanda Dutta 2016- present; Winifred Rossi 2012-2016; Sherry Sherman 1994 – 2012; Marcia Ory 1994 – 2001; National Institute of Nursing Research, Bethesda, MD – Program Officers. Central Laboratory: University of Michigan, Ann Arbor – Daniel McConnell (Central Ligand Assay Satellite Services). Coordinating Center: University of Pittsburgh, Pittsburgh, PA – Maria Mori Brooks, PI 2012 - present; Kim Sutton-Tyrrell, PI 2001 – 2012; New England Research Institutes, Watertown, MA – Sonja McKinlay, PI 1995 – 2001. Steering Committee: Susan Johnson, Current Chair, Chris Gallagher, Former Chair. We thank the study staff at each site and all the women who participated in SWAN.

## LONGITUDINAL STAIR CLIMB IN MIDLIFE WOMEN

**Disclosures:** Dr. Kravitz reports grants from National Institutes of Health/National Institute on Aging during the conduct of the study. No other authors have any conflicts to report.

### References

1. Sowers MF, Tomey K, Jannausch M, et al. Physical functioning and menopause states. *Obstet Gynecol.* 2007;110(6):1290-1296.
2. Tseng LA, El Khoudary SR, Young EZ, et al. The association of menopause status with physical function: The study of women's health across the nation. *Menopause.* 2012;19(11):1186-1192.
3. Avis NE, Colvin A, Bromberger JT, et al. Change in health-related quality of life over the menopausal transition in a multiethnic cohort of middle-aged women: Study of Women's Health Across the Nation. *Menopause.* 2009;16(5):860-869.
4. El Khoudary SR, McClure CK, VoPham T, et al. Longitudinal assessment of the menopausal transition, endogenous sex hormones, and perception of physical functioning: The Study of Women's Health across the Nation. *J Gerontol A Biol Sci Med Sci.* 2014;69(8):1011-1017.
5. Sowers M, Pope S, Welch G, Sternfeld B, Albrecht G. The association of menopause and physical functioning in women at midlife. *J Am Geriatr Soc.* 2001;49(11):1485-1492.
6. Ylitalo KR, Herman WH, Harlow SD. Performance-based physical functioning and peripheral neuropathy in a population-based cohort of women at midlife. *Am J Epidemiol.* 2013;177(8):810-817.
7. Sowers M, Jannausch ML, Gross M, et al. Performance-based physical functioning in African-American and Caucasian Women at Midlife: Considering body composition, quadriceps strength, and knee osteoarthritis. *Am J Epidemiol.* 2006;163(10):950-958.
8. Tomey K, Sowers MF, Harlow SD, Jannausch M, Zheng H, Bromberger J. Physical functioning among mid-life women: Associations with trajectory of depressive symptoms. *Social Science & Medicine.* 2010;71(2010):1259-1267.
9. Ylitalo KR, Karvonen-Gutierrez CA, Fitzgerald N, et al. Relationship of race-ethnicity, body mass index, and economic strain with longitudinal self-report of physical functioning: The Study of Women's Health Across the Nation. *Ann Epidemiol.* 2013;23(7):401-408.
10. Kirkness CS, Ren J. Race differences: Use of walking speed to identify community-dwelling women at risk for poor health outcomes--Osteoarthritis Initiative Study. *Phys Ther.* 2015;95(7):955-965.
11. Andresen EM, Brownson RC. Disability and health status: Ethnic differences among women in the United States. *J Epidemiol Community Health.* 2000;54(3):200-206. doi: 10.1136/jech.54.3.200.
12. Colbert CJ, Almagor O, Chmiel JS, et al. Excess body weight and four-year function outcomes: Comparison of African Americans and whites in a prospective study of osteoarthritis. *Arthritis Care Res (Hoboken).* 2013;65(1):5-14.
13. Sternfeld B, Colvin A, Stewart A, et al. The effect of a healthy lifestyle on future physical functioning in midlife women. *Med Sci Sports Exerc.* 2017;49(2):274-282.
14. Dugan SA, Everson-Rose SA, Karavolos K, Sternfeld B, Wesley D, Powell LH. The impact of physical activity level on SF-36 role-physical and bodily pain indices in midlife women. *J Phys Act Health.* 2009;6(1):33-42.
15. Pettee Gabriel K, Sternfeld B, Colvin A, et al. Physical activity trajectories during midlife and subsequent risk of physical functioning decline in late mid-life: The Study of Women's Health across the Nation (SWAN). *Prev Med.* 2017;105:287-294.
16. Adams PF, Marano MA. Current estimates from the National Health Interview Survey, 1994. *Vital Health Stat 10.* 1995;(193 Pt 1):1-260.
17. Wittink H, Rogers W, Sukiennik A, Carr DB. Physical functioning: Self-report and performance measures are related but distinct. *Spine (Phila Pa 1976).* 2003;28(20):2407-2413.
18. Bean JF, Olveczky DD, Kiely DK, LaRose SI, Jette AM. Performance-based versus patient-reported physical function: What are the underlying predictors? *Phys Ther.* 2011;91(12):1804-1811.
19. Guralnik JM, Branch LG, Cummings SR, Curb JD. Physical performance measures in aging research. *J Gerontol.* 1989;44(5):M141-6.
20. Cress ME, Schechtman KB, Mulrow CD, Fiatarone MA, Gerety MB, Buchner DM. Relationship between physical performance and self-perceived physical function. *J Am Geriatr Soc.* 1995;43(2):93-101.
21. Oh-Park M, Wang C, Verghese J. Stair negotiation time in community-dwelling older adults: Normative values and association with functional decline. *Arch Phys Med Rehabil.* 2011;92(12):2006-2011.
22. Bendayan R, Cooper R, Wloch EG, Hofer SM, Piccinin AM, Muniz-Terrera G. Hierarchy and speed of loss in physical functioning: A comparison across older U.S. and English men and women. *J Gerontol A Biol Sci Med Sci.* 2017;72(8):1117-1122.
23. Chen HY, Wang CY, Lee MY, Tang PF, Chu YH, Suen MW. A hierarchical categorization of tasks in mobility disability. *Disabil Rehabil.* 2010;32(19):1586-1593.
24. Weiss CO, Fried LP, Bandeen-Roche K. Exploring the hierarchy of mobility performance in high-functioning older women. *J Gerontol A Biol Sci Med Sci.* 2007;62(2):167-173.
25. Sowers MF, Crawford SL, Sternfeld B, Morganstein D, Gold EB, Greendale GA. SWAN: A multi-center, multi-ethnic, community-based cohort study of women and the menopausal transition. In: Lobo RA, Kelsey J, Marcus R, eds. *Menopause: Biology and pathobiology.* San Diego, CA: Academic Press; 2000:175-188.
26. Kurina LM, Gulati M, Everson-Rose SA, et al. The effect of menopause on grip and pinch strength: results from the Chicago, Illinois, site of the Study of Women's Health across the Nation. *Am J Epidemiol.* 2004;160:484-491.
27. Sternfeld B, Ainsworth BE, Quesenberry CP. Physical activity patterns in a diverse population of women. *Prev Med.* 1999;28(3):313-323.
28. Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr.* 1982;36(5):936-42.
29. Radloff L. The CES-D scale: A self-report depression scale for research in the general population. *Appl Psychol Meas.* 1977;1(3):385-401.
30. Boyd JH, Weissman MM, Thompson WD, Myers JK. Screening for depression in a community sample. understanding the discrepancies between depression symptom and diagnostic scales. *Arch Gen Psychiatry.* 1982;39(10):1195-1200.
31. Mendes de Leon CF, Beckett LA, Fillenbaum GG, et al. Black-white differences in risk of becoming disabled and recovering from disability in old age: A longitudinal analysis of two EPESE populations. *Am J Epidemiol.* 1997;145(6):488-497.
32. Mendes de Leon CF, Barnes LL, Bienias JL, Skarupski KA, Evans DA. Racial disparities in disability: Recent evidence from self-reported and performance-based disability measures in a population-based study of older adults. *J Gerontol B Psychol Sci Soc Sci.* 2005;60(5):S263-71.
33. Thorpe RJ Jr, Koster A, Kritchevsky SB, et al. Race, socioeconomic resources, and late life mobility and decline: Findings from the Health, Aging, and Body Composition Study. *J Gerontol A Biol Sci Med Sci.* 2011;66(10):1114-1123.
34. Thorpe RJ Jr, Kasper JD, Szanton SL, Frick KD, Fried LP, Simonsick EM. Relationship of race and poverty to lower extremity function and decline: Findings from the Women's Health and Aging Study. *Soc Sci Med.* 2008;66(4):811-821.
35. Lange-Maia B, Strotmeyer E, Harris T, et al. Physical activity and change in long distance corridor walk performance in the Health, Aging and Body Composition Study. *J Am Geriatr Soc.* 2015;63(7):1348-54.
36. Fleg J, Morrell C, Bos A, et al. Accelerated longitudinal decline of aerobic capacity in healthy older adults. *Circulation.* 2005;112:674-682.
37. Chiu CJ, Wray LA. Physical disability trajectories in older Americans with and without diabetes: The role of age, gender, race or ethnicity, and education. *Gerontologist.* 2011;51(1):51-63.
38. Houston DK, Ding J, Nicklas BJ, et al. Overweight and obesity over the adult life course and incident mobility limitation in older adults: The Health, Aging and Body Composition Study. *Am J Epidemiol.* 2009;169(8):927-936.
39. Nightingale EJ, Pourkazemi F, Hiller CE. Systematic review of timed stair tests. *J Rehabil Res Dev.* 2014;51(3):335-350.
40. Oh-Park M, Perera S, Verghese J. Clinically meaningful change in stair negotiation performance in older adults. *Gait Posture.* 2012;36(3):532-536.
41. Bean JF, Kiely DK, LaRose S, Alian J, Frontera WR. Is stair climb power a clinically relevant measure of leg power impairments in at-risk older adults? *Arch Phys Med Rehabil.* 2007;88(5):604-9.
42. Ni M, Brown LG, Lawler D, Bean JF. Reliability, validity, and minimal detectable change of four-step stair climb power test in community-dwelling older adults. *Phys Ther.* 2017;97(7):767-773.
43. Silva AG, Queiros A, Cerqueira M, Rocha NP. Pain intensity is associated with both performance-based disability and self-reported disability in a sample of older adults attending primary health care centers. *Disabil Health J.* 2014;7(4):457-465.
44. Scudds RJ, Østbye T. Pain and pain-related interference with function in older Canadians: The Canadian Study of Health and Aging. *Disabil Rehabil.* 2001;23(15):654-664.
45. Weiner DK, Haggerty CL, Kritchevsky SB, et al. How does low back pain impact physical function in independent, well-functioning older adults? Evidence from the Health ABC cohort and implications for the future. *Pain Med.* 2003;4(4):311-320.