



ERRATUM

Erratum to: Same-day discharge after craniotomy for supratentorial tumour surgery: a retrospective observational single-centre study

Lashmi Venkatraghavan, MD · Suparna Bharadwaj, MD · Karolyn Au, MD ·
Mark Bernstein, MD · Pirjo Manninen, MD

Published online: 4 January 2017
© Canadian Anesthesiologists' Society 2017

**Erratum to: Can J Anesth/J Can Anesth (2016)
63:1245–1257
DOI 10.1007/s12630-016-0717-8**

In the article entitled: “Same-day discharge after craniotomy for supratentorial tumour surgery: a retrospective observational single-centre study” published in the November 2016 issue of the *Journal*, Can J Anesth 2016; 63: 1245-1257, in Table 3, next to *Osmotic therapy*, the values between *Inpatient* and *Same-day discharge* have been corrected and are highlighted in bold. The publisher apologizes most sincerely for this error.

The online version of the original article can be found under
doi:[10.1007/s12630-016-0717-8](https://doi.org/10.1007/s12630-016-0717-8).

L. Venkatraghavan, MD · S. Bharadwaj, MD ·
P. Manninen, MD
Department of Anesthesia and Pain Management, Toronto
Western Hospital, University of Toronto, Toronto, ON, Canada

L. Venkatraghavan, MD (✉)
Department of Anesthesia, Toronto Western Hospital University
Health Network, University of Toronto, 399 Bathurst St,
Toronto, ON M5T 2S8, Canada
e-mail: lashmi.venkatraghavan@uhn.on.ca

K. Au, MD · M. Bernstein, MD
Division of Neurosurgery, Toronto Western Hospital, University
of Toronto, Toronto, ON, Canada

Table 3 Perioperative management

Perioperative management	Inpatient (<i>n</i> =154)	Same-day discharge (<i>n</i> =175)	<i>P</i> value
Monitoring	103(67%)	23(13%)	<0.001
Invasive arterial blood pressure	6(3%)	0(0%)	0.01
Central venous pressure	89(58%)	11(6%)	<0.001
Urinary catheter			
Intraoperative fentanyl use (μg)	177(89)	81(56)	<0.001
Osmotic therapy	15(10%)	1(1%)	<0.001
Total fluid administered (mL)	2,370 (430)	1,345 (665)	<0.001
Hypotension (>20% from baseline)	106(69%)	114(65%)	0.48
Use of vasopressors	85(55%)	39(22%)	<0.001
Use of anti-hypertensive medication	84(55%)	8(5%)	<0.001
Estimated blood loss >300 mL	54(35%)	6(3%)	<0.001
Use of colloid	16(10%)	1(1%)	<0.001
Blood transfusion	7(4%)	0(0%)	0.004
Postoperative opioid consumption (morphine equivalent) (mg)	23(16)	16(17)	<0.001
Antiemetic use in PACU	84(55%)	39(22%)	<0.001

All values are *n* (%) or mean (SD), where indicated; *n* = number; PACU = postanesthesia care unit