

Erratum to: Total tumor volume predicts survival following liver resection in patients with hepatocellular carcinoma

Mu-xing Li¹ · Hong Zhao¹ · Xin-yu Bi¹ · Zhi-yu Li¹ · Zhen Huang¹ · Yue Han² ·
Jian-guo Zhou¹ · Jian-jun Zhao¹ · Ye-fan Zhang¹ · Wen-qiang Wei³ · Dong-bin Zhao¹ ·
Jian-qiang Cai¹

Published online: 4 March 2016

© International Society of Oncology and BioMarkers (ISOBM) 2016

Erratum to: Tumor Biol.

DOI 10.1007/s13277-016-4794-7

The original version of this article contained a mistake.

The same figures were published twice in the online version namely, figure 4E and figure 4F.

The corrected Figure 4 image is shown in the next page.

The online version of the original article can be found at <http://dx.doi.org/10.1007/s13277-016-4794-7>.

✉ Jian-qiang Cai
caijianqiang@cicams.ac.cn

¹ Department of Abdominal Surgical Oncology, Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College (CAMS and PUMC), No. 17, Panjiayuan Nanli, Beijing 100021, People's Republic of China

² Department of Interventional Therapies, Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College (CAMS and PUMC), No. 17, Panjiayuan Nanli, Beijing 100021, People's Republic of China

³ Department of Cancer Epidemiology, Cancer Hospital, Chinese Academy of Medical Sciences and Peking Union Medical College (CAMS and PUMC), No. 17, Panjiayuan Nanli, Beijing 100021, People's Republic of China




Fig. 4 Comparison of the survival distribution of the TTV-CLIP scoring system (a), CLIP scoring system (b), TNM staging system (c), BCLC staging system (d), Okuda staging system (e), and Milan criteria (f)