

Videophone-based multimodal home telecare support system for patients with diabetes

Masakazu Kubota · Kiminori Hosoda ·
Kyoko Eguchi · Ayana Furuya · Yuki Nishijima ·
Kazuwa Nakao · Ayae Kinoshita

Received: 17 April 2012 / Accepted: 4 September 2012 / Published online: 14 November 2012
© The Japan Diabetes Society 2012

Abstract

Aims We conducted an intervention study to clarify how effectively a telemedicine system using a multi-functional videophone could be used in lifestyle guidance, with special focus on diet for diabetic patients cared for at home.

Methods Patients were assigned to a 3-month intervention group or to a usual care group. In the intervention group, patients and a medical professional communicated bi-directionally through a videophone-based communication system for 30 min, once a week. The participants were encouraged to send pictures of each meal online, through the videophone system, in real time.

Results In intra-subjective comparison, the 3-month intervention program resulted in a significant decrease in body weight (BW) ($p < 0.0005$) and average HbA1C level ($p < 0.005$) compared with before the intervention program. Three months after the conclusion of the intervention program, average HbA1C levels returned to almost the same levels as before intervention ($p < 0.05$). In the usual care group, average HbA1C level did not change significantly for 6 months.

Conclusions 3-month intensive communication using the multimodal videophone system led to a significant decrease in BW and average HbA1C level. This novel bi-directional communication is useful for improving conditions such as diabetes, BW, and hyperglycemia for homecare diabetes patients, thus, reducing cardiovascular risk.

Keywords Videophone · Multimodal · Telecare · Homecare

Introduction

It is estimated that throughout the world the number of adults with diabetes will rise from 135 million in 1995 to 300 million in 2025 [1]. In Japan also, the prevalence of type 2 diabetes and prediabetes has been increasing in recent decades. In 2007, approximately 8.9 million people were estimated to have diabetes. Also, there were approximately 13.2 million people with possibilities of developing diabetes [2]. The pathogenesis of type 2 diabetes is characterized by increased peripheral insulin resistance and reduced insulin secretion, which leads to complications such as diabetic neuropathy, nephropathy, and retinopathy. Thus, it is important for patients with diabetes to prevent complications of diabetes by improving glycemic control through sequential therapy. To date, a sedentary lifestyle and a Westernized high-calorie diet, which result in overweight, have been considered to be associated with an increased risk of development of diabetes [3–5]. Therefore, the introduction of weight management, an energy-restricted diet, and a physically active lifestyle, is thought to be the primary approach to treatment of diabetes. Many studies have demonstrated that lifestyle intervention, including an energy-restricted diet, exercise

M. Kubota (✉) · K. Hosoda · A. Furuya · Y. Nishijima ·
A. Kinoshita
Department of Human Health Sciences, Graduate School of
Medicine, Kyoto University, 53 Shogoin Kawahara-cho,
Sakyo-ku, Kyoto 606-8507, Japan
e-mail: kubota.masakazu.5w@kyoto-u.ac.jp

K. Hosoda · K. Nakao
Department of Medicine and Clinical Science, Graduate School
of Medicine, Kyoto University, Kyoto 606-8507, Japan

K. Eguchi
Department of Geriatric Nursing, School of Nursing,
Osaka Prefecture University, Habikino 583-8555, Japan

habit, and cognitive behavior therapy, yield significant behavioral improvements, in addition to weight loss and better glycemic control [6–10].

In Japan, lifestyle education for diabetes is mainly conducted in hospitals. Patients with diabetes are hospitalized for education for a short time and given self-management training, which includes dietary intervention, exercise therapy, and drug therapy, under the direction of professionals such as doctors, nurses, and nutritionists in hospitals. After the in-hospital training program, the patients with diabetes are expected to observe the self-care program at home also. However, the brief training session is not sufficient to rectify their dietary behavior problems, with the hospital visits being only once a month or once in 2 months [11]. It can be challenging for patients to accept that it is worthwhile making significant lifestyle changes to tackle problems such as glycemic control, unbalanced diet, and sedentary lifestyle.

Recently, with the spread of internet-based technology, the number of people using information communication technology (ICT) devices has increased dramatically. ICT devices are gradually being introduced in the field of healthcare as a telemedicine system tool. Previous studies from our laboratory obtained evidence that videophone-based intensive communication is useful for cognitive rehabilitation of dementia patients and for the psychological care of their caregivers [12]. Because of their portability, ICT devices are now used in telemedicine to manage chronic diseases, including diabetes. Studies have evaluated weight maintenance and glycemic control programs, delivered via the Internet, telephone, and e-mail intervention [13–16]. Of these, two recent studies by Harvey-Berino et al. [13] and Noh et al. [16] have investigated whether medical intervention using ICT devices were effective in treating patients with diabetes. The former report indicated that the Internet was an effective vehicle for promoting long-term clinically significant levels of weight loss. The latter report showed that HbA1c and postprandial glucose levels were significantly reduced by web-based intervention 6 months post-initiation in patients with type 2 diabetes. Thus, ICT devices have tremendous potential as tools that offer health benefits. Therefore, we have to pursue the most effective ICT-based approach and learn how to maximize its use to help diabetes patients regain and maintain a balanced lifestyle. However, some of the patients, particularly elderly people, are likely to have difficulty in operating computer-based ICT devices. As type 2 diabetes is common in older adults [17], it may be expected that many may express great anxiety about using ICT devices—compared to younger adults [18]. Recently, innovative multi-functional devices have been developed as a new communication tool, equipped with not only a telephone but also an Internet-based videophone and other

multimodal functions. The main characteristic of multimodal videophone equipment is its user-friendliness. It can be easily operated by use of a touch panel, has a relatively large display, and requires no knowledge of computers. Considering the advantages of these innovative devices, we felt that an effective approach utilizing these tools in healthcare should be thoroughly evaluated.

We therefore conducted an intervention study to clarify how effectively a bi-directional communication system with multimodal functions could be used in lifestyle guidance, with special focus on diet, for diabetic patients cared for at home. On the basis of data collected by use of the real time videophone system, which can transmit information on body weight, diet record, and pictures of meals, we conclude that this new ICT-based intervention had a significant effect on control of diabetes, thus reducing the number of regular hospital visits. The interactive nature of this Internet-based communication will surely be an appealing mode of telemedicine in treating homecare patients in the future.

Materials and methods

Subjects

Participants diagnosed with type 2 diabetes were recruited from an outpatient section at the Department of Medicine and Clinical Science, Kyoto University Hospital. Twenty patients with diabetes were randomly assigned to a 3-month intervention group or to a usual care group (Table 1). Ten subjects with type 2 diabetes (63 ± 2 years, four males and six females) were invited to participate in this interventional study. The other ten patients (65 ± 3 years, four males and six females) were treated on an outpatient basis once every 1 or 2 months, as usual. Eligibility was assessed on the basis of medical history. Both intervention group and control group were presented to a diabetes specialist once every 1 or 2 months. Only telephonic participants were prompted by us to receive the ICT-based intervention. Before the recruitment and before they gave their voluntary consent to participate in the study, they were informed about the nature, purpose, and possible risks of the study. All the subjects with diabetes were treated with oral hypoglycemic agents and medical nutritional therapy. All the subjects had had stable BW for at least 3 months before their participation in the study. Exclusion criteria were: psychological problems, disordered recognition, impaired visual acuity, disabled hands restricting or prohibiting the use of the videophone system, impaired hearing. Participants who did not have a Nippon Telegraph and Telephone West Corporation's local cable system in their home were also excluded, because this was a requirement for videophone installation. The study protocol was

Table 1 Characteristics of the patients at baseline

	Age (years)	Sex (M/F)	BW (kg)	BMI (kg/m ²)	DM duration (years)	FBG (mg/dl)	HbA1C (%)	TG (mg/dl)	Total-CHO (mg/dl)	HDL-CHO (mg/dl)	LDL-CHO (mg/dl)
Intervention group	63 ± 2	4/6	69.0 ± 5.6	26.8 ± 1.5	14 ± 3	134.7 ± 12.5	7.4 ± 0.3	134.6 ± 24.9	183.7 ± 12.0	51.1 ± 4.5	100.1 ± 7.5
Usual care group	65 ± 3	4/6	61.5 ± 2.4	24.6 ± 0.9	16 ± 3	157.0 ± 18.7	7.5 ± 0.2	106.6 ± 20.7	185.3 ± 9.1	51.9 ± 4.0	101.8 ± 7.8

Values are mean ± SE for subjects with type 2 diabetes

approved by the ethics committee of Kyoto University Graduate School of Medicine (approval number E661) and all individuals provided written informed consent after the study purpose and design had been explained to them.

Telemedicine system

The telemedicine system used in this study consisted of a central hospital unit and peripheral units (at patients' homes), connected by multimodal videophones (Flet's phone VP1000; Nippon Telegraph and Telephone West Corporation, Japan) via optical cable. The multimodal videophone was installed at each patient's home before the nursing intervention was started. The main characteristic of the multimodal videophone equipment was its user-friendliness. It could be easily operated by use of a touch panel and had a relatively large display. Thus, it could be easily used even by elderly people. It could be operated without difficulty after a 30-min guidance session. Furthermore, this multimodal system had a function which was able to send and display pictures of each meal taken by the patients and a graph file of the BW data on a screen. The price of the multimodal videophone is 200 dollars and monthly telecommunication costs only a flat rate of 40 dollars.

Study design

Nguyen et al. [19] suggested that weight loss and obesity prevention are important medical intervention targets in the effort to reduce the effects of diabetes on health-care. According to the recommendations of the Japan Diabetes Society (JDS) and the Japan Society for Study of Obesity (JASSO), ideal daily calorific intake (kcal/day) is estimated from the formula: ideal body weight (IBW, kg) × 25–35 (kcal) depending on daily activity. IBW is defined as the weight that is equal to 22 times the square of the height (m) [20, 21]. Nakajima et al. [22] concluded that the diet therapy recommended by JDS and JASSO can be a useful and simple program for control of BW and hyperglycemia in daily life. On the basis of their instructions, we encouraged diabetic patients to eat properly, by communicating via videophone, with visual and live messages, and also using examples based on recommended calorific intake. We also encouraged physical activity.

For a period of 3 months, the patients in the intervention group were instructed to record their BW, contents of meals, and exercise every day. They were instructed to take and save digital photographs of every meal. During the intervention, they communicated bi-directionally on a one-to-one basis with a staff nurse, by use of the videophone system, for 30 min, once a week (Fig. 1). The nurse facilitated the patient's problem-solving abilities from the perspective of diabetes self-management, based on a graph

of weight history, drug compliance, and weekly exercise habit. Additionally, taking advantage of a multimodal system which can send and display pictures and a graph file of the BW data, the staff nurse checked the attached pictures of every meal; this was done jointly with the patients by use of the videophone system in real time. After assessing the adequacy of dietary intake, the staff nurse advised the patients on the amount and types of food that they should eat by drawing marks, noting the approximate calorie content of the meal, and by writing a few comments directly on the display (Fig. 2). The period of intervention was followed by 3 months of withheld intervention. The patients were instructed to return to their usual lifestyle in self-care while receiving outpatient treatment on a monthly basis during this period. During this time control patients were treated on an outpatient basis once every 1 or 2 months as usual.

Measurement

Medical diagnosis, duration of diabetes, height, fasting blood glucose level (FBG), HbA1c, total cholesterol, triglycerides (TG), low density lipoprotein-cholesterol (LDL-



Fig. 1 Situation of intercommunication between a patient with diabetes and a nurse through the videophone system. The patients in the intervention group communicated bi-directionally on a one-to-one basis with a staff nurse through the videophone system for 30 min, once a week

Fig. 2 A real screen with writing marks, calories, and a few words of caution written by a nurse so that a patient with diabetes learns naturally to have an appropriate diet. The staff nurse advised the patients on the amount and types of food that they should eat by drawing marks, noting the approximate calorie content of the meal, and by writing a few comments directly on the display



cholesterol), and high-density lipoprotein-cholesterol (HDL-cholesterol) at baseline were obtained from the patients' medical records. BMI was calculated by measuring weight in kilograms and height in meters, then using the formula weight (kg)/height (m²). Blood was collected after an overnight fast of at least 12 h, when the participants returned for their follow-up visit 1, 2, 3, and 6 months after the study initiation. Daily weight was measured before breakfast, without shoes or heavy clothing and using the same weighing scale in the patient's home.

Statistical analysis

All data were expressed as mean \pm SE. Data were analyzed by use of the Statistical Package for the Social Sciences for Windows, version 16.0 (SPSS Japan, Tokyo, Japan). All results were evaluated by use of the paired *t* test. A *p* value less than 0.05 was considered to be statistically significant.

Results

Baseline characteristics

All the patients, who were diagnosed with diabetes by the attending doctor, participated in this study. The patient information is listed in Table 1. All the patients had uncontrolled diabetes with high FBG levels (intervention, 134.7 ± 12.5 mg/dl; control, 157.0 ± 18.7 mg/dl) and HbA1c levels (intervention, 7.4 ± 0.3 %; control, 7.5 ± 0.2 %) despite being educated on the diet to be followed, exercise therapy, and the use of oral anti-diabetes drugs, as an outpatient, over the years. Most of the participants had normal TG levels (intervention, 134.6 ± 24.9 mg/dl; control, 106.6 ± 20.7 mg/dl), total cholesterol levels (intervention, 183.7 ± 12.0 mg/dl; control, 185.3 ± 9.1 mg/dl), HDL-cholesterol levels (intervention, 51.1 ± 4.5 mg/dl; control, 51.9 ± 4.0 mg/dl), and LDL-cholesterol levels (intervention, 100.1 ± 7.5 mg/dl; control, 101.8 ± 7.8 mg/dl) as shown in Table 1.

Effects of videophone-based intervention

During the session, once a week the staff nurse communicated with the patients on a one-to-one basis. Taking advantage of multimodal functions of the touch panel system of a videophone, the nurse examined and remediated contents of meals and food intake through the videophone system in real time. To provide nutritional advice, the nurse calculated the numbers of calories in the meals during the session and wrote them directly on the display (Fig. 2). On the basis of calorific intake and nutrition, the nurse advised the patients on the amount and types of food they should eat. In addition, the nurse showed a line graph of the change in their BW on the display and evaluated the effect of dietary and exercise habits. If the weight change was not satisfactory, the nurse analyzed the reason for this and encouraged the patient to follow adequate dietary and




Fig. 3 Effects of the bi-directional communication via the videophone system on average body weight. The 3-month intervention program resulted in a significant decrease in the body weight ($-3.3 \pm 0.6 \%$, $p < 0.0005$). Three months after termination of the intervention program, body weight had returned to almost the same levels as before intervention ($p < 0.005$). Values are mean \pm SE for ten subjects with type 2 diabetes. ** $p < 0.0005$ versus before intervention. * $p < 0.005$ versus 3 months after intervention

fluid intake. During the session, the patient was given advice on exercise also.

In intra-subject comparison, the 3-month intervention program resulted in a significant decrease in BW ($-3.3 \pm 0.6 \%$, $p < 0.0005$, Fig. 3), BMI (from 26.8 ± 1.5 to $25.9 \pm 1.5 \text{ kg/m}^2$, $p < 0.0005$, Table 2), and average HbA1c level (from 7.4 ± 0.3 to $6.8 \pm 0.2 \%$, $p < 0.005$, Fig. 4). Three months after termination of the intervention program (6 months after study initiation), average HbA1c level returned to almost the same levels as it was at the pre-intervention stage (from 6.8 ± 0.2 to $7.2 \pm 0.3 \%$, $p < 0.05$, Fig. 4). In usual care group, the average HbA1c level of the patients did not change significantly for 6 months, as shown in Fig. 4, which confirmed that this intervention program was effective in reducing HbA1c level.

The average FBG level had a tendency to decrease with intervention (from 134.7 ± 12.5 to $111.1 \pm 7.7 \text{ mg/dl}$, 18 % decrease, Table 2), although not significantly. The total plasma cholesterol and HDL-cholesterol levels did not change significantly before and after intervention. TG levels tended to decrease with intervention (from 134.6 ± 24.9 to $116.2 \pm 14.4 \text{ mg/dl}$, 14 % decrease, Table 2), although not statistically significantly.

Discussion

In this study, we used a telemedicine system using multimodal videophones to communicate bi-directionally, to create an environment in which diabetes patients can live without anxiety. The purpose of the study was to test the feasibility and efficacy of an intervention program delivered via a multimodal videophone. After a 3-month intervention program, BW, BMI, and average HbA1c level decreased significantly. From these results we conclude that our novel telemedicine system is a useful, effective,

Table 2 Characteristics of sample at baseline, 3 months after intervention, and 3 months after termination of intervention in the intervention group

	Before intervention	3 months after intervention	<i>p</i> value (vs. before intervention)	3 months after termination of intervention	<i>p</i> value (vs. 3 months after intervention)
Body weight (kg)	69 ± 5.6	66.9 ± 5.6	<0.0005	67.9 ± 5.7	<0.005
BMI (kg/m^2)	26.8 ± 1.5	25.9 ± 1.5	<0.0005	26.4 ± 1.5	<0.005
Fasting glucose (mg/dl)	134.7 ± 12.5	111.1 ± 7.7	0.1288	120.2 ± 15	0.5969
HbA1C (%)	7.4 ± 0.3	6.8 ± 0.2	<0.005	7.2 ± 0.3	<0.05
Triglyceride (mg/dl)	134.6 ± 24.9	116.2 ± 14.4	0.4133	158.2 ± 30.9	0.0637
Total-CHO (mg/dl)	183.7 ± 12	190.5 ± 12	0.2213	184.7 ± 12.4	0.5012
HDL-CHO (mg/dl)	51.1 ± 4.5	54.8 ± 5.1	0.2349	54.7 ± 5.1	0.9714
LDL-CHO (mg/dl)	100.1 ± 7.5	106.1 ± 8.2	0.2269	101.1 ± 8.5	0.3786

Values are mean \pm SE for ten subjects with type 2 diabetes




Fig. 4 Effects of the bi-directional communication via the videophone system on average HbA1c level. *Filled diamonds*, the 3-month intervention program resulted in a significant decrease in average HbA1c level (from 7.4 ± 0.3 to 6.8 ± 0.2 %, $p < 0.005$). Three months after termination of the intervention program, average HbA1c levels returned to almost the same levels as before intervention (from 6.8 ± 0.2 to 7.2 ± 0.3 %, $p < 0.05$). *Filled squares*, changes in HbA1c level of the control patients who were treated on an outpatient basis as usual. The average HbA1c level of the patients did not change significantly for 6 months. Values are mean \pm SE. ** $p < 0.005$ versus before intervention. * $p < 0.05$ versus 3 months after intervention

and low-cost intervention that can improve the quality of life of patients with diabetes.

All participants in this study had diabetes that had been poorly controlled for many years despite repeated hospitalization for patient education about the condition under the direction of medical professionals. Thus, it is plausible that this novel intervention led to more effective glycemic control than the conventional approach, because of the reasons discussed below.

One reason for the effectiveness of this study was that the program led to increased awareness of a good lifestyle among participants by making them record BW, compliance checks, walk counts, and remarks every day. Additionally, sharing the information with a health care professional once a week led to a decrease in BW, BMI, and HbA1c level. This is in line with literature showing that raising awareness of the disease and self-monitoring could have a beneficial effect on glycemic control [23, 24]. Our multimodal videophone system, which is able to illustrate change in BW graphically, also led the participants to success in self-management, because they could recognize the benefits and merits from visual feedback. We speculate that this intervention program was effective in increasing awareness of a good lifestyle—not only by making participants document their daily records but also by convincing them that their lifestyle strongly correlates with their health status by showing a graph every time.

Another reason for the effectiveness of the program was the use of photographs of the meals the participants actually ate. The videophone was able to display attached photographs of the meals and to add a note of caution on the touch

panel in real time. Because we advised a patient with diabetes on the amount and types of food he or she ate by giving marks, calories, and adding a few words of caution on the screen directly, we could take realistic and specific approaches in dietary intervention. The patients could easily understand how to select and combine the food to control calorie intake, by receiving individual guidance using the attached pictures of the meals. Thus, they could set a realistic objective for weight loss. Providing patients with personalized and specific advice from diabetes care specialists gives rise to a beneficial result [25]. In this regard, use of the videophone system, which focuses on patients' education for self-management, provided an appropriate opportunity for patients with diabetes who have difficulty maintaining adequate glucose levels at home.

The third reason for the success of the program was the continuous intensive communication with a member of the medical staff for 3 months. There is no easy way to change lifestyle over the years, and maintenance of desirable blood glucose levels is left to the self-management ability of the individual. Considering that type 2 diabetes patients return for follow-up visits once every 1 or 2 months, they must monitor their blood glucose levels without medical professionals' help in the interval between visits. Previous studies have already revealed the effectiveness of frequent intervention. Pimazoni-Netto et al. [26] showed that within a short time there is rapid improvement of glycemic control in diabetics using weekly intensive multifactorial intervention. Our current results are consistent with this research. However, regarding the frequency of hospital visits, the usual care group presented had several visits to a clinic during the study, whereas the intervention group had weekly videophone sessions. Thus, it may be considered that the usual care group did not serve as an appropriate control, which is a limitation of our study. In the literature, similar comparative studies in which control groups were treated differently from the intervention group [26] have been reported. Therefore, our comparison can be regarded as standard. Further, the different frequency of intervention is a major aspect of our study, in which the novelty of the research is that home-care diabetics can receive helpful guidance from a member of the medical staff without having to make frequent hospital visits. As we would like to show the possibility that the multimodal videophone system is helpful as a means of providing lifestyle guidance at outpatient clinics, we selected diabetics with regular hospital visits as controls. The videophone-based intensive communication made it possible for the patients with diabetes who only visit the hospital once a month or once in 2 months to continuously acquire sufficient knowledge and encouragement by communicating with a health care professional. The ICT-based videophone system made frequent counseling available to home-care diabetes patients.

In this study, we made a special effort to instruct participants to adhere to an appropriate diet because diet therapy is an important approach for patients with diabetes, especially obese patients. The risk of developing diabetes seems to increase with increased BMI [27, 28]. Thus, keeping a daily calorie count and any decrease in BMI toward the recommended range are expected to improve glycemic control. Our results support the theory of a correlation between reduced BMI and improved glycemic control. For participants in the intervention group, the average HbA1c level improved in parallel with reduced BMI, compared with readings at the start of the study. Additionally, the average HbA1c level 3 months after termination of the intervention program (6 months after study initiation), returned to almost the same level as it was at the pre-intervention stage. Taken together, these results reveal that encouraging diabetic patients to become more aware of weight reduction is a realistic and effective approach.

After the intervention, many participants expressed a feeling of satisfaction with this videophone system. Some of the opinions include: “talking to the medical staff frequently through the videophone system played an important role in resolving issues regarding the diet, especially those regarding dairy products, on the spot.” “We could easily understand how to select and combine the food to control calorie intake in our real life by receiving individual guidance using the attached pictures of the meals.” “This videophone system is a helpful tool because it is hard for elderly people to visit the hospital frequently to receive outpatient treatment.” From these comments, we believe that this videophone system can be useful from viewpoint of the patients.

There was no significant reduction of the average FBG level, although average HbA1c level decreased significantly. This may be because HbA1c is believed to reflect average blood glucose over the previous 12-week period better than FBG [29, 30]. Alternatively, this gives rise to the possibility that postprandial hyperglycemia may be improved by this intervention program. As stated above, we made a special effort to instruct participants to adhere to an appropriate well-balanced low-carbohydrate and low-fat diet which included many vegetables. It is known that an appropriate diet, especially low-carbohydrate, improves postprandial blood glucose control [5, 31]. However, it is not clear whether the intervention actually affected postprandial hyperglycemia which was not measured in this study. This possibility should be pursued in any future study.

TG levels tended to decrease after the 3-month intervention program, although not significantly. We suppose that longer intervention periods might induce a significant decrease in this.

The system used in this study may or may not be useful for a long-term educational approach, because in this study we did not focus on factors which may have long-term effects. To obtain long-term favorable effects, we suggest that this videophone system should be continued for a longer time even though the duration and frequency of communication are yet to be decided. We should consider using it to maintain a favorable effect for a longer period; we are now planning to implement the next intervention program.

In the future, the telemedicine approach will contribute to cost savings if this system is covered by insurance and used in daily practice. Currently, because there are approximately 550,000 latent nurses in Japan, we assume they may be able to work as health care professionals who stay at home and communicate with patients through the Internet.

In conclusion, we showed that 3-month bi-directional intensive communication using the videophone-based system resulted in a significant decrease in BW, BMI, and average HbA1c level. Bi-directional communication using multimodal videophones can be useful as an auxiliary therapy for switching to a good lifestyle.

The results of the study show a potential strategy for diabetes care, which can be used as an alternative approach to conventional hospital visits in the near future.

Acknowledgments We thank Hideyo Tsutsui from the Department of Sports Medicine, the University of Nagoya, for her valuable comments. We also thank the patients and their families for participating in this study. A part of this study was supported by the Grant-in-Aid for Young Scientists (B) (grant no. 21792305) from the Japanese Ministry of Education, Culture, Sports, Science, and Technology (MEXT) and a research grant from the UNIVERS foundation.

References

1. King H, Aubert RE, Herman WH. Global burden of diabetes, 1995–2025: prevalence, numerical estimates, and projections. *Diabetes Care*. 1998;21(9):1414–31.
2. Ministry of Health LaW. Outline of results from 2007 National Health and Nutrition Survey. Tokyo.
3. Chan JCN, Malik V, Jia WP, Kadowaki T, Yajnik CS, Yoon KH, et al. Diabetes in Asia epidemiology, risk factors, and pathophysiology. *J Am Med Assoc*. 2009;301(20):2129–40.
4. Schmidt MI, Duncan BB. Diabesity: an inflammatory metabolic condition. *Clin Chem Lab Med*. 2003;41(9):1120–30.
5. Kubota M, Koshinaka K, Kawata Y, Koike T, Oshida Y. Effects of continuous low-carbohydrate diet after long-term exercise on GLUT-4 protein content in rat skeletal muscle. *Horm Metab Res*. 2008;40(1):24–8.
6. Kubota M, Nagasaki M, Tokudome M, Shinomiya Y, Ozawa T, Sato Y. Mechanical horseback riding improves insulin sensitivity in elder diabetic patients. *Diabetes Res Clin Pract*. 2006;71(2): 124–30.
7. Klein S, Sheard NF, Pi-Sunyer X, Daly A, Wylie-Rosett J, Kulkarni K, et al. Weight management through lifestyle modification for the prevention and management of type 2 diabetes:

- rationale and strategies. *Diabetes Care*. 2004;27(8):2067–73.
8. Matvienko OA, Hoehns JD. A lifestyle intervention study in patients with diabetes or impaired glucose tolerance: translation of a research intervention into practice. *J Am Board Fam Med*. 2009;22(5):535–43.
 9. Wing RR, Blair EH, Bononi P, Marcus MD, Watanabe R, Bergman RN. Caloric restriction per se is a significant factor in improvement in glycemic control and insulin sensitivity during weight-loss in obese NIDDM patients. *Diabetes Care*. 1994;17(1):30–6.
 10. Pi-Sunyer FX, Maggio CA, McCarron DA, Reusser ME, Stern JS, Haynes RB, et al. Multicenter randomized trail of a comprehensive prepared meal program in type 2 diabetes. *Diabetes Care*. 1999;22(2):191–7.
 11. Shibayama T, Kobayashi K, Takano A, Kadowaki T, Kazuma K. Effectiveness of lifestyle counseling by certified expert nurse of Japan for non-insulin-treated diabetic outpatients: a 1-year randomized controlled trial. *Diabetes Res Clin Pract*. 2007;76(2):265–8.
 12. Hori M, Kubota M, Ando K, Kihara T, Takahashi R, Kinoshita A. The effect of videophone communication (with Skype and webcam) for elderly patients with dementia and their caregivers. *Gan To Kagaku Ryoho*. 2009;35(1):36–8.
 13. Harvey-Berino J, Pintauro S, Buzzell P, Gold EC. Effect of Internet support on the long-term maintenance of weight loss. *Obes Res*. 2004;12(2):320–9.
 14. Eakin EG, Reeves MM, Marshall AL, Dunstan DW, Graves N, Healy GN, et al. Living well with diabetes: a randomized controlled trial of a telephone-delivered intervention for maintenance of weight loss, physical activity and glycaemic control in adults with type 2 diabetes. *BMC Public Health*. 2010;10.
 15. Sherwood NE, Jeffery RW, Pronk NP, Boucher JL, Hanson A, Boyle R, et al. Mail and phone interventions for weight loss in a managed-care setting: weigh-to-be 2-year outcomes. *Int J Obes*. 2006;30(10):1565–73.
 16. Noh JH, Cho YJ, Nam HW, Kim JH, Kim DJ, Yoo HS, et al. Web-based comprehensive information system for self-management of diabetes mellitus. *Diabetes Technol Ther*. 2010;12(5):333–7.
 17. Ueno M, Takeuchi S, Oshiro A, Shinada K, Ohara S, Kawaguchi Y. Association between diabetes mellitus and oral health status in Japanese adults. *Int J Oral Sci*. 2010;2(2):82–9.
 18. Czaja SJ, Charness N, Fisk AD, Hertzog C, Nair SN, Rogers WA, et al. Factors predicting the use of technology: findings from the Center for Research and Education on Aging and Technology Enhancement (CREATE). *Psychol Aging*. 2006;21(2):333–52.
 19. Nguyen NT, Nguyen XMT, Lane J, Wang P. Relationship between obesity and diabetes in a US adult population: findings from the National Health and Nutrition Examination Survey, 1999–2006. *Obes Surg*. 2011;21(3):351–5.
 20. Society JD. Guideline for treatment of diabetes mellitus (http://www.jds.or.jp/jds_or_jp0/data/guideline/021-027.pdf).
 21. Obesity JSFTSO. Guideline for treatment of obesity 2006, Japan society for the study of obesity, Tokyo, 2006.
 22. Nakajima Y, Sato K, Sudo M, Nagao M, Kano T, Harada T, et al. Practical dietary calorie management, body weight control and energy expenditure of diabetic patients in short-term hospitalization. *J Atheroscler Thromb*. 2010;17(6):558–67.
 23. Reynolds LR, Anderson JW. Practical office strategies for weight management of the obese diabetic individual. *Endocr Pract*. 2004;10(2):153–9.
 24. Chan WB, Chan JC, Chow CC, Yeung VT, So WY, Li JK, et al. Glycaemic control in type 2 diabetes: the impact of body weight, beta-cell function and patient education. *QJM*. 2000;93(3):183–90.
 25. Iizuka Y, Ohashi K, Tsukamoto K, Ueki K, Kadowaki T. Education and management of diabetes at The University of Tokyo Hospital. *J Diabetes*. 2011;3(2):104–8.
 26. Pimazoni-Netto A, Rodbard D, Zanella MT, Grp DEC. Rapid improvement of glycemic control in type 2 diabetes using weekly intensive multifactorial interventions: structured glucose monitoring, patient education, and adjustment of therapy—a randomized controlled trial. *Diabetes Technol Ther*. 2011;13(10):997–1004.
 27. Colditz GA, Willett WC, Stampfer MJ, Manson JE, Hennekens CH, Arky RA, et al. Weight as a risk factor for clinical diabetes in women. *Am J Epidemiol*. 1990;132(3):501–13.
 28. Knowler WC, Pettitt DJ, Savage PJ, Bennett PH. Diabetes incidence in Pima Indians: contributions of obesity and parental diabetes. *Am J Epidemiol*. 1981;113(2):144–56.
 29. Rohlfing CL, Wiedmeyer HM, Little RR, England JD, Tennill A, Goldstein DE. Defining the relationship between plasma glucose and HbA(1c): analysis of glucose profiles and HbA(1c) in the Diabetes Control and Complications Trial. *Diabetes Care*. 2002;25(2):275–8.
 30. Goldstein DE, Little RR, Wiedmeyer HM, England JD, McKenzie EM. Glycated hemoglobin: methodologies and clinical applications. *Clin Chem*. 1986;32(10 Suppl):B64–70.
 31. De Natale C, Annuzzi G, Bozzetto L, Mazzarella R, Costabile G, Ciano O, et al. Effects of a plant-based high-carbohydrate/high-fiber diet versus high-monounsaturated fat/low-carbohydrate diet on postprandial lipids in type 2 diabetic patients. *Diabetes Care*. 2009;32(12):2168–73.