



Correction to: A candidate gene study of intermediate histopathological phenotypes in HIV-associated neurocognitive disorders

Andrew J. Levine¹ · Virawudh Soontornniyomkij² · Eliezer Masliah³ · Janet S. Sinsheimer^{4,5} · Sarah S. Ji⁵ · Steve Horvath^{5,6} · Elyse J. Singer¹ · Asha Kallianpur⁷ · David J. Moore²

Published online: 6 July 2020
© Journal of NeuroVirology, Inc. 2020

Correction to: Journal of NeuroVirology
<https://doi.org/10.1007/s13365-020-00846-z>

Due to a production error data in Table 1 were not presented correctly. The corrected table is shown below.

The online version of the original article can be found at <https://doi.org/10.1007/s13365-020-00846-z>

✉ Andrew J. Levine
ajlevine@mednet.ucla.edu

¹ Department of Neurology, David Geffen School of Medicine at the University of California, Los Angeles, CA, USA

² Department of Psychiatry, University of California San Diego School of Medicine, San Diego, CA, USA

³ Departments of Neurosciences and Pathology, University of California San Diego School of Medicine, San Diego, CA, USA

⁴ Departments of Human Genetics and Computational Biology, David Geffen School of Medicine at the University of California, Los Angeles, CA, USA

⁵ Department of Biostatistics, UCLA Fielding School of Public Health, Los Angeles, CA, USA

⁶ Department of Human Genetics, David Geffen School of Medicine at the University of California, Los Angeles, CA, USA

⁷ Department of Molecular Medicine, Genomic Medicine, Medicine, & Pediatrics, Cleveland Clinic/Lerner Research Institute, Cleveland Clinic Lerner College of Medicine of Case Western Reserve University, Cleveland, OH, USA

Age at death	47.4 years ^a (SD = 9.2)
Length of HIV infection	12 years (SD = 6.4)
CD4+ T cell count	122 (SD = 168)
Median plasma HIV-1 RNA viral load	15,406 copies/mL
CPE	8.96 (SD = 4.08)
Global clinical rating	5.31 (SD = 1.87)
	Percent ^b of sample (<i>n</i>)
Detectable (>50 copies/mL) plasma HIV-1 RNA viral load	84% (30)
Male	81.4% (162)
Race/ethnicity	
Caucasian	50.8% (100)
African American	24.9% (49)
Hispanic	22.3% (44)
Asian/Native Alaskan	2% (4)
Major depression	
Current	23.4% (32)
Past	16.8% (23)
Substance use disorder	
Current	20.3% (31)
Past	29.4% (45)
Alcohol use disorder	
Current	9% (14)
Past	26% (40)
HIV encephalitis	9.2% (16)
HAND	74.2% (135)
ANI	14.2% (26)
MND	31% (56)
HAD	29% (53)

SD standard deviation, *CPE* central nervous system penetration or effectiveness, *ANI* asymptomatic

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.