



Correction: Productivity and biological N₂-fixation in cereal-cowpea intercropping systems in sub-Saharan Africa. A review

Talent Namatsheve¹ · Rémi Cardinael^{1,2,3}  · Marc Corbeels^{2,3,4} · Regis Chikowo^{1,5}

Accepted: 30 May 2024 / Published online: 11 July 2024
© INRAE and Springer-Verlag France SAS, part of Springer Nature 2024

Correction: Agronomy for Sustainable Development (2020) 40:30
<https://doi.org/10.1007/s13593-020-00629-0>

The original online version of this article was revised: In Figure 2, right panel, for the sorghum-cowpea and millet-cowpea systems, the yields of the cereal instead of the yields of the cowpea were plotted.

The original article can be found online at <https://doi.org/10.1007/s13593-020-00629-0>.

✉ Rémi Cardinael
remi.cardinael@cirad.fr

¹ Crop Science Department, University of Zimbabwe, Box MP167, Mt. Pleasant, Harare, Zimbabwe


² CIRAD, UPR AIDA, Harare, Zimbabwe

³ AIDA, CIRAD, Université Montpellier, Montpellier, France


⁴ CIMMYT, Sustainable Intensification Program (SIP), P.O. Box 1041–00621, Gigiri, Nairobi, Kenya

⁵ Plant, Soil and Microbial Sciences Department, Michigan State University, East Lansing, MI 48824, USA

Incorrect Figure 2:



Correct Figure 2:



The original article has been corrected.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.