

Derrisisoflavones H–K and One Isoflavan Derivative from *Derris robusta*




Guo-Zhu Wei · Mei-Fen Mao · Xiang-Mei Li ·
Fu-Cai Ren · Fei Wang

Received: 7 January 2016 / Accepted: 31 January 2016 / Published online: 19 February 2016
© The Author(s) 2016. This article is published with open access at Springerlink.com

Abstract Four hitherto unknown prenylated isoflavonoids, named derrisisoflavones H–K (**1–4**) and one new isoflavan, namely 6-hydroxyisosativan (**5**), were isolated from the ethanol extract of *Derris robusta*. Their structures were elucidated on the basis of extensive spectroscopic studies. To our knowledge, derrisisoflavones J (**3**) and K (**4**) are the first examples of hydroxyethylated isoflavonoid.

Graphical Abstract



Keywords *Derris robusta* · Isoflavonoid · Isoflavan · Derrisisoflavone

1 Introduction

Derris is a genus belonging to the Leguminosae family with about 800 species that are widely distributed in

Electronic supplementary material The online version of this article (doi:10.1007/s13659-016-0090-x) contains supplementary material, which is available to authorized users.

G.-Z. Wei · M.-F. Mao · X.-M. Li · F.-C. Ren · F. Wang (✉)
BioBioPha Co., Ltd., Kunming 650201,
People's Republic of China
e-mail: f.wang@mail.biobioph.com

tropical, subtropical areas of Asia and Africa [1]. Published studies have shown that the genus is a rich source of pterocarpan, flavonoids, particularly prenylated isoflavonoids and flavonoids [2–5] and these phytochemicals are associated with a broad spectrum of biological activities, including insecticidal, antimicrobial, cytotoxic, and antioxidant activities [3–7]. As part of a BioBioPha [http://www.chemlib.cn/] objective to assemble a large-scale natural product library valuable in the discovery of new drug leads from nature [8–10], the phytochemical investigation on the twigs and leaves of *Derris robusta* led to the

isolation of four new prenylated isoflavonoids, named derrisisoflavones H–K (**1–4**), and a new isoflavan, namely 6-hydroxyisosativan (**5**). This paper describes the isolation and structural elucidation of five new compounds (Fig. 1).

2 Results and Discussion

Compound **1** was obtained as a yellow amorphous powder and had a molecular formula $C_{26}H_{26}O_7$ determined by its HRESIMS, showing a negative molecular ion peak at m/z 449.1608 $[M - H]^-$ (calcd. for $C_{26}H_{25}O_7$, 449.1606). The UV spectrum of **1** with a set of absorption maxima at 268, 297 (sh), 359 nm suggested that it had an isoflavone skeleton as the chromophore [11]. This inference was further supported by characteristic proton singlet at δ_H 8.17 (H-2) and sp^2 methine

carbon at δ_C 155.6 (C-2). The 1H NMR spectrum (Table 1) showed a set of signals at δ_H 7.05 (br. s), 6.87 (br. d, $J = 8.0$ Hz) and 6.82 (d, $J = 8.0$ Hz) due to a 1,3,4-trisubstituted benzene ring, two aromatic or olefinic protons at δ_H 6.86 (s) and 5.30 (t, $J = 6.8$ Hz), one methylene signal at δ_H 3.64 (d, $J = 6.8$ Hz), and five methyl singlets at δ_H 3.11, 1.85, 1.67, 1.60 and 1.60. The ^{13}C NMR (DEPT) spectrum (Table 2) displayed a total of 26 carbon resonances, including five methyls, one methylene, six methines and 14 quaternary carbons. The above NMR spectroscopic features were very similar to those of 5,4'-dihydroxy-8-(3,3-dimethylallyl)-2''-methoxyisopropylfuran[4,5:6,7]isoflavone [12], and the most dramatic difference was the presence of an additional hydroxy group in **1**. The hydroxy group was located at C-3', on the basis of the HMBC correlations from the protons at δ_H 7.05 (br. s, H-2') and 6.87 (br. d, $J = 8.0$ Hz, H-6') to the




Fig. 1 Structures of compounds **1–5**

Table 1 ^1H NMR spectroscopic data for derrisoflavones H–K (**1–4**) and 6-hydroxyisositivan (**5**)

No.	1 ^a	2 ^b	3 ^a	4 ^a	5 ^a
2	8.17 (s)	8.51 (s)	7.97 (s)	8.00 (s)	4.17 (ddd, 10.2, 3.3, 1.9, H _{eq}) 3.90 (t, 10.2, H _{ax})
3					3.43 (dddd, 10.7, 10.2, 5.3, 3.3, H _{ax})
4					2.91 (dd, 15.9, 10.7, H _{ax}) 2.73 (ddd, 15.9, 5.3, 1.9, H _{eq})
5					6.51 (s)
8			6.37 (s)	6.36 (s)	6.37 (s)
2'	7.05 (br. s)	7.05 (d, 1.6)	7.19 (d, 2.0)	7.25 (d, 2.3)	
3'					6.38 (d, 2.5)
5'	6.82 (d, 8.0)	6.81 (d, 8.0)	6.79 (d, 8.2)	6.82 (d, 8.2)	6.36 (dd, 8.4, 2.5)
6'	6.87 (br. d, 8.0)	6.85 (dd, 8.0, 1.6)	7.15 (dd, 8.2, 2.0)	7.21 (dd, 8.2, 2.3)	6.95 (d, 8.4)
1''	6.86 (s)	3.48 (d, 7.3)	2.92 (t, 7.3)	3.30 (overlapped)	
2''		5.29 (t, 7.3)	3.68 (t, 7.3)	5.22 (t, 7.2)	
4''	1.60 (s)	1.62 (s)		1.65 (s)	
5''	1.60 (s)	1.79 (s)		1.77 (s)	
1'''	3.64 (d, 6.8)	7.07 (s)	3.32 (d, 7.3)	2.87 (t, 7.0)	
2'''	5.30 (t, 6.8)		5.33 (t, 7.3)	3.78 (t, 7.0)	
4'''	1.67 (s)	1.56 (s)	1.72 (s)		
5'''	1.85 (s)	1.56 (s)	1.72 (s)		
5-OH		13.21 (s)			
3'-OH		9.14 (br. s)			
4'-OH		9.14 (br. s)			
7-OCH ₃					3.78 (s)
4'-OCH ₃					3.71 (s)
3''-OCH ₃	3.11 (s)				
3'''-OCH ₃		3.00 (s)			

^a Measured in CD₃OD (δ_{H} 3.30 ppm)

^b Measured in DMSO-*d*₆ (δ_{H} 2.50 ppm)

carbon at δ_{C} 123.7 (s, C-3) (Fig. 2). The HMBC correlations from the protons at δ_{H} 3.64 (2H, d, $J = 6.8$ Hz, H-1''') to the carbons at δ_{C} 158.6 (s, C-7), 105.2 (s, C-8), and 152.4 (s, C-9) verified the location of the prenyl group [δ_{H} 1.67, 1.85 (each s), 3.64 (d, $J = 6.8$ Hz), and 5.30 (t, $J = 6.8$ Hz)] at C-8. Furthermore, the correlations from the proton at δ_{H} 6.86 (s, H-1'') to the carbons at δ_{C} 154.2 (s, C-5), 114.1 (s, C-6), and 158.6 (s, C-7) confirmed that the furan ring, derived from a prenyl group, was fused along the C-6 to C-7 bond. Consequently, the structure of **1** was determined and named as derrisoflavone H.


Compound **2**, was purified as a yellow amorphous powder and had the same molecular formula as **1** based on its HRESIMS (neg.): m/z 449.1607 [$\text{M} - \text{H}$]⁻ (calcd. for C₂₆H₂₅O₇, 449.1606). The NMR spectroscopic data of **2** (Tables 1, 2) were very similar to those of **1**. The structural discrepancy was only from the switch positions of prenyl group and furan ring on ring A. The positions were verified by the HMBC correlations from the protons at δ_{H} 3.48 (2H, d, $J = 7.3$ Hz, H-1'') and 13.21 (s, 5-OH) to the carbon at

δ_{C} 154.6 (s, C-5), and from the protons at δ_{H} 7.07 (s, H-1''') and 8.51 (s, H-2) to the carbon at δ_{C} 147.7 (s, C-9), respectively. Accordingly, the structure of **2** was elucidated as shown and given the name derrisoflavone I.

Compound **3**, was isolated as a white amorphous powder, with a molecular formula of C₂₂H₂₂O₆ according to its HRESIMS (pos.): m/z 405.1305 [$\text{M} + \text{Na}$]⁺ (calcd. for C₂₂H₂₂O₆Na, 405.1309). The general features of NMR spectra (Tables 1, 2) of **3** were similar to those of lupalbigenin, a diprenylated isoflavone [13], except for the signals of a hydroxyethyl moiety [δ_{H} 2.92 (t, $J = 7.3$ Hz), 3.68 (t, $J = 7.3$ Hz); δ_{C} 26.8 (t), 61.9 (t)] instead of one of prenyl group in the latter. The hydroxyethyl group was linked to C-6 on the basis of the following HMBC correlations: from the protons at δ_{H} 2.92 (t, $J = 7.3$ Hz, H-1'') to the carbons at δ_{C} 161.3 (s, C-5), 109.9 (s, C-6) and 164.2 (s, C-7), and from the protons at δ_{H} 7.97 (s, H-2) and 6.37 (s, H-8) to the carbon at δ_{C} 157.9 (s, C-9). Similarly, the connection of the prenyl group to C-3' was established by the correlation from the proton at δ_{H} 7.19 (d, $J = 2.0$ Hz,

Table 2 ^{13}C NMR spectroscopic data for derrisisoflavones H–K (1–4) and 6-hydroxyisositivan (5)

No.	1 ^a	2 ^b	3 ^a	4 ^a	5 ^a
2	155.6 (d)	154.1 (d)	154.5 (d)	154.6 (d)	71.0 (t)
3	123.7 (s)	123.5 (s)	125.0 (s)	124.7 (s)	33.1 (d)
4	184.2 (s)	181.4 (s)	182.4 (s)	182.3 (s)	31.4 (t)
5	154.2 (s)	154.6 (s)	161.3 (s)	160.5 (s)	116.4 (d)
6	114.1 (s)	107.6 (s)	109.9 (s)	113.1 (s)	141.0 (s)
7	158.6 (s)	157.0 (s)	164.2 (s)	163.7 (s)	148.1 (s)
8	105.2 (s)	107.9 (s)	94.2 (d)	93.9 (d)	101.4 (d)
9	152.4 (s)	147.7 (s)	157.9 (s)	157.6 (s)	148.8 (s)
10	107.6 (s)	107.4 (s)	106.2 (s)	106.1 (s)	115.0 (s)
1'	123.7 (s)	121.5 (s)	123.4 (s)	123.4 (s)	121.4 (s)
2'	117.5 (d)	116.8 (d)	131.4 (d)	132.8 (d)	157.2 (s)
3'	146.2 (s)	145.0 (s)	129.5 (s)	126.7 (s)	102.3 (d)
4'	146.8 (s)	145.8 (s)	156.5 (s)	157.0 (s)	160.8 (s)
5'	116.3 (d)	115.5 (d)	115.8 (d)	116.1 (d)	105.6 (d)
6'	121.8 (d)	120.3 (d)	128.7 (d)	129.5 (d)	128.8 (d)
1''	102.7 (d)	21.6 (t)	26.8 (t)	22.3 (t)	
2''	161.3 (s)	120.9 (d)	61.9 (t)	123.4 (d)	
3''	74.7 (s)	132.1 (s)		132.1 (s)	
4''	25.5 (q)	25.5 (q)		26.0 (q)	
5''	25.5 (q)	17.6 (q)		17.9 (q)	
1'''	22.9 (t)	101.2 (d)	29.3 (t)	35.1 (t)	
2'''	122.2 (d)	159.3 (s)	123.9 (d)	63.0 (t)	
3'''	133.8 (s)	72.8 (s)	133.1 (s)		
4'''	25.9 (q)	24.9 (q)	25.9 (q)		
5'''	18.0 (q)	24.9 (q)	17.9 (q)		
7-OCH ₃					56.3 (q)
4'-OCH ₃					55.5 (q)
3''-OCH ₃	51.4 (q)				
3'''-OCH ₃		50.4 (q)			

^a Measured in CD₃OD (δ_{C} 49.0 ppm)^b Measured in DMSO-*d*₆ (δ_{C} 39.5 ppm)**Fig. 2** Selected HMBC (↔) correlations of derrisisoflavone H (1)

H-2') to the carbon at 29.3 (t, C-1'''). Therefore, the structure of **3** was characterized and named as derrisisoflavone J.

Compound **4** was afforded as a white amorphous powder and possessed the same molecular formula as **3** according to its HRESIMS (pos.): m/z 405.1307 [$\text{M} + \text{Na}$]⁺ (calcd.

for C₂₂H₂₂O₆Na, 405.1309). The NMR data (Tables 1, 2) were very similar to those of **3**, which allowed us to infer that their structural discrepancy may result from the different substitution patterns of the hydroxyethyl and prenyl groups. This deduction was confirmed by the HMBC correlations from the protons at 3.30 (overlapped, H-1'') to δ_{C} 160.5 (s, C-5), 113.1 (s, C-6) and 163.7 (s, C-7), and from the proton at δ_{H} 7.25 (d, $J = 2.3$ Hz, H-2') to δ_{C} 35.1 (t, C-1'''). Therefore, the structure of **4** was established as shown and given the name derrisisoflavone K.

Compound **5**, a white amorphous powder, had a molecular formula of C₁₇H₁₈O₅ by its HRESIMS (pos.): m/z 325.1031 [$\text{M} + \text{Na}$]⁺ (calcd. for C₁₇H₁₈O₅Na, 325.1046). Its ¹H NMR spectrum (Table 1) displayed an ABX-type aromatic proton system [δ_{H} 6.38 (d, $J = 2.5$ Hz), 6.36 (dd, $J = 8.4, 2.5$ Hz), and 6.95 (d, $J = 8.4$ Hz)], two aromatic singlets at δ_{H} 6.51 and 6.37, two methoxy signals at δ_{H} 3.78 and 3.71, and a set of signals [δ_{H} 4.17 (ddd, $J = 10.2, 3.3, 1.9$ Hz, H_{eq}-2), 3.90 (t, $J = 10.2$ Hz, H_{ax}-2), 3.43 (dddd, $J = 10.7, 10.2, 5.3, 3.3$ Hz, H_{ax}-3), 2.91 (dd, $J = 15.9, 10.7$ Hz, H_{ax}-4), and 2.73 (ddd, $J = 15.9, 5.3, 1.9$ Hz, H_{eq}-4)] due to ring C protons of an isoflavan. The above NMR signals were similar to those of isositivan (also called 7-*O*-methylvestitol) [14], and a prominent difference was two aromatic singlets at δ_{H} 6.51 and 6.37 instead of one of ABX-type system of isositivan. By careful analysis of the MS and NMR data, the isoflavan was inferred as a hydroxylated derivative of isositivan. The additional hydroxy group was located at C-6 by the HMBC correlations from the proton singlet at δ_{H} 6.51 (H-5) to the carbon at δ_{C} 31.4 (t, C-4), and from the methoxy signal at δ_{H} 3.78 (7-OMe) to the carbons at δ_{C} 148.1 (s, C-7) and 101.4 (d, C-8). Thereupon, the structure of **5** was established and named 6-hydroxyisositivan. The absolute configuration at C-3 was postulated as being *R*-form in the light of a negative specific rotation value (−11.7, MeOH), consistent with those of (3*R*)-vestitol derivatives [15].

3 Experimental Section

3.1 General Experimental Procedures

Optical rotation was measured on a Jasco P-1020 automatic digital polarimeter. UV data were obtained from HPLC online analysis. NMR spectra were carried out on a Bruker AV-400, Bruker DRX-500 or Bruker AV-600 instrument with deuterated solvent signals used as internal standards. ESI and HRESIMS were performed with a Shimadzu LC-IT-TOF mass spectrometer equipped with an ESI interface (Shimadzu, Kyoto, Japan). Silica gel 200–300 mesh (Qingdao Marine Chemical Inc., Qingdao, China),

Chromatorex C-18 (40–75 μm , Fuji Silysia Chemical Ltd., Japan) and Sephadex LH-20 (Amersham Biosciences, Uppsala, Sweden) were used for normal pressure column chromatography (CC). Fractions were monitored and analyzed by TLC, in combination with Agilent 1200 series HPLC system equipped by Extend-C18 column (5 μm , 4.6×150 mm).

3.2 Plant Material

The twigs and leaves of *D. robusta* were collected from the Pu'er region of Yunnan Province, People's Republic of China, in May 2011, and identified by Mr. Yu Chen of Kunming Institute of Botany, Chinese Academy of Sciences. A voucher specimen (BBP0350022DR) was deposited at BioBioPha Co., Ltd.

3.3 Extraction and Isolation

The air-dried and powdered twigs and leaves (12.0 kg) of *D. robusta* were extracted with EtOH-H₂O (95:5, v/v; 3×20 L, each 4 days) at room temperature, and the combined filtrates were concentrated under reduced pressure to give crude extract (ca. 870 g), which was further fractionated by silica gel CC successively eluted with a gradient of increasing acetone in petroleum ether (PE) (10:1 \rightarrow 0:1, v/v) and then MeOH to obtain nine fractions A–I. Fraction D (PE/acetone, 5:1, v/v) was subjected to silica gel CC (CHCl₃/MeOH, 100:0 \rightarrow 100:1, v/v) and Sephadex LH-20 (CHCl₃/MeOH, 1:1, v/v) to give **5** (14 mg). Fraction E (PE/acetone, 4:1, v/v) was isolated on silica gel CC (CHCl₃/MeOH, 100:1 \rightarrow 5:1, v/v), RP-18 (30 % MeOH/H₂O, v/v), and Sephadex LH-20 (MeOH) to yield **1** (7 mg), **2** (11 mg), and **4** (14 mg). Fraction H (PE/acetone, 1:1) was purified by silica gel CC (CHCl₃/MeOH, 10:1 \rightarrow 2:1, v/v) and repeated Sephadex LH-20 (MeOH) to afford **3** (10 mg). The retention times (t_R) of **1**–**5** on an analytical HPLC Extend-C18 column (20 % \rightarrow 100 % MeOH in H₂O over 8.0 min followed by 100 % MeOH to 13.0 min, 1.0 ml/min, 25 °C) were 10.32, 10.12, 9.13, 9.16, and 7.76 min, respectively.

3.4 Derrisisoflavone H (**1**)

Yellow amorphous powder; UV (MeOH) λ_{max} (log ϵ): 268 (4.75), 297 (sh) (4.23), 359 (3.60) nm; ¹H NMR data: see Table 1; ¹³C NMR data: see Table 2; ESIMS (neg.): m/z 449 [M – H][–]; HRESIMS (neg.): 449.1608 [M – H][–] (calcd. for C₂₆H₂₅O₇, 449.1606).

3.5 Derrisisoflavone I (**2**)

Yellow amorphous powder; UV (MeOH) λ_{max} (log ϵ): 263 (4.75), 302 (sh) (4.35), 357 (sh) (3.68) nm; ¹H NMR data:

see Table 1; ¹³C NMR data: see Table 2; ESIMS (neg.): m/z 449 [M – H][–]; HRESIMS (neg.): m/z 449.1607 [M – H][–] (calcd. for C₂₆H₂₅O₇, 449.1606).

3.6 Derrisisoflavone J (**3**)

White amorphous powder; UV (MeOH) λ_{max} (log ϵ): 213 (4.59), 267 (4.56), 338 (sh) (3.59) nm; ¹H NMR data: see Table 1; ¹³C NMR data: see Table 2; ESIMS (pos.): m/z 405 [M + Na]⁺; HRESIMS (pos.): m/z 405.1305 [M + Na]⁺ (calcd. for C₂₂H₂₂O₆Na, 405.1309).

3.7 Derrisisoflavone K (**4**)

White amorphous powder; UV (MeOH) λ_{max} (log ϵ): 214 (4.59), 268 (4.57), 336 (sh) (3.67) nm; ¹H NMR data: see Table 1; ¹³C NMR data: see Table 2; ESIMS (pos.): m/z 405 [M + Na]⁺; HRESIMS (pos.): m/z 405.1307 [M + Na]⁺ (calcd. for C₂₂H₂₂O₆Na, 405.1309).

3.8 6-Hydroxyisotivan (**5**)

White amorphous powder; UV (MeOH) λ_{max} (log ϵ): 226 (sh) (4.22), 287 (3.89) nm; [α]_D²³ –11.7 (*c* 0.2, MeOH); ¹H NMR data: see Table 1; ¹³C NMR data: see Table 2; ESIMS (pos.): m/z 325 [M + Na]⁺; HRESIMS (pos.): m/z 325.1031 [M + Na]⁺ (calcd. for C₁₇H₁₈O₅Na, 325.1046).

Acknowledgments This work was financially supported by the “Large-scale Compound Library” project of National Development and Reform Commission of China.

Compliance with Ethical Standards

Conflict of Interest The authors declare no conflict of interest.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. Editorial Committee of Flora Reipublicae Popularis Sinicae, Flora Reipublicae Popularis Sinicae. Academic Press: Beijing **40**, 194–195 (1994)
2. X.M. Li, M.F. Mao, F.C. Ren, X.J. Jiang, P. Hai, F. Wang, Nat. Prod. Bioprospect. **5**, 287–291 (2015)
3. S. Morel, J.J. Helesbeux, D. Séraphin, S. Derbré, J. Gatto, M.C. Aumond, Y. Abatuci, P. Grellier, M.A. Beniddir, P. Le Pape, F. Pagniez, M. Litaudon, A. Landreau, P. Richomme, Phytochem. Lett. **6**, 498–503 (2013)

4. T. Sekine, M. Inagaki, F. Ikegami, Y. Fujii, N. Ruangrunsi, *Phytochemistry* **52**, 87–94 (1999)
5. W. Mahabusarakam, S. Deachathai, S. Phongpaichit, C. Jansakul, W.C. Taylor, *Phytochemistry* **65**, 1185–1191 (2004)
6. P. Decharchoochart, J. Suthiwong, P. Samatiwat, V. Kukongviriyapan, C. Yenjai, *J. Nat. Med.* **68**, 730–736 (2014)
7. S.A. Rao, P.V. Srinivas, A.K. Tiwari, U.M.S. Vanka, R.V.S. Rao, K.R. Dasari, M.J. Rao, *J. Chromatogr. B* **855**, 166–172 (2007)
8. F. Wang, Y. Gao, L. Zhang, J.K. Liu, *Org. Lett.* **12**, 2354–2357 (2010)
9. F. Wang, Y.J. Li, F.C. Ren, G.Z. Wei, J.K. Liu, *Chem. Pharm. Bull.* **59**, 484–487 (2011)
10. F. Wang, X.L. Li, G.Z. Wei, F.C. Ren, J.K. Liu, *Nat. Prod. Bioprospect.* **3**, 238–242 (2013)
11. G.R. Wang, W.Z. Tang, Q.Q. Yao, H. Zhong, Y.J. Liu, *J. Nat. Med.* **64**, 358–361 (2010)
12. X.L. Li, N.L. Wang, M.S. Wong, A.S.C. Chen, X.S. Yao, *Chem. Pharm. Bull.* **54**, 570–573 (2006)
13. L. Pistelli, K. Spera, G. Flamini, S. Mele, I. Morelli, *Phytochemistry* **42**, 1455–1458 (1996)
14. A.L. Piccinelli, M.C. Fernandez, O. Cuesta-Rubio, I.M. Hernández, F. De Simone, L. Rastrelli, *J. Agric. Food Chem.* **53**, 9010–9016 (2005)
15. S. Yahara, T. Ogata, R. Saijo, R. Konishi, J. Yamahara, K. Miyahara, T. Nohara, *Chem. Pharm. Bull.* **37**, 979–987 (1989)