

# The beneficial role of lutein and zeaxanthin in cataracts

Samanta Maci, Rafaela Santos



Correspondence to:

Samanta Maci

samanta.maci@kemin.com

**Keywords:**

*lutein*  
*zeaxanthin*  
*age-related cataracts*  
*antioxidants*  
*eye*

Received: 4 February 2015 / Accepted: 9 April 2015

© Springer – CEC Editore 2015

## Abstract

Cataracts remain the leading cause of visual loss and blindness worldwide. Cataract surgery is the most common surgical procedure in several countries and its occurrence is expected to increase substantially due to the ageing of the population and the reduced visual acuity threshold required for considering it. Cataracts are characterised by lens opacity and are caused by the precipitation of proteins induced by oxidative damage. The fact that oxidation of the lens is a contributing cause for cataracts suggests that antioxidants may play a positive role in cataract prevention. Lutein and zeaxanthin, two dietary xanthophylls exclusively deposited in the macula lutea and known for their beneficial role in age-related macular degeneration, are the only carotenoids deposited in the lens and may have an important role in reducing the ROS-

induced damage of the lens contributing to cataract development. This paper will review the current science derived from epidemiological and intervention studies addressing the association between lutein and zeaxanthin intake, their serum level, and the possible benefits for primary prevention of age-related cataract. Additional insights into the potential health cost savings from the increased intake of these dietary ingredients originating from a U.S. study are also provided.

## Introduction

The incidence of age-related cataracts, responsible for 51% of world blindness (approximately 20 million people in 2010), is expected to rise due to the increase in life expectancy [1], escalating the already large public health expenditures and resulting in billions of dollars in healthcare costs. In developing countries, cataracts are the principal cause of blindness among people over 40 years of age due to improper nutrition and infectious diseases [2].

Cataracts are characterised by lens opacity induced by cumulative or excessive exposure to light, oxygen and metals that generate reactive oxygen species (ROS) leading to peroxidation of membrane lipids and subsequent destruction and precipitation of lens protein crystalline(s). Currently, cataracts are successfully treated by surgical removal of the

Samanta Maci (✉)

Kemin Human Nutrition and Health

a division of Kemin Foods L.C.

Campo Grande 35- 8<sup>o</sup>D, 1700-087 Lisbon, Portugal

tel +351214157500; fax +351211412172

samanta.maci@kemin.com

Rafaela Santos

Kemin Human Nutrition and Health

a division of Kemin Foods L.C.

Rua Ettore Soliane, 471 13347-394, Indaiatuba - SP, Brazil

lens, followed by implantation of an artificial lens at the time of surgery. The surgical procedure provides excellent improvement in patient visual acuity and quality of life; however, patients living in developing countries or not covered by health insurance can have limited or delayed access to it. In addition, the costs of cataract extraction represent an important public health burden; thus, strategies to prevent cataracts or delay their progression should be followed [3].

The fact that oxidation of the lens is a major cause for cataracts suggests that antioxidants may play a positive role in cataract prevention. Because most of the enzymatic cofactors and other constituents which contribute to the lens antioxidant defence mechanism originate from diet, it is likely that diet has an important role in reducing the ROS-induced damage of the lens [4]. Evidence has emerged suggesting that healthy lifestyle and proper nutrition may have a beneficial effect on the onset of cataracts [5, 6] and special attention has been paid to two dietary antioxidants in particular: lutein and zeaxanthin.

The purpose of this paper is to provide an overview of the science addressing the beneficial relationship between lutein and zeaxanthin intake and cataracts.

### Lutein and zeaxanthin: Focus on prevention

Lutein and zeaxanthin, two dietary xanthophylls exclusively deposited in the macula lutea and key components of the macular pigment, are the only carotenoids deposited in the lens. Here they can be found in all the lens layers although they are more concentrated in the epithelial and cortical layers than the nuclear layer [7].

The rationale for the protective role of lutein and zeaxanthin in both the retina and the lens stems from their ability (1) to filter harmful short-wave blue light, (2) to function as antioxidants and (3) to stabilise membrane integrity. These biological functions are believed to play an important role in helping to reduce light-induced oxidative damage caused by ROS, which contributes to the patho-

genesis of age-related macular degeneration (AMD) and cataracts [8].

Studies conducted in patients with AMD or healthy subjects show that macular pigment optical density (MPOD), which provides an indication of lutein and zeaxanthin in the macula and is measured with non-invasive technologies, can be augmented with increased intake of lutein and zeaxanthin. The scientific literature also suggests that in individual 50+ years old, there is an inverse relationship between MPOD and lens opacity quantified in terms of lens optical density (LOD) [9, 10]. Since increased LOD provides an indication of lens senescence, which is strongly related to cataracts [9], the fact that high MPOD is associated with decreased LOD suggests that lutein and zeaxanthin in the eye may retard lens ageing [9, 10].

The protective role of these two carotenoids and the underlying mechanism of action is supported by (1) their exclusive presence in the lens and (2) observations from *in vitro* studies indicating that pretreatment of human lens epithelial (HLE) cells with lutein and zeaxanthin: (i) induces their accumulation in the lens, (ii) decreases UVB-induced lipid peroxidation and stress signalling activation [4] and (iii) reduces H<sub>2</sub>O<sub>2</sub>-mediated damage of lens protein, lipid and DNA [11].

An *in vivo* intervention study conducted in quails confirmed that lens carotenoids can be increased by dietary intake of zeaxanthin [12]. Additionally, a study performed in streptozotocin-induced diabetic rats showed that lutein supplementation was effective in preventing the development and progression of cataracts and an increase in lipid peroxidation [13].

Epidemiological studies provide much of the scientific data on lutein and zeaxanthin and indicate the possible protective effect of dietary intake of lutein and zeaxanthin and/or their serum levels against cataract incidence, mainly nuclear cataracts, or risk of cataract extraction [14–29], although some publications report a null or negative effect [14, 28–31]. A summary of some of the papers addressing

Study	Cohort	Age	Lutein (L) and/or zeaxanthin (Z) daily intake ( $\mu\text{g}$ )		Key outcome
			Lowest	Highest	
Brown et al., 1999 [15]	36,344 men	45–75 years	1.300	6.871	<ul style="list-style-type: none"> <li>Risk of cataract extraction was 19% lower in the group with the highest quintile of consumption of L and Z when compared to the lowest quintile of intake (RR: 0.81, 95% CI: 0.65–1.01; <math>p</math> for trend=0.03).</li> <li>As cataracts develop over many years, frequent consumption of foods high in carotenoids early in life or for long periods may be associated with a lower risk of senile cataract extractions.</li> </ul>
Chasan Taber et al., 1999 [16]	77,466 women	45–71 years	1.172	11.685	<ul style="list-style-type: none"> <li>Subjects with the highest quintile of intake of L and Z had a 22% decreased risk of cataract extraction compared with those in the lowest quintile (RR: 0.78; 95% CI: 0.63–0.95; <math>p</math> for trend=0.04).</li> </ul>
Lyle et al., 1999 [23]	1,354	43–84 years	298	1.245	<ul style="list-style-type: none"> <li>Subjects in the highest quintile of L intake in the distant past were half as likely to have an incident cataract as persons in the lowest quintile of intake (95% CI 0.3–0.8; <math>p</math> for trend=0.002).</li> <li>High intake of L at baseline was associated with a non-significant OR of 0.6 (95% CI 0.4–1.1; <math>p</math> for trend=0.09).</li> <li>Lutein was the only carotenoid out of five examined that was associated with nuclear cataracts.</li> <li>Consumption of spinach and other dark leafy greens (concentrated source of lutein) was inversely associated with nuclear cataracts for persons with the highest intakes (OR: 0.6; 95% CI: 0.4–0.9; <math>p</math> for trend=0.02).</li> </ul>
Jaques et al., 2001 [18]	478 women	53–73 years	–	5.600	<ul style="list-style-type: none"> <li>The prevalence of nuclear opacification was significantly lower in the upper four L/Z quintiles of intake categories relative to the lowest quintile category (<math>p</math> for trend=0.03), although the association did not remain statistically significant after adjustment for other nutrients.</li> </ul>
Rodriguez-Rodriguez et al., 2006 [19]	177	>65 years	<256 (5th %)	>3290 (95th %)	<ul style="list-style-type: none"> <li>Subjects whose daily L intake was &gt;3290 <math>\mu\text{g}</math> were less likely to have cataracts (OR=0.086; CI: 0.007–1.084; <math>p</math>&lt;0.05) than those whose consumption was &lt;256 <math>\mu\text{g}</math>.</li> <li>In men, high intakes of Z seemed to have a protective effect (OR=0.96; CI: 0.91–0.99; <math>p</math>&lt;0.05).</li> </ul>
Vu et al., 2006 [20]	2,322	>40 years	472	1037	<ul style="list-style-type: none"> <li>Inverse association between high dietary LZ intake and prevalence of nuclear cataracts. The odds ratio (95% CI) for those in the top quintile of energy-adjusted LZ intake was 0.64 (CI: 0.40–1.03; <math>p</math> for trend=0.018).</li> <li>In the analysis with LZ as continuous variable, the odds ratio for every 1 mg increase in energy-adjusted daily LZ intake was 0.60 (0.40–0.90) for nuclear cataracts.</li> <li>Cortical and posterior subcapsular cataracts were not significantly associated with LZ intake.</li> </ul>
Christen et al., 2008 [44]	35,551 women	>45 years	1.177	6.716	<ul style="list-style-type: none"> <li>Women in the highest quintiles of lutein/zeaxanthin per day (corresponding to more than 6 mg/day) had 18% lower risk of cataracts compared to women with median lutein/zeaxanthin intake approximating 1 mg per day (RR: 0.82; 95% CI: 0.71–0.95; <math>p</math> for trend =0.04).</li> <li>A borderline significant inverse association between higher intake of green leafy vegetables such as spinach and risk of cataracts was also observed.</li> </ul>
Moeller et al., 2008 [22]	1802 women	50–79 years	600	3.300	<ul style="list-style-type: none"> <li>Women in the highest quintiles of dietary lutein and zeaxanthin intake had a 32% lower prevalence of nuclear cataract (OD: 0.68; 95% CI: 0.48–0.97; <math>p</math>=0.04) and 37% lower prevalence of nuclear sclerosis score <math>\geq 0.4</math> (OD: 0.63; CI: 0.43–0.94; <math>p</math>=0.04 for trend) compared with those with low levels.</li> </ul>

**Table 1** Published observational studies addressing the relationship between dietary consumption of lutein (L)/zeaxanthin (Z) and cataracts

the positive relationship between cataract incidence or extraction and high lutein and zeaxanthin in the diet or serum is presented in Tables 1 and 2. Three recently published meta-analyses examining the association between dietary or blood lutein and zeaxanthin and age-related cataract and referring to the studies listed in Tables 1 and 2 support the efficacy of these carotenoids in reducing the risk

of this eye condition, in particular nuclear cataracts [32–34]. More specifically, the dose-response analysis published by Ma et al. in 2013 showed a statistically significant 3% decrease in nuclear cataract risk for each 300  $\mu\text{g}$  increase in daily dietary lutein and zeaxanthin intake [33].

Only two human intervention studies [35, 36] have been conducted up to the present date to explore

Study	Cohort	Age	Serum level ( $\mu\text{mol/l}$ )		Key outcome
			Lutein	Zeaxanthin	
Lyle et al., 1999 [17]	325	50–86 years	Lutein+zeaxanthin combined Low: <0.18 High: $\geq$ 0.38		<ul style="list-style-type: none"> <li>Serum lutein/zeaxanthin were not significantly associated with 5 y incidence of severe nuclear cataract, however a marginal inverse association with lutein (OR: 0.3; 95% CI: 0.1–1.2; <math>p=0.13</math> for linear trend) was suggested for people <math>\geq</math>65 y of age.</li> </ul>
Gale et al., 2001 [14]	372 men and women	66–75 years	Low: 0.14 High: >0.20	Low: 0.03 High: >0.05	<ul style="list-style-type: none"> <li>Risk of posterior subcapsular cataract was lowest in those with higher concentrations of lutein (OR: 0.5; 95% CI: 0.2–1.0, <math>p</math> for trend: 0.012).</li> <li>No statistical significant association was found between lutein or zeaxanthin serum levels and risk of nuclear or cortical cataracts.</li> </ul>
Delcourt et al., 2006 [24]	644	>60 years	Low: 0.17 High: $\geq$ 0.38	Low: 0.04 High: $\geq$ 0.09	<ul style="list-style-type: none"> <li>Highest tertiles of plasma zeaxanthin were significantly associated with 75% reduced risk of nuclear cataract (OR: 0.23; 95% CI: 0.08–0.68; <math>p</math> for trend 0.003) and 43% reduced risk for any cataract (OR: 0.53; 95% CI: 0.31–0.89; <math>p</math> for trend 0.01).</li> <li>No association with plasma lutein levels was observed, while a statistically significant 66% reduction of risk for nuclear cataract was found with high plasma dehydro-lutein.</li> <li>No statistically significant association with cortical or posterior subcapsular cataract was found for any of the carotenoids.</li> </ul>
Dherani et al., 2008 [25]	1112	>50 years	Low: <0.0762 High: $\geq$ 0.1137	Low: <0.0155 High: $\geq$ 0.0214	<ul style="list-style-type: none"> <li>Highest tertiles of plasma zeaxanthin or lutein were associated with decreased odds of cataract (OR: 0.66; 95% CI: 0.45–0.96; <math>p=0.03</math> for zeaxanthin and OR: 0.73; 95% CI: 0.43–1.02; <math>p=0.06</math> for lutein).</li> <li>Inverse associations were found between cataract and blood antioxidants in an antioxidant-depleted study sample.</li> </ul>
Moeller et al., 2008 [22]	1802 women	50–79 years	Lutein+zeaxanthin combined Low: 0.15 High: 0.5		<ul style="list-style-type: none"> <li>Women in the highest quintile category of serum levels of lutein and zeaxanthin were 32% less likely to have nuclear cataract (OR: 0.68; 95% CI, 0.47–0.98; <math>p</math> for trend=0.01) compared with those in the lowest quintile category.</li> <li>A 21% reduction on nuclear sclerosis (score <math>\geq</math>4) was also observed, although it did not reach statistical significance in the multivariable adjusted analysis.</li> </ul>
Karppi et al., 2012 [26]	1689	61–80 years	Low: <0.2 High: >0.27	Low: <0.030 High: >0.041	<ul style="list-style-type: none"> <li>Subjects in the highest tertile of plasma concentrations of lutein had 42% lower risks of nuclear cataract compared with those in the lowest tertile (relative risk (RR)=0.58; 95% CI 0.35–0.98; <math>p=0.041</math>).</li> <li>Subjects in the highest tertile of plasma concentrations of zeaxanthin had 41% lower risks of nuclear cataract compared with those in the lowest tertile (RR=0.59; 95% CI 0.35–0.99; <math>p=0.046</math>).</li> </ul>
Ravindran et al., 2011 [27]	5638 people	>60 years	Q1: 0.125 Q4: 0.460	Q1: 0.021 Q2: 0.066	<ul style="list-style-type: none"> <li>Lutein and zeaxanthin were significantly inversely associated with cataract, but the associations were weaker and not consistently observed by type of cataract.</li> </ul>

**Table 2** Published observational studies addressing the relationship between the L/Z serum/plasma concentrations and cataracts

the cause–effect relationship between lutein and zeaxanthin supplementation and benefits for cataracts, providing additional support for the positive findings collected from observational studies and addressing the importance of supplementation, especially when the dietary intake of these carotenoids is low. The Age-Related Eye Disease Study 2 (AREDS2), a large intervention trial primarily conducted to evaluate the effect of nutritional intervention with lutein and zeaxanthin on reduction in the risk of progression to advanced AMD [37, 38], also assessed the effect of these carotenoids

on cataracts or progression to cataract surgery. Although the results of the primary analysis comparing the progression of cataract with or without lutein and zeaxanthin supplementation resulted in a non-significant hazard ratio of 0.96 (95% CI 0.84–1.10), in subjects in the lowest quintiles of lutein and zeaxanthin dietary intake, supplementation with 10 mg lutein and 2 mg zeaxanthin for 5 years resulted in a 30% reduction of any cataract and 32% reduction of cataract surgery when compared to no lutein and zeaxanthin supplementation [36]. It should be noted that the AREDS2 study popula-

Study	Cohort	Age	Lutein/zeaxanthin dose	Period of supplementation	Key outcome
The AREDS2 research group, 2013 [36]	3159	50–85 years	10 mg L+2 mg Z (daily)	5 years	<ul style="list-style-type: none"> <li>Subjects in the lowest quintile of lutein and zeaxanthin intake benefit from the addition of lutein/zeaxanthin from the AREDS supplement: When comparing L+Z vs. no L+Z, in these subjects a significant reduction in risk of progression to cataract surgery, any cataract or any severe cataract of 32%, 30% and 36%, respectively (<math>p&lt;0.05</math>) was observed.</li> <li>No overall effect of lutein/zeaxanthin supplementation was observed in the study. The hazard ratios for progression to cataract surgery for patients receiving the carotenoids was 0.96 (95% CI: 0.84–1.10; <math>p=0.54</math>).</li> </ul>
Olmedilla et al., 2003 [35]	15	55–73 years	12 mg of all-trans lutein (3 times a week)	2 years	<ul style="list-style-type: none"> <li>Supplementation with lutein increased serum concentrations, whereas no changes were observed in the placebo group.</li> <li>Lutein supplementation significantly improved visual acuity in cataract patients.</li> <li>Glare sensitivity at low, medium and high thresholds also improved in the lutein group compared to the baseline values.</li> <li>Cataract progression was not observed in 80% of the patients in the lutein group compared to just 20% of subjects in the placebo group.</li> </ul>

**Table 3** Interventional studies addressing the effect of supplementation with L and Z on visual function and cataract progression in cataract patients

tion was well nourished compared to the general U.S. adult population, with above-average intake of dietary nutrients, including much higher intake and mean serum levels of lutein and zeaxanthin. The intake of the lowest quintile is more representative of what is typical in the adult population, therefore the outcomes of the subgroup analysis may be what is more relevant. In the study published by Olmedilla in 2003, 2-year supplementation of cataract patients with lutein resulted in improvements in visual function (visual acuity and glare) [35]. A summary of these two human intervention studies is shown in Table 3.

These observed beneficial effects are important in light of a possible reduction of medical expenditures associated with the condition. A report issued in 2013 by Frost and Sullivan and commissioned by the Council for Responsible Nutrition (CRN) Foundation [39] provided insights into the potential net savings that could be achieved in four different diseases, including age-related eye diseases (AMD and cataract), by using dietary supplements as a preventive health-care measure to reduce the rate of medical events in the U.S. high-risk population over 55 years. By reviewing published studies directly assessing the direct causal relationship be-

tween intake of lutein and zeaxanthin and the relative risk of the disease events, the report showed that the relative risk reductions for AMD and cataract associated with the use of lutein and zeaxanthin were 23% and 15.3%, respectively, overall offering a potential net health-care cost savings (after accounting for the cost of lutein and zeaxanthin supplements) between 2013 and 2020 of nearly \$1 billion/year.

### Conclusion

Lutein and zeaxanthin are the only dietary carotenoids deposited in the macula lutea and are well known for their ability to act as antioxidants and as a filter of blue light. Lutein and zeaxanthin are also the only dietary carotenoids deposited in the lens. The selective uptake of these carotenoids in this structure combined with their biological function support their potential beneficial role also in age-related cataracts, thus expanding the positive effect observed in patients with AMD and in healthy subjects. The science originating from observational studies suggests that adequate dietary intake or serum lutein and zeaxanthin may help to reduce the risk of cataract incidence as well as the need for cataract extraction. The results of intervention stud-

ies further support the importance of an adequate intake of lutein and zeaxanthin and reinforce the benefits of supplementation with these carotenoids when proper dietary intake is not achieved, as occurs in the modern diet [40–43].

#### Conflict of interest

Samanta Maci and Rafaela Pontes are employees of Kemin Human Nutrition and Health.

#### Human and Animal rights

This article does not contain any studies with human or animal subjects performed by any of the authors.

#### References

1. WHO. Prevention of blindness and visual impairment. World Health Organization. <http://www.who.int/blindness/causes/priority/en/index1.html>. Accessed January 2015
2. Abdel-Aal EM, Akhtar H, Zaheer K, Ali R (2013) Dietary sources of lutein and zeaxanthin carotenoids and their role in eye health. *Nutrients* 5:1169–1185
3. Fernandez MM, Afshari NA (2011) Cataracts: we have perfected the surgery, but is it time for prevention? *Curr Opin Ophthalmol* 22:2–3
4. Chitchumroonchokchai C, Bomser JA, Glamm JE, Failla ML (2004) Xanthophylls and  $\alpha$ -tocopherol decrease UVB-induced lipid peroxidation and stress signaling in human lens epithelial cells. *J Nutr* 134:3225–3232
5. Seddon JM (2007) Multivitamin-multimineral supplements and eye disease: age-related macular degeneration and cataract. *Am J Clin Nutr* 85:304S–307S
6. Mares JA, Volland R, Adler R, Tinker L, Millen AE, Moeller SM, Blodi B, Gehrs KM, Wallace RB, Chappell RJ et al (2010) Healthy diets and the subsequent prevalence of nuclear cataract in women. *Arch Ophthalmol* 128:738–749
7. Yeum KJ, Shang FM, Schalch WM, Russell RM, Taylor A (1999) Fat-soluble nutrient concentrations in different layers of human cataractous lens. *Curr Eye Res* 19:502–505
8. Krinsky NI, Landrum JT, Bone RA (2003) Biologic mechanisms of the protective role of lutein and zeaxanthin in the eye. *Annu Rev Nutr* 23:171–201
9. Berendschot TT, Broekmans WM, Klopping-Ketelaars IA, Kardinaal AF, Van Poppel G, Van Norren D (2002) Lens aging in relation to nutritional determinants and possible risk factors for age-related cataract. *Arch Ophthalmol* 120:1732–1737
10. Hammond BR, Wooten BR, Snodderly DM (1997) Density of the human crystalline lens is related to the macular pigment carotenoids, lutein and zeaxanthin. *Optom Vis Sci* 74:499–504
11. Gao S, Qin T, Liu Z, Caceres MA, Ronchi CF, Chen CO, Yeum K, Taylor A, Blumberg JB, Liu Y, Shang F (2011) Lutein and zeaxanthin supplementation reduces H<sub>2</sub>O<sub>2</sub>-induced oxidative damage in human lens epithelial cells. *Mol Vis* 17:3180–3190
12. Dorey CK, Granata L, Nichols CR, Cheng KM, Craft NE (2005) Dietary modulation of lens zeaxanthin in quail. *Exp Eye Res* 81:464–477
13. Arnal E, Miranda M, Almansa I, Muriach M, Barcia JM, Romero FJ, Diaz-Llopis M, Bosch-Morell F (2009) Lutein prevents cataract development and progression in diabetic rats. *Graefes Arch Clin Exp Ophthalmol* 247:115–120
14. Gale CR, Hall NF, Phillips DI, Martyn CN (2001) Plasma antioxidant vitamins and carotenoids and age-related cataract. *Ophthalmology* 108:1992–1998
15. Brown L, Rimm EB, Seddon JM et al (1999) A prospective study of carotenoid intake and risk of cataract extraction in US men. *Am J Clin Nutr* 70:517–524
16. Chasan-Taber L, Willett WC, Seddon JM et al (1999) A prospective study of carotenoid and vitamin A intakes and risk of cataract extraction in US women. *Am J Clin Nutr* 70:509–516
17. Lyle BJ, Mares-Perlman JA, Klein BE, Klein R, Palta M, Bowen PE, Greger JL (1999) Serum carotenoids and tocopherols and incidence of age-related nuclear cataract. *Am J Clin Nutr* 69:272–277
18. Jacques PF, Chylack LT Jr, Hankinson SE, Khu PM, Rogers G, Friend J, Tung W, Wolfe JK, Padhye N, Willett WC et al (2001) Long-term nutrient intake and early age-related nuclear lens opacities. *Arch Ophthalmol* 119:1009–1019
19. Rodriguez-Rodriguez E, Ortega RM, Lopez-Sobaler AM, Aparicio A, Bermejo LM, Marin-Arias LI (2006) The relationship between antioxidant nutrient intake and cataracts in older people. *Int J Vit Nutr Res* 76:359–366
20. Vu HT, Robman L, Hodge A, McCarty CA, Taylor H (2006) Lutein and zeaxanthin and the risk of cataract: the Melbourne visual impairment project. *Invest Ophthalmol Vis Sci* 47:3783–3786
21. Christen WG, Liu S, Glynn RJ et al (2008) Dietary carotenoids, vitamin C and E, and risk of cataract in women. *Arch Ophthalmol* 126:102–109
22. Moeller SM, Volland R, Tinker L, Blodi BA, Klein ML, Gehrs KM, Johnson EJ, Snodderly DM, Wallace RB, Chappell RJ et al (2008) Associations between age-related nuclear cataract and lutein and zeaxanthin in the diet and serum in the Carotenoids in the Age-Related Eye Disease Study, an Ancillary Study of the Women's Health Initiative. *Arch Ophthalmol* 126:354–364
23. Lyle BJ, Mares-Perlman JA, Klein BE, Klein R, Greger JL (1999) Antioxidant intake and risk of incident age-related nuclear cataracts in the Beaver Dam Eye Study. *Am J Epidemiol* 149:801–809
24. Delcourt C, Carrière I, Delage M, Barberger-Gateau P, Schalch W (2006) Plasma lutein and zeaxanthin and other



- carotenoids as modifiable risk factors for age-related maculopathy and cataract: the POLA Study. *Invest Ophthalmol Vis Sci* 47:2329–2335
25. Dherani M, Murthy GV, Gupta SK, Young IS, Maraini G, Camparini M, Price GM, John N, Chakravarthy U, Fletcher AE (2008) Blood levels of vitamin C, carotenoids and retinol are inversely associated with cataract in a North Indian population. *Invest Ophthalmol Vis Sci* 49:3328–3335
  26. Karppi J, Laukkanen JA, Kurl S (2012) Plasma lutein and zeaxanthin and the risk of age-related nuclear cataract among the elderly Finnish population. *Br J Nutr* 14:148–154
  27. Ravindran RD, Vashist P, Gupta SK, Young IS, Maraini G, Camparini M, Jayanthi R, John N, Fitzpatrick KE, Chakravarthy U et al (2011) Inverse association of vitamin C with cataract in older people in India. *Ophthalmology* 118:1958–1965
  28. Chiu CJ, Taylor A (2007) Nutritional antioxidants and age-related cataract and maculopathy. *Exp Eye Res* 84:229–245
  29. Weikel KA, Garber C, Baburins A, Taylor A (2014) Nutritional modulation of cataract. *Nutr Rev* 72:30–47
  30. Vitale S, West SK, Halfrisch J, Allston C, Wang F, Moorman C, Muller D, Singh V, Taylor H (1993) Plasma antioxidants and risk of cortical and nuclear cataract. *Epidemiology* 4:195–203
  31. Mares-Perlman JA, Brady WE, Klein BE, Klein R, Palta M, Bowen P, Stacewicz-Sapuntzakis M (1995) Serum carotenoids and tocopherols and severity of nuclear and cortical opacities. *Invest Ophthalmol Vis Sci* 36:276–288
  32. Cui YH, Jing CX, Pan HW (2013) Association of blood antioxidants and vitamins with risk of age-related cataract: a meta-analysis of observational studies. *Am J Clin Nutr* 98:778–786
  33. Ma L, Hao ZX, Liu RR, Yu RB, Shi Q, Pan JP (2014) A dose-response meta-analysis of dietary lutein and zeaxanthin intake in relation to risk of age-related cataract. *Graefes Arch Clin Exp Ophthalmol* 252:63–70
  34. Liu XH, Yu RB, Liu R, Hao ZX, Han CC, Zhu ZH, Ma L (2014) Association between lutein and zeaxanthin status and the risk of cataract: a meta-analysis. *Nutrients* 6:452–465
  35. Olmedilla B, Granado F, Blanco I et al (2003) Lutein, but not alpha-tocopherol, supplementation improves visual function in patients with age-related cataracts: a 2-y double-blind, placebo-controlled pilot study. *Nutrition* 19:21–24
  36. Age-related Eye Disease Study 2 (AREDS2) Research Group (2013) Lutein/zeaxanthin for the treatment of age-related cataract: AREDS2 randomized trial report no. 4. *JAMA Ophthalmol* 131:843–850
  37. Age-related Eye Disease Study 2 Research Group (2013) Lutein + zeaxanthin and omega-3 fatty acids for age-related macular degeneration: the Age-Related Eye Disease Study 2 (AREDS2) randomized clinical trial. *JAMA* 309:2005–2015
  38. Age-related Eye Disease Study 2 (AREDS2) Research Group (2014) Secondary analyses of the effects of lutein/zeaxanthin on age-related macular degeneration progression. AREDS2 Report No. 3. *JAMA Ophthalmol* 132:142–149
  39. Shanahan C, de Lorimier R; Frost and Sullivan (2013) Smart prevention: health care cost savings resulting from the targeted use of dietary supplements. <http://www.crnusa.org/CRNFoundation/HCCS/chapters/CRNFrostSullivan-fullreport0913.pdf>
  40. Granado F, Blázquez S, Olmedilla B (2007) Changes in carotenoid intake from fruit and vegetables in the Spanish population over the period 1964–2004. *Public Health Nutr* 10:1018–1023
  41. Giuliano AR, Siegel EM, Roe DJ, Ferreira S, Baggio ML, Galan L, Duarte-Franco E, Villa LL, Rohan TE, Marshall JR et al (2003) Dietary intake and risk of persistent human papillomavirus (HPV) infection: the Ludwig-McGill HPV Natural History Study. *J Infect Dis* 188:1508–1516
  42. O'Neill ME, Carroll Y, Corridan B, Olmedilla B, Granado F, Blanco I, Van Den Berg H, Hininger I, Rousell AM, Chopra M, Southon S, Thurnham DI (2001) A European carotenoid database to assess carotenoid intakes and its use in a five-country comparative study. *Br J Nutr* 85:499–507
  43. Johnson EJ, Maras JE, Rasmussen HM, Tucker KL (2010) Intake of lutein and zeaxanthin differ with age, sex, and ethnicity. *J Am Diet Assoc* 110:1357–1362
  44. Christen WG, Liu S, Glynn RJ, Gaziano JM, Buring JE (2008) Dietary carotenoids, vitamins C and E, and risk of cataract in women: a prospective study. *Arch Ophthalmol* 126:102–109