

# Formaldehyde degradation by a newly isolated fungus *Aspergillus* sp. HUA

D. S. Yu · G. Song · L. L. Song · W. Wang ·  
C. H. Guo

Received: 8 September 2013 / Accepted: 3 November 2013 / Published online: 27 November 2013  
© Islamic Azad University (IAU) 2013

**Abstract** Formaldehyde is widely used in chemical manufacturing industry and classified as a human carcinogen. Discharging wastewater containing formaldehyde without treatment can cause serious risk to the water environment. In this study, a formaldehyde-resistant fungal strain was isolated from sewerage of a furniture factory. Isolate strain was identified based on the morphological and phylogenetic analyses. Formaldehyde-degrading fungus was determined by characterizing the mycelia growth in culture media, formaldehyde-resistant, formaldehyde-degrading efficiencies, and specific enzyme activity involved in formaldehyde removal. Isolate strain HUA was identified as a member of *Aspergillus sydowii*. The strain HUA showed a growth in the presence of formaldehyde up to 2,400 mg l<sup>-1</sup> at an optimum temperature of 25 °C and optimum pH of 7. The specific activity of formaldehyde dehydrogenase and formate dehydrogenase could reach up to 5.02 and 1.06 U mg<sup>-1</sup>, respectively. It indicated that isolated formaldehyde-resistant *A. sydowii* HUA strain would be potential used for removing formaldehyde from industrial wastewater.

**Keywords** Biodegradation · Formaldehyde · Formaldehyde-degrading fungus · *Aspergillus sydowii* · Wastewater

## Introduction

Formaldehyde, a highly reactive chemical material, is widely used in chemical manufacturing industry. It is often released from wastewater of construction, textile, wood processing, furniture, and pharmacy industry (Tang et al. 2009). Formaldehyde is a water-soluble compound, which can diffuse into many tissues rapidly, react with different macromolecules such as proteins and nucleic acids, cause DNA–DNA, protein–DNA, and protein–protein cross-links (Metz et al. 2004; Merk and Speit 1998; Solomon and Varshavsky 1985). Therefore, it has a toxic effect on all organisms, the IARC (2006) (International Agency for Research on Cancer) has classified formaldehyde as a human carcinogen that causes nasopharyngeal cancer and probably leukemia.

Discharging wastewater containing formaldehyde generated by organic synthesis cause environmental worsening and human disease (Tişler and Zagorc-Koncan 1997). Much more researches have been done to wastewater treatment (Moussavi and Heidarizad 2010). Biological control, such as microbial degradation, is the most commonly applied method for treatment of wastewaters containing biodegradable compounds (Bhakta et al. 2012; Kanmani et al. 2012; Ashraf et al. 2011). Many microorganisms have been shown to degrade formaldehyde in the sewage, and bacterial degradability has been the research hotspot on formaldehyde degradation (Di Maiuta et al. 2009; Arutchelvan et al. 2005; Hidalgo et al. 2002). Nevertheless, there is fewer fungal resources that are able to degrade formaldehyde have been reported (Sawada et al. 2006). Filamentous fungus plays an important role in bioremediation in wastewater treatment (Mani and Kumar 2013; Usharani and Muthukumar 2013; More et al. 2010). It is shown that filamentous fungus *A. versicolor* and *A.*

D. S. Yu · G. Song · L. L. Song · W. Wang · C. H. Guo (✉)  
Key Laboratory of Molecular and Cytogenetics, Heilongjiang  
Province, College of Life Science and Technology,  
Harbin Normal University, Harbin 150025, China  
e-mail: kaku\_2008@163.com

*niger* has bio-accumulate capacity of heavy metal from industrial effluent (Taştan et al. 2010; Das et al. 2007; Srivastava and Thakur 2006). The effects of biodegrading behavior of polymers and utilization of organophosphate pesticides by genus *Aspergillus* have been reported (Zhao et al. 2005).

Formaldehyde presents as a natural product endogenous biological compound in most living systems (Tang et al. 2009). Formaldehyde is situated at the branch point between dissimilation and assimilation pathway of formaldehyde (Yurimoto et al. 2005). Organisms have several biological forms of formaldehyde metabolism to deal with the hazards of formaldehyde. One of which is the combination of formaldehyde with cofactors, the combination product then is oxidized into CO<sub>2</sub> and generated energy (Hanson and Roje 2001). The glutathione (GSH)-dependent formaldehyde dehydrogenase (FADH) and formate dehydrogenase (FDH) participate in the above strategy (Tada and Kidu 2011). In addition, formaldehyde participate the C<sub>1</sub> assimilation pathways, the xylulose monophosphate pathway in yeasts, and the ribulose monophosphate pathway in methylotrophic bacteria (Yurimoto et al. 2009).

There are few reports on fungal degradation of formaldehyde and filamentous fungi formaldehyde metabolism. The aims of this study were to isolate and identify formaldehyde-degrading fungi and to determine the effects of the degradation of formaldehyde and the activities of key enzymes in formaldehyde metabolism.

## Materials and methods

### Fungal isolation and cultivation

The sludge samples were collected from sewerage of a furniture factory (Harbin, China). The samples were stored at 4 °C in sterile plastic bags. The sludge (10 g) was transferred into 90 ml of the sterilized liquid basal medium (1 l) contained: sucrose 3 %; NaNO<sub>3</sub> 0.3 %; K<sub>2</sub>HPO<sub>4</sub> 0.1 %; MgSO<sub>4</sub>·7H<sub>2</sub>O 0.05 %; KCl 0.05 %; FeSO<sub>4</sub>·7H<sub>2</sub>O 0.001 % (adjusted to pH 7) supplemented with 150 mg formaldehyde and incubated at 30 °C shaking with 180 rpm. After cultivation for 5 days, the culture was inoculated to plates containing basal medium with 2 % agar and 150 mg l<sup>-1</sup> formaldehyde incubated at 30 °C. Single colony was picked up and inoculated into another new plate. Above operation was repeated several times until the pure cultures were obtained.

### Identification and characterization of isolates

The fungal isolates were incubated on basal medium agar plates in the dark at 25 °C for 15 days. At the end of the

cultivation, morphology (colony growth, color of conidia, mycelia, exudate, and colony reverse) was characterized using light microscope (Leica DM1000, Leica Microsystems Germany). The genomic DNA was obtained from the mycelia according to the method of Zhang et al. (1996). Partial fragments of ITS region (internal transcribed spacer region of the ribosomal DNA containing ITS1, ITS2, and 5.8S rRNA gene) and β-tubulin gene of strain HUA were amplified for sequence analysis. The primers and conditions used to amplify the ITS region and the β-tubulin gene have been described previously (Glass and Donaldson 1995). The β-tubulin gene was sequenced to help clarify interspecific relationships when ITS region was inadequate for definitive identification. The amplified products were sequenced at Sangon Biotech (Shanghai, China). Alignment of the determined sequences and those of related ITS regions and β-tubulin sequences from the GenBank database by searching with BLAST program was aligned using ClustalX with manual adjustment (Thompson et al. 1997). Phylogenesis analysis was performed with MEGA4, and the evolutionary distances between the sequences were calculated using Kimura's two-parameter model (Tamura et al. 2007). The phylogenetic trees were constructed by the neighbor-joining method with bootstrap analysis (Saitou and Nei 1987).

The optimal temperature for isolates growth was determined by measuring the dry cell mass of mycelia under different temperatures. Flasks containing 100 ml of liquid basal medium inoculated with almost 10 mg (fresh weight) mycelia, which previously cultured aerobically in 100 ml basal medium on a shaker (180 rpm) at 30 °C for 5 days. The isolates were incubated at 20, 25, 30, 35, and 40 °C, the cultivation was carried out with shaking at 180 rpm for 7 days to investigate temperature profile. Mycelia were collected by filtration through filters and measured after drying to a constant weight at 105 °C.

### Formaldehyde resistance and degradation

Formaldehyde resistance and biodegradation of isolates with different initial formaldehyde concentrations were screened in liquid medium. Pre-culture of isolates, as described for characterization above, was inoculated in basal medium (50 ml) supplemented with 600–2,400 mg l<sup>-1</sup> of formaldehyde. In order to clarify the connection between formaldehyde degradation curve and fungal growth, fungal initial inoculation amount control in this experiment was 10 mg of fresh weight. Control of formaldehyde solution with same amount of the autoclaved mycelia was run in parallel. All cultivations were repeated five times. The flasks were incubated at 25 °C with shaking at 180 rpm for 7 days. The residual formaldehyde was determined as previous description (Nash 1953). Growth rates were determined by



measuring the fresh weight in time course. The experiments were performed in three replications.

#### The characterization of formaldehyde degradation


To characterize the formaldehyde degradation under aerobic condition efficiency of fungal strains, the effects of initial pH and supplemental carbon sources were investigated. The pre-culture (10 mg) of fungal strains was inoculated in autoclaved liquid basal medium supplemented with  $1,500 \text{ mg l}^{-1}$  formaldehyde. The pH of medium was adjusted from pH 4 to pH 8. Mycelia were collected as describe above after incubation at  $25 \text{ }^\circ\text{C}$  for 7 days. Formaldehyde-degrading abilities of the isolates under different carbon conditions were characterized by 4 different carbon media supplemented with  $1,500 \text{ mg l}^{-1}$  formaldehyde. Different carbon media were set by adding (w/v) 3 % glucose, 3 % lactose, and 3 % soluble starch replaced 3 % sucrose in basal medium. Basal medium with same amount of the autoclaved cells was set as control. In addition, basal medium without carbon source addition to


$1,500 \text{ mg l}^{-1}$  formaldehyde was set to detect formaldehyde degradation ability of HUA-utilizing formaldehyde as the sole carbon source. All cultivations were repeated five times. Cultivation was carried out with shaking at 180 rpm at  $25 \text{ }^\circ\text{C}$  for 7 days. Formaldehyde concentration was analyzed with methods described before. Fresh weight of mycelia and residual formaldehyde was measured every day.

#### Enzyme activity assay

A basal medium supplemented with  $300 \text{ mg l}^{-1}$  formaldehyde was set as enzyme-inducing medium, and a basal medium without formaldehyde was used as the control. Mycelia were then collected by filtration and washed well with sodium phosphate buffer (pH 7.5). Crude enzyme extract was obtained through sonication (4 V, 50 Hz) at  $4 \text{ }^\circ\text{C}$  for 30 s and centrifuged 13,000 rpm for 10 min. Enzyme activities of GSH-dependent FADH and FDH were assayed at 340 nm spectrophotometrically by monitoring the formation of NADH using a photometer (Hitachi

**Fig. 1** Characterization of HUA: **a** fungal colony grown on basal medium agar, **b** conidiogenous structures, **c** neighbor-joining phylogenetic tree based on  $\beta$ -tubulin sequences of HUA and other related taxa





**Fig. 2** Formaldehyde degradation of HUA with different initial formaldehyde concentration: **a** 1,500 mg l<sup>-1</sup> and **b**, 2,400 mg l<sup>-1</sup>

U-2800, Japan) (Schutte et al. 1976). The protein concentration of crude enzyme extracts was estimated according to Bradford method (1976).

#### Statistical analysis

Statistical analysis was performed using SPSS for Windows version 16.0 (SPSS Inc., USA) and Excel (Microsoft, USA). Graphical representations were presented using Sigma Plot for Windows version 11.0 (SPSS Inc., USA). Data were statistically evaluated using one-way analysis of variance, and significant differences between means were determined by Duncan's multiple range tests ( $p < 0.05$ ).

## Results and discussion

### Identification and characterization of fungal strain HUA

A fungal strain HUA was isolated on basal medium agar plate supplemented with 150 mg l<sup>-1</sup> formaldehyde through

a simple procedure. The colony color was white at first and then turned to olive, the reverse color was white initially and then turned to orange (Fig. 1a). The color of exudates and soluble pigments were brown and orange, respectively. Radiation pattern was found on both surface and medium. Sporangium and conidiogenous cells were globose-shaped (Fig. 1b). The ITS region of strain HUA was selected as the specific sequence for phylogenetic analysis. When comparing with public ITS region sequences in the GenBank database, HUA exhibited the highest ITS gene sequence similarity to *Aspergillus sydowii* (99 %) and *Aspergillus versicolor* (99 %), both belonged to *Aspergillus* section *versicolors*. The  $\beta$ -tubulin conserved sequence confirmed that strain HUA belonged to *A. sydowii* (Fig. 1c). Temperature had substantial effect on the dry cell mass of strain HUA. The strain could grow in the range of temperature 20–35 °C, and the optimum temperature for strain was 25 °C, whereas HUA could not grow at 40 °C.

### Formaldehyde resistance and degradation in liquid medium



The pre-culture strain HUA could completely degrade formaldehyde with initial concentration of 600, 900, 1,200, and 1,500 mg l<sup>-1</sup> within 4, 5, 6, 7 days, respectively. With an increase in mycelia mass, the formaldehyde concentration (1,500 mg l<sup>-1</sup>) reduced gradually in medium (Fig. 2a). As shown in Fig. 2a, in the first 2 days, the fungal mycelia mass and the concentration of formaldehyde were almost unchanged. However, mycelia mass increased rapidly after the second day. At the same time, the formaldehyde concentration decreased rapidly, the amounts of formaldehyde degraded by the strains HUA per gram of mycelia and per hour was 8.9 mg beyond 6.3 mg as previously reported about a formaldehyde-degrading fungus, *Trichoderma virens* BDF002 (Sawada et al. 2006). The strain HUA was able to grow in the presence of formaldehyde up to 2,400 mg l<sup>-1</sup> (Fig. 2b). Many fungi are resistant to formaldehyde (Kondo et al. 2008, 2002). According to Sawada et al. (2006), *Trichoderma virens* was identified as formaldehyde-resistant strains based on morphological and phylogenetic analyses.

### Effect of initial pH on formaldehyde degradation

The formaldehyde degradation was dependent upon pH values. The experiments were performed in the range of pH 4 to pH 8 for formaldehyde degrading. Formaldehyde removal was 40.04 % at pH 4, 89.49 % at pH 5, 82.12 % at pH 6, 100 % at pH 7, and 38.96 % at pH 8 after 7 days. It was observed that the biodegrading of formaldehyde



**Fig. 3** Formaldehyde degradation of HUA with initial formaldehyde concentration of 1500 mg l<sup>-1</sup> in different carbon sources: **a** fungal mass and **b** formaldehyde degradation



**Fig. 4** Enzyme activities of HUA grown on inducing medium supplemented with formaldehyde

increased between pH ranges from pH 5 to pH 7. Moreover, the mycelia mass also increased between pH 5 and pH 7. It indicates that fungal mycelia growth state affects

formaldehyde-degrading rates. In the previous paper on the effect of initial pH on the growth and formaldehyde-degrading rates of fungi, it was shown that the formaldehyde-degrading capacity affected by the pH values of the culture medium, where the optimum pH values were found to be between 4 and 9 in fungus cultures (Sawada et al. 2006). Fungi within the same genus have similar growth conditions. Similar results were observed, an optimum pH value range for heavy metal adsorption of fungi belonging to genus *Aspergillus* has been found from pH 5 to pH 7 (Taştan et al. 2010; Das et al. 2007).

Formaldehyde degradation with different carbon source

Strain HUA could grow in medium with sucrose, glucose, lactose, soluble starch, and formaldehyde as carbon source. These carbon sources had impact on fungal growth. Fresh weight of mycelia grew around the time from 0.01 to 0.24, 0.28, 0.18, 0.16, and 0.08 g under the conditions of sucrose, glucose, lactose, soluble starch, and formaldehyde after 7 days (Fig. 3a), respectively. The

degradation rates of formaldehyde were 100, 100, 99.5, 91.3, and 22.8 % (Fig. 3b), respectively. The amount of evaporation of formaldehyde was 4.8 % in control sample. The result showed strain HUA was able to grow and degrade formaldehyde effectively in carbon sources mentioned above.

*Aspergillus sydowii* spreads widely in ocean water (Martins et al. 2011; Rocha et al. 2010). *A. sydowii* could be used for field biodegradation of organophosphate pesticides (Hasan 1999). Recent studies have focused on the application of xylanase from *A. sydowii* (Nair et al. 2008). Closely related fungus *A. versicolor* has been found to play a considerable role in reducing polymer materials and heavy metals such as chromium, copper, and nickel and decolorizing Remazol Blue reactive dye (Taştan et al. 2010). The degradation of formaldehyde by *Aspergillus* section *versicolors* has not yet been reported.

#### Enzyme activity analysis

Degradation of formaldehyde by strain HUA would occur through the action of FADH and FDH. The activities of FADH and FDH were measured under formaldehyde-induced and non-induced conditions. FADH activity and FDH activity could only be detected under formaldehyde-induced condition. After 12 h of culturing with formaldehyde, FADH activity can be detected at a lower level that closely resembles FDH (Fig. 4). The activity of FADH reached a maximum after 24 h induced by formaldehyde, followed by a gradual decrease until to the undetectable levels at the fourth day. FADH activity can reach up to  $5.02 \text{ U mg}^{-1}$ . After induction of 24 h, FDH was possible to achieve the maximum, and after that, the activity was maintained to formaldehyde completely consumed. Activity of FDH achieved the highest  $1.06 \text{ U mg}^{-1}$ . The activity of formaldehyde dehydrogenase was higher than that of formate dehydrogenase during the entire formaldehyde metabolism.

Formaldehyde degradation pathways have been studied in certain prokaryotic and eukaryotic microorganisms. FADH has been found in methylotrophic bacteria, as well as in mammals, plants, and yeasts, where it plays a general role in formaldehyde detoxification and antioxidation (Vorholt 2002). In the present study, fungal FADH and FDH activities were induced by  $300 \text{ mg l}^{-1}$  formaldehyde and the enzyme activities were also change over time. The two enzymes were gradually induced and accumulated at the first 24 h. At this point, the strain HUA had the strongest formaldehyde degradation (data not shown), and formaldehyde concentration was reduced rapidly. After 48 h, the formaldehyde was completely consumed, the

FADH and FDH accumulated in strain HUA gradually restored to the initial level. The results indicate that there was a connection between formaldehyde degradation curve, fungal growth curves, and enzyme activities. Achkor et al. (2003) reported that transgenic *Arabidopsis* overexpressing *AtFADH* show a 25 % increase in the detoxification rate compared with the wild-type plants. Recently, FADH genes from *Arabidopsis thaliana*, golden pothos, and rice were isolated and characterized (Tada and Kidu 2011). Studies on gene cloning of this fungus were in progress to elucidate the gene regulation of formaldehyde metabolism. In addition, when glutathione or NAD was not added in reaction solution, the FADH activity could not be measured. This result clearly showed that formaldehyde dehydrogenase in strain HUA requires glutathione and NAD for activity, which is the same as *Candida boidinii* (Schutte et al. 1976). It indicates that FADH is one of the key enzymes in formaldehyde metabolism of strain HUA, and the strain HUA has potential applications for contaminated water treatment.

#### Conclusion

In this study, a formaldehyde-resistant fungal strain HUA was identified as a member of *A. sydowii*. The optimal temperature, pH, and carbon source conditions of culture preparation were investigated, and formaldehyde degradation characteristics were determined. The formaldehyde dehydrogenase could play an important role in metabolic formaldehyde process in strain HUA. From the above results, it can be concluded that *A. sydowii* HUA can degrade formaldehyde efficiently, and it could be successfully used for the treatment of industrial wastewater containing formaldehyde. Further experiment should be studied in natural condition to ascertain the exact formaldehyde degradation capacity.

**Acknowledgments** This study was supported by the Science Foundation for Distinguished Young Scholars of Heilongjiang Province, China (JC201109), the Graduate Innovation Fund of Heilongjiang Province (YJSCX2012-149HLJ), and Aid program for Science and Technology Innovative Research Team in Higher Educational Institutions of Heilongjiang Province (2010TD10) and Harbin Normal University (KJTD2011-2).

#### References

- Achkor H, Díaz M, Fernández MR, Biosca JA, Parés X, Martínez MC (2003) Enhanced formaldehyde detoxification by overexpression of glutathione-dependent formaldehyde dehydrogenase from *Arabidopsis*. *Plant Physiol* 132:2248–2255

- Arutchelvan V, Kanakasabai V, Nagarajan S, Muralikrishnan V (2005) Isolation and identification of novel high strength phenol degrading bacterial strains from phenol–formaldehyde resin manufacturing industrial wastewater. *J Hazard Mater* 127:238–243
- Ashraf MA, Maah MJ, Yusoff I (2011) Heavy metals accumulation in plants growing in ex tin mining catchment. *Int J Environ Sci Technol* 8:401–416
- Bhakta J, Munekage Y, Ohnishi K, Jana B (2012) Isolation and identification of cadmium- and lead-resistant lactic acid bacteria for application as metal removing probiotic. *Int J Environ Sci Technol* 9:433–440
- Bradford M (1976) A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
- Das SK, Das AR, Guha AK (2007) A study on the adsorption mechanism of mercury on *Aspergillus versicolor* biomass. *Environ Sci Technol* 41:8281–8287
- Di Maiuta N, Hubschmid S, Giuliani N, Schwarzenhuber P, Dow CS (2009) Microbial degradation of formaldehyde in white mineral dispersions preserved with formaldehyde-releasing biocides. *Int Biodeter Biodegr* 63:769–777
- Glass NL, Donaldson GC (1995) Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol* 61:1323–1330
- Hanson AD, Roje S (2001) One-carbon metabolism in higher plants. *Ann Rev Plant Physiol Plant Mol Biol* 52:119–137
- Hasan H (1999) Fungal utilization of organophosphate pesticides and their degradation by *Aspergillus flavus* and *A. sydowii* in soil. *Folia Microbiol* 44:77–84
- Hidalgo A, Lopategi A, Prieto M, Serra J, Llama M (2002) Formaldehyde removal in synthetic and industrial wastewater by *Rhodococcus erythropolis* UPV-1. *Appl Microbiol Biotechnol* 58:260–264
- IARC (2006) Formaldehyde, 2-Butoxyethanol, and 1-tert-Butoxy-2-Propanol. International Agency for Research on Cancer, Lyon
- Kanmani P, Aravind J, Preston D (2012) Remediation of chromium contaminants using bacteria. *Int J Environ Sci Technol* 9:183–193
- Kondo T, Morikawa Y, Hayashi N, Kitamoto N (2002) Purification and characterization of formate oxidase from a formaldehyde-resistant fungus. *FEMS Microbiol Lett* 214:137–142
- Kondo T, Morikawa Y, Hayashi N (2008) Purification and characterization of alcohol oxidase from *Paecilomyces variotii* isolated as a formaldehyde-resistant fungus. *Appl Microbiol Biotechnol* 77:995–1002
- Mani D, Kumar C (2013) Biotechnological advances in bioremediation of heavy metals contaminated ecosystems: an overview with special reference to phytoremediation. *Int J Environ Sci Technol* 10:1–30
- Martins M, Mouad A, Boschini L, Regali Selegim M, Sette L, Meleiro Porto A (2011) Marine fungi *Aspergillus sydowii* and *Trichoderma* sp. catalyze the hydrolysis of benzyl glycidyl ether. *Mar Biotechnol* 13:314–320
- Merk O, Speit G (1998) Significance of formaldehyde-induced DNA–protein crosslinks for mutagenesis. *Environ Mol Mutagen* 32:260–268
- Metz B, Kersten GFA, Hoogerhout P, Brugghe HF, Timmermans HAM, de Jong A, Meiring H, Hove JT, Hennink WE, Crommelin DJA, Jiskoot W (2004) Identification of formaldehyde-induced modifications in proteins. *J Biol Chem* 279:6235–6243
- More TT, Yan S, Tyagi RD, Surampalli RY (2010) Potential use of filamentous fungi for wastewater sludge treatment. *Bioresour Technol* 101:7691–7700
- Moussavi G, Heidarizad M (2010) Biodegradation of mixture of phenol and formaldehyde in wastewater using a single-basin MSCR process. *J Biotechnol* 150:240–245
- Nair S, Sindhu R, Shashidhar S (2008) Purification and biochemical characterization of two xylanases from *Aspergillus sydowii* SBS 45. *Appl Biochem Biotech* 149:229–243
- Nash T (1953) The colorimetric estimation of formaldehyde by means of the Hantzsch reaction. *Biochem J* 55:416–421
- Rocha L, Ferreira H, Pimenta E, Berlinck R, Rezende M, Landgraf M, Selegim M, Sette L, Porto A (2010) Biotransformation of  $\alpha$ -bromoacetophenones by the marine fungus *Aspergillus sydowii*. *Mar Biotechnol* 12:552–557
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sawada A, Ikeda R, Tamiya E, Yoshida T, Oyabu T, Nanto H (2006) A novel formaldehyde-degrading fungus, *Trichoderma virens*: isolation and some properties. *IEICE Trans Electron* 89:1786–1791
- Schutte H, Flossdorf J, Sahn H, Kula M-R (1976) Purification and properties of formaldehyde dehydrogenase and formate dehydrogenase from *Candida boidinii*. *Eur J Biochem* 62:151–160
- Solomon MJ, Varshavsky A (1985) Formaldehyde-mediated DNA–protein crosslinking: a probe for in vivo chromatin structures. *PNAS* 82:6470–6474
- Srivastava S, Thakur I (2006) Biosorption potency of *Aspergillus niger* for removal of chromium (VI). *Curr Microbiol* 53:232–237
- Tada Y, Kidu Y (2011) Glutathione-dependent formaldehyde dehydrogenase from golden pothos (*Epipremnum aureum*) and the production of formaldehyde detoxifying plants. *Plant Biotechnol* 28:373–378
- Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0. *Mol Biol Evol* 24:1596–1599
- Tang X, Bai Y, Duong A, Smith MT, Li L, Zhang L (2009) Formaldehyde in China: production, consumption, exposure levels, and health effects. *Environ Int* 35:1210–1224
- Taştan BE, Ertuğrul S, Dönmez G (2010) Effective bioremoval of reactive dye and heavy metals by *Aspergillus versicolor*. *Bioresour Technol* 101:870–876
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Tišler T, Zagorc-Končan J (1997) Comparative assessment of toxicity of phenol, formaldehyde, and industrial wastewater to aquatic organisms. *Water Air Soil Poll* 97:315–322
- Usharani K, Muthukumar M (2013) Optimization of aqueous methylparathion biodegradation by *Fusarium* sp. in batch scale process using response surface methodology. *Int J Environ Sci Technol* 10:591–606
- Vorholt J (2002) Cofactor-dependent pathways of formaldehyde oxidation in methylotrophic bacteria. *Arch Microbiol* 178:239–249
- Yurimoto H, Kato N, Sakai Y (2005) Assimilation, dissimilation, and detoxification of formaldehyde, a central metabolic intermediate of methylotrophic metabolism. *Chem Rec* 5:367–375
- Yurimoto H, Kato N, Sakai Y (2009) Genomic organization and biochemistry of the ribulose monophosphate pathway and its application in biotechnology. *Appl Microbiol Biotechnol* 84:407–416



- Zhang D, Yang Y, Castlebury LA, Cerniglia CE (1996) A method for the large scale isolation of high transformation efficiency fungal genomic DNA. *FEMS Microbiol Lett* 145:261–265
- Zhao J, Wang X, Zeng J, Yang G, Shi F, Yan Q (2005) Biodegradation of poly(butylene succinate-co-butylene adipate) by *Aspergillus versicolor*. *Polym Degrad Stab* 90:173–179

