

Hiram H. López · Rafael H. Villarreal · Leticia Zárate

# Complete intersection vanishing ideals on degenerate tori over finite fields

Received: 2 July 2012 / Accepted: 20 December 2012 / Published online: 30 January 2013  
© The Author(s) 2013. This article is published with open access at Springerlink.com

**Abstract** We study the complete intersection property and the algebraic invariants (index of regularity, degree) of vanishing ideals on degenerate tori over finite fields. We establish a correspondence between vanishing ideals and toric ideals associated to numerical semigroups. This correspondence is shown to preserve the complete intersection property, and allows us to use some available algorithms to determine whether a given vanishing ideal is a complete intersection. We give formulae for the degree, and for the index of regularity of a complete intersection in terms of the Frobenius number and the generators of a numerical semigroup.

**Mathematics Subject Classification** 13F20 · 13P25 · 14H45 · 11T71 · 94B25

## المخلص

ندرس خاصية التقاطع التام واللامتغيرات الجبرية (مؤشر الانتظام، الدرجة) لمثاليات متلاشية على طارات منحلة معرفة على حقول منتهية. نرسخ تقابلاً بين مثاليات متلاشية ومثاليات موطرة مرتبطة بأنصاف زمر عددية. تم إثبات أن هذا التقابل يحافظ على خاصية التقاطع التام ويسمح لنا باستخدام بعض الخوارزميات المتاحة لتحديد فيما إذا كان مثالي متلاش معطى تقاطعاً تاماً. نعطي صيغة للدرجة، ومؤشر الانتظام لتقاطع تام بدلالة عدد فروبينيس ومولدات نصف الزمرة العددية.

## 1 Introduction

Let  $K = \mathbb{F}_q$  be a finite field with  $q$  elements and let  $v_1, \dots, v_n$  be a sequence of positive integers. Consider the *degenerate projective torus*

$$X := \{[(x_1^{v_1}, \dots, x_n^{v_n})] \mid x_i \in K^* \text{ for all } i\} \subset \mathbb{P}^{n-1},$$

parameterized by the monomials  $x_1^{v_1}, \dots, x_n^{v_n}$ , where  $K^* = \mathbb{F}_q \setminus \{0\}$  and  $\mathbb{P}^{n-1}$  is a projective space over the field  $K$ . This set is a multiplicative group under componentwise multiplication. If  $v_i = 1$  for all  $i$ ,  $X$  is just a *projective torus*.

H. H. López · R. H. Villarreal (✉)  
Departamento de Matemáticas, Centro de Investigación y de Estudios Avanzados del IPN,  
Apartado Postal 14-740, 07000 Mexico City, DF, Mexico  
E-mail: vila@math.cinvestav.mx

H. H. López  
E-mail: hlopez@math.cinvestav.mx

L. Zárate  
CEFyMAP, Universidad Autónoma de Chiapas, 4a. Oriente Norte No. 1428. Entre 13a. y 14a. Norte,  
Col. Barrio La Pimienta, 29000 Tuxtla Gutiérrez, Chiapas, Mexico  
E-mail: leticia@math.cinvestav.mx



Let  $S = K[t_1, \dots, t_n] = \bigoplus_{d=0}^{\infty} S_d$  be a polynomial ring over the field  $K$  with the standard grading. Recall that the *vanishing ideal* of  $X$ , denoted by  $I(X)$ , is the ideal of  $S$  generated by the homogeneous polynomials that vanish on  $X$ . To study  $I(X)$ , we will associate with this a semigroup  $\mathcal{S}$  and a toric ideal  $P$  that depend on  $v_1, \dots, v_n$  and the multiplicative group of  $\mathbb{F}_q$ .

In what follows  $\beta$  denotes a generator of the cyclic group  $(K^*, \cdot)$ ,  $d_i$  denotes  $o(\beta^{v_i})$ , the order of  $\beta^{v_i}$  for  $i = 1, \dots, n$ , and  $\mathcal{S}$  denotes the semigroup  $\mathbb{N}d_1 + \dots + \mathbb{N}d_n$ . If  $d_1, \dots, d_n$  are relatively prime,  $\mathcal{S}$  is called a *numerical semigroup*. As is seen in Sect. 3, the algebra of  $I(X)$  is closely related to the algebra of the *toric ideal* of the semigroup ring

$$K[\mathcal{S}] = K[y_1^{d_1}, \dots, y_n^{d_n}] \subset K[y_1],$$

where  $K[y_1]$  is a polynomial ring. Recall that the *toric ideal* of  $K[\mathcal{S}]$ , denoted by  $P$ , is the kernel of the following epimorphism of  $K$ -algebras

$$\varphi: S = K[t_1, \dots, t_n] \longrightarrow K[\mathcal{S}], \quad f \longmapsto f(y_1^{d_1}, \dots, y_n^{d_n}).$$

Thus,  $S/P \simeq K[\mathcal{S}]$ . Since  $K[y_1]$  is integral over  $K[\mathcal{S}]$  we have  $\text{ht}(P) = n - 1$ . The ideal  $P$  is graded if one gives degree  $d_i$  to variable  $t_i$ . For  $n = 3$ , the first non-trivial case, this type of toric ideals were studied by Herzog [14]. For  $n \geq 4$ , these toric ideals have been studied by many authors [2, 4, 5, 8, 9, 23].

In this paper, we relate some of the algebraic invariants and properties of  $I(X)$  with those of  $P$  and  $\mathcal{S}$ . We are especially interested in the degree and the regularity index, and in the complete intersection property.

One of the most well known properties that  $P$  and  $I(X)$  have in common is that both are Cohen–Macaulay graded lattice ideals of dimension 1 [14, 19].

The contents of this paper are as follows. In Sect. 2, we introduce some of the notions that will be needed throughout the paper.

A key fact that allows us to link the properties of  $P$  and  $I(X)$  is that the homogeneous lattices of these ideals are closely related (Proposition 3.2). If  $g_1, \dots, g_m$  is a set of generators for  $P$  consisting of binomials, then  $h_1, \dots, h_m$  is a set of generators for  $I(X)$ , where  $h_k$  is the binomial obtained from  $g_k$  after substituting  $t_i$  by  $t_i^{d_i}$  for  $i = 1, \dots, n$  (Proposition 3.3). As a consequence if  $n = 3$ , then  $I(X)$  is minimally generated by 2 or 3 binomials (Corollary 3.4). If  $I(X)$  is a complete intersection, one of our main results shows that a minimal generating set for  $I(X)$  consisting of binomials corresponds to a minimal generating set for  $P$  consisting of binomials and viceversa (Theorem 3.6). As a consequence  $I(X)$  is a complete intersection if and only if  $P$  is a complete intersection (Corollary 3.7).

We show a formula for the degree of  $S/I(X)$  (Lemma 3.11). The *Frobenius number* of a numerical semigroup is the largest integer not in the semigroup. For complete intersections, we give a formula that relates the index of regularity of  $S/I(X)$  with the Frobenius number of the numerical semigroup generated by  $o(\beta^{rv_1}), \dots, o(\beta^{rv_n})$ , where  $r$  is the greatest common divisor of  $d_1, \dots, d_n$  (Corollary 3.13).

The Frobenius number occurs in many branches of mathematics and is one of the most studied invariants in the theory of semigroups. A great deal of effort has been directed at the effective computation of this number, see the monograph of Ramírez-Alfonsín [18].

The complete intersection property of  $P$  has been nicely characterized, using the notion of a binary tree [2, 4] and the notion of *suites distinguées* [5]. For  $n = 3$ , there is a classical result of [14] showing an algorithm to construct a generating set for  $P$ . Thus, using our results, one can obtain various classifications of the complete intersection property of  $I(X)$ . Furthermore, in [2] an effective algorithm is given to determine whether  $P$  is a complete intersection. This algorithm has been implemented in the distributed library `cimonom.lib` [3] of *Singular* [11]. Thus, using our results, one can use this algorithm to determine whether  $I(X)$  is a complete intersection (see Example 3.14). If  $I(X)$  is a complete intersection, this algorithm returns the generators of  $P$  and the Frobenius number. As a byproduct, we can construct interesting examples of complete intersection vanishing ideals (see Example 3.16).

We show how to compute the vanishing ideal  $I(X)$  using the notion of saturation of an ideal with respect to a polynomial (Proposition 3.18).

It is worth mentioning that our results could be applied to coding theory. The algebraic invariants of  $S/I(X)$  occur in algebraic coding theory as we now briefly explain. An *evaluation code* over  $X$  is a linear code obtained by evaluating the linear space of homogeneous  $d$ -forms of  $S$  on the set of points  $X \subset \mathbb{P}^{n-1}$ . A linear code obtained in this way, denoted by  $C_X(d)$ , has *length*  $|X|$  and *dimension*  $\dim_K(S/I(X))_d$ . The computation of the index of regularity of  $S/I(X)$  is important for applications to coding theory: for  $d \geq \text{reg } S/I(X)$  the code  $C_X(d)$  coincides with the underlying vector space  $K^{|X|}$  and has, accordingly, minimum distance equal

to 1. Thus, potentially good codes  $C_X(d)$  can occur only if  $1 \leq d < \text{reg}(S/I(X))$ . The length, dimension and minimum distance of evaluation codes  $C_X(d)$  arising from complete intersections have been studied in [6, 10, 12, 15, 16, 20, 21].

For all unexplained terminology and additional information, we refer to [7] (for the theory of lattice ideals), [22, 24] (for commutative algebra and the theory of Hilbert functions).

## 2 Preliminaries

We continue to use the notation and definitions used in Sect. 1. In this section, we introduce the notions of degree and regularity via Hilbert functions, and the notion of a lattice ideal.

The *Hilbert function* of  $S/I(X)$  is given by  $h_X(d) := \dim_K(S_d/I(X) \cap S_d)$ , and the *Krull-dimension* of  $S/I(X)$  is denoted by  $\dim(S/I(X))$ . The unique polynomial

$$h_X(t) = \sum_{i=0}^{k-1} c_i t^i \in \mathbb{Q}[t]$$

of degree  $k - 1 = \dim(S/I(X)) - 1$  such that  $h_X(d) = H_X(d)$  for  $d \gg 0$  is called the *Hilbert polynomial* of  $S/I(X)$ . The integer  $c_{k-1}(k - 1)!$ , denoted by  $\text{deg}(S/I(X))$ , is called the *degree* of  $S/I(X)$ . According to [13, Lecture 13],  $h_X(d) = |X|$  for  $d \geq |X| - 1$ . Hence

$$|X| = h_X(d) = c_0 = \text{deg}(S/I(X))$$

for  $d \geq |X| - 1$ . Thus,  $|X|$  is the degree of  $S/I(X)$ .

**Definition 2.1** The *index of regularity* of  $S/I(X)$ , denoted by  $\text{reg}(S/I(X))$ , is the least integer  $\ell \geq 0$  such that  $h_X(d) = H_X(d)$  for  $d \geq \ell$ .

The index of regularity of  $S/I(X)$  is equal to the Castelnuovo Mumford regularity of  $S/I(X)$  because this ring is Cohen–Macaulay of dimension 1.

*Remark 2.2* The Hilbert series of  $S/I(X)$  can be written as

$$F_X(t) := \sum_{i=0}^{\infty} H_X(i)t^i = \frac{h_0 + h_1t + \dots + h_r t^r}{1 - t},$$

where  $h_0, \dots, h_r$  are positive integers. The number  $r$  is the regularity index of  $S/I(X)$  and  $h_0 + \dots + h_r$  is the degree of  $S/I(X)$  (see [24, Corollary 4.1.12]). The same observation holds for any graded Cohen–Macaulay ideal  $I \subset S$  of height  $n - 1$ .

Recall that a binomial in  $S$  is a polynomial of the form  $t^a - t^b$ , where  $a, b \in \mathbb{N}^n$  and where, if  $a = (a_1, \dots, a_n) \in \mathbb{N}^n$ , we set

$$t^a = t_1^{a_1} \dots t_n^{a_n} \in S.$$

A *binomial ideal* is an ideal generated by binomials.

Given  $c = (c_i) \in \mathbb{Z}^n$ , the set  $\text{supp}(c) = \{i \mid c_i \neq 0\}$  is the *support* of  $c$ . The vector  $c$  can be written as  $c = c^+ - c^-$ , where  $c^+$  and  $c^-$  are two non-negative vectors with disjoint support. If  $t^a$  is a monomial, with  $a = (a_i) \in \mathbb{N}^n$ , the set  $\text{supp}(t^a) = \{t_i \mid a_i > 0\}$  is called the *support* of  $t^a$ .

**Definition 2.3** A subgroup  $\mathcal{L}$  of  $\mathbb{Z}^n$  is called a *lattice*. A *lattice ideal* is an ideal of the form

$$I(\mathcal{L}) = (\{t^{\alpha^+} - t^{\alpha^-} \mid \alpha \in \mathcal{L}\}) \subset S$$

for some lattice  $\mathcal{L}$  in  $\mathbb{Z}^n$ . A lattice  $\mathcal{L}$  is called *homogeneous* if there is an integral vector  $\omega$  with positive entries such that  $\langle \omega, a \rangle = 0$  for  $a \in \mathcal{L}$ .

**Definition 2.4** An ideal  $I \subset S$  is called a *complete intersection* if there exists  $g_1, \dots, g_m$  such that  $I = (g_1, \dots, g_m)$ , where  $m$  is the height of  $I$ .

*Remark 2.5* A graded binomial ideal  $I \subset S$  is a complete intersection if and only if  $I$  is generated by a set of homogeneous binomials  $g_1, \dots, g_m$ , with  $m = \text{ht}(I)$ , and any such set of homogeneous generators is already a regular sequence (see [24, Proposition 1.3.17, Lemma 1.3.18]).

**Lemma 2.6** *Let  $S = K[t_1, \dots, t_n]$  be a polynomial ring with the standard grading. If  $I$  is a graded ideal of  $S$  generated by a homogeneous regular sequence  $f_1, \dots, f_{n-1}$ , then*

$$\text{reg}(S/I) = \sum_{i=1}^{n-1} (\text{deg}(f_i) - 1) \quad \text{and} \quad \text{deg}(S/I) = \text{deg}(f_1) \cdots \text{deg}(f_{n-1}).$$

*Proof* We set  $\delta_i = \text{deg}(f_i)$ . By Villarreal [24, p. 104], the Hilbert series of  $S/I$  is given by

$$F_I(t) = \frac{\prod_{i=1}^{n-1} (1 - t^{\delta_i})}{(1 - t)^n} = \frac{\prod_{i=1}^{n-1} (1 + t + \cdots + t^{\delta_i - 1})}{(1 - t)}. \tag{2.1}$$

Thus, by Remark 2.2,  $\text{reg}(S/I) = \sum_{i=1}^{n-1} (\delta_i - 1)$  and  $\text{deg}(S/I) = \delta_1 \cdots \delta_{n-1}$ . □

### 3 Complete intersections and algebraic invariants

We continue to use the notation and definitions used in Sects. 1 and 2. In this section, we study vanishing ideals over degenerate projective tori. We study the complete intersection property and the algebraic invariants of vanishing ideals. We will establish a correspondence between vanishing ideals and toric ideals associated to semigroups of  $\mathbb{N}$ .

Let  $D$  be the non-singular matrix  $D = \text{diag}(d_1, \dots, d_n)$ . Consider the homomorphisms of  $\mathbb{Z}$ -modules:

$$\begin{aligned} \psi: \mathbb{Z}^n &\rightarrow \mathbb{Z}, & e_i &\mapsto d_i, \\ D: \mathbb{Z}^n &\rightarrow \mathbb{Z}^n, & e_i &\mapsto d_i e_i. \end{aligned}$$

If  $c = (c_i) \in \mathbb{R}^n$ , we set  $|c| = \sum_{i=1}^n c_i$ . Notice that  $|D(c)| = \psi(c)$  for any  $c \in \mathbb{Z}^n$ . There are two homogeneous lattices that will play a role here:

$$\mathcal{L}_1 = \ker(\psi) \quad \text{and} \quad \mathcal{L} = D(\ker(\psi)).$$

The map  $D$  induces a  $\mathbb{Z}$ -isomorphism between  $\mathcal{L}_1$  and  $\mathcal{L}$ . It is well known [24] that the toric ideal  $P$  is the lattice ideal of  $\mathcal{L}_1$ . Below, we show that  $I(X)$  is the lattice ideal of  $\mathcal{L}$ .

**Lemma 3.1** *The map  $t^a - t^b \mapsto t^{D(a)} - t^{D(b)}$  induces a bijection between the binomials  $t^a - t^b$  of  $P$  whose terms  $t^a, t^b$  have disjoint support and the binomials  $t^{a'} - t^{b'}$  of  $I(X)$  whose terms  $t^{a'}, t^{b'}$  have disjoint support.*

*Proof* If  $f = t^a - t^b$  is a binomial of  $P$  whose terms have disjoint support, then  $a - b \in \mathcal{L}_1$  and the terms of  $g = t^{D(a)} - t^{D(b)}$  have disjoint support because

$$\text{supp}(t^a) = \text{supp}(t^{D(a)}) \quad \text{and} \quad \text{supp}(t^b) = \text{supp}(t^{D(b)}).$$

Thus,  $|D(a)| = \psi(a) = \psi(b) = |D(b)|$ . This means that  $g = t^{D(a)} - t^{D(b)}$  is homogeneous in the standard grading of  $S$ . As  $(\beta^{v_i})^{d_i} = 1$  for all  $i$ , it is seen that  $g$  vanishes at all points of  $X$ . Hence,  $g \in I(X)$  and the map is well defined.

The map is clearly injective. To show that the map is onto, take a binomial  $f' = t^{a'} - t^{b'}$  in  $I(X)$  with  $a' = (a'_i), b' = (b'_i)$  and such that  $t^{a'}$  and  $t^{b'}$  have disjoint support. Then,  $(\beta^{v_i})^{a'_i - b'_i} = 1$  for all  $i$  because  $f'$  vanishes at all points of  $X$ . Hence, since the order of  $\beta^{v_i}$  is  $d_i$ , there are integers  $c_1, \dots, c_n$  such that  $a'_i - b'_i = c_i d_i$  for all  $i$ . Since  $f'$  is homogeneous, one has  $|a'| = |b'|$ . It follows readily that  $c \in \mathcal{L}_1$  and  $a' - b' = D(c)$ . We can write  $c = c^+ - c^-$ . As  $a'$  and  $b'$  have disjoint support, we get  $a' = D(c^+)$  and  $b' = D(c^-)$ . Thus, the binomial  $f = t^{c^+} - t^{c^-}$  is in  $P$  and maps to  $t^{a'} - t^{b'}$ . □

**Proposition 3.2**  $P = I(\mathcal{L}_1)$  and  $I(X) = I(\mathcal{L})$ .

*Proof* As mentioned above, the first equality is well known [24]. Since  $I(X)$  is a lattice ideal [19], it is generated by binomials of the form  $t^{a^+} - t^{a^-}$  (this follows using that  $t_i$  is a non-zero divisor of  $S/I(X)$  for all  $i$ ). To show the second equality, take  $t^{a^+} - t^{a^-}$  in  $I(X)$ . Then, by Lemma 3.1,  $a^+ - a^- \in \mathcal{L}$  and  $t^{a^+} - t^{a^-}$  is in  $I(\mathcal{L})$ . Thus,  $I(X) \subset I(\mathcal{L})$ . Conversely, take  $f = t^{a^+} - t^{a^-}$  in  $I(\mathcal{L})$  with  $a^+ - a^-$  in  $\mathcal{L}$ . Then, there is  $c \in \mathcal{L}_1$  such that  $a^+ - a^- = D(c^+ - c^-)$ . Then,  $t^{c^+} - t^{c^-}$  is in  $P$  and maps, under the map of Lemma 3.1, to  $f$ . Thus,  $f \in I(X)$ . This proves that  $I(\mathcal{L}) \subset I(X)$ .  $\square$

**Proposition 3.3** *If  $P = (\{t^{a_i} - t^{b_i}\}_{i=1}^m)$ , then  $I(X) = (\{t^{D(a_i)} - t^{D(b_i)}\}_{i=1}^m)$ .*

*Proof* We set  $g_i = t^{a_i} - t^{b_i}$  and  $h_i = t^{D(a_i)} - t^{D(b_i)}$  for  $i = 1, \dots, n$ . Notice that  $h_i$  is equal to  $g_i(t_1^{d_1}, \dots, t_n^{d_n})$ , the evaluation of  $g_i$  at  $(t_1^{d_1}, \dots, t_n^{d_n})$ . By Lemma 3.1, one has the inclusion  $(h_1, \dots, h_m) \subset I(X)$ . To show the reverse inclusion take a binomial  $0 \neq f \in I(X)$ . We may assume that  $f = t^{a^+} - t^{a^-}$ . Then, by Lemma 3.1, there is  $g = t^{c^+} - t^{c^-}$  in  $P$  such that  $f = t^{D(c^+)} - t^{D(c^-)}$ . By hypothesis we can write  $g = \sum_{i=1}^m f_i g_i$  for some  $f_1, \dots, f_m$  in  $S$ . Then, evaluating both sides of this equality at  $(t_1^{d_1}, \dots, t_n^{d_n})$ , we get

$$f = t^{D(c^+)} - t^{D(c^-)} = g(t_1^{d_1}, \dots, t_n^{d_n}) = \sum_{i=1}^m f_i(t_1^{d_1}, \dots, t_n^{d_n})g_i(t_1^{d_1}, \dots, t_n^{d_n}) = \sum_{i=1}^m f_i' h_i,$$

where  $f_i' = f_i(t_1^{d_1}, \dots, t_n^{d_n})$  for all  $i$ . Then,  $f \in (h_1, \dots, h_m)$ .  $\square$

**Corollary 3.4** *If  $n = 3$ , then  $I(X)$  is minimally generated by at most 3 binomials.*

*Proof* By a classical theorem of Herzog [14],  $P$  is generated by at most 3 binomials. Hence, by Proposition 3.3,  $I(X)$  is generated by at most 3 binomials.  $\square$

Given a subset  $I \subset S$ , its *variety*, denoted by  $V(I)$ , is the set of all  $a \in \mathbb{A}_K^n$  such that  $f(a) = 0$  for all  $f \in I$ , where  $\mathbb{A}_K^n$  is the affine space over  $K$ . Given a binomial  $g = t^a - t^b$ , we set  $\widehat{g} = a - b$ . If  $B$  is a subset of  $\mathbb{Z}^n$ ,  $\langle B \rangle$  denotes the subgroup of  $\mathbb{Z}^n$  generated by  $B$ .

**Proposition 3.5** [4, Proposition 2.5] *Let  $\mathcal{B} = \{g_1, \dots, g_{n-1}\}$  be a set of binomials in  $P$ . Then,  $P = \langle \mathcal{B} \rangle$  if and only if the following two conditions hold:*

- (i')  $\mathcal{L}_1 = \langle \widehat{g}_1, \dots, \widehat{g}_{n-1} \rangle$ , where  $\mathcal{L}_1 = \ker(\psi)$ .
- (ii')  $V(g_1, \dots, g_{n-1}, t_i) = \{0\}$  for  $i = 1, \dots, n$ .

We come to the main result of this section.

**Theorem 3.6** (a) *If  $I(X)$  is a complete intersection generated by binomials  $h_1, \dots, h_{n-1}$ , then  $P$  is a complete intersection generated by binomials  $g_1, \dots, g_{n-1}$  such that  $h_i$  is equal to  $g_i(t_1^{d_1}, \dots, t_n^{d_n})$  for all  $i$ .* (b) *If  $P$  is a complete intersection generated by binomials  $g_1, \dots, g_{n-1}$ , then  $I(X)$  is a complete intersection generated by binomials  $h_1, \dots, h_{n-1}$ , where  $h_i$  is equal to  $g_i(t_1^{d_1}, \dots, t_n^{d_n})$  for all  $i$ .*

*Proof* (a) Since  $t_k$  is a non-zero divisor of  $S/I(X)$  for all  $k$ , it is not hard to see that the monomials of  $h_i$  have disjoint support for all  $i$ , i.e., we can write  $h_i = t^{a_i^+} - t^{a_i^-}$  for  $i = 1, \dots, n - 1$ . We claim that the following two conditions hold.

- (i)  $\mathcal{L} = \langle a_1, \dots, a_{n-1} \rangle$ , where  $a_i = a_i^+ - a_i^-$  and  $\mathcal{L}$  is the lattice that defines  $I(X)$ .
- (ii)  $V(h_1, \dots, h_{n-1}, t_i) = \{0\}$  for  $i = 1, \dots, n$ .

As  $I(X)$  is generated by  $h_1, \dots, h_{n-1}$ , by López and Villarreal [17, Lemma 2.5], condition (i) holds. The binomial  $t_i^{q-1} - t_n^{q-1}$  is in  $I(X)$  for all  $i$  because  $\mathbb{F}_q^*$  is a group of order  $q - 1$ . Thus,  $V(I(X), t_i) = \{0\}$  for all  $i$ . From the equality  $(h_1, \dots, h_{n-1}, t_i) = (I(X), t_i)$ , we get

$$V(h_1, \dots, h_{n-1}, t_i) = V(I(X), t_i) = \{0\}.$$

Thus, (ii) holds. This completes the proof of the claim.

By (i) and Proposition 3.2, there are  $b_1, \dots, b_{n-1}$  in  $\mathcal{L}_1 = \ker(\psi)$  such that  $a_i = D(b_i)$  for all  $i$ . Accordingly  $a_i^+ = D(b_i^+)$  and  $a_i^- = D(b_i^-)$  for all  $i$ . We set  $g_i = t^{b_i^+} - t^{b_i^-}$  for all  $i$ . Clearly, all the  $g_i$ 's are in  $P$  and  $h_i$  is equal to  $g_i(t_1^{d_1}, \dots, t_n^{d_n})$  for all  $i$ . Next, we prove that  $P$  is generated by  $g_1, \dots, g_{n-1}$ . By Proposition 3.5 it suffices to show that the following two conditions hold:

- (i')  $\mathcal{L}_1 = \langle b_1, \dots, b_{n-1} \rangle$ , where  $\mathcal{L}_1 = \ker(\psi)$ .
- (ii')  $V(g_1, \dots, g_{n-1}, t_i) = \{0\}$  for  $i = 1, \dots, n$ .

First we show (i'). Since  $b_1, \dots, b_{n-1}$  are in  $\mathcal{L}_1$ , we need only show the inclusion “ $\subset$ ”. Take  $\gamma \in \ker(\psi)$ , then  $D(\gamma) \in \mathcal{L}$ , and by (i) it follows that  $\gamma \in \langle b_1, \dots, b_{n-1} \rangle$ .

Next we show (ii'). For simplicity of notation, we may assume that  $i = n$ . Take  $c$  in the variety  $V(g_1, \dots, g_{n-1}, t_n)$  and write  $c = (c_1, \dots, c_n)$ . Then,  $c_n = 0$  and  $g_i(c) = c^{b_i^+} - c^{b_i^-} = 0$  for all  $i$ , were  $c^{b_i^+}$  means to evaluate the monomial  $t^{b_i^+}$  at the point  $c$ . Let  $i$  be a fixed but arbitrary integer in  $\{1, \dots, n - 1\}$ . We can write

$$b_i = b_i^+ - b_i^- = (b_{i1}^+, \dots, b_{in}^+) - (b_{i1}^-, \dots, b_{in}^-)$$

and  $a_i = a_i^+ - a_i^- = (a_{i1}^+, \dots, a_{in}^+) - (a_{i1}^-, \dots, a_{in}^-)$ . Then

$$\begin{aligned} h_i(c_1^{v_1}, \dots, c_n^{v_n}) &= (c_1^{v_1})^{a_{i1}^+} \dots (c_n^{v_n})^{a_{in}^+} - (c_1^{v_1})^{a_{i1}^-} \dots (c_n^{v_n})^{a_{in}^-} \\ &= c_1^{v_1 d_1 b_{i1}^+} \dots c_n^{v_n d_n b_{in}^+} - c_1^{v_1 d_1 b_{i1}^-} \dots c_n^{v_n d_n b_{in}^-}. \end{aligned} \tag{3.1}$$

We claim that  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$ . To show this we consider two cases.

Case (I):  $b_{in}^+ > 0$ . Then, as  $g_i(c) = c^{b_i^+} - c^{b_i^-} = 0$  and  $c^{b_i^+} = 0$ , one has  $c^{b_i^-} = 0$ . Hence, there is  $j$  such that  $b_{ij}^- > 0$  and  $c_j = 0$ . Thus, by Eq. (3.1),  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$ .

Case (II):  $b_{in}^+ = 0$ . If  $c_j = 0$  for some  $b_{ij}^+ > 0$ , then  $c^{b_i^-} = 0$  because  $g_i(c) = 0$ . Hence, there is  $k$  such that  $c_k = 0$  and  $b_{ik}^- > 0$ . Thus, by Eq. (3.1),  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$ . Similarly, if  $c_j = 0$  for some  $b_{ij}^- > 0$ , then  $c^{b_i^+} = 0$  because  $g_i(c) = 0$ . Hence, there is  $k$  such that  $c_k = 0$  and  $b_{ik}^+ > 0$ . Thus, by Eq. (3.1),  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$ . We may now assume that  $c_j \neq 0$  if  $b_{ij}^+ > 0$ , and  $c_m \neq 0$  if  $b_{im}^- > 0$ . Let  $\beta$  be a generator of the cyclic group  $(\mathbb{F}^*, \cdot)$ . Any  $c_j \neq 0$  has the form  $c_j = \beta^{j\epsilon}$ . Thus, using that  $(\beta^{v_j})^{d_j} = 1$ , we get that  $(c_j^{v_j})^{d_j b_{ij}^+} = 1$  if  $b_{ij}^+ > 0$  and  $(c_j^{v_j})^{d_j b_{ij}^-} = 1$  if  $b_{ij}^- > 0$ . Hence, by Eq. (3.1),  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$ , as required. This completes the proof of the claim.

As  $h_i(c_1^{v_1}, \dots, c_n^{v_n}) = 0$  for all  $i$ , the point  $c' = (c_1^{v_1}, \dots, c_n^{v_n})$  is in  $V(h_1, \dots, h_{n-1}, t_n)$ . By (ii), the point  $c'$  is zero. Hence,  $c = 0$  as required. This completes the proof of (ii'). Hence,  $P$  is a complete intersection generated by  $g_1, \dots, g_{n-1}$ .

(b) It follows from Proposition 3.3. □

Using the notion of a binary tree, a criterion for complete intersection toric ideals of affine monomial curves is given in [4]. In [2] an effective algorithm is given to determine whether  $P$  is a complete intersection. If  $P$  is a complete intersection, this algorithm returns the generators of  $P$  and the Frobenius number.

In our situation, the next result allows us to: (A) use the results of [4,5,14] to give criteria for complete intersection vanishing ideals over a finite field, (B) use the effective algorithms of [2] to recognize complete intersection vanishing ideals over finite fields and to compute its invariants (see Example 3.14).

**Corollary 3.7**  *$I(X)$  is a complete intersection if and only if  $P$  is a complete intersection.*

*Proof* Assume that  $I(X)$  is a complete intersection. By Remark 2.5, there are binomials  $h_1, \dots, h_{n-1}$  that generate  $I(X)$ . Hence,  $P$  is a complete intersection by Theorem 3.6. The converse follows by similar reasons. □

**Lemma 3.8** *If  $r = \gcd(d_1, \dots, d_n)$  and  $d'_i = o(\beta^{rv_i})$ , then  $d_i = rd'_i$  and  $\gcd(d'_1, \dots, d'_n) = 1$ .*

*Proof* It follows readily by recalling that  $o(\beta^{rv_i}) = o(\beta^{v_i}) / \gcd(r, o(\beta^{v_i}))$ . □

In what follows  $X'$  will denote the degenerate torus in  $\mathbb{P}^{n-1}$  parameterized by  $x_1^{v'_1}, \dots, x_n^{v'_n}$ , where  $v'_i = rv_i$  and  $r = \gcd(d_1, \dots, d_n)$ . Below, we relate  $I(X)$  and  $I(X')$ .

**Proposition 3.9** *The vanishing ideal  $I(X)$  is a complete intersection if and only if  $I(X')$  is a complete intersection.*

*Proof* Let  $P$  and  $P'$  be the toric ideals of  $K[y_1^{d_1}, \dots, y_n^{d_n}]$  and  $K[y_1^{d'_1}, \dots, y_n^{d'_n}]$ , respectively, where  $d'_i = o(\beta^{rv_i})$  for all  $i$ . It is not hard to see that  $P = P'$ . Then, by Theorem 3.6,  $P$  is a complete intersection if and only if  $I(X)$  is a complete intersection and  $P'$  is a complete intersection if and only if  $I(X')$  is a complete intersection. Thus,  $I(X)$  is a complete intersection if and only if  $I(X')$  is a complete intersection.  $\square$

**Definition 3.10** The set  $X^* := \{(x_1^{v_1}, \dots, x_n^{v_n}) \mid x_i \in K^* \text{ for all } i\} \subset K^n$  is called an *affine degenerate torus* parameterized by  $x_1^{v_1}, \dots, x_n^{v_n}$ .

**Lemma 3.11**  $|X^*| = d_1 \cdots d_n$  and  $\deg(S/I(X)) = |X| = d_1 \cdots d_n / \gcd(d_1, \dots, d_n)$ .

*Proof* Let  $S_i = \langle \beta^{v_i} \rangle$  be the cyclic group generated by  $\beta^{v_i}$ . The set  $X^*$  is equal to the Cartesian product  $S_1 \times \cdots \times S_n$ . Hence, to show the first equality, it suffices to recall that  $|S_i|$  is  $o(\beta^{v_i})$ , the order of  $\beta^{v_i}$ . Notice that any element of  $X^*$  can be written as  $((\beta^{i_1})^{v_1}, \dots, (\beta^{i_n})^{v_n})$  for some integers  $i_1, \dots, i_n$ . The kernel of the epimorphism of groups  $X^* \mapsto X, x \mapsto [x]$ , is equal to

$$\{(\gamma, \dots, \gamma) \in (K^*)^n : \gamma \in \langle \beta^{v_1} \rangle \cap \cdots \cap \langle \beta^{v_n} \rangle\}.$$

Hence,  $|X^*|/|\cap_{i=1}^n \langle \beta^{v_i} \rangle| = |X|$ . Since  $\langle \beta^{v_i} \rangle$  is a subgroup of  $K^*$  for all  $i$  and  $K^*$  is a cyclic group, one has  $|\cap_{i=1}^n \langle \beta^{v_i} \rangle| = \gcd(d_1, \dots, d_n)$  (see for instance [1, Theorem 4, p. 4]). Thus, the second equality follows.  $\square$

**Definition 3.12** If  $\mathcal{S}$  is a numerical semigroup of  $\mathbb{N}$ , the *Frobenius number* of  $\mathcal{S}$ , denoted by  $g(\mathcal{S})$ , is the largest integer not in  $\mathcal{S}$ .

Consider the semigroup  $\mathcal{S}' = \mathbb{N}d'_1 + \cdots + \mathbb{N}d'_n$ , where  $d'_i = o(\beta^{rv_i})$  for  $i = 1, \dots, n$ . By Lemma 3.8, one has  $\gcd(d'_1, \dots, d'_n) = 1$ , i.e.,  $\mathcal{S}'$  is a numerical semigroup. Thus,  $g(\mathcal{S}')$  is finite. If the toric ideal of  $K[\mathcal{S}']$  is a complete intersection, then  $g(\mathcal{S}')$  can be expressed entirely in terms of  $d'_1, \dots, d'_n$  [4, Remark 4.5].

**Corollary 3.13** (i)  $\deg(S/I(X)) = d_1 \cdots d_n / \gcd(d_1, \dots, d_n)$ .  
 (ii) If  $I(X)$  is a complete intersection, then

$$\text{reg} S/I(X) = \gcd(d_1, \dots, d_n)g(\mathcal{S}') + \sum_{i=1}^n d_i - (n - 1).$$

*Proof* Part (i) follows at once from Lemma 3.11. Next, we prove (ii). Let  $P$  and  $P'$  be as in the proof of Proposition 3.9. With the notation above, by Lemma 3.8, we get that  $d_i = rd'_i$  for all  $i$ . The toric ideals  $P$  and  $P'$  are equal but they are graded differently. Recall that  $P$  and  $P'$  are graded with respect to the gradings induced by assigning  $\deg(t_i) = d_i$  and  $\deg(t_i) = d'_i$  for all  $i$ , respectively. Let  $g_1, \dots, g_{n-1}$  be a generating set of  $P = P'$  consisting of binomials. Then, by Theorem 3.6,  $I(X)$  is generated by  $h_1, \dots, h_{n-1}$ , where  $h_i$  is  $g_i(t_1^{d_1}, \dots, t_n^{d_n})$  for all  $i$ . Accordingly,  $I(X')$  is generated by  $h'_1, \dots, h'_{n-1}$ , where  $h'_i$  is  $g_i(t_1^{d'_1}, \dots, t_n^{d'_n})$  for all  $i$ . If  $D_i = \deg(h_i)$  and  $D'_i = \deg(h'_i)$ , then  $D_i = rD'_i$  for all  $i$ . As  $P'$  is a complete intersection generated by  $g_1, \dots, g_{n-1}$  and  $\deg_{P'}(g_i) = D'_i$  for all  $i$ , using [4, Remark 4.5], we get

$$g(\mathcal{S}') = \sum_{i=1}^{n-1} D'_i - \sum_{i=1}^n d'_i = \sum_{i=1}^{n-1} (D_i/r) - \sum_{i=1}^n (d_i/r).$$

Therefore, using the equality  $\text{reg} S/I(X) = \sum_{i=1}^{n-1} (D_i - 1)$  (see Lemma 2.6), the formula for the regularity follows.  $\square$

*Example 3.14* To illustrate how to use the algorithm of [2] we consider the degenerate torus  $X$ , over the field  $\mathbb{F}_q$ , parameterized by  $x_1^{v_1}, \dots, x_5^{v_5}$ , where  $v_1 = 1,500, v_2 = 1,000, v_3 = 432, v_4 = 360, v_5 = 240$ , and  $q = 54,001$ . In this case, one has

$$d_1 = 36, \quad d_2 = 54, \quad d_3 = 125, \quad d_4 = 150, \quad d_5 = 225.$$

Using [2, Algorithm CI, p. 981], we get that  $P$  is a complete intersection generated by the binomials

$$g_1 = t_1^3 - t_2^2, \quad g_2 = t_4^3 - t_5^2, \quad g_3 = t_3^3 - t_4t_5, \quad g_4 = t_1^8t_2^3 - t_4^3,$$

and we also get that the Frobenius number of  $S$  is 793. Hence, by our results, the vanishing ideal  $I(X)$  is a complete intersection generated by the binomials

$$h_1 = t_1^{108} - t_2^{108}, \quad h_2 = t_4^{450} - t_5^{450}, \quad h_3 = t_3^{375} - t_4^{150}t_5^{225}, \quad h_4 = t_1^{288}t_2^{162} - t_4^{450},$$

the index of regularity and degree of  $S/I(X)$  are 1,379 and 8,201,250,000, respectively.

The next example is interesting because if one chooses  $v_1, \dots, v_n$  at random, it is likely that  $I(X)$  will be generated by binomials of the form  $t_i^m - t_j^m$ .

*Example 3.15* Let  $\mathbb{F}_q$  be the field with  $q = 211$  elements. Consider the sequence  $v_1 = 42, v_2 = 35, v_3 = 30$ . In this case, one has  $d_1 = 5, d_2 = 6, d_3 = 7$ . By a well known result of Herzog [14], one has

$$P = (t_2^2 - t_1t_3, t_1^4 - t_2t_3^2, t_1^3t_2 - t_3^3).$$

Hence, by our results,  $I(X) = (t_2^{12} - t_1^5t_3^7, t_1^{20} - t_2^6t_3^{14}, t_1^{15}t_2^6 - t_3^{21})$  and this ideal is not a complete intersection. The index of regularity and the degree of  $S/I(X)$  are 25 and 210, respectively. The Frobenius number of  $S$  is equal to 9. Notice that the toric relations  $t_1^{30} - t_2^{30}, t_1^{35} - t_3^{35}, t_2^{42} - t_3^{42}$  do not generate  $I(X)$ .

The next example was found using Theorem 3.6. Without using this theorem it is very difficult to construct examples of complete intersection vanishing ideals not generated by binomials of the form  $t_i^m - t_j^m$ .

*Example 3.16* Let  $\mathbb{F}_q$  be the field with  $q = 271$  elements. Consider the sequence  $v_1 = 30, v_2 = 135, v_3 = 54$ . In this case, one has  $d_1 = 9, d_2 = 2, d_3 = 5$ . The ideals  $P$  and  $I(X)$  are complete intersections given by

$$P = (t_1 - t_2^2t_3, t_2^5 - t_3^2) \quad \text{and} \quad I(X) = (t_1^9 - t_2^4t_3^5, t_2^{10} - t_3^{10}).$$

By Lemma 2.6, the index of regularity of  $S/I(X)$  is 17 and by Corollary 3.13 the Frobenius number of  $S$  is 3.

*The computation of the vanishing ideal.* In this part we show how to compute the vanishing ideal using the notion of saturation of an ideal with respect to a polynomial.

The next lemma is easy to show.

**Lemma 3.17** *If  $c_{ij} := \text{lcm}\{d_i, d_j\} = \text{lcm}\{o(\beta^{v_i}), o(\beta^{v_j})\}$ , then  $t_i^{c_{ij}} - t_j^{c_{ij}} \in I(X)$ .*

The set of toric relations  $\mathcal{T} = \{t_i^{c_{ij}} - t_j^{c_{ij}} : 1 \leq i, j \leq n\}$  does not generate  $I(X)$ , as is seen in Example 3.15. If  $v_i = 1$  for all  $i$ , then  $c_{ij} = q - 1$  for all  $i, j$  and  $I(X)$  is generated by  $\mathcal{T}$ .

For an ideal  $I \subset S$  and a polynomial  $h \in S$  the saturation of  $I$  with respect to  $h$  is the ideal

$$(I : h^\infty) := \{f \in S \mid fh^k \in I \text{ for some } k \geq 1\}.$$

**Proposition 3.18** *Let  $I'$  be the ideal  $(t_i^{c_{ij}} - t_j^{c_{ij}} \mid 1 < i < j \leq n)$ , where  $c_{ij} = \text{lcm}\{d_i, d_j\}$ . If  $\text{gcd}(d_1, \dots, d_n) = 1$ , then  $I(X) = (I' : (t_1 \cdots t_n)^\infty)$ .*

*Proof* We claim that  $\mathcal{L} = \langle c_{ij}e_i - c_{ij}e_j \mid 1 \leq i < j \leq n \rangle$ . By Villarreal [24, Proposition 10.1.8], we get

$$\mathcal{L}_1 = \langle (d_j / \text{gcd}(d_i, d_j))e_i - (d_i / \text{gcd}(d_i, d_j))e_j \mid 1 \leq i < j \leq n \rangle.$$

Thus, the claim follows from the equality  $\mathcal{L} = D(\mathcal{L}_1)$ . The inclusion “ $\supset$ ” follows readily using that  $t_i$  is a non-zero divisor of  $S/I(X)$  for all  $i$  because  $I(X)$  is a lattice ideal containing  $I'$  (see Lemma 3.17). To show the inclusion “ $\subset$ ”, take a binomial  $f = t^a - t^b \in I(X)$ . By Proposition 3.2,  $I(X) = I(\mathcal{L})$ . Thus,  $a - b \in \mathcal{L}$ . Using the previous claim and [17, Lemma 2.3], there is  $\delta \in \mathbb{N}^n$  such that  $t^\delta f \in I'$ . Hence,  $f \in (I' : (t_1 \cdots t_n)^\infty)$ . □

**Acknowledgements** R. H. Villarreal and L. Zárate were partially supported by SNI.

**Open Access** This article is distributed under the terms of the Creative Commons Attribution License which permits any use, distribution, and reproduction in any medium, provided the original author(s) and the source are credited.



## References

1. Alperin, J.L.; Bell, R.B.: Groups and Representations. Graduate Texts in Mathematics, vol. 162. Springer, Berlin (1995)
2. Bermejo, I.; García-Marco, I.; Salazar-González, J.J.: An algorithm for checking whether the toric ideal of an affine monomial curve is a complete intersection. *J. Symb. Comput.* **42**(10), 971–991 (2007)
3. Bermejo, I.; García-Marco, I.; Salazar-González, J.J.: cimonom.lib, A SINGULAR 3.0.3 library for determining whether the toric ideal of an affine monomial curve is a complete intersection (2007)
4. Bermejo, I.; Gimenez, P.; Reyes, E.; Villarreal, R.H.: Complete intersections in affine monomial curves. *Bol. Soc. Mat. Mex.* **11**(3), 191–203 (2005)
5. Delorme, C.: Sous-monoïdes d'intersection complète de  $\mathbb{N}$ . *Ann. Sci. École Norm. Sup.* **9**, 145–154 (1976)
6. Duursma, I.M.; Rentería, C.; Tapia-Recillas, H.: Reed–Muller codes on complete intersections. *Appl. Algebra Eng. Commun. Comput.* **11**(6), 455–462 (2001)
7. Eisenbud, D.; Sturmfels, B.: Binomial ideals. *Duke Math. J.* **84**, 1–45 (1996)
8. Eliahou, S.: Idéaux de définition des courbes monomiales. In: Greco, S.; Strano, R. (eds.) Complete Intersections. Lecture Notes in Mathematics, vol. 1092, pp. 229–240. Springer, Heidelberg (1984)
9. Eliahou, S.; Villarreal, R.H.: On systems of binomials in the ideal of a toric variety. *Proc. Am. Math. Soc.* **130**, 345–351 (2002)
10. Gold, L.; Little, J.; Schenck, H.: Cayley–Bacharach and evaluation codes on complete intersections. *J. Pure Appl. Algebra* **196**(1), 91–99 (2005)
11. Greuel, G.M.; Pfister, G.; Schoenemann, H.: *Singular*, A computer algebra system for polynomial computations. <http://www.singular.uni-kl.de> (2007)
12. Hansen, J.: Linkage and codes on complete intersections. *Appl. Algebra Eng. Commun. Comput.* **14**(3), 175–185 (2003)
13. Harris, J.: Algebraic Geometry. A First Course. Graduate Texts in Mathematics, vol. 133. Springer, New York (1992)
14. Herzog, J.: Generators and relations of abelian semigroups and semigroup rings. *Manuscr. Math.* **3**, 175–193 (1970)
15. López, H.H.; Rentería, C.; Villarreal, R.H.: Affine cartesian codes. *Des. Codes Cryptogr.* doi:10.1007/s10623-012-9714-2
16. López, H.H.; Sarmiento, E.; Vaz Pinto, M.; Villarreal, R.H.: Parameterized affine codes. *Stud. Sci. Math. Hung.* **49**(3), 406–418 (2012)
17. López, H.H.; Villarreal, R.H.: Complete intersections in binomial and lattice ideals. Preprint (2012). arXiv:1205.0772
18. Ramírez Alfonsín, J.L.: The Diophantine Frobenius Problem. Oxford Lecture Series in Mathematics and Its Applications, vol. 30. Oxford University Press, Oxford (2005)
19. Rentería, C.; Simis, A.; Villarreal, R.H.: Algebraic methods for parameterized codes and invariants of vanishing ideals over finite fields. *Finite Fields Appl.* **17**(1), 81–104 (2011)
20. Sarmiento, E.; Vaz Pinto, M.; Villarreal, R.H.: The minimum distance of parameterized codes on projective tori. *Appl. Algebra Eng. Commun. Comput.* **22**(4), 249–264 (2011)
21. Sarmiento, E.; Vaz Pinto, M.; Villarreal, R.H.: On the vanishing ideal of an algebraic toric set and its parameterized linear codes. *J. Algebra Appl.* **11**(4), 1250072 (2012)
22. Stanley, R.: Hilbert functions of graded algebras. *Adv. Math.* **28**, 57–83 (1978)
23. Thoma, A.: On the set-theoretic complete intersection problem for monomial curves in  $\mathbb{A}^n$  and  $\mathbb{P}^n$ . *J. Pure Appl. Algebra* **104**, 333–344 (1995)
24. Villarreal, R.H.: Monomial Algebras. Monographs and Textbooks in Pure and Applied Mathematics, vol. 238. Marcel Dekker, New York (2001)

