



# Patient Preferences in the Medical Product Lifecycle

Jennifer A. Whitty<sup>1,2</sup> · Esther W. de Bekker-Grob<sup>3,4</sup> · Nigel S. Cook<sup>5</sup> · Fern Terris-Prestholt<sup>6</sup> · Michael Drummond<sup>7</sup> · Rocco Falchetto<sup>8</sup> · Hans L. Hillege<sup>9</sup>

Published online: 29 November 2019  
© Springer Nature Switzerland AG 2019

There is an emerging consensus that evidence on patients' perspectives provides crucial guidance for decision making throughout the medical product lifecycle (MPLC). Considering the patient voice has become increasingly important for developers of new medical products and for authorities that assess, regulate and decide which products are effective, safe, well tolerated and cost effective [1, 2]. However, as key stakeholders, we lack a shared understanding about how to consider patient preferences in the MPLC [3].

This commentary captures opinions of thought leaders presenting at the symposium on “Patient preferences in the medical product lifecycle” in Basel, Switzerland in July 2019. The symposium was jointly sponsored by the International Academy of Health Preference Research (<http://www.iahpr.org>) and the research project “Patient Preferences in Benefit and Risk Assessments during the Treatment Life Cycle” (PREFER; <http://www.imi-prefer.eu>). This commentary is one of a series of papers reporting on health preference research symposiums held alongside International Academy of Health Preference Research meetings [4–6].

---

Jennifer A. Whitty and Esther W. de Bekker-Grob are joint first authors for this commentary.

---

This article is part of the topical collection “From the International Academy of Health Preference Research”. Please refer the previous topical collection article (<https://doi.org/10.1007/s40271-018-0346-0>).

---

✉ Jennifer A. Whitty  
Jennifer.Whitty@uea.ac.uk

<sup>1</sup> Norwich Medical School, University of East Anglia, Norwich, UK

<sup>2</sup> National Institute for Health Research Applied Research Collaboration East of England, Cambridge, UK

<sup>3</sup> Erasmus School of Health Policy & Management, Erasmus University Rotterdam, Rotterdam, The Netherlands

<sup>4</sup> Erasmus Choice Modelling Centre, Erasmus University Rotterdam, Rotterdam, The Netherlands

At the symposium, a varied panel shared their experience as researchers, policy makers, industry leaders and patients on the use of patient preferences in the MPLC and spoke to the following six questions.

## 1 Does Patient Preference Information in Reimbursement Decision-Making Lead to Higher Quality Decisions and Increased Public Acceptance of Decisions Regarding Allocation of Healthcare Resources?

Although it is widely acknowledged that patient preferences contribute important evidence for reimbursement decisions, there are challenges in assessing its impact on decision-making. First, there have not been enough published examples where preference studies have been submitted as part of the reimbursement process on which to judge their impact. Second, even where examples are available, the complexity of decision-making means it is difficult to single out the impact of preference information from the impact of other decision ‘inputs’ such as clinical and economic evidence and societal factors. Third, preference studies are mostly undertaken late in the product development process. When this occurs, there may be a tendency to retro-fit the attributes to those which the product in question addresses, or which have been measured in the trials. There is the risk that such attributes are

<sup>5</sup> Decision Support, Global Patient Access, Novartis Pharma AG, Basel, Switzerland

<sup>6</sup> Department of Global Health and Development, London School of Hygiene and Tropical Medicine, London, UK

<sup>7</sup> Centre for Health Economics, University of York, York, UK

<sup>8</sup> International Porphyria Patient Network, Zurich, Switzerland

<sup>9</sup> Department of Cardiology, Department of Epidemiology, University Medical Center Groningen, Groningen, The Netherlands

not addressing the issues that matter most to the patient. The health technology assessment (HTA) body may well recognise this discrepancy [7]. Likewise, the patient voice might be dampened if what matters to them is not addressed by the new product, with or without a preference study to support it. Fourth, where a preference study is sponsored by a manufacturer, there may be a perceived conflict of interest resulting in a high level of scrutiny, but also a potentially high and possibly inappropriate level of scepticism in their findings.

Undertaking preference studies early in the development process is an opportunity to strengthen their impact on both the development process and reimbursement. There are examples where preference evidence has impacted decision-making and therefore potentially improved the ‘quality’ of decisions [8, 9]. As such, the panel were generally of the view that preference information will improve the quality of decisions, although less decided on whether the consideration of patient preference information increased public acceptance of decisions regarding the allocation of health-care resources. At least it has the potential to increase the transparency of the decision-making process, and there was general agreement that broader stakeholder involvement in the decision-making process was desirable.

## 2 What Should Happen First to Achieve Successful Integration of Patient Preference Information into the Medical Product Lifecycle?

To date, successful integration of patient preference information into the MPLC has been slow, unsystematic or very limited [10, 11]. To promote successful integration, it is necessary first to address the concerns of users of preference information. Publication of examples where preference information has been beneficial and creating projects that involve collaboration across stakeholders would also promote successful integration. For some conditions, there are multiple companies with similar interests. Therefore, collaboration between competing companies when undertaking studies would allow companies to pool resources and expertise, and reduce any perception of biased interests.

Who should fund preference studies presents a dilemma. Regulators look to industry to generate good-quality preference studies to support their decision making [12]. However, pharmaceutical and medical technology companies may be averse to spending research resources to undertake studies that are not seen to influence regulators’ and/or HTA body decisions. The uptake of preference studies and examples of impact might be expected to break this deadlock, even if progress is gradual initially. Finally, industry standards, research standards, and quality guidelines for undertaking preference studies are needed to support the use of patient

preferences in regulatory and HTA decisions. Textbooks in the field (such as that currently being developed by the International Academy of Health Preference Research), consensus guidance, and a more consistent approach to developing protocols before undertaking a preference study should all assist the perceived rigour and therefore integration of preference studies. Rigour can be improved by adhering to an a priori study protocol; for example, via the Health Preference Study and Technology Registry database [13], and reporting according to appropriate peer-review standards such as those recommended by the International Society for Pharmacoeconomics and Outcomes Research review task force [14] and through publication in a peer-reviewed journal.

## 3 Which Preference Methods are Most Promising or Acceptable to be Used in Regulatory Assessments or Reimbursement Decision Making?

As with any research area, the method needs to suit the research question. Currently, discrete choice experiments are perceived as the gold standard for evaluating preferences and are widely used. However, they are not suitable in all scenarios: when there are many attributes to consider (which may be the case in a new disease area in particular) or when patients might not be able to process a large amount of information being given to them. Furthermore, if the question relates to a rare disease, preference studies via discrete choice experiments may not reach a sufficient sample size to allow for valid statistical analyses and reliable preference data. In such circumstances, alternative approaches might be more suitable.

The importance of using appropriate qualitative methods in the early stages of undertaking a preference study is now well established [9, 15, 16]. The development of consensus guidance to aid the selection of appropriate methods for different research questions was encouraged by the panel.

## 4 At What Point in the Medical Product Lifecycle is Patient Preference Information Most Useful, and for What Purpose?

Whilst funding can be challenging to obtain for preference studies, it may be easiest to obtain funding to elicit patient preference information during product development to support marketing, or for reimbursement decisions. Typically, funding availability is greatest in the 12–24 months approaching product approval and launch, and is likely to be quite restricted early in the MPLC. However, consideration of patient preference evidence early and if feasible continuously in the product lifecycle can provide strong benefits for stakeholders. The net present value of engaging patients in

the MPLC is high in the early phases of development [17]. The challenge is to secure the funding at this early stage of development. Once again, good examples of the impact of such early preference initiatives are needed in the literature to foster this shift of attention to earlier investment. Moreover, patient preference data may play an important role in designing trials, to help understand what risks and benefits should be measured and what effect size is meaningful for patients. It has also been shown that trials can achieve faster recruitment and completion when patient input had been solicited during the design [18].

Resources to undertake preference studies would be best directed to those products where preferences are most likely to influence the regulatory or reimbursement decisions. We need to understand the conditions and factors that make these decisions most sensitive to preferences [12], and should promote further research to help understand this. It is important for companies to have an early discussion with regulators and HTA bodies to both identify products for which patient preference information might be of particular benefit to decision making and to identify relevant effects for which risk–benefit trade-offs might be elicited, and to determine what constitutes patient value. Of note, the nuance between regulators and HTA agencies and the purpose for which they might use preference information also need to be considered.

## 5 Is There a Role for the Use of Preference Information Elicited from People Other than Patients to Inform the Medical Product Lifecycle?

The decision problem and context are important considerations when deciding whose preferences are relevant. Where the general public are the payers for care, their preferences might be relevant for informing priorities about health and social care provision, and resource allocation at a population level (and this is typically achieved, for example, through the use of the EQ-5D and similar preference-based outcome measures).

Once resources are allocated to a specific area, patient preferences on how to develop or prioritise treatments or products within that area become most relevant. Where patients are able to provide preferences, every effort should be made to obtain preferences from patients themselves, or in some cases, their caregivers, and the use of a proxy might not be acceptable. It is important to also remember that the relevant individuals to seek preferences from may differ in different cultural or geographical settings. This also raises the question of who is the ‘patient’? A patient preference study needs to clearly define the characteristics of the ‘patient’ of interest, using inclusion and exclusion criteria. Within a disease, people may have different characteristics, some of whom may be candidates for the drug based on the marketing

authorisation or reimbursement criteria, others not. Further, participants within a clinical trial may be a very narrow subset of patients affected by the disease. For products targeting prevention, it may be service users or the general public rather than ‘patients’ that are most relevant. There is a continuum of ‘patients’, at one extreme is the general public, who are all potential service users, and at the other extreme is an individual patient with a specific condition making an individual decision about their treatment [19]. The patient population of interest will lie somewhere on this continuum.

## 6 How Might Patient Preference Information Best be Elicited from Patients and When? For Example, is it Acceptable (or even Preferable) to Elicit Information Alongside Clinical Trials?

Eliciting preferences alongside a trial may be opportunistic. However, there are also arguments against eliciting preferences alongside a trial. Participants in a trial might be more willing to accept risk than other patients with the same condition. Whilst a trial may be helpful scientifically to accommodate randomisation and investigate comparative effects, it may not be the best setting for investigating preferences that would ideally be generalisable to the wider patient population. Preferences elicited alongside a clinical trial are also unlikely to provide information on the uptake by the broader patient population, which is useful particularly in the public health setting. If preferences are to be elicited alongside a trial, the timing of elicitation is important. Patients may be naïve to the product at the start of the trial, but some will be more familiar with the product by the end (although they may be blinded to allocation). Product preferences may also be associated with trial drop-out, introducing a selection bias if preferences are elicited from participants later on in a trial.

## 7 Concluding Remarks

To conclude, the symposium was unanimous about the important role of patient preference information and that collecting preference data from an early stage of product development has clear benefits. There is a role for qualitative approaches to gather evidence on patient experiences and to inform the development of preference instruments. Qualitative evidence complements the quantified weights and trade-offs that preference information provides. Finally, it is important to recognise patient heterogeneity, in terms not only of defining the characteristics of the patient group of interest, but also of considering when in the patient’s journey is the most relevant time point to elicit preferences.

**Acknowledgements** The authors thank all the symposium panellists (Nigel S. Cook, Michael Drummond, Fern Terris-Prestholt, Irina Cleemput, Hans L. Hillege, and Rocco Falchetto) and participants for joining the conversation and sharing their opinions and ideas. They also thank Irina Cleemput for providing input into an initial draft of this paper and Samare P.I. Huls for taking notes to assist with the commentary preparation during the symposium.

**Author Contributions** JW and EBG conceived the topic and framework for the symposium panel and paper, chaired the panel and prepared a joint draft of the paper. NSC, FTP, MD, RF and HLH presented at the panel and contributed their academic ideas to the paper. All authors reviewed and revised the draft for academic content and approved the final version for publication.

## Compliance with Ethical Standards

**Funding** Jennifer A. Whitty is supported by the National Institute for Health Research (NIHR) Applied Research Collaboration East of England (NIHR200177). Esther W. de Bekker-Grob is funded through a personal grant from The Netherlands Organisation for Scientific Research (NWO-Talent-Scheme-Veni-Grant No. 451-15-039) and supported by the EU/EFPIA Innovative Medicines Initiative [2] Joint Undertaking PREFER grant no. 115966. Nigel S. Cook is an employee of Novartis Pharma AG and the work presented was funded by Novartis. RF is funded through out-of-pocket private donations to the IPPN. The views expressed are those of the authors and not necessarily those of the National Health Service, the NIHR, or the Department of Health and Social Care.

**Conflict of Interest** Jennifer A. Whitty, Rocco Falchetto, Hans L. Hillege, Fern Terris-Prestholt and Michael Drummond are members of the International Academy of Health Preference Research (IAHPR). Esther W. de Bekker-Grob is a member of IAHPR and IMI-PREFER. Nigel S. Cook is a member of IAHPR and IMI-PREFER and is an employee of Novartis Pharma AG. None of the authors have any conflicts of interest that are directly relevant to the content of this article.

## References

1. Hoos A, Anderson J, Boutin M, Dewulf L, Geissler J, Johnston G, et al. Partnering with patients in the development and lifecycle of medicines: a call for action. *Ther Innov Regul Sci*. 2015;49(6):929–39. <https://doi.org/10.1177/2168479015580384>.
2. Smith MY, Hammad TA, Metcalf M, Levitan B, Noel R, Wolka AM, et al. Patient engagement at a tipping point: the need for cultural change across patient, sponsor, and regulator stakeholders: insights from the DIA Conference, “Patient Engagement in Benefit Risk Assessment Throughout the Life Cycle of Medical Products”. *Ther Innov Regul Sci*. 2016;50(5):546–53. <https://doi.org/10.1177/2168479016662902>.
3. de Bekker-Grob EW, Berlin C, Levitan B, Raza K, Christoforidi K, Cleemput I, et al. Giving patients’ preferences a voice in medical treatment life cycle: the PREFER Public-Private Project. *Patient*. 2017;10(3):263–6. <https://doi.org/10.1007/s40271-017-0222-3>.
4. Gonzalez JM, Johnson FR, Levitan B, Noel R, Peay H. Symposium title: preference evidence for regulatory decisions. *Patient*. 2018;11(5):467–73. <https://doi.org/10.1007/s40271-018-0311-y>.
5. Norman R, Craig BM, Hansen P, Jonker MF, Rose J, Street DJ, et al. Issues in the design of discrete choice experiments. *Patient*. 2019;12(3):281–5. <https://doi.org/10.1007/s40271-018-0346-0>.
6. Ostermann J, Brown DS, van Til JA, Bansback N, Légaré F, Marshall DA, et al. Support tools for preference-sensitive decisions in healthcare: where are we? Where do we go? How do we get there? *Patient*. 2019;12(5):439–43. <https://doi.org/10.1007/s40271-019-00372-z>.
7. Berglas S, Jutai L, MacKean G, Weeks L. Patients’ perspectives can be integrated in health technology assessments: an exploratory analysis of CADTH Common Drug Review. *Res Invol Engagem*. 2016;2(1):21. <https://doi.org/10.1186/s40900-016-0036-9>.
8. Cleemput I, Devriese S, Kohn L, Westhovens R. A multi-criteria decision approach for ranking unmet needs in healthcare. *Health Policy*. 2018;122(8):878–84. <https://doi.org/10.1016/j.healthpol.2018.06.010>.
9. Cook NS, Cave J, Holtorf A-P. Patient preference studies during early drug development: aligning stakeholders to ensure development plans meet patient needs. *Front Med (Lausanne)*. 2019;6:82. <https://doi.org/10.3389/fmed.2019.00082>.
10. van Overbeeke E, Whichello C, Janssens R, Veldwijk J, Cleemput I, Simoens S, et al. Factors and situations influencing the value of patient preference studies along the medical product lifecycle: a literature review. *Drug Discov Today*. 2019;24(1):57–68. <https://doi.org/10.1016/j.drudis.2018.09.015>.
11. Janssens R, Russo S, van Overbeeke E, Whichello C, Harding S, Kübler J, et al. Patient preferences in the medical product life cycle: what do stakeholders think? Semi-structured qualitative interviews in Europe and the USA. *Patient*. 2019;12(5):513–26. <https://doi.org/10.1007/s40271-019-00367-w>.
12. Benz HL, Lee T-H, Tsai J-H, Bridges JFP, Eggers S, Moncur M, et al. Advancing the use of patient preference information as scientific evidence in medical product evaluation: a summary report of the patient preference workshop. *Patient*. 2019. <https://doi.org/10.1007/s40271-019-00396-5>.
13. Health Preference Study and Technology Register (HPSTR). International Academy of Health Preference Research (IAHPR). <http://www.hpstr.org>. Accessed 23 Nov 2019.
14. Bridges JF, Hauber AB, Marshall D, Lloyd A, Prosser LA, Regier DA, et al. Conjoint analysis applications in health—a checklist: a report of the ISPOR Good Research Practices for Conjoint Analysis Task Force. *Value Health*. 2011;14(4):403–13. <https://doi.org/10.1016/j.jval.2010.11.013>.
15. Vass C, Rigby D, Payne K. The role of qualitative research methods in discrete choice experiments. *Med Decis Making*. 2017;37(3):298–313. <https://doi.org/10.1177/0272989X16683934>.
16. Coast J, Al-Janabi H, Sutton EJ, Horrocks SA, Vosper AJ, Swancutt DR, et al. Using qualitative methods for attribute development for discrete choice experiments: issues and recommendations. *Health Econ*. 2012;21(6):730–41. <https://doi.org/10.1002/hec.1739>.
17. Levitan B, Getz K, Eisenstein E, Goldberg M, Harker M, Hesterlee S, et al. Assessing the financial value of patient engagement: a quantitative approach from CTTI’s Patient Groups and Clinical Trials Project. *Ther Innov Regul Sci*. 2018;52:220–9. <https://doi.org/10.1177/2168479017716715>.
18. Economist Intelligence Unit Limited. The innovation imperative: the future of drug development. Part I: research methods and findings. A report by the Economist Intelligence Unit (commissioned by Paraxel). London: The Economist Intelligence Unit; 2018. [https://druginnovation.eiu.com/wp-content/uploads/2018/10/Paraxel-innovations-in-drug-development\\_V10.pdf](https://druginnovation.eiu.com/wp-content/uploads/2018/10/Paraxel-innovations-in-drug-development_V10.pdf)
19. Whitty JA, Fraenkel L, Saigal CS, Groothuis-Oudshoorn CGM, Regier DA, Marshall DA. Assessment of individual patient preferences to inform clinical practice. *Patient*. 2017;10:519–21. <https://doi.org/10.1007/s40271-017-0254-8>.