

# Psychometric properties of the Italian Food Cravings Questionnaire-Trait-reduced (FCQ-T-r)

Marco Innamorati · Claudio Imperatori · Adrian Meule ·  
Dorian A. Lamis · Anna Contardi · Michela Balsamo ·  
Antonino Tamburello · Stella Tamburello · Mariantonietta Fabbriatore

Received: 27 May 2014 / Accepted: 13 July 2014 / Published online: 29 July 2014  
© Springer International Publishing Switzerland 2014

## Abstract

**Purpose** The aim of the present study was to investigate dimensionality and psychometric properties of the Italian Food Cravings Questionnaire-Trait-reduced (FCQ-T-r) in a sample of obese and overweight patients seeking weight loss treatment.

**Methods** Participants were 504 (416 women and 88 men) overweight and obese patients ( $BMI \geq 25 \text{ kg/m}^2$ ), and 289 (215 women and 74 men) Italian adults not currently seeking weight loss treatment. All participants were administered the Food Cravings Questionnaire-Trait (FCQ-T) and the Binge Eating Scale.

**Results** The fifteen items included in the FCQ-T-r explained 93 % of the variance of the 39-item FCQ-T total score ( $R^2 = 0.93$ ). A principal axis factoring analysis indicated a one-factor solution, explaining 55.6 % of the variance of the data. The FCQ-T-r had high internal consistency and was also able to differentiate between individuals with various severities of binge eating behavior.

**Conclusions** The FCQ-T-r may be considered a useful instrument for measuring trait food craving, when time constraints impede the use of the 39-item FCQ-T.

**Keywords** Food Cravings Questionnaire · Obesity · Overweight · Validity · Exploratory factor analysis

M. Innamorati · C. Imperatori (✉) · A. Contardi ·  
A. Tamburello · S. Tamburello · M. Fabbriatore  
Università Europea di Roma, via degli Aldobrandeschi 190,  
00163 Rome, Italy  
e-mail: imperatori.c@libero.it

M. Innamorati · C. Imperatori · A. Contardi · A. Tamburello ·  
S. Tamburello · M. Fabbriatore  
Istituto Skinner, via Nazionale 163, 00184 Rome, Italy

A. Meule  
Department of Psychology I, University of Würzburg,  
Würzburg, Germany

A. Meule  
Hospital for Child and Adolescent Psychiatry, LWL University  
Hospital of the Ruhr University Bochum, Hamm, Germany

D. A. Lamis  
Department of Psychiatry and Behavioral Sciences, Emory  
University School of Medicine, Atlanta, GA, USA

M. Balsamo  
DISPUTer, “G. d’Annunzio” University of Chieti-Pescara,  
Chieti, Italy

## Introduction

Food craving (FC) can be defined as an intense desire to consume a specific food that is difficult to resist [1, 2]. Several studies have reported that FC is an important feature of eating and weight disorders, especially Binge Eating Disorder (BED) and obesity [3–7]. Indeed, FC appears to be a proximal factor in precipitating episodes of binge eating [8–10] and it is predictive of high caloric food intake [11]. More intense or frequent FC is also associated with higher body mass index (BMI) [2, 12–15], lower dieting success and more dropouts from weight loss treatment programs [14, 16, 17], and a higher number of lifetime weight loss attempts [18].

The Food Cravings Questionnaire-Trait (FCQ-T; [9, 19]) is the most extensively validated and internationally used FC measure [15, 20]. The FCQ-T is a multidimensional questionnaire which assesses several aspects of FC such as, anticipation of positive/negative reinforcement from eating, thoughts about and

preoccupation with food, or lack of control over eating, among others.

Although the multidimensionality of the FCQ-T may help clinicians and investigators assess all the major components of FC, it is a relatively long self-report measure (39 items). Thus, Meule et al. [15] developed and validated the Food Cravings Questionnaire-Trait-reduced (FCQ-T-r), a short form of the FCQ-T consisting of 15 items, which showed the highest item-total correlations in a study in which the full version was used [14]. Psychometric properties of the FCQ-T-r were also investigated, with satisfactory results, among Cuban adults [20]. However, to date there are no studies investigating the psychometric properties of an Italian version of the FCQ-T-r. Thus, the aim of the present study was to investigate dimensionality and psychometric properties of the Italian FCQ-T-r in a sample of obese and overweight patients seeking weight loss treatment. Furthermore, to assess whether the FCQ-T-r item and total scores could discriminate obese and overweight patients from the general population, a control group was enrolled. We hypothesized that overweight patients seeking weight loss treatment would report higher food craving severity on the FCQ-T-r.

## Methods

### Participants

All patients were consecutively admitted to three private medical centers in Rome (Italy) that specialized in the treatment of obesity and shared the same therapeutic approach [21], and who met inclusion criteria were enrolled in the study. Approximately 89.6 % of the patients agreed to participate in the study, and 95.5 % of those who agreed to participate in the study completed the assessment. The final sample was composed of 504 (416 women and 88 men) overweight and obese patients. Inclusion criteria were a minimum age of 18 years and a minimum BMI of 25 kg/m<sup>2</sup>. Exclusion criteria were the presence of major disorders of the central nervous system (e.g., epilepsy, dementia, or Parkinson's disease), any psychotic symptoms (hallucinations, delusions, or negative symptoms) or major mood disorders (e.g., bipolar disorder, major depressive disorder), and the presence of any condition affecting the ability to complete the assessment, including the denial of informed consent. Participant characteristics are displayed in Table 1.

Four hundred and ninety-seven patients were part of the sample of a recent study investigating the psychometric properties of the FCQ-T [22], and more detailed information about the sample and the recruitment process can be found in Innamorati et al. [22].

**Table 1** Participant characteristics and group differences between patients and controls

Variable	Patients ( <i>n</i> = 504)	Controls ( <i>n</i> = 289)	Test statistics
Women, <i>n</i> (%)	416 (82.5)	215 (74.4)	<i>p</i> = 0.01*
Age (years), M (SD)	44.00 (12.14)	32.67 (13.49)	<i>t</i> <sub>(791)</sub> = 11.80, <i>p</i> < 0.001
Employed, <i>n</i> (%)	390 (77.4)	207 (71.6)	<i>p</i> = 0.37*
Education (>13 years), <i>n</i> (%)	445 (88.3)	238 (82.4)	<i>p</i> = 0.64*
Married or in a stable relationship, <i>n</i> (%)	314 (62.3)	195 (67.5)	<i>p</i> = 0.27*
Body mass index (kg/m <sup>2</sup> ), M (SD)	30.53 (5.35)	22.83 (3.13)	<i>t</i> <sub>(791)</sub> = 25.55, <i>p</i> < 0.001
Binge eating, <i>n</i> (%)			$\chi^2_{(2)}$ = 48.84, <i>p</i> < 0.001
No binge eating	326 (64.6)	259 (89.5)	
Probable binge eating	127 (25.2)	27 (9.5)	
Clinical-level binge eating	51 (10.3)	3 (0.9)	

\* One-way Fisher exact tests

Controls included 289 (215 women and 74 men) adults from the Italian general population (see Table 1 for participant characteristics). The same inclusion/exclusion criteria were used for controls with the exception of BMI limits. The controls were recruited through advertisements posted in established community groups (e.g., universities and community organizations operating in Rome and in Central Italy).

Groups differed in sex distribution, age, and BMI (Table 1). Patients and controls voluntarily participated in the study, did not receive payment, provided written informed consent, and completed the protocol anonymously. The research was approved by the Università Europea's ethics review board.

### Measures

All participants completed the 39-item Italian version of the FCQ-T [22] and the Binge Eating Scale (BES) [23]. Clinical and socio-demographic information was obtained from medical records for patients, while controls completed a socio-demographic form during the assessment.

The FCQ-T is a 39-item questionnaire with items rated on a 6-point Likert scale (from 1 = never to 6 = always) for the assessment of the frequency of FC experiences. Several studies [9, 19, 24, 25] indicated a nine-factor structure: (1) anticipation of positive reinforcement from

eating (Ant+); (2) anticipation of relief from negative states and feelings from eating (Ant-); (3) intentions and plans to consume food (Intent); (4) cues that may trigger food cravings (Cues); (5) thoughts or preoccupation with food (Thoughts); (6) craving as hunger (Hunger); (7) lack of control over eating (Control); (8) emotions that may be experienced before or during food cravings or eating (Emotions); and (9) guilt from cravings and/or for giving into them (Guilt). Previous studies conducted both in samples of healthy individuals and patients with eating disorders reported that the scale has high internal consistency (Cronbach's alphas between 0.92 and 0.97); however, alphas for some subscales were not always satisfactory ranging from 0.57 to 0.84, most likely due to the low number of items included on some factors [3, 9, 13, 14, 19, 24, 25]. Test-retest reliability at 3 weeks was high for the English and the Spanish Versions ( $r = 0.88$ ; [9, 19]), and for the German version ( $r = 0.84$ ; [14]). Cronbach's alpha in the present sample was 0.96.

The FCQ-T-r is composed of 15 items from the FCQ-T. The items belonged to five dimensions of the original FCQ-T: Control (items 2, 3, 25, 26, 29), Thoughts (items 6, 8, 27, 32, 33), Intent (items 5, 18), Emotions (items 20, 34), and Cues (item 36). Meule et al. [15] reported unidimensionality and high internal consistency (Cronbach's alpha = 0.94). Validity was demonstrated by, for example, positive correlations between scores on the FCQ-T-r, BMI and cue-elicited FC [15].

The BES is a 16-item questionnaire assessing binge eating severity as well as the feelings and thoughts associated with such behavior. Thus, it assesses both behavioral and cognitive/emotional manifestations of binge eating. Each item has three or four weighted statements, and the respondent is directed to choose one. In the validation sample, the BES successfully discriminated among persons judged by trained interviewers to have or not have binge eating problems [26]. The scores range from 0 to 46. Individuals with scores <18 were considered not to be engaging in significant binge eating; individuals with scores in the 18–26 range likely experience binge eating; and individuals with scores of >27 engage in clinically significant binge eating behaviors [26]. Using these categories resulted in a 98 % concordance rate with diagnosis made in a semi-structured interview [27] and, thus, we also used these cut-offs in the present study. Furthermore, the BES has been shown to have good internal consistency (Cronbach's alpha of 0.85) [26]. Cronbach's alpha in the present sample was 0.88.

#### Statistical analyses

To investigate dimensionality of the FCQ-T-r in obese and overweight patients we used principal axis factoring

analysis. The number of factors to retain was determined by means of Kaiser criterion (only factors with eigenvalues >1 were retained), and parallel analysis [28]. To investigate psychometric properties of the FCQ-T-r at the item level, we report corrected item-total correlations and descriptive statistics (means and standard deviations), and analyzed differences between patients and controls by means of independent  $t$  tests. At the scale level, we report Cronbach's alpha and mean inter-item correlation as measures of reliability, and descriptive statistics (mean and standard deviation). We also calculated the proportion of variance ( $R^2$ ) of the full FCQ-T that is explained by the FCQ-T-r by means of a linear regression analysis with the FCQ-T as the criterion and the FCQ-T-r as the independent variable. Scores on the FCQ-T-r were compared between patients and controls, and scores were also compared with those reported in previous studies [15, 20]. Relationships between FCQ-T-r scores and socio-demographic characteristics were tested with independent  $t$  test (sex) and Pearson correlation coefficients (age, BMI, and BES). Cohen's  $d$  coefficients are reported as measures of effect size. To assess the performance of the FCQ-T-r in categorizing individuals based on the severity of their current binge eating, we performed the Receiver Operating Characteristic (ROC) test procedures. A ROC curve is a two-dimensional depiction of test performance [29, 30]. The main outcome variable is the area under the ROC curve (AUC), which is interpreted as the probability that a randomly sampled respondent will be correctly assigned to the appropriate group. The AUC directly represents the overall accuracy of the instrument in screening for binge eating. ROC curve analyses were performed both on the whole sample, which included patients and controls, and on the sample of patients. In line with the validation study of the Italian version of the FCQ-T [22], we tested the ability of FCQ-T-r to discriminate respondents with clinical-level binge eating (BES scores  $\geq 27$ ) from those with probable binge eating (BES scores between 18 and 26). We analyzed the accuracy of the FCQ-T-r in differentiating only respondents with clinical-level binge eating from those with probable binge eating because this may be the most challenging situation in clinical settings, due to the fact that the use of the BES only may produce too many false positive and negative. All analyses were performed with the Statistical Package for the Social Sciences (SPSS) 19.0 for Windows (IBM Corp., Armonk, NY).

#### Results

In Table 2, statistics for the FCQ-T-r in obese and overweight patients are reported. Corrected item-total correlations for the FCQ-T-r items ranged between 0.60 and 0.79

**Table 2** Psychometric properties of the FCQ-T-r in obese and overweight patients and differences in FCQ-T-r item and total scores compared with controls

Item #	Obese and overweight patients ( <i>n</i> = 504)			Controls ( <i>n</i> = 289)	Cohen's <i>d</i> <sup>a</sup>
	Factor loadings	Corrected item-total correlation	Item M (SD)	Item M (SD)	
2	0.63	0.61	3.52 (1.51)	2.64 (1.39)	0.61
3	0.70	0.69	3.63 (1.46)	2.69 (1.24)	0.69
5	0.68	0.66	2.72 (1.47)	2.12 (1.22)	0.44
6	0.77	0.74	2.52 (1.46)	1.79 (1.13)	0.55
8	0.76	0.73	2.25 (1.35)	1.52 (0.90)	0.64
18	0.70	0.67	2.49 (1.44)	2.07 (1.15)	0.32
20	0.60	0.59	3.55 (1.47)	2.79 (1.31)	0.55
25	0.71	0.70	3.18 (1.55)	2.12 (1.20)	0.76
26	0.84	0.82	2.91 (1.55)	1.97 (1.13)	0.69
27	0.83	0.80	2.24 (1.44)	1.51 (0.93)	0.60
29	0.82	0.79	2.38 (1.55)	1.55 (0.89)	0.66
32	0.76	0.73	2.32 (1.36)	1.88 (1.06)	0.35
33	0.72	0.70	1.78 (1.16)	1.37 (0.74)	0.42
34	0.66	0.64	2.66 (1.42)	2.09 (1.63)	0.37
36	0.64	0.62	3.54 (1.45)	3.00 (1.33)	0.39
		FCQ-T-r sum score	41.69 (16.09)	31.00 (11.26)	0.77

<sup>a</sup> All tests on group differences in FCQ-T-r item and total scores were significant at  $p < 0.001$


(mean = 0.70). The FCQ-T-r score explained 93 % of the variance of the 39-item FCQ-T total score ( $R^2 = 0.93$ ), and was correlated strongly with the sum score of the 24 items excluded from the FCQ-T-r ( $r = 0.89$ ;  $p < 0.01$ ). In the principal axis factoring, two factors had eigenvalues  $>1$  (Factor 1 = 8.35, 55.6 % of the variance explained; factor 2 = 1.13, 7.5 % of the variance explained), but a parallel analysis (eigenvalues of 1.30 and 1.23 for simulated data at the parallel analysis) suggested to retain only 1 factor (factor loadings for the FCQ-T-r are reported in Table 2). Internal consistency for the 15-item FCQ-T-r was  $\alpha = 0.94$  with a mean inter-item correlation of 0.52. All items scores differed between patients and controls with weak to moderate effects (Table 2). Also, the FCQ-T-r sum score differed between obese patients and controls (Table 2). Mean scores for our sample of obese patients were higher than those reported by Meule et al. [15] in German students ( $38.47 \pm 13.11$ ;  $t_{825} = 3.01$ ;  $p < 0.01$ ; Cohen's  $d = 0.22$ ), and lower than those reported for Cuban adults ( $45.02 \pm 19.74$ ;  $t_{1,743} = 3.36$ ;  $p < 0.001$ ; Cohen's  $d = -0.18$ ) [20]. Mean scores for controls were lower than those reported by German students ( $t_{610} = 7.52$ ;  $p < 0.001$ ; Cohen's  $d = -0.61$ ) and Cuban adults ( $t_{1,528} = 11.64$ ;  $p < 0.001$ ; Cohen's  $d = -0.87$ ). FCQ-T-r scores were weakly but significantly associated with BMI ( $r = 0.15$ ;  $p < 0.01$ ) and strongly correlated with BES scores ( $r = 0.77$ ;  $p < 0.01$ ). They were not correlated with age ( $r = -0.04$ ;  $p = 0.34$ ), and were significantly associated with sex ( $t_{141.90} = 4.65$ ;  $p < 0.001$ ; Cohen's  $d = 0.52$ ), with female patients reporting higher scores than males ( $43.06 \pm 16.21$  vs.  $35.24 \pm 13.89$ ).

Approximately 25 % of obese patients reported probable binge eating as indicated by BES scores (vs. 9.5 % of the controls), and 10.3 % reported clinical-level binge eating (vs. 0.9 % of the controls) (Table 1). A ROC curve procedure indicated that FCQ-T-r scores (area under the ROC curve = 0.78,<sup>1</sup> 95 % CI = 0.71/0.85; SE = 0.03;  $p < 0.001$ ) may categorize individuals with different severity of binge eating well (Fig. 1). A score of 57.5 or higher on the FCQ-T-r categorized individuals with a sensitivity of 0.69 (69 % of all the patients with clinical-level binge eating were correctly identified) and a specificity of 0.69 (31 % of all individuals with only probable binge eating were incorrectly identified to have clinical-level binge eating).

## Discussion

The aim of the present study was to investigate dimensionality and psychometric properties of the Italian FCQ-T-r in a sample of obese and overweight patients seeking weight loss treatment. Results showed that FCQ-T-r scores explained more than 90 % of the variance of the original 39-item version of the FCQ-T, that the FCQ-T-r is unidimensional and has high internal consistency, comparable to that of the longer version. Thus, results suggest that the

<sup>1</sup> Results were similar when only overweight and obese patients were considered in the analysis (area under the ROC curve = 0.75, 95 % CI = 0.69/0.83; SE = 0.04;  $p < 0.001$ ).



**Fig. 1** ROC curve graph for the ability of the FCQ-T-r to discriminate patients with clinical-level binge eating (BES scores  $\geq 27$ ) from those with probable binge eating (BES scores between 18 and 26)

short form may be used as a substitute of the long version without sacrificing much information.

Obese participants had higher FCQ-T-r scores than controls. Obese patients also reported higher mean FCQ-T-r scores than German students [15]. Unexpectedly, they had lower scores compared with those reported by Cuban adults [20]. This difference could reflect some social and cultural factors. After a period of 5 years in which difficulties caused a decrease of food availability for the majority of the Cuban population [20, 31], obesity rates began to rise [32], indicating changes in food-consuming habits among the Cuban population. Likewise, previous studies have reported cultural differences in FC [33, 34]. However, some characteristics among our sample of obese patients may explain the results. In a previous study [22] in which we compared scores on the FCQ-T in obese/overweight patients who made up the majority of the current sample, their mean scores were intermediate among those reported in other studies [3, 4, 35, 36], indicating that FC severity only was moderate.

Inline with previous data [3, 5–7, 12–15, 22], in our sample, FC was strongly associated with binge eating and weakly but significantly associated with BMI. The small correlation observed between FCQ-T-r and BMI is not unexpected; previous studies [13, 14] reported small correlation between BMI and the FCQ-T ( $< 0.30$ ) and the exact correlation coefficient ( $r = 0.15$ ) found in the present study was found for the German FCQ-T-r [15]. Our results also suggest that FCQ-T-r was able to discriminate between people with clinical-level binge eating (BES scores  $\geq 27$ ) and those with probable binge eating (BES scores between 18 and 26). The ability of the FCQ-T-r in discriminating people with different levels of binge eating

was comparable to that of the 39-item FCQ-T. A randomly chosen individual with clinical-level binge eating had a 78 % of probability of having higher scores on the FCQ-T-r than a randomly chosen individual with probable binge eating, versus 76 % of probability of having higher scores on the 39-item FCQ-T [22].

Finally, consistent with previous findings [15, 24, 37, 38], we observed gender differences in FC, with female patients reporting higher scores on the FCQ-T-r than males. It is possible that women are more weight and shape concerned and that they more often try to restrict their eating [39], exposing themselves to the risk of inducing in FC [40]. At the biological level, these differences could reflect differences observed in women's reward system [41, 42] which could be secondary to modifications associated with the menstruation cycle [43, 44].

The study's contributions should be considered in light of its limitations. First, our clinical sample mostly consisted of slightly obese female patients; thus, results may be different in samples of patients with severe obesity, and/or in male patients. Second, obese patients and controls differed in major socio-demographic characteristics, raising the possibility that other covariates may explain the differences between groups. Third, we only administered self-report measures, which may have been biased due to social desirability. Moreover, this study was a reanalysis of existing data, that means the FCQ-T and not the FCQ-T-r was administered. Thus, future studies on the validation of the FCQ-T-r need to be done in which only the FCQ-T-r items are used.

Nevertheless, our results suggest that the FCQ-T-r may be considered a useful instrument for assessing food cravings, when time constraints impede the use of the 39-item version of the FCQ-T. In fact, the FCQ-T-r explained above 90 % of the variance of FCQ-T scores, correlated strongly with FCQ-T's items excluded from this short version, and correlations with, for example, BMI and binge eating severity were comparable to those reported in studies in which the full version was used [13, 22, 36]. However, it should be noted that the FCQ-T-r may only be able to capture a subset of aspects related to FC that are evaluated by the 39-item version. Thus, the full version should be administered when assessing the impact that different facets of FC may have on eating patterns. Future studies are needed to further evaluate the psychometric properties of the FCQ-T-r. For example, these studies should investigate test–retest reliability of the FCQ-T-r or further examine the utility of the cut-off score to successfully differentiate people with clinical-level binge eating versus those with probable binge eating. Lastly, investigating the invariance of the FCQ-T-r structure among samples from different countries could be interesting.

**Acknowledgements** This study was not supported financially by any grants.

**Conflict of interests** The authors report no conflicts of interest.

## References

- Weingarten HP, Elston D (1991) Food cravings in a college population. *Appetite* 17:167–175
- White MA, Whisenhunt BL, Williamson DA, Greenway FL, Netemeyer RG (2002) Development and validation of the food-craving inventory. *Obes Res* 10:107–114. doi:10.1038/oby.2002.17
- Vander Wal JS, Johnston KA, Dhurandhar NV (2007) Psychometric properties of the State and Trait Food Cravings Questionnaires among overweight and obese persons. *Eat Behav* 8:211–223. doi:10.1016/j.eatbeh.2006.06.002
- Fabbricatore M, Imperatori C, Morgia A, Contardi A, Tamburello S, Tamburello A et al (2011) Food craving and personality dimensions in overweight and obese patients attending low energy diet therapy. *Obe Metab* 7:e28–e34
- Abiles V, Rodriguez-Ruiz S, Abiles J, Mellado C, Garcia A, Perez de la Cruz A et al (2010) Psychological characteristics of morbidly obese candidates for bariatric surgery. *Obes Surg* 20:161–167. doi:10.1007/s11695-008-9726-1
- White MA, Grilo CM (2005) Psychometric properties of the Food Craving Inventory among obese patients with binge eating disorder. *Eat Behav* 6:239–245. doi:10.1016/j.eatbeh.2005.01.001
- Ng L, Davis C (2013) Cravings and food consumption in Binge Eating Disorder. *Eat Behav* 14:472–475. doi:10.1016/j.eatbeh.2013.08.011
- van der Ster Wallin G, Norring C, Holmgren S (1994) Binge eating versus nonpurged eating in bulimics: is there a carbohydrate craving after all? *Acta Psychiatr Scand* 89:376–381
- Cepeda-Benito A, Gleaves DH, Fernandez MC, Vila J, Williams TL, Reynoso J (2000) The development and validation of Spanish versions of the State and Trait Food Cravings Questionnaires. *Behav Res Ther* 38:1125–1138 S0005-7967(99)00141-2
- Waters A, Hill A, Waller G (2001) Internal and external antecedents of binge eating episodes in a group of women with bulimia nervosa. *Int J Eat Disord* 29:17–22. doi:10.1002/1098-108
- Martin CK, O'Neil PM, Tollefson G, Greenway FL, White MA (2008) The association between food cravings and consumption of specific foods in a laboratory taste test. *Appetite* 51:324–326. doi:10.1016/j.appet.2008.03.002
- Delahanty LM, Meigs JB, Hayden D, Williamson DA, Nathan DM (2002) Psychological and behavioral correlates of baseline BMI in the diabetes prevention program (DPP). *Diabetes Care* 25:1992–1998
- Franken IH, Muris P (2005) Individual differences in reward sensitivity are related to food craving and relative body weight in healthy women. *Appetite* 45:198–201. doi:10.1016/j.appet.2005.04.004
- Meule A, Lutz A, Vögele C, Kübler A (2012) Food cravings discriminate differentially between successful and unsuccessful dieters and non-dieters. Validation of the Food Cravings Questionnaires in German. *Appetite* 58:88–97. doi:10.1016/j.appet.2011.09.010
- Meule A, Hermann T, Kübler A (2014) A short version of the Food Cravings Questionnaire-Trait: the FCQ-T-reduced. *Front Psychol* 5:190. doi:10.3389/fpsyg.2014.00190
- Sitton SC (1991) Role of craving for carbohydrates upon completion of a protein-sparing fast. *Psychol Rep* 69:683–686
- Meule A, Westenhofer J, Kubler A (2011) Food cravings mediate the relationship between rigid, but not flexible control of eating behavior and dieting success. *Appetite* 57:582–584. doi:10.1016/j.appet.2011.07.013
- Fabbricatore M, Imperatori C, Contardi A, Tamburello A, Innamorati M (2013) Food craving is associated with multiple weight loss attempts. *Mediterr J Nutr Metab* 6:79–83. doi:10.1007/s12349-012-0115-x
- Cepeda-Benito A, Gleaves DH, Williams TL, Erath SA (2000) The development and validation of the State and Trait Food-Cravings Questionnaires. *Behav Ther* 31:151–173
- Rodriguez-Martin BC, Molerio-Perez O (2014) Exploring the factor structure of the Food Cravings Questionnaire-Trait in Cuban adults. *Front Psychol* 5:214. doi:10.3389/fpsyg.2014.00214
- Atkinson RL, Dietz WH, Foreyt JP, Goodwin NJ, Hill JO, Hirsch J et al (1993) Very low-calorie diets. National Task Force on the Prevention and Treatment of Obesity, National Institutes of Health. *JAMA* 270:967–974
- Innamorati M, Imperatori C, Balsamo M, Tamburello S, Murri MB, Contardi A et al (2014) Food Cravings Questionnaire-Trait (FCQ-T) discriminates between obese and overweight patients with and without binge eating tendencies: the Italian version of the FCQ-T. *J Pers Assess* (in press)
- Di Bernardo M, Barciulli E, Ricca V, Manucci E, Moretti S, Cabras PL et al (1998) Validazione della versione italiana della Binge Eating Scale in pazienti obesi. *Minerva Psichiatria* 39:125–130
- Cepeda-Benito A, Fernandez MC, Moreno S (2003) Relationship of gender and eating disorder symptoms to reported cravings for food: construct validation of state and trait craving questionnaires in Spanish. *Appetite* 40:47–54 S0195666302001459
- Moreno S, Rodriguez S, Fernandez MC, Tamez J, Cepeda-Benito A (2008) Clinical validation of the trait and state versions of the Food Craving Questionnaire. *Assessment* 15:375–387. doi:10.1177/1073191107312651
- Gormally J, Black S, Daston S, Rardin D (1982) The assessment of binge eating severity among obese persons. *Addict Behav* 7:47–55
- Marcus MD, Wing RR, Hopkins J (1988) Obese binge eaters: affect, cognitions, and response to behavioural weight control. *J Consult Clin Psychol* 56:433–439
- O'Connor BP (2000) SPSS and SAS programs for determining the number of components using parallel analysis and Velicer's MAP test. *Behav Res Methods Instrum Comput* 32:396–402. doi:10.3758/BF03200807
- Balsamo M, Imperatori C, Sergi MR, Belvederi Murri M, Continisio M, Tamburello A et al (2013) Cognitive vulnerabilities and depression in young adults: an ROC curves analysis. *Depress Res Treat* 2013:407602. doi:10.1155/2013/407602
- Fawcett T (2006) An introduction to ROC analysis. *Pattern Recogn Lett* 27:861–874. doi:10.1016/j.patrec.2005.10.010
- Jiménez-Acosta S, Porrata-Maury C, Pérez M (1998) Evolución de algunos indicadores alimentario-nutricionales en Cuba a partir de 1993. *Rev Cubana Med Trop* 50:270–272
- Jiménez-Acosta S, Díaz-Sánchez ME, García-Roche RG, Bonet-Gorbea M, Wong-Ordóñez I (2012) Cambios en el estado nutricional de la población cubana adulta de diferentes regiones de Cuba. *Rev Cubana Hig Epidemiol* 50:4–13
- Komatsu S (2008) Rice and sushi cravings: a preliminary study of food craving among Japanese females. *Appetite* 50:353–358. doi:10.1016/j.appet.2007.08.012
- Osman JL, Sobal J (2006) Chocolate cravings in American and Spanish individuals: biological and cultural influences. *Appetite* 47:290–301. doi:10.1016/j.appet.2006.04.008
- Jarosz PA, Dobal MT, Wilson FL, Schram CA (2007) Disordered eating and food cravings among urban obese African American women. *Eat Behav* 8:374–381. doi:10.1016/j.eatbeh.2006.11.014

36. Meule A, Kübler A (2012) Food cravings in food addiction: the distinct role of positive reinforcement. *Eat Behav* 13:252–255. doi:[10.1016/j.eatbeh.2012.02.001](https://doi.org/10.1016/j.eatbeh.2012.02.001)
37. Lafay L, Thomas F, Mennen L, Charles MA, Eschwege E, Borys JM et al (2001) Gender differences in the relation between food cravings and mood in an adult community: results from the fleurbaix laventie ville sante study. *Int J Eat Disord* 29:195–204. doi:[10.1002/1098-108](https://doi.org/10.1002/1098-108)
38. Imperatori C, Innamorati M, Tamburello S, Continisio M, Contardi A, Tamburello A et al (2013) Gender differences in food craving among overweight and obese patients attending low energy diet therapy: a matched case-control study. *Eat Weight Disord* 18:297–303. doi:[10.1007/s40519-013-0054-7](https://doi.org/10.1007/s40519-013-0054-7)
39. Wardle J, Haase AM, Steptoe A (2006) Body image and weight control in young adults: international comparisons in university students from 22 countries. *Int J Obes (Lond)* 30:644–651. doi:[10.1038/sj.ijo.0803050](https://doi.org/10.1038/sj.ijo.0803050)
40. Polivy J, Coleman J, Herman CP (2005) The effect of deprivation on food cravings and eating behavior in restrained and unrestrained eaters. *Int J Eat Disord* 38:301–309. doi:[10.1002/eat.20195](https://doi.org/10.1002/eat.20195)
41. Geliebter A, Pantazatos SP, McOuatt H, Puma L, Gibson CD, Atalayer D (2013) Sex-based fMRI differences in obese humans in response to high vs. low energy food cues. *Behav Brain Res* 243:91–96. doi:[10.1016/j.bbr.2012.12.023](https://doi.org/10.1016/j.bbr.2012.12.023)
42. Wang GJ, Volkow ND, Telang F, Jayne M, Ma Y, Pradhan K et al (2009) Evidence of gender differences in the ability to inhibit brain activation elicited by food stimulation. *Proc Natl Acad Sci USA* 106:1249–1254. doi:[10.1073/pnas.0807423106](https://doi.org/10.1073/pnas.0807423106)
43. Rozin P, Levine E, Stoess C (1991) Chocolate craving and liking. *Appetite* 17:199–212. doi:[10.1016/0195-6663\(91\)90022](https://doi.org/10.1016/0195-6663(91)90022)
44. Dye L, Warner P, Bancroft J (1995) Food craving during the menstrual cycle and its relationship to stress, happiness of relationship and depression; a preliminary enquiry. *J Affect Disord* 34:157–164 016503279500013