



Approach to hyponatremia according to the clinical setting: Consensus statement from the Italian Society of Endocrinology (SIE), Italian Society of Nephrology (SIN), and Italian Association of Medical Oncology (AIOM)

E. Sbardella¹ · A. M. Isidori¹ · G. Arnaldi² · M. Arosio³ · C. Barone⁴ · A. Benso⁵ · R. Berardi⁶ · G. Capasso⁷ · M. Caprio^{8,9} · F. Ceccato¹⁰ · G. Corona¹¹ · S. Della Casa¹² · L. De Nicola¹³ · M. Faustini-Fustini¹⁴ · E. Fiaccadori¹⁵ · L. Gesualdo¹⁶ · S. Gori¹⁷ · A. Lania¹⁸ · G. Mantovani³ · P. Menè¹⁹ · G. Parenti²⁰ · C. Pinto²¹ · R. Pivonello²² · P. Razzore²³ · G. Regolisti¹⁵ · C. Scaroni¹⁰ · F. Trepiccione⁷ · A. Lenzi¹ · A. Peri²⁴ · On behalf of the: Fluid and Electrolyte Disorders Club of the Italian Society of Endocrinology; Italian Society of Nephrology; and Italian Association of Medical Oncology

Received: 25 July 2017 / Accepted: 13 October 2017 / Published online: 20 November 2017
© Italian Society of Endocrinology (SIE) 2017

Introduction

Hyponatremia (hypoNa, serum sodium levels < 135 mEq/L) is the most frequently observed electrolyte disorder in clinical practice, affecting up to 15–30% of hospitalized patients [1]. HypoNa is characterized by an excess of water relative to exchangeable total body sodium that can be normal, increased, or decreased. As a consequence, hypoNa can be classified by the fluid volume status of the patient (euvolemic, hypovolemic, and hypervolemic hypoNa) or by plasma tonicity, i.e., the effective osmolality (isotonic, hypertonic, and hypotonic hypoNa). Hypotonic hypoNa is the most commonly observed form in daily clinical practice [2]. Severe hypoNa, especially if acutely developed (i.e., in less than 48 h), may determine major neurological symptoms due to brain edema, a potentially life-threatening complication if not promptly recognized and treated [3]. However, even mild hypoNa (130–134 mEq/L) may also be associated with other strictly related clinical problems, often insidious and scarcely symptomatic, such as bone demineralization or gait instability and attention deficits, which may increase the risk of falls and bone fractures, especially in the elderly [4–8]. Accordingly, recent meta-analyses have shown that even mild forms of hypoNa are associated with an increased risk of mortality in different clinical settings [9], along with

prolonged hospital stay, increased readmission rates, and higher hospital costs [10].

Although the main mechanisms of renal sodium and water handling, especially for what concerns the fine regulation by the distal nephron, have been fully elucidated [11], several clinical issues make the approach to hypoNa a complex task. In fact, on one hand, it should be timely diagnosed and appropriately managed according to the severity of the neurological status, but on the other hand, an overly rapid correction may cause neurological damage possibly leading to the Osmotic Demyelination Syndrome (ODS) [12–14]. Conditions that may be associated with a risk of hypercorrection of hypoNa and their related mechanisms are shown in Table 1.

Nevertheless, despite its high prevalence, especially among hospitalized patients, and its clinical impact, hypoNa is often neglected, or under- or mistreated [15]. This is mainly due to both an empirical approach (i.e., not pathophysiology-driven) and to the high degree of heterogeneity of the clinical settings, where hypoNa is encountered.

In recent years, a new class of drugs, namely, the vasopressin receptor antagonists or vaptans, has become available for the treatment of hypoNa secondary to the Syndrome of Inappropriate Antidiuresis (SIAD), one of the most frequent causes of hypoNa [16]. Tolvaptan is the only licensed vaptan in Europe, so far [12]. However, there is no agreement between the European Guidelines [17] and the recommendations of an US Expert Group [14] concerning the use of vaptans in clinical practice [18].

On this basis, a task force generated by the Fluid and Electrolyte Disorders Club of the Italian Society of

E. Sbardella and A. M. Isidori contributed equally to this work.

✉ A. Peri
alessandro.peri@unifi.it

Extended author information available on the last page of the article

Table 1 Conditions at risk for hypoNa overcorrection

Condition	Mechanism of hypercorrection
Hypovolemia	Elimination of the stimulus for ADH secretion due to baroreceptor activation by volume expansion by crystalloid
Low solute diet (beer potomania, tea and toast diet, etc.)	Diet correction increases dietary solute load → increased renal free water clearance
Thiazide diuretic therapy	Discontinuation of the drug directly restores renal diluting capacity
SSRI antidepressant drug therapy	Discontinuation of the drug reduces the serotonergic stimulus on ADH secretion
Hypopituitarism	Restoration of physiologic suppression of ADH secretion by cortisol replacement therapy
Hypoxemia	Elimination of non osmotic stimulus on ADH by normalization of blood gases
Stress, pain, nausea	Elimination of transient stimuli on ADH secretion
Hypokalemia, potassium depletion	With K administration, sodium leaves the cell in exchange with potassium entrance, to maintain electro-neutrality

Endocrinology, the Italian Society of Nephrology, and the Italian Association of Medical Oncology has joined together, to prepare a practical guide to recognize and manage hypoNa in different clinical settings.

In the intention of the task force, the present paper should be considered nor as a formal Guideline nor as an all-inclusive in-depth review on the topic. Rather, this paper should be viewed as a pocket guide to support the practical approach to hyponatremic patients by different specialists.

We propose a simplified diagnostic algorithm for hypoNa (Fig. 1, Table 2) and a treatment algorithm for hypoNa secondary to SIAD (Fig. 2). The treatment strategies for hypovolemic (rehydration) or hypervolemic hypoNa (fluid restriction, hypertonic saline solution, furosemide) are very well established, and for a detailed description, we redirect the reader to the already mentioned recommendations/guidelines [14, 17]. Here, we would like to remind that fluid restriction is not very effective, and in several clinical situations this approach aiming to correct hypoNa is going to fail [19].

Hyponatremia in oncology

Prevalence and etiology

HypoNa in patients with cancer is a common finding, because three major pathogenetic factors may concur to its development: the tumor, through the ectopic secretion of the antidiuretic hormone (ADH), also named vasopressin, the anti-neoplastic treatments, and again the tumor itself, through non-hormonal mechanisms.

About 14% of all cases of hypoNa occur in oncological patients [3]. SIAD is one of the leading causes of hypoNa in inpatients with cancer, affecting 1–2% of the entire cancer population [20, 21]. The likelihood that SIAD is the cause of hypoNa in cancer patients is > 30% [22].

SIAD is commonly reported in small-cell lung cancer. However, hypoNa has also been reported in other tumors, such as gastrointestinal, genitourinary, breast, prostate, or hematological malignancies. SIAD in these patients may also be caused by pharmacological treatments, e.g., by a number of chemotherapeutic agents, opioid analgesics, antidepressants, including tricyclics and selective serotonin reuptake inhibitors (SSRI), as well as phenothiazines used as antiemetic agents (Table 3) [23]. Of notice, hypoNa in oncology patients may be secondary to other conditions besides SIAD (Table 4) [22], and for this reason, a careful differential diagnosis is needed. Finally, hypoNa can be precipitated by fluid and salt losses due to emesis or diarrhea, with severely symptomatic acute hypoNa that can be superimposed to a relatively stable chronic electrolyte imbalance.

Mortality

HypoNa significantly contributes to both morbidity and mortality in cancer patients, and it is an independent prognostic marker: in a large study, the hazard ratio risk of 90-day mortality for mild, moderate, and severe hypoNa was 2.04, 4.74, and 3.46, respectively [24].

Notes on treatment

In general, the treatment strategy of hypoNa in oncologic patients is not different from that suggested by the available recommendations [12–14].

However, specific situations may occur. For instance, in patients with mild hypoNa secondary to SIAD, fluid restriction may be problematic because of the need of parenteral hydration used during chemotherapy.

The use of urea for the treatment of hypoNa, especially in cases of SIAD, has also been proposed since the 1980s [25, 26]. The rationale of this approach is based on the capability of urea to increase the free water clearance by the kidney. The urea dosages usually used in patients with SIAD to




Fig. 1 Simplified diagnostic algorithm for hyponatremia

Table 2 Diagnosis of SIAD. Modified from Ref. [120]

Essential features

Decreased effective osmolality (< 275 mOsm/kg of water)

Urine osmolality > 100 mOsm/kg of water

Clinical euvoemia

No clinical signs of volume depletion of extracellular fluid (tachycardia, decreased skin turgor, or dry mucous membranes)

No clinical signs of excessive volume of extracellular fluid (edema or ascites)

Urinary sodium > 40 mmol/L with normal dietary salt intake

Normal thyroid and adrenal function

No recent use of diuretic agents

Supplemental features

Plasma uric acid < 4 mg/dL

Blood urea nitrogen < 10 mL/dL

Fractional sodium excretion > 1%; fractional urea excretion > 55%

Failure to correct hyponatremia after 0.9% saline infusion

Correction of hyponatremia through fluid restriction

Abnormal results on test of water load (< 80% excretion of 20 mL of water per kilogram of body weight over a period of 4 h), or inadequate urinary dilution (< 100 mOsm/kg of water)

Elevated plasma AVP levels, despite the presence of hypo-tonicity and clinical euvoemia




Fig. 2 Proposed treatment algorithm for hyponatremia secondary to SIAD

correct serum sodium range between 15 and 30 g/day taken orally after a meal in one or two doses [25].

While a few non-controlled studies [27, 28] have reported that urea is effective in normalizing hypoNa, hypercorrection with hyponatremia has also been reported in the same studies. In fact, the urea-induced increase in serum sodium concentration is not easily predicted, as it depends on both hydration status and urine osmolality. Thus, while the European Guidelines [13] recommend urea as the treatment of choice in patients with SIAD when water restriction is ineffective or not feasible, poor palatability, scarce clinical experience, and the risk of hypercorrection suggest that the advantages and disadvantages of urea should be balanced against the possible use of vasopressin receptor antagonists in this clinical setting.

Therefore, a valuable option in cancer patients with SIAD could be represented by vaptans. In a recent prospective study on small-cell lung cancer patients with severe SIAD,

tolvaptan led to an effective correction and stabilization of the serum sodium levels, also enabling patients to receive chemotherapy without any delay [29]. In addition, the use of vaptans may avoid withdrawal of hypoNa-inducing chemotherapies.

The duration of treatment for hypoNa is largely dependent on the cause. In drug-induced hypoNa, the electrolyte alteration is usually reverted within days after the cessation of the involved drug. Conversely, in ADH secreting tumors, hypoNa usually requires a longer and somewhat unpredictable duration of therapy, which is also dependent on the response to anti-tumoral treatments [30].

In summary, we suggest that hypoNa should be carefully taken into account and timely corrected in oncology patients, preferably avoiding severe fluid restriction or agents that may increase nausea (urea), taking into account that the normalization of sodium levels has been found to have a positive effect on the prognosis and length of in-hospital stay [31].

Table 3 Drugs possibly used in oncological patients that may induce hyponatremia

DRUGS	Indication for use	Mechanism involved	References
Vinca alkaloids vincristine, vinblastine	Chemotherapy	Increase ADH secretion	[23]
Platinum compounds Cisplatin, carboplatin		Increase ADH secretion and renal waist syndrome	[121, 122]
Alkylating agents ev cyclophosphamide, melphalan, ifosfamide		Increase ADH secretion and increase renal sensitivity	[123]
Anthracyclines		Hypovolemic hyponatremia	[122, 123]
TK inhibitors and monoclonal antibodies Afatinib Brivanib Cetuximab Gefitinib Linifanib Pazopanib Sorafenib Vorinostat		Direct natriuretic effect and interference in Na pathway and Increase ADH secretion Possible role for iatrogenic hypothyroidism Increase ADH secretion and possible fluid redistribution	[121, 122, 124] [122, 125] [23, 121–123]
Others Methotrexate IFN α - γ Pentostatina IL2			
Opioid Acetaminophen Non-steroidal anti-inflammatory drugs	Pain control	Increased renal sensitivity, indirect increase in ADH secretion secondary to nausea or hypotension	[23, 122, 123]
Tricyclic antidepressant Amitriptyline Protriptyline Desipramine SSRI MAO inhibitors Others Duloxetine, venlafaxine, mirtazapine	Antidepressant	Increase ADH secretion Reset osmostat	[23, 122, 123] [23]
Carbamazepine, oxcarbazepine Sodium valproate Lamotrigine	Antiepileptic	Increase ADH secretion and potentiation ADH effect Reset osmostat	[23, 123, 126]
Phenothiazine	Antiemetic	Drug induced polydipsia	[122]
Corticosteroid	Anti-edema, nausea	Hyperglycemia—pseudohyponatremia	[122, 123]
First antidiabetic generation Chlorpropamide, tolbutamide	Diabetes	potentiation ADH effect	[123]
Antibiotics Ciprofloxacin Trimethoprim/sulfamethoxazole Linezolid Cefoperazone sulbactam	Infections	Increase ADH secretion Hypovolemic hyponatremia	[23] [127, 128]
Proton pump inhibitor Omeprazole, esomeprazole	Prevention gastric ulceration stress or drug related	Increase ADH secretion	[23]
Hypotensive drug Diuretic loop Furosemide Angiotensin converting enzyme inhibitors Thiazide	Hypertension therapy	Hypovolemic hyponatremia Increase osmotic renal losses Increase ADH secretion Increase of thirst	[23]
Mannitol	Anti-edema	Pseudohyponatremia	[122, 123]
Hypotonic solution Isotonic solution	Hydratation	Dilutional	[122, 123]

Table 4 Causes of hypoNa in cancer patients. Modified from Ref. [22]

Causes	%
SIAD	30.4
Dehydration	28.7
Diuretic use	14.0
Hypervolemia	7.8
Kidney failure	3.5
Hypotonic solutions	1.7
Miscellaneous	5.2
Not defined	1.7
False positive	7.0
Mixed causes	9.6
Total	100

Hyponatremia in the elderly

Prevalence and etiology

The prevalence of hypoNa is increased in elderly patients compared with that in the general population, reaching almost 50% of all acute geriatric admissions [32, 33].

In the elderly, the etiology of hypoNa is multifactorial in 50–75% of cases [34, 35]. SIAD is the most common cause, even if a risk of over-diagnosis has been claimed [34]. Other frequent causes are congestive heart failure, water and sodium homeostasis alterations, renal and hepatic dysfunction, and especially drug-induced hypoNa, because older people often receive multiple pharmacological treatments (Table 5) [35].

In elderly patients, alterations of electrolyte and water balance are favoured by age-related reduction in total body water, reduced renal function [36], decreased cortical blood flow and glomerular filtration rate, impaired responsiveness to sodium balance changes [37], osmoreceptors hypersensitivity, and higher ADH release [38]. In addition, the ability to excrete free water is reduced [39].

Mortality

HypoNa is associated with increased all-cause mortality in elderly subjects: a recent study showed that the adjusted

hazards ratio (95% CI) in hyponatremic men without chronic kidney disease (CKD), stroke, or heart failure was 1.30 (confidence interval 1.02–1.66) [40].

Specific notes on clinical features and diagnosis

HypoNa in the elderly is mostly mild, chronic, and apparently asymptomatic, but is often associated with bone demineralization and cognitive impairment, increased risk of falls and fractures [7, 34].

Conversely, acute hypoNa in the elderly is characterized by confusion, irritability, lethargy, anorexia, and nausea, but pre-existing cognitive and sensory impairment might interfere with timely identification of symptoms [34, 38].

The diagnosis in older people may be challenging, due to polypharmacy, difficult assessment of fluid volume status by clinical examination, presence of several confounding co-morbidities, and difficulties in obtaining a reliable clinical history [34, 41].

Notes on treatment

Treatment of both acute and chronic hypoNa in the elderly does not differ from that of younger patients [14, 34]. Vaptans could represent an option in hypoNa secondary to SIAD also in the elderly. The use of low doses—at least initially—may reduce the risk of overtreatment. Appropriate hydration should be strictly monitored. In the case of hypovolemic hypoNa, rehydration should be provided with special caution, especially when cardiac function is reduced and/or chronic kidney disease coexists [36].

Hyponatremia in congestive heart failure

Prevalence and etiology

The prevalence of hypoNa among patients with heart failure (HF) is about 20–25% [42–46]; however, it may be higher in patients admitted for acutely decompensated HF (ADHF): 38% at hospital admission and 28% as new-onset hypoNa during hospital stay [47].

Table 5 Examples of drugs commonly used in elderly patients that may cause or worsen hypoNa [129–133]

Drug classes	Principal drugs involved in the class
Diuretic drugs	Loop diuretics Thiazides
Second-generation antidepressants	Citalopram, escitalopram, paroxetine, fluoxetine, fluvoxamine, venlafaxine, duloxetine, mirtazapine, or sertraline
Proton pump inhibitors	Omeprazole, esomeprazole
Hypotensive drugs	Angiotensin converting enzyme inhibitor

In this clinical setting, effective arterial blood volume (EABV) is decreased, due to low cardiac output and systemic venous congestion. The decreased EABV releases the tonic baroreceptor-dependent inhibition on efferent sympathetic tone and vasopressin release. The ensuing hyperactivation of the sympathetic nervous system, together with renal hypoperfusion, is associated with decreased glomerular filtration rate, increased proximal sodium reabsorption, and reduced sodium delivery to distal nephron segments. These latter mechanisms and the high circulating vasopressin levels—always disproportionate to the reduced plasma tonicity—are mainly responsible for decreased free water clearance and development of hypoNa. On the other hand, secondary hyperaldosteronism due to increased renin release and elevated circulating angiotensin II favors increased sodium reabsorption at the distal nephron and increased total body sodium balance, edema formation, and hypervolemic hypoNa [48].

Mortality

A correlation between hypoNa and overall mortality was first documented 30 years ago in HF. Moreover, in patients with ADHF, hypoNa is associated with an increased mortality and risk of re-hospitalization [47, 49]. In particular, a recent meta-analysis of the available data documented that hypoNa doubled the risk of mortality in patients with HF [9]. In addition, even in patients admitted for ADHF and normal serum sodium values, it has been shown that the development and worsening of hypoNa during hospital stay are strongly correlated with an increase in overall and cardiovascular mortality [50].

Finally, a recent retrospective study in patients admitted for ADHF with hypoNa at admission reported that even persistent hypoNa at the time of hospital discharge is associated with a significant increase of re-hospitalization or mortality at 30 days [51].

Notes on treatment

Clearly, sodium and fluid restriction, diuretics, blockers of the renin–angiotensin–aldosterone system, and beta-blockers are the mainstay of treatment in patients with HF. While hypoNa bears a clear negative prognostic impact, few data are currently available in the literature to clearly ascertain whether correction of hypoNa per se may ameliorate outcomes in patients with HF [13]. As a matter of fact, long-term treatment with tolvaptan in patients with HF was not associated with decreased mortality or risk of re-hospitalization compared with placebo, notwithstanding greater weight loss, better dyspnea relief, and a significant increase in serum sodium values at discharge [52]. However, a post hoc analysis of the Acute and Chronic Therapeutic

Impact of a Vasopressin antagonist (ACTIV) study [53] suggested a possible correlation between increased serum sodium levels and increased survival. Furthermore, a post hoc analysis of the Efficacy of Vasopressin Antagonism in Heart Failure Outcome Study With Tolvaptan (EVEREST) trial showed decreased incidence of the combined endpoint of cardiovascular mortality and cardiovascular morbidity in tolvaptan-treated patients with serum sodium values $\text{Na} < 130 \text{ mEq/L}$ [54]. Thus, expert consensus by US investigators [14] suggested that vaptans (tolvaptan, and possibly conivaptan, not licensed in Europe, so far) may represent a useful therapeutic tool in patients with CHF (chronic heart failure) and mild-to-moderate hyponatremia. On the other hand, based on the results of an extended meta-analysis indicating a non-significant trend towards increased mortality in hyponatremic patients with expanded extracellular fluid volume, European guidelines [13] are against the use of vaptans in conditions where hyponatremia is associated with expanded extracellular fluid volume. A faster decongestion with dyspnea relief represents a desirable goal in the treatment of patients with ADHF, and no cases of osmotic myelinolysis have been reported either in the ACTIV and EVEREST trials or in the subgroup of patients with CHF enrolled in the Study of Ascending Levels of Tolvaptan in Hyponatremia 1 and 2 (SALT-1 and SALT-2) trials [55]. For these reasons, the use of tolvaptan may be envisaged as a potentially useful add-on treatment strategy in patients with CHF and mild-to-moderate hypoNa. However, very recently, the Targeting Acute Congestion with Tolvaptan in Congestive Heart Failure (TACTICS-HF) trial [56] reported no significant differences in dyspnea relief and in-hospital or post-discharge clinical outcomes in patients with ADHF treated with tolvaptan 30 mg given at 0, 24, and 48 h on top of fixed-dose furosemide compared with patients receiving placebo, despite greater weight loss and net fluid loss in tolvaptan-treated patients. Thus, at the present time, no firm recommendation about the use of vaptans in CHF can be supported by the available literature data.

Hyponatremia in decompensated liver cirrhosis

Prevalence and etiology

The prevalence of hypoNa in patients admitted for decompensated liver cirrhosis reaches 57% [57]. Between 21 and 28% of patients have serum sodium values $< 130 \text{ mEq/L}$ [57–59], whereas severe hypoNa (serum $\text{Na} \leq 120 \text{ mEq/L}$) is relatively infrequent ($< 1.2\%$) in this setting [57].

Post-sinusoidal capillary hypertension, hypoalbuminemia, and splanchnic vasodilation play a pivotal role in ascites accumulation; specifically, overproduction of nitric oxide,

mainly due to circulating endotoxin associated with bacterial translocation, maintains splanchnic vasodilation [60]. In this clinical setting, pathophysiological mechanisms triggered by decreased EABV are essentially the same as in HF. Thus, while total body sodium balance is increased, the development of hypoNa is facilitated by reduced sodium delivery to the distal nephron and high circulating ADH levels [60].

Mortality

The negative prognostic role of hypoNa in patients with liver cirrhosis has been clearly documented [59, 60]. In a study performed in 6769 patients with liver failure waiting for liver transplantation, of whom 422 had deceased within 90 days since entering the waiting list, the investigators found an increased risk of death associated with hypoNa, even independent of the MELD score [61]. Accordingly, a recent meta-analysis of the available data documented that hypoNa was associated with more than threefold increased risk of mortality in patients with cirrhosis [9].

Notes on treatment

In patients with liver failure, treatment with vaptans has been shown to ameliorate fluid balance in some studies [62–64]. Moreover, treatment with satavaptan was associated with greater increase in serum sodium values and decreased ascites formation in cirrhotic patients receiving either diuretics [65] and spironolactone [59]. However, one of those studies [65] also found an increase in the risk of death due to complication of cirrhosis in patients treated with both satavaptan and diuretics, which subsequently lead to drug withdrawal from commerce. Lixivaptan, on top of standard treatment with spironolactone, proved to be effective in increasing free water clearance and serum sodium values in patients with decompensated liver cirrhosis [66].

Tolvaptan, so far the only vaptan approved in Europe and allowed only in patient with hypoNa secondary to SIAD, has also been tested on top of the standard treatment with furosemide and spironolactone in patients with decompensated liver cirrhosis. Although the drug proved to be effective, also at low doses (e.g., 3.5 and 7.5 mg/day), in reducing ascites volume and body weight [67], as well as in increasing serum sodium values [68, 69], both the European guidelines [13] and the US expert consensus document [14] recommend against the use of vaptans in patients with liver disease. Anyway, as hypoNa may complicate the use of high-dose loop diuretics in oliguric patients with refractory ascites and may predispose them to hepatic encephalopathy, a cautious use of tolvaptan in combination with diuretics may represent a treatment strategy that should be explored in future research [60]. In any case, liver function should be closely monitored,

and the drug should be discontinued if worsening of it is detected.

Hyponatremia in chronic kidney disease

Prevalence and etiology

Prevalence and incidence values of hypoNa in chronic kidney disease (CKD) are, respectively, 13.5 and 26% (mean follow-up 5.5 years) [70]. HypoNa is especially common among patients with stage 5 CKD on dialysis (End-Stage Renal Disease, ESRD): 29.3% prevalence in hemodialysis and 14.5% incidence in peritoneal dialysis [71].

Three main pathogenetic mechanisms may lead to hypoNa in renal patients [72]:

- Direct renal sodium loss, such as in salt-losing nephropathies (chronic pyelonephritis, chronic drug-associated, or toxic tubule interstitial), characterized by reduced renal sodium reabsorption, sodium and potassium depletion, reduced concentration ability, hypovolemia, and ADH stimulation [73]. The consequence is hypoNa with volume depletion.
- Reduced urinary dilution capacity, due to severe impairment of glomerular filtration rate with ensuing lower availability of preurine at the diluting segments of the distal nephron and reduced ability to generate free water. In this case, hypoNa will be euvolemic or hypervolemic [72, 74].
- Oligoanuria or anuria along with free water/hypotonic fluid intakes exceeds losses, such as the case of ESRD on chronic dialysis or in acute kidney injury. These forms of hypoNa are usually hypervolemic [74].

Mortality

Low sodium levels are associated with increased mortality risk, both in CKD patients on conservative treatment [70] and in ESRD patients on dialysis [75–77].

Notes on treatment

There is paucity of data in the literature concerning the treatment of hypoNa in patients with CKD on conservative therapy. Tolvaptan has been tested in small studies performed in CKD patients with or without congestive heart failure [78]. On the whole, a significant increase in urine volume was observed, together with an increase in serum sodium concentration. Moreover, treatment with tolvaptan

was not associated with deterioration of kidney function in these patients.

Issues associated with the treatment of hypoNa during renal replacement therapy (RRT), and the inherent risks of overcorrection and osmotic demyelization syndrome, have been addressed mainly in the critically ill patients. Specifically, reducing serum sodium concentration in the substitution fluids has been advocated as the best approach to avoid the risk of overcorrection during continuous veno-venous hemofiltration or continuous veno-venous hemodialysis [79]. When the standard intermittent hemodialysis is chosen, sodium concentration in the dialysate should be reduced to a minimum of 130 mEq/L, and blood flow rate as low as 50 mL/min together with short duration (e.g., 3 h) of dialysis session should be prescribed [80].

Hyponatremia in neurology

Prevalence and etiology

HypoNa is a frequent complication of traumatic brain injury and meningitis [81].

Limited information is available for other neurological disorders. A large Swedish registry study documented that epilepsy and stroke accounted for about 10% of all cases of hypoNa [82].

In subjects with stroke, many factors including dietary sodium restriction for hypertension control, use of thiazide diuretics, and infections might precipitate hypoNa [83].

Several antiepileptic drugs (AEDs) and in particular carbamazepine, oxcarbazepine, eslicarbazepine, and levetiracetam may cause asymptomatic or mildly symptomatic hypoNa secondary to SIAD, which in turn may exacerbate seizures (Table 3) [81].

HypoNa can occur in Guillain–Barré syndrome (GBS) as a consequence of SIAD caused by the intravenous immunoglobulin therapy, or of renal salt wasting syndrome as part of GBS-related dysautonomia [84].

Mortality

HypoNa is associated with an increased risk of mortality in patients with neurological diseases: in a Danish cohort study, the adjusted 30-day relative risk of death among hyponatremic patients compared to patients with normonatremia was 1.5 (0.9–2.5) [85].

Serum sodium evaluation should be mandatory in the presence of neurological symptoms. Routine sodium monitoring for patients receiving AEDs is not usually necessary, except in elderly subjects or in those receiving AED polytherapy or sodium-depleting drugs [81].

Notes on treatment

HypoNa in neurological patients should be managed according to the general recommendations. Treatment mainly depends on etiology; it has been shown, for instance, that traumatic brain injury treatment usually lasts 0.5–2 years [30]. In SIAD caused by AEDs, the possibility to reduce the dose, to switch to a different drug or stop treatment, should be evaluated together with the neurologist.

Hyponatremia in neurosurgery

Prevalence and etiology

Hypotonic hypoNa is a frequent finding in neurosurgical patients, with the highest rate (20–50%) in some series among patients with subarachnoid hemorrhage (SAH) [86, 87]. Observational studies have shown that brain tumors, during their course, may be associated with hypoNa in about 15–20% of cases [86]. The occurrence of hypoNa as a result of transsphenoidal surgery varies a lot in the different series, depending on the selection criteria. Symptomatic hypoNa was much less frequent (4–7%) than asymptomatic hypoNa, which in some series occurred in up to 20–35% of patients, according to serum sodium measured every day for at least 12 days after surgery [88–90].

Some neurosurgical disorders, such as acute and chronic SAH, subdural hematoma, hemorrhagic stroke, tumors, cysts, metastases, and inflammatory diseases of the brain, pituitary, or hypothalamus, become harder to manage when hypoNa develops [86, 87, 90–92]. Such a complication may occur both before and after surgery. HypoNa may also be observed at presentation in patients with pituitary apoplexy yet much less frequently than hypernatremia due to diabetes insipidus (DI).

Mortality

Besides hypoNa, several other factors may contribute to increase the mortality risk in the neurosurgery setting. A recent systematic review, aimed at characterizing the effect of hypoNa on morbidity and mortality after SAH, included 13 studies with a total number of 2387 patients and showed that hypoNa was associated with increased morbidity (especially due to vasospasm), but it did not influence mortality [87]. Interestingly, a recent retrospective observational study reviewed 198 consecutive patients with SAH and indicated sodium fluctuation, rather than hypoNa per se, as a significant factor associated with worse neurologic outcome [91].

Specific notes on clinical features and diagnosis

Most observational studies have shown that SIAD is the commonest cause of hypotonic hypoNa in neurosurgical patients [86]. However, in this setting, it is essential to differentiate SIAD and cerebral salt wasting syndrome (CSWS) as a possible cause of hypoNa, especially in pediatric series and in patients with SAH [86, 92, 93]. The differential diagnosis between CSWS and SIAD may not be easy in clinical practice, the former having hypovolemia as a crucial point for the proper diagnosis [94].

In the evaluation of the hyponatremic patient after neurosurgery, it is essential to consider the possible occurrence of DI with a triphasic pattern [95] and the possible late occurrence of hypoNa due to SIAD, which can occur after the patient has been discharged [88–90].

To improve the outcome in neurosurgical patients, we suggest careful monitoring of serum sodium on admission and during the hospital stay. Whenever hypoNa is observed, a proper workup has to be instructed to elucidate the underlying cause, bearing in mind that the evaluation of extracellular fluid volume status is mandatory. In addition, in case of early discharge, the patients should receive clear instructions on what to do if hypoNa-compatible symptoms appear.

Hyponatremia in the patient with trauma and polytrauma

Prevalence and etiology

Little is known about hypoNa in patients with polytrauma (PT). In several studies, hypoNa has been reported in up to 15% of patients after trauma or PT [96, 97]. After PT, the occurrence of hypoNa can be related to the event per se (fluid depletion, hemorrhage), to the immediate treatment at the site of trauma or Emergency Department (hypotonic intravenous fluids), or to a pre-existing comorbidity, especially in the elderly.

Mortality

HypoNa is associated with poor prognosis and increased mortality in patients with crush syndrome: in a retrospective study conducted in Chinese reference hospitals during the Wenchuan earthquake the presence of hypoNa was common, up to 50% of patients were affected and 15% of them died. However, here, the hypoNa was mainly correlated with the development of acute kidney failure [96].

Specific notes on clinical features and diagnosis

Hip fracture is the commonest cause of traumatic death in Europe: immediate surgery has been associated with higher rates of independent living, lower mortality rates, improved patient outcomes by reducing pain scores, and lowering the risk of decubitus ulcers. The occurrence of hypoNa in the course of the pre-surgical planning was the main medical pre-operative risk factor for surgery delay after 36 h from trauma [98].

Rather than considering hypoNa always as a consequence of PT, this electrolyte disorder could be the cause of trauma: even mild hypoNa in fact has been associated with unsteady gait, falls, impaired concentration, and risk of fractures, especially hip and femur fractures [6, 99]. In an extensive series of elderly adults, fracture risk incidence was higher in patients with hypoNa, also after adjusting for osteoporosis. Patients with moderate–severe hypoNa ($[\text{Na}]^+ < 130 \text{ mEq/L}$) presented an 11-fold risk of fractures [100], and fragility fractures increased incrementally with a categorical decrease in median serum sodium levels in multivariate logistic regression models [101].

Notes on treatment

Specific consensus about the treatment of hypoNa targeted to PT patients has not been developed, yet. In general, we suggest to follow currently available guidelines and recommendations for the management of hypoNa.

HypoNa is common in PT patients without neurological involvement: however, larger studies are needed to investigate the relationship between trauma and serum sodium levels, and hypotonic intravenous fluids should be supplied carefully to patients with PT.

HypoNa is not always recognized in patients with PT: we suggest to pay attention to sodium balance in such patients. Population studies with a large number of participants have to be performed.

Hyponatremia in the outpatient setting

Prevalence and etiology

The prevalence of hypoNa in this setting greatly depends on the age of the population considered [102–105]. In a young and ethnically diverse population, the prevalence of hypoNa was 6.3% [105], but with aging, the potential risk of developing hypoNa increases [103, 106].

In the outpatient setting, hyponatremic individuals are more likely to be smokers, to have black ethnicity, a history of diabetes mellitus, congestive HF, or cirrhosis and to use thiazide diuretics, antiepileptic drugs, or SSRI [103, 105].

However, in the Dallas Heart Study, among hyponatremic individuals with no predisposing medical conditions, 20% of them had criteria indicating a diagnosis of SIAD [105].

Mortality

There are limited data in the literature regarding hypoNa in the outpatient setting, but similar to hospitalized patients, also in the outpatient studies hypoNa has been reported to be an independent mortality risk factor [9, 102, 105, 107, 108]. In a recent survey, hypoNa was found to be associated with a nearly twofold increase in deaths, even after adjusting for major risk factors [105].

Specific notes on clinical features and diagnosis

HypoNa in the outpatient is more likely to be mild, chronic, and asymptomatic [103, 105, 107].

Actually, in clinical practice, hypoNa very often represents an incidental finding during an outpatient visit for another reason and it is difficult for the physician to formulate promptly a correct etiological diagnosis, which greatly depends on additional laboratory tests that may not be readily available.

Notes on treatment

In the case of moderately or severely symptomatic hypoNa, hospitalization should be considered.

In the case of hypoNa secondary to SIAD, when vaptans use is indicated, patients should be hospitalized because of the need of close initial monitoring and to identify the appropriate dose [9, 109]. A day-hospital admission may be suitable if there are no other serious concomitant disorders [110].

A regular outpatient follow-up is recommended, to evaluate the effectiveness of the therapy as well as the possibility of discontinuing it.

We suggest that hypoNa in this setting should be taken into consideration even if mild-to-moderate in order to timely correct it, particularly in the elderly and in patients assuming drugs, thus likely limiting the consequences of persistent low sodium levels.

Life-threatening hyponatremia

Prevalence and etiology

Acute and severely symptomatic hypoNa is rare. However, if not rapidly recognized and correctly treated, it may carry a high morbidity and mortality rate, even in previously healthy subjects, such as, for example, marathon runners, in which an incidence of 13% has been documented [111]. Other causes can be represented by the rapid ingestion of large amounts of water, for example, in psychiatric patients, or of other hypotonic liquids, such as in beer potomania or tea and toast diet. Other conditions associated with acute and potentially life-threatening hypoNa are the post-operative period, in particular after prostate transurethral resection or post uterine endoscopic surgery due to the use of hypotonic irrigant solutions, colonoscopy preparation, the use of some drugs such as oxytocin or cyclophosphamide, or a recent prescription of thiazides or desmopressin, and use of recreational drugs such as ecstasy (MDMA) [13, 112]. The severity of the picture correlates both with the magnitude and with the rate of sodium decrease.

Mortality

Mortality in this setting has been noted to be as high as 55% [113]. However, the estimate from a broad-based literature survey gives much lower values [114].

Specific notes on clinical features and diagnosis

HypoNa may be itself the direct cause of death because of brain stem herniation due to cerebral edema for serum hypotonicity. Risk factors for brain edema are both the rate and the depth of sodium fall.

The risk of death as a consequence of brain edema is increased in the presence of an intracranial disease, in the case of post-operative hypoNa or acute water intoxication.

Notes on treatment

Prompt infusion of hypertonic saline, independent of volume status, may save lives in life-threatening hypoNa.

In this emergency setting, hypertonic 3% NaCl saline solution is administered as a 100/150 mL bolus (or 2 mL/kg of body weight) given over 10–20 min, strictly monitoring sodium levels (every 20 min), and repeating the bolus administration, as needed, up to a maximum of three times. According to the European guidelines, this protocol is recommended until a serum sodium increase of 5 mEq/L is achieved [13].

In the case of symptoms improvement and/or after a 5–6 mEq/L increase in serum sodium (symptoms relief can take longer), 3% NaCl should be stopped, but the i.v. access kept. Meanwhile, a diagnosis-specific process should be initiated, and appropriate management performed [13, 14].

Hyponatremia overcorrection: conditions at risk, prevention, and treatment

Excessive correction of hypoNa (i.e., too much and/or too rapid increase of serum sodium levels) is associated with an increased risk of negative neurologic outcomes (i.e., the ODS), especially in the chronic forms of hypoNa. On this regard, specific attention has to be paid to the fact that during sodium correction an adequate renal response to hypotonicity (i.e., a hypotonic polyuria) is often spontaneously (and rapidly) restored, even in the first 8–12 h from starting of treatment. This usually happens when pathogenetic factors responsible for the electrolyte derangement are promptly taken away, such as, for example, by volume expansion with 0.9% saline in hypovolemic hypoNa, or by ceasing the trigger mechanism for inappropriate secretion/response to ADH (i.e., drugs or inflammation). Thus, since the most frequent cause of hypoNa overcorrection is actually the reactivation of the normal renal physiological response (increased free water clearance), special attention should be paid to hypoNa settings characterized by rapidly reversible causes (Table 1) [18]. A hypotonic polyuria with maximally diluted urine output may in fact further increase the programmed/estimated rate of correction.

In such a circumstance, administration of a hypotonic solution should be started, as intravenous 5% dextrose or free water by a nasogastric tube, initially at 10 mL/kg/h over 1 h [13] or in repeated 3 mL/kg infusions [14] and then matched to urinary output in terms of rate and tonicity; desmopressin (i.v. or s.c.) at 2–4 µg every 8 h can be associated, to bring back the rate of correction to below 12 mEq/L/24 h (or better to a target of 6–8 mEq/L in the first 24 h) [13]. It is mandatory that specific measures to blunt overcorrection of hypoNa must be implemented by/or under the direction of experienced medical personnel [13].

Another possible (and underrated) cause of overcorrection of hypoNa is represented by the administration of potassium salts along with NaCl, aiming at correcting coexisting hypokalemia/potassium depletion. Based on the original Edelman equation, potassium and sodium salts are equivalent in terms of tonicity effects [115]. In fact, in case of cellular potassium depletion, the administered potassium enters the cells, with ensuing Na exit to maintain the electrical equilibrium. Thus, serum sodium values increase [116].

The risk of overcorrection is not significantly reduced by the use of specific formulas aimed at estimating the rate and

the temporal trajectory of serum sodium during correction [117]. Formulas may be useful to set the start of therapy, but they do not completely avoid the risk of overcorrection due to their inherent limitations: they do not take into account the possibility of a rapidly restored diluting capacity by the kidney. Ongoing losses and other electrolyte supplements are difficult to be integrated in the calculation, as it is the case of potassium administration in potassium depletion [117]. More conservatively, in these cases, it is better to frequently check (at least every 4 h in the first 24 h) the actual serum sodium levels.

Finally, it should be kept in mind that some conditions are associated with a higher risk of ODS due to overly rapid correction of hypoNa: serum sodium levels less than 105 mEq/L, alcoholism, malnutrition, and advanced liver disease [14] (Table 1). These conditions must be recognized even before the start of treatment and a great caution should be used in these situations.

Conclusions

Despite being the most common electrolyte disorder encountered in clinical practice, hypoNa is frequently underdiagnosed and/or not appropriately treated. This may be due to a lack of awareness of the implications of this condition on patient outcomes, particularly when hospital-acquired and mildly or moderately symptomatic. Appropriate workup and treatment in the various clinical settings associated with hypoNa require a multidisciplinary approach. In such a need, this task force has provided the above-outlined suggestions and warnings. Ineffective management of hypoNa may negatively affect patient prognosis. New therapeutic options for the correction of hypoNa, particularly vaptans, the vasopressin receptor antagonists, represent an effective tool to safely treat this disorder and improve outcomes among a wide range of patients with hypoNa secondary to SIAD. However, the different clinical scenarios in which hypoNa may occur suggest that a thoughtful and personalized management should be individuated. This scenario is even more complex when we consider that not all the hospitals are properly equipped to perform an accurate differential diagnosis of hypoNa. As an example, the infrequent availability of osmometers in the medium/small hospital facilities is a limiting factor for the diagnosis of SIAD. Thus, we propose that clinicians may refer to calculated serum and/or urinary osmolality according to recently reviewed formulae [118].

A rapid recognition and optimal treatment of hypoNa can reduce the risk of death [119], also reducing the length of hospitalization and associated costs, and improving the quality of life.

Acknowledgements We would like to thank all the members of the *Fluid and Electrolyte Disorders Club of the Italian Society of Endocrinology*, for the scientific support during the writing of the manuscript.

Compliance with ethical standards

Conflict of interest A.M.I. has been an occasional consultant for Shire and Novartis; R.B. has been an occasional consultant for Otsuka; G.M. has been an occasional consultant for Shire and Novartis; A.L. has been an occasional consultant for Shire and Novartis; A.P. has been an occasional consultant for Otsuka. All other authors have nothing to disclose.

Funding The study was not funded.

Ethical approval This consensus statement does not contain any studies with human participants or animals performed by any of the authors.

Informed consent No informed consent needed.

References

- Upadhyay A, Jaber BL, Madias NE (2006) Incidence and prevalence of hyponatremia. *Am J Med* 119(7 Suppl 1):S30–S35
- Adroge HJ, Madias NE (2000) Hyponatremia. *N Engl J Med* 342(21):1581–1589
- Gill G, Huda B, Boyd A, Skagen K, Wile D, Watson I et al (2006) Characteristics and mortality of severe hyponatraemia—a hospital-based study. *Clin Endocrinol (Oxf)* 65(2):246–249
- Renneboog B, Musch W, Vandemergel X, Manto MU, Decaux G (2006) Mild chronic hyponatremia is associated with falls, unsteadiness, and attention deficits. *Am J Med.* 119(1):71e1–71e8
- Gankam Kengne F, Andres C, Sattar L, Melot C, Decaux G (2008) Mild hyponatremia and risk of fracture in the ambulatory elderly. *QJM* 101(7):583–588
- Kinsella S, Moran S, Sullivan MO, Molloy MG, Eustace JA (2010) Hyponatremia independent of osteoporosis is associated with fracture occurrence. *Clin J Am Soc Nephrol* 5(2):275–280
- Verbalis JG, Barsony J, Sugimura Y, Tian Y, Adams DJ, Carter EA et al (2010) Hyponatremia-induced osteoporosis. *J Bone Miner Res* 25(3):554–563
- Barsony J, Sugimura Y, Verbalis JG (2011) Osteoclast response to low extracellular sodium and the mechanism of hyponatremia-induced bone loss. *J Biol Chem* 286(12):10864–10875
- Corona G, Giuliani C, Parenti G, Norello D, Verbalis JG, Forti G et al (2013) Moderate hyponatremia is associated with increased risk of mortality: evidence from a meta-analysis. *PLoS One* 8(12):e80451
- Corona G, Giuliani C, Parenti G, Colombo GL, Sforza A, Maggi M et al (2016) The economic burden of hyponatremia: systematic review and meta-analysis. *Am J Med.* 129(8):823e4–835e4
- Chambrey R, Trepiccione F (2015) Relative roles of principal and intercalated cells in the regulation of sodium balance and blood pressure. *Curr Hypertens Rep* 17(4):538
- Cuesta M, Garrahy A, Thompson CJ (2016) SIAD: practical recommendations for diagnosis and management. *J Endocrinol Invest* 39(9):991–1001
- Spasovski G, Vanholder R, Allolio B, Annane D, Ball S, Bichet D et al (2014) Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Intensive Care Med* 40(3):320–331
- Verbalis JG, Goldsmith SR, Greenberg A, Korzelius C, Schrier RW, Sterns RH et al (2013) Diagnosis, evaluation, and treatment of hyponatremia: expert panel recommendations. *Am J Med* 126(10 Suppl 1):S1–S42
- Hoorn EJ, Lindemans J, Zietse R (2006) Development of severe hyponatraemia in hospitalized patients: treatment-related risk factors and inadequate management. *Nephrol Dial Transplant* 21(1):70–76
- Rondon-Berrios H, Berl T (2016) Vasopressin receptor antagonists: characteristics and clinical role. *Best Pract Res Clin Endocrinol Metab* 30(2):289–303
- Spasovski G, Vanholder R, Allolio B, Annane D, Ball S, Bichet D et al (2014) Clinical practice guideline on diagnosis and treatment of hyponatraemia. *Nephrol Dial Transplant* 29(Suppl 2):i1–i39
- Regolisti G, Cabassi A, Antonucci E, Brusasco I, Cademartiri C, Pistolesi V et al (2015) Hyponatremia in clinical practice. *G Ital Nefrol* 32(1)
- Aylwin S, Burst V, Peri A, Runkle I, Thatcher N (2015) ‘Dos and don’ts’ in the management of hyponatremia. *Curr Med Res Opin* 31(9):1755–1761
- Glover DJ, Glick JH (1987) Metabolic oncologic emergencies. *CA Cancer J Clin* 37(5):302–320
- Silverman P, Distelhorst CW (1989) Metabolic emergencies in clinical oncology. *Semin Oncol* 16(6):504–515
- Berghmans T, Paesmans M, Body JJ (2000) A prospective study on hyponatraemia in medical cancer patients: epidemiology, aetiology and differential diagnosis. *Support Care Cancer* 8(3):192–197
- Liamis G, Milionis H, Elisaf M (2008) A review of drug-induced hyponatremia. *Am J Kidney Dis* 52(1):144–153
- Doshi SM, Shah P, Lei X, Lahoti A, Salahudeen AK (2012) Hyponatremia in hospitalized cancer patients and its impact on clinical outcomes. *Am J Kidney Dis* 59(2):222–228
- Sterns RH, Silver SM, Hix JK (2015) Urea for hyponatremia? *Kidney Int* 87(2):268–270
- Decaux G, Unger J, Brimiouille S, Mockel J (1982) Hyponatremia in the syndrome of inappropriate secretion of antidiuretic hormone. Rapid correction with urea, sodium chloride, and water restriction therapy. *JAMA* 247(4):471–474
- Decaux G, Andres C, Gankam Kengne F, Soupart A (2010) Treatment of euvolemic hyponatremia in the intensive care unit by urea. *Crit Care* 14(5):R184
- Soupart A, Coffernils M, Couturier B, Gankam-Kengne F, Decaux G (2012) Efficacy and tolerance of urea compared with vaptans for long-term treatment of patients with SIADH. *Clin J Am Soc Nephrol* 7(5):742–747
- Petereit C, Zaba O, Teber I, Luders H, Grohe C (2013) A rapid and efficient way to manage hyponatremia in patients with SIADH and small cell lung cancer: treatment with tolvaptan. *BMC Pulm Med* 13:55
- Verbalis JG (2010) Managing hyponatremia in patients with syndrome of inappropriate antidiuretic hormone secretion. *Endocrinol Nutr* 57(Suppl 2):30–40
- Berardi R, Caramanti M, Castagnani M, Guglielmi S, Marcucci F, Savini A et al (2015) Hyponatremia is a predictor of hospital length and cost of stay and outcome in cancer patients. *Support Care Cancer* 23(10):3095–3101
- Berl T (2013) An elderly patient with chronic hyponatremia. *Clin J Am Soc Nephrol* 8(3):469–475
- Mannesse CK, Vondeling AM, van Marum RJ, van Solinge WW, Egberts TC, Jansen PA (2013) Prevalence of hyponatremia on geriatric wards compared to other settings over four decades: a systematic review. *Ageing Res Rev* 12(1):165–173

34. Soiza RL, Cumming K, Clarke JM, Wood KM, Myint PK (2014) Hyponatremia: special considerations in older patients. *J Clin Med* 3(3):944–958
35. Shapiro DS, Sonnenblick M, Galperin I, Melkonyan L, Munter G (2010) Severe hyponatraemia in elderly hospitalized patients: prevalence, aetiology and outcome. *Intern Med J* 40(8):574–580
36. Beck LH (1998) Changes in renal function with aging. *Clin Geriatr Med* 14(2):199–209
37. Lindeman RD, Tobin J, Shock NW (1985) Longitudinal studies on the rate of decline in renal function with age. *J Am Geriatr Soc* 33(4):278–285
38. Moran D, Fronk C, Mandel E (2014) Managing hyponatremia in adults. *JAAPA* 27(4):23–29 (quiz 30)
39. Kugler JP, Husted T (2000) Hyponatremia and hypernatremia in the elderly. *Am Fam Physician* 61(12):3623–3630
40. Wannamethee SG, Shaper AG, Lennon L, Papacosta O, Whincup P (2016) Mild hyponatremia, hypernatremia and incident cardiovascular disease and mortality in older men: a population-based cohort study. *Nutr Metab Cardiovasc Dis* 26(1):12–19
41. Hoyle GE, Chua M, Soiza RL (2011) Volaemic assessment of the elderly hyponatraemic patient: reliability of clinical assessment and validation of bioelectrical impedance analysis. *QJM* 104(1):35–39
42. De Luca L, Klein L, Udelson JE, Orlandi C, Sardella G, Fedele F et al (2005) Hyponatremia in patients with heart failure. *Am J Cardiol* 96(12A):19L–23L
43. Gheorghide M, Abraham WT, Albert NM, Gattis Stough W, Greenberg BH, O'Connor CM et al (2007) Relationship between admission serum sodium concentration and clinical outcomes in patients hospitalized for heart failure: an analysis from the OPTIMIZE-HF registry. *Eur Heart J* 28(8):980–988
44. Friedewald VE, Emmett M, Gheorghide M, Roberts WC (2011) The editor's roundtable: pathophysiology and management of hyponatremia and the role of vasopressin antagonists. *Am J Cardiol* 107(9):1357–1364
45. Farmakis D, Filippatos G, Parissis J, Kremastinos DT, Gheorghide M (2009) Hyponatremia in heart failure. *Heart Fail Rev* 14(2):59–63
46. Gheorghide M, Rossi JS, Cotts W, Shin DD, Hellkamp AS, Pina IL et al (2007) Characterization and prognostic value of persistent hyponatremia in patients with severe heart failure in the ESCAPE Trial. *Arch Intern Med* 167(18):1998–2005
47. Wald R, Jaber BL, Price LL, Upadhyay A, Madias NE (2010) Impact of hospital-associated hyponatremia on selected outcomes. *Arch Intern Med* 170(3):294–302
48. Urso C, Brucculeri S, Caimi G (2015) Acid-base and electrolyte abnormalities in heart failure: pathophysiology and implications. *Heart Fail Rev* 20(4):493–503
49. Bettari L, Fiuzat M, Shaw LK, Wojdyla DM, Metra M, Felker GM et al (2012) Hyponatremia and long-term outcomes in chronic heart failure—an observational study from the Duke Databank for Cardiovascular Diseases. *J Card Fail* 18(1):74–81
50. Konishi M, Haraguchi G, Ohgashi H, Sasaoka T, Yoshikawa S, Inagaki H et al (2012) Progression of hyponatremia is associated with increased cardiac mortality in patients hospitalized for acute decompensated heart failure. *J Card Fail* 18(8):620–625
51. De Vecchis R, Di Maio M, Di Biase G, Ariano C (2016) Effects of hyponatremia normalization on the short-term mortality and rehospitalizations in patients with recent acute decompensated heart failure: a retrospective study. *J Clin Med* 5(10):92
52. Konstam MA, Gheorghide M, Burnett JC Jr, Grinfeld L, Maggioni AP, Swedberg K et al (2007) Effects of oral tolvaptan in patients hospitalized for worsening heart failure: the EVEREST Outcome Trial. *JAMA* 297(12):1319–1331
53. Rossi J, Bayram M, Udelson JE, Lloyd-Jones D, Adams KF, O'Connor CM et al (2007) Improvement in hyponatremia during hospitalization for worsening heart failure is associated with improved outcomes: insights from the Acute and Chronic Therapeutic Impact of a Vasopressin Antagonist in Chronic Heart Failure (ACTIV in CHF) trial. *Acute Card Care* 9(2):82–86
54. Hauptman PJ, Burnett J, Gheorghide M, Grinfeld L, Konstam MA, Kostic D et al (2013) Clinical course of patients with hyponatremia and decompensated systolic heart failure and the effect of vasopressin receptor antagonism with tolvaptan. *J Card Fail* 19(6):390–397
55. Schrier RW, Gross P, Gheorghide M, Berl T, Verbalis JG, Czerwiec FS et al (2006) Tolvaptan, a selective oral vasopressin V₂-receptor antagonist, for hyponatremia. *N Engl J Med* 355(20):2099–2112
56. Felker GM, Mentz RJ, Adams KF, Cole RT, Egnaczyk GF, Patel CB et al (2015) Tolvaptan in patients hospitalized with acute heart failure: rationale and design of the TACTICS and the SECRET of CHF Trials. *Circ Heart Fail* 8(5):997–1005
57. Angeli P, Wong F, Watson H, Gines P, Investigators C (2006) Hyponatremia in cirrhosis: results of a patient population survey. *Hepatology* 44(6):1535–1542
58. Gines P, Berl T, Bernardi M, Bichet DG, Hamon G, Jimenez W et al (1998) Hyponatremia in cirrhosis: from pathogenesis to treatment. *Hepatology* 28(3):851–864
59. Gines P, Guevara M (2008) Hyponatremia in cirrhosis: pathogenesis, clinical significance, and management. *Hepatology* 48(3):1002–1010
60. Fukui H (2015) Do vasopressin V₂ receptor antagonists benefit cirrhotics with refractory ascites? *World J Gastroenterol* 21(41):11584–11596
61. Kim WR, Biggins SW, Kremers WK, Wiesner RH, Kamath PS, Benson JT et al (2008) Hyponatremia and mortality among patients on the liver-transplant waiting list. *N Engl J Med* 359(10):1018–1026
62. Guyader D, Patat A, Ellis-Grosse EJ, Orczyk GP (2002) Pharmacodynamic effects of a nonpeptide antidiuretic hormone V₂ antagonist in cirrhotic patients with ascites. *Hepatology* 36(5):1197–1205
63. Gines P, Wong F, Watson H, Terg R, Bruha R, Zarski JP et al (2010) Clinical trial: short-term effects of combination of sata-vaptan, a selective vasopressin V₂ receptor antagonist, and diuretics on ascites in patients with cirrhosis without hyponatraemia—a randomized, double-blind, placebo-controlled study. *Aliment Pharmacol Ther* 31(8):834–845
64. Wong F, Gines P, Watson H, Horsmans Y, Angeli P, Gow P et al (2010) Effects of a selective vasopressin V₂ receptor antagonist, satavaptan, on ascites recurrence after paracentesis in patients with cirrhosis. *J Hepatol* 53(2):283–290
65. Wong F, Watson H, Gerbes A, Vilstrup H, Badalamenti S, Bernardi M et al (2012) Satavaptan for the management of ascites in cirrhosis: efficacy and safety across the spectrum of ascites severity. *Gut* 61(1):108–116
66. Gerbes AL, Gulberg V, Gines P, Decaux G, Gross P, Gandjini H et al (2003) Therapy of hyponatremia in cirrhosis with a vasopressin receptor antagonist: a randomized double-blind multicenter trial. *Gastroenterology* 124(4):933–939
67. Okita K, Kawazoe S, Hasebe C, Kajimura K, Kaneko A, Okada M et al (2014) Dose-finding trial of tolvaptan in liver cirrhosis patients with hepatic edema: a randomized, double-blind, placebo-controlled trial. *Hepatol Res* 44(1):83–91
68. Sakaida I, Kawazoe S, Kajimura K, Saito T, Okuse C, Takaguchi K et al (2014) Tolvaptan for improvement of hepatic edema: a phase 3, multicenter, randomized, double-blind, placebo-controlled trial. *Hepatol Res* 44(1):73–82

69. Watkins PB, Lewis JH, Kaplowitz N, Alpers DH, Blais JD, Smotzer DM et al (2015) Clinical pattern of tolvaptan-associated liver injury in subjects with autosomal dominant polycystic kidney disease: analysis of clinical trials database. *Drug Saf* 38(11):1103–1113
70. Kovcsdy CP, Lott EH, Lu JL, Malakauskas SM, Ma JZ, Molnar MZ et al (2012) Hyponatremia, hypernatremia, and mortality in patients with chronic kidney disease with and without congestive heart failure. *Circulation* 125(5):677–684
71. Dimitriadis C, Sekercioglu N, Pipili C, Oreopoulos D, Bargman JM (2014) Hyponatremia in peritoneal dialysis: epidemiology in a single center and correlation with clinical and biochemical parameters. *Perit Dial Int* 34(3):260–270
72. Khan S, Floris M, Pani A, Rosner MH (2016) Sodium and volume disorders in advanced chronic kidney disease. *Adv Chronic Kidney Dis* 23(4):240–246
73. Negro A, Regolisti G, Perazzoli F, Davoli S, Sani C, Rossi E (1998) Ifosfamide-induced renal Fanconi syndrome with associated nephrogenic diabetes insipidus in an adult patient. *Nephrol Dial Transplant* 13(6):1547–1549
74. Combs S, Berl T (2014) Dysnatremias in patients with kidney disease. *Am J Kidney Dis* 63(2):294–303
75. Waikar SS, Curhan GC, Brunelli SM (2011) Mortality associated with low serum sodium concentration in maintenance hemodialysis. *Am J Med* 124(1):77–84
76. Nigwekar SU, Wenger J, Thadhani R, Bhan I (2013) Hyponatremia, mineral metabolism, and mortality in incident maintenance hemodialysis patients: a cohort study. *Am J Kidney Dis* 62(4):755–762
77. Hecking M, Karaboyas A, Saran R, Sen A, Inaba M, Rayner H et al (2012) Dialysate sodium concentration and the association with interdialytic weight gain, hospitalization, and mortality. *Clin J Am Soc Nephrol* 7(1):92–100
78. Tanaka A, Katsuno T, Ozaki T, Sakata F, Kato N, Suzuki Y et al (2015) The efficacy of tolvaptan as a diuretic for chronic kidney disease patients. *Acta Cardiol* 70(2):217–223
79. Dangoisse C, Dickie H, Tovey L, Ostermann M (2014) Correction of hyper- and hyponatraemia during continuous renal replacement therapy. *Nephron Clin Pract* 128(3–4):394–398
80. Wendland EM, Kaplan AA (2012) A proposed approach to the dialysis prescription in severely hyponatremic patients with end-stage renal disease. *Semin Dial* 25(1):82–85
81. Ball SG, Iqbal Z (2016) Diagnosis and treatment of hyponatraemia. *Best Pract Res Clin Endocrinol Metab* 30(2):161–173
82. Gisby M, Lundberg J, Landin M, O'Reilly K, Robinson P, Sobocki P et al (2016) The burden of illness in patients with hyponatraemia in Sweden: a population-based registry study. *Int J Clin Pract* 70(4):319–329
83. Rodrigues B, Staff I, Fortunato G, McCullough LD (2014) Hyponatremia in the prognosis of acute ischemic stroke. *J Stroke Cerebrovasc Dis* 23(5):850–854
84. Hiew FL, Winer JB, Rajabally YA (2016) Hyponatraemia in Guillain–Barre syndrome revisited. *Acta Neurol Scand* 133(4):295–301
85. Holland-Bill L, Christiansen CF, Heide-Jorgensen U, Ulrichsen SP, Ring T, Jorgensen JO et al (2015) Hyponatremia and mortality risk: a Danish cohort study of 279 508 acutely hospitalized patients. *Eur J Endocrinol* 173(1):71–81
86. Sherlock M, O'Sullivan E, Agha A, Behan LA, Owens D, Finucane F et al (1002) Incidence and pathophysiology of severe hyponatraemia in neurosurgical patients. *Postgrad Med J* 2009(85):171–175
87. Mapa B, Taylor BE, Appelboom G, Bruce EM, Claassen J, Connolly ES Jr (2016) Impact of hyponatremia on morbidity, mortality, and complications after aneurysmal subarachnoid hemorrhage: a systematic review. *World Neurosurg* 85:305–314
88. Kelly DF, Laws ER Jr, Fossett D (1995) Delayed hyponatremia after transphenoidal surgery for pituitary adenoma. Report of nine cases. *J Neurosurg* 83(2):363–367
89. Kristof RA, Rother M, Neuloh G, Klingmuller D (2009) Incidence, clinical manifestations, and course of water and electrolyte metabolism disturbances following transphenoidal pituitary adenoma surgery: a prospective observational study. *J Neurosurg* 111(3):555–562
90. Olson BR, Rubino D, Gumowski J, Oldfield EH (1995) Isolated hyponatremia after transphenoidal pituitary surgery. *J Clin Endocrinol Metab* 80(1):85–91
91. Bales J, Cho S, Tran TK, Korab GA, Khandelwal N, Spiekerman CF et al (2016) The effect of hyponatremia and sodium variability on outcomes in adults with aneurysmal subarachnoid hemorrhage. *World Neurosurg* 96:340–349
92. Williams C, Simon TD, Riva-Cambrin J, Bratton SL (2012) Hyponatremia with intracranial malignant tumor resection in children. *J Neurosurg Pediatr* 9(5):524–529
93. Kao L, Al-Lawati Z, Vavao J, Steinberg GK, Katznelson L (2009) Prevalence and clinical demographics of cerebral salt wasting in patients with aneurysmal subarachnoid hemorrhage. *Pituitary* 12(4):347–351
94. Palmer BF (2003) Hyponatremia in patients with central nervous system disease: SIADH versus CSW. *Trends Endocrinol Metab* 14(4):182–187
95. Loh JA, Verbalis JG (2007) Diabetes insipidus as a complication after pituitary surgery. *Nat Clin Pract Endocrinol Metab* 3(6):489–494
96. Zhang L, Fu P, Wang L, Cai G, Zhang L, Chen D et al (2013) Hyponatraemia in patients with crush syndrome during the Wenchuan earthquake. *Emerg Med J* 30(9):745–748
97. Roquilly A, Mahe PJ, Seguin P, Guitton C, Floch H, Tellier AC et al (2011) Hydrocortisone therapy for patients with multiple trauma: the randomized controlled HYPOLYTE study. *JAMA* 305(12):1201–1209
98. Aqil A, Hossain F, Sheikh H, Aderinto J, Whitwell G, Kapoor H (2016) Achieving hip fracture surgery within 36 hours: an investigation of risk factors to surgical delay and recommendations for practice. *J Orthop Traumatol* 17(3):207–213
99. Sharif S, Dominguez M, Imbriano L, Mattana J, Maesaka JK (2015) Recognition of hyponatremia as a risk factor for hip fractures in older persons. *J Am Geriatr Soc* 63(9):1962–1964
100. Murthy K, Koshkina O, Marcantonio AJ, Pala N, Breeze JL, Paulus J et al (2015) Hyponatremia and fracture risk: a hospital-based case-control study. *J Am Geriatr Soc* 63(8):1699–1701
101. Usala RL, Fernandez SJ, Mete M, Cowen L, Shara NM, Barsony J et al (2015) hyponatremia is associated with increased osteoporosis and bone fractures in a large US health system population. *J Clin Endocrinol Metab* 100(8):3021–3031
102. Hawkins RC (2003) Age and gender as risk factors for hyponatremia and hypernatremia. *Clin Chim Acta* 337(1–2):169–172
103. Tasdemir V, Oguz AK, Sayin I, Ergun I (2015) Hyponatremia in the outpatient setting: clinical characteristics, risk factors, and outcome. *Int Urol Nephrol* 47(12):1977–1983
104. Liamis G, Rodenburg EM, Hofman A, Zietse R, Stricker BH, Hoorn EJ (2013) Electrolyte disorders in community subjects: prevalence and risk factors. *Am J Med* 126(3):256–263
105. Gankam-Kengne F, Ayers C, Khera A, de Lemos J, Maalouf NM (2013) Mild hyponatremia is associated with an increased risk of death in an ambulatory setting. *Kidney Int* 83(4):700–706
106. Liamis G, Filippatos TD, Elisaf MS (2016) Thiazide-associated hyponatremia in the elderly: what the clinician needs to know. *J Geriatr Cardiol* 13(2):175–182

107. Sajadieh A, Binici Z, Mouridsen MR, Nielsen OW, Hansen JF, Haugaard SB (2009) Mild hyponatremia carries a poor prognosis in community subjects. *Am J Med* 122(7):679–686
108. Hoorn EJ, Rivadeneira F, van Meurs JB, Ziere G, Stricker BH, Hofman A et al (2011) Mild hyponatremia as a risk factor for fractures: the Rotterdam study. *J Bone Miner Res* 26(8):1822–1828
109. Verbalis JG, Adler S, Schrier RW, Berl T, Zhao Q, Czerwiec FS et al (2011) Efficacy and safety of oral tolvaptan therapy in patients with the syndrome of inappropriate antidiuretic hormone secretion. *Eur J Endocrinol* 164(5):725–732
110. De las Penas R, Ponce S, Henao F, Camps Herrero C, Carcereny E, Escobar Alvarez Y et al (2016) SIADH-related hyponatremia in hospital day care units: clinical experience and management with tolvaptan. *Support Care Cancer*. 24(1):499–507
111. Almond CS, Shin AY, Fortescue EB, Mannix RC, Wypij D, Bin-stadt BA et al (2005) Hyponatremia among runners in the Boston Marathon. *N Engl J Med* 352(15):1550–1556
112. Ben-Abraham R, Szold O, Rudick V, Weinbroum AA (2003) ‘Ecstasy’ intoxication: life-threatening manifestations and resuscitative measures in the intensive care setting. *Eur J Emerg Med* 10(4):309–313
113. Arieff AI (1984) Central nervous system manifestations of disordered sodium metabolism. *Clin Endocrinol Metab* 13(2):269–294
114. Berl T (1990) Treating hyponatremia: what is all the controversy about? *Ann Intern Med* 113(6):417–419
115. Edelman IS, Leibman J, O’Meara MP, Birkenfeld LW (1958) Interrelations between serum sodium concentration, serum osmolarity and total exchangeable sodium, total exchangeable potassium and total body water. *J Clin Invest* 37(9):1236–1256
116. Sterns RH, Hix JK, Silver S (2010) Treating profound hyponatremia: a strategy for controlled correction. *Am J Kidney Dis* 56(4):774–779
117. Hanna RM, Yang WT, Lopez EA, Riad JN, Wilson J (2016) The utility and accuracy of four equations in predicting sodium levels in dysnatremic patients. *Clin Kidney J* 9(4):530–539
118. Trepiccione F, Capasso G, Lippi G (2014) Serum and urine osmolality: clinical and laboratory features. *G Ital Nefrol* 31(5)
119. Corona G, Giuliani C, Verbalis JG, Forti G, Maggi M, Peri A (2015) Hyponatremia improvement is associated with a reduced risk of mortality: evidence from a meta-analysis. *PLoS One* 10(4):e0124105
120. Ellison DH, Berl T (2007) Clinical practice. The syndrome of inappropriate antidiuresis. *N Engl J Med* 356(20):2064–2072
121. Grohe C, Berardi R, Burst V (2015) Hyponatraemia–SIADH in lung cancer diagnostic and treatment algorithms. *Crit Rev Oncol Hematol* 96(1):1–8
122. Onitilo AA, Kio E, Doi SA (2007) Tumor-related hyponatremia. *Clin Med Res* 5(4):228–237
123. Berardi R, Rinaldi S, Caramanti M, Grohe C, Santoni M, Morgese F et al (2016) Hyponatremia in cancer patients: time for a new approach. *Crit Rev Oncol Hematol* 102:15–25
124. Berardi R, Santoni M, Rinaldi S, Nunzi E, Smerilli A, Caramanti M et al (2016) Risk of hyponatraemia in cancer patients treated with targeted therapies: a systematic review and meta-analysis of clinical trials. *PLoS One* 11(5):e0152079
125. Illouz F, Braun D, Briet C, Schweizer U, Rodien P (2014) Endocrine side-effects of anti-cancer drugs: thyroid effects of tyrosine kinase inhibitors. *Eur J Endocrinol* 171(3):R91–R99
126. Lu X, Wang X (2017) Hyponatremia induced by antiepileptic drugs in patients with epilepsy. *Expert Opin Drug Saf* 16(1):77–87
127. Tsapepas D, Chiles M, Babayev R, Rao MK, Jaitly M, Salerno D et al (2016) Incidence of hyponatremia with high-dose trimethoprim-sulfamethoxazole exposure. *Am J Med* 129(12):1322–1328
128. Tanaka R, Suzuki Y, Takumi Y, Iwao M, Sato Y, Hashinaga K et al (2016) A retrospective analysis of risk factors for linezolid-associated hyponatremia in Japanese patients. *Biol Pharm Bull* 39(12):1968–1973
129. Spital A (1999) Diuretic-induced hyponatremia. *Am J Nephrol* 19(4):447–452
130. Gandhi S, Shariff SZ, Al-Jaishi A, Reiss JP, Mamdani MM, Hackam DG et al (2017) Second-generation antidepressants and hyponatremia risk: a population-based cohort study of older adults. *Am J Kidney Dis* 69(1):87–96
131. Buon M, Gaillard C, Martin J, Fedrizzi S, Mosquet B, Coquerel A et al (2013) Risk of proton pump inhibitor-induced mild hyponatremia in older adults. *J Am Geriatr Soc* 61(11):2052–2054
132. Izzedine H, Fardet L, Launay-Vacher V, Dorent R, Petitclerc T, Deray G (2002) Angiotensin-converting enzyme inhibitor-induced syndrome of inappropriate secretion of antidiuretic hormone: case report and review of the literature. *Clin Pharmacol Ther* 71(6):503–507
133. Yamada H, Asano T, Aoki A, Ikoma A, Yoshida M, Kusaka I et al (2014) Combination therapy of angiotensin II receptor blocker and thiazide produces severe hyponatremia in elderly hypertensive subjects. *Intern Med* 53(7):749–752

Affiliations

E. Sbardella¹ · A. M. Isidori¹ · G. Arnaldi² · M. Arosio³ · C. Barone⁴ · A. Benso⁵ · R. Berardi⁶ · G. Capasso⁷ · M. Caprio^{8,9} · F. Ceccato¹⁰ · G. Corona¹¹ · S. Della Casa¹² · L. De Nicola¹³ · M. Faustini-Fustini¹⁴ · E. Fiaccadori¹⁵ · L. Gesualdo¹⁶ · S. Gori¹⁷ · A. Lania¹⁸ · G. Mantovani³ · P. Menè¹⁹ · G. Parenti²⁰ · C. Pinto²¹ · R. Pivonello²² · P. Razzore²³ · G. Regolisti¹⁵ · C. Scaroni¹⁰ · F. Trepiccione⁷ · A. Lenzi¹ · A. Peri²⁴ · On behalf of the: Fluid and Electrolyte Disorders Club of the Italian Society of Endocrinology; Italian Society of Nephrology; and Italian Association of Medical Oncology

- ¹ Department of Experimental Medicine, Sapienza University of Rome, Rome, Italy
- ² Clinica di Endocrinologia e Malattie del Metabolismo, Università Politecnica delle Marche Azienda Ospedaliero-Universitaria, Ospedali Riuniti Umberto I-GM Lancisi-G Salesi, Ancona, Italy
- ³ Endocrinology and Diabetology Unit, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Department of Clinical Sciences and Community Health, University of Milan, Milan, Italy
- ⁴ UOC di Oncologia Medica, Università Cattolica del Sacro Cuore, Rome, Italy
- ⁵ Division of Endocrinology, Diabetes and Metabolism, Department of Medical Sciences, University of Turin, Turin, Italy
- ⁶ Clinica Oncologica, Università Politecnica delle Marche Azienda Ospedaliero-Universitaria; Ospedali Riuniti Umberto I-GM Lancisi-G Salesi, Ancona, Italy
- ⁷ Dipartimento di Scienze Cardio-Toraciche e Respiratorie, Università della Campania "Luigi Vanvitelli", Caserta, Italy
- ⁸ Laboratory of Cardiovascular Endocrinology, IRCCS San Raffaele Pisana, Rome, Italy
- ⁹ Department of Human Sciences and Promotion of the Quality of Life, San Raffaele Roma Open University, Rome, Italy
- ¹⁰ Endocrinology Unit, Department of Medicine DIMED, University-Hospital of Padova, Padua, Italy
- ¹¹ Endocrinology Unit, Medical Department, Azienda UsI Bologna Maggiore-Bellaria Hospital, Bologna, Italy
- ¹² Endocrinology and Metabolic Diseases Unit, Catholic University of the Sacred Heart, Rome, Italy
- ¹³ Nephrology, Medical School, University of Campania Luigi Vanvitelli, Naples, Italy
- ¹⁴ Pituitary Unit, IRCCS Institute of Neurological Sciences, Bellaria Hospital, Bologna, Italy
- ¹⁵ Renal Unit, Parma University Medical School, Parma, Italy
- ¹⁶ Nephrology Dialysis and Transplantation, Bari University Medical School, Bari, Italy
- ¹⁷ UOC Oncologia Medica, Ospedale Sacro Cuore Don Calabria, Negrar, Verona, Italy
- ¹⁸ Endocrine Unit, Department of Biomedical Sciences, Humanitas Research Hospital, Humanitas University, Rozzano (MI), Italy
- ¹⁹ Nephrology, Sapienza University of Rome, Rome, Italy
- ²⁰ Endocrine Unit, Careggi Hospital, Florence, Italy
- ²¹ Oncologia Medica IRCCS Arcispedale S. Maria Nuova, Reggio Emilia, Italy
- ²² Dipartimento di Medicina Clinica e Chirurgia, Sezione di Endocrinologia, Università "Federico II" di Napoli, Naples, Italy
- ²³ Endocrine Unit, AO Ordine Mauriziano, Turin, Italy
- ²⁴ Endocrine Unit, Department of Experimental and Clinical Biomedical Sciences "Mario Serio", AOU Careggi, University of Florence, Viale Pieraccini, 6, 50139 Florence, Italy