

RESEARCH



# Conic stability of polynomials

Thorsten Jörgens and Thorsten Theobald\* 

\*Correspondence:  
theobald@math.uni-frankfurt.de  
FB 12 – Institut für Mathematik,  
Goethe-Universität, Postfach 11  
19 32, 60054 Frankfurt am Main,  
Germany

## Abstract

We introduce and study the notion of conic stability of multivariate complex polynomials in  $\mathbb{C}[\mathbf{z}]$ , which naturally generalizes the stability of multivariate polynomials. In particular, we generalize Borcea’s and Brändén’s multivariate version of the Hermite–Kakeya–Obreschkoff Theorem to the conic stability and provide a characterization in terms of a directional Wronskian. And we generalize a major criterion for stability of determinantal polynomials to stability with respect to the positive semidefinite cone.

## 1 Introduction

Stable polynomials have a rich history (see, e.g., [22]) and attracted a lot of interest in recent years. Prominent research directions include the generalization of classical results on univariate stable polynomials to multivariate stable polynomials (see, e.g., [3, 4, 6, 16, 26]) as well as applications of stable polynomials to various areas of mathematics and theoretical computer science, see [1, 5, 10, 19–21, 24] and the references therein. A polynomial  $f = f(\mathbf{z}) = f(z_1, \dots, z_n) \in \mathbb{C}[\mathbf{z}] = \mathbb{C}[z_1, \dots, z_n]$  is called *stable* if every root  $\mathbf{z}$  satisfies  $\text{Im}(z_j) \leq 0$  for some  $j$ . A stable polynomial  $f$  with real coefficients is called *real stable*.

In [15], the authors and de Wolff introduced a geometric approach to stability phenomena introducing the imaginary projection of a polynomial as the set

$$\mathcal{I}(f) = \{\text{Im}(\mathbf{z}) = (\text{Im}(z_1), \dots, \text{Im}(z_n)) : f(\mathbf{z}) = 0\}, \quad (1.1)$$

where  $\text{Im}(\cdot)$  denotes the imaginary part of a complex number. Using this notion, stability of  $f$  is equivalent to  $\mathcal{I}(f) \cap (\mathbb{R}_{>0})^n = \emptyset$ .

This geometric view upon stability of polynomials naturally suggests to extend the results of the (usual) stability notion to more general real cones. In this article, given a cone  $K \subset \mathbb{R}^n$ , a polynomial  $f$  is called *K-stable* if  $\mathcal{I}(f) \cap \text{int } K = \emptyset$ , where  $\text{int } K$  denotes the interior of  $K$ . Note that  $(\mathbb{R}_{\geq 0})^n$ -stability coincides with the usual stability. And note that setting  $\Omega = \mathbb{R}^n + iK$ , our  $K$ -stability also falls into the more general class of stability notions which forbid zeroes in an arbitrarily given complex set  $\Omega \subset \mathbb{C}^n$ —however, as pointed out in [26, p. 81], little can be said on a class of that generality. For polynomials with matrix variables, we consider the special case where  $K = \mathcal{S}_n^+$  is the cone of positive semidefinite matrices.

In the paper, we initiate to develop a theory of  $K$ -stability of multivariate polynomials. To begin with, we extend the well-known characterization of stable polynomials in terms

of hyperbolic polynomials to the conic case, see Lemma 3.4. Our main contribution is the generalization of three core results on multivariate stable polynomials to the conic stability. Firstly, we show that the classical Theorem of Hermite–Takeya–Obreschkoff, which has been generalized from the univariate to the multivariate case by Borcea and Brändén [6, Theorem 6.3.8], can be further generalized to  $K$ -stability for multivariate polynomials; see Theorem 4.3. Secondly, we characterize conic stability with respect to polyhedral cones and non-polyhedral cones in terms of a directional Wronskian; see Theorem 4.4. Thirdly, we show that Borcea’s and Brändén’s prominent criterion for stability of determinantal polynomials in [3, Theorem 2.4] can be generalized to stability with respect to the positive semidefinite cone  $\mathcal{S}_n^+$ ; see Theorem 5.3.

Our statements and their proofs apply conic duality, and the generalization of the stability criterion for determinantal polynomials is given in terms of the Khatri–Rao product of matrices.

While our work was mainly motivated by the intrinsic relevance and the structure of stable polynomials, we note that the case  $K = \mathcal{S}_n^+$  is naturally related to the Siegel upper half-spaces in the theory of modular forms, see Sect. 2.3.

Beside the actual statements themselves, we think that these extensions pinpoint that the conic stability offers a very natural generalized framework for studying stability issues of multivariate polynomials.

The paper is structured as follows. In Sect. 2, we collect some known statements and the connection to Siegel upper half-spaces in the theory of modular forms. In Sect. 3, we provide some basic results on  $K$ -stable polynomials. Then, in Sect. 4, we generalize the multivariate Hermite–Takeya–Obreschkoff Theorem to the conic setting and study conic stability by means of the directional Wronskian. Section 5 contains the generalization of the characterization of stability for determinantal polynomials.

## 2 Preliminaries

Throughout the text, bold letters denote  $n$ -dimensional vectors unless noted otherwise.

### 2.1 Stability theorems

As general references on stable polynomials, we refer to [6, 22, 26]. Note that in our definition of stability, the zero polynomial is not stable, in consistency with the convention in [6].

For univariate, real stable polynomials  $f, g \in \mathbb{R}[z]$ , let  $W(f, g) = f'g - g'f$  denote the *Wronskian* of  $f$  and  $g$  and write  $f \ll g$  if  $W(f, g) \leq 0$  on  $\mathbb{R}$ . Note that univariate, real stable polynomials are real-rooted. In the context of univariate stable polynomials, the following concept of interlacing roots naturally appears.

**Definition 2.1** Let  $f, g \in \mathbb{R}[z]$  be two univariate, real-rooted polynomials with roots  $\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_{\deg f}$  and  $\beta_1 \leq \beta_2 \leq \dots \leq \beta_{\deg g}$ . We say that  $f$  and  $g$  *interlace* if their roots alternate, i.e.,  $\alpha_1 \leq \beta_1 \leq \alpha_2 \leq \beta_2 \leq \dots$  or  $\beta_1 \leq \alpha_1 \leq \beta_2 \leq \alpha_2 \leq \dots$ . If all inequalities are strict,  $f$  and  $g$  *interlace strictly*.

We say that  $f$  *interlaces  $g$  properly* (or:  $f$  is a *proper interlacing* of  $g$ ), if

- $\beta_{\deg g} \geq \alpha_{\deg f} \geq \beta_{\deg g - 1} \geq \alpha_{\deg f - 1} \geq \dots$ , when the leading coefficients of  $f$  and  $g$  have the same sign,

- $\alpha_{\deg f} \geq \beta_{\deg g} \geq \alpha_{\deg f-1} \geq \beta_{\deg g-1} \geq \dots$ , when the leading coefficients of  $f$  and  $g$  have different signs.

For interlacing polynomials  $f$  and  $g$ , the degrees of  $f$  and  $g$  can only differ by at most 1. We collect two classical theorems on univariate stable polynomials (see [22, 26]).

**Proposition 2.2** (Hermite–Biehler) *For  $f, g \in \mathbb{R}[z] \setminus \{0\}$ , the following are equivalent:*

- (1)  $g + if$  is stable.
- (2)  $f, g$  are real stable and  $f \ll g$ .
- (3)  $f, g$  are real stable and  $f$  interlaces  $g$  properly.

Extending the definition of  $\ll$  and of interlacing to arbitrary  $f, g \in \mathbb{R}[x]$  by requiring real stability of  $f$  and  $g$ , condition (2) can be written shortly as  $f \ll g$  and (3) can be written shortly as:  $f$  interlaces  $g$  properly.

**Proposition 2.3** (Hermite–Kakeya–Obreschkoff (HKO, for short)) *Let  $f, g \in \mathbb{R}[z]$ . Then  $\lambda f + \mu g$  is stable or the zero polynomial for all  $\lambda, \mu \in \mathbb{R}$  if and only if  $f$  and  $g$  interlace or  $f \equiv g \equiv 0$ .*

Moreover, the following theorem will be useful.

**Proposition 2.4** (Hurwitz, see Theorem 1.3.8 in [22]) *Let  $\{f_j\}_{j \in \mathbb{N}} \subset \mathbb{C}[z]$  be a sequence of polynomials non-vanishing in a connected open set  $U \subset \mathbb{C}^n$ , and assume it converges to a function  $f$  uniformly on compact subsets of  $U$ . Then,  $f$  is either non-vanishing on  $U$  or it is identically 0.*

For multivariate polynomials  $f, g \in \mathbb{R}[\mathbf{z}]$ , one writes  $f \ll g$  if  $g + if$  is stable (see, e.g., [6, 26]; note that this makes the multivariate Hermite–Biehler statement a definition rather than a theorem). The multivariate version of the HKO Theorem then has the same format as the univariate version. The multivariate theorem was shown in [6, Theorem 1.6], see also [4, Theorem 2.9], [26, Theorem 2.9].

**Proposition 2.5** (Multivariate HKO of Borcea and Brändén) *Let  $f, g \in \mathbb{R}[\mathbf{z}]$ . Then,  $\lambda f + \mu g$  is stable or the zero polynomial for all  $\lambda, \mu \in \mathbb{R}$  if and only if  $f \ll g$  or  $g \ll f$  or  $f \equiv g \equiv 0$ .*

An important class of stable polynomials comes from determinantal representations ([3, Theorem 2.4], see also [7, 11, 17]).

**Proposition 2.6** (Borcea, Brändén) *Let  $A_1, \dots, A_n$  be positive semidefinite  $d \times d$ -matrices and  $B$  be a Hermitian  $d \times d$ -matrix, then*

$$f(\mathbf{x}) = \det \left( \sum_{j=1}^n x_j A_j + B \right)$$

*is real stable or the zero polynomial.*

Determinantal representations of this kind are relevant in the context of the Lax conjecture (proven by Lewis et al. [17], see [6, Corollary 6.7] for a formulation on stable polynomials) as well as its variations and generalizations (see, for example, [3, Section 5]).

### 2.2 Imaginary projections and hyperbolicity cones

Set  $A^c = \mathbb{R}^n \setminus A$  for the complement of a set  $A \subseteq \mathbb{R}^n$ , and write  $\text{cl}A$  for its closure. For a polynomial  $f \in \mathbb{C}[\mathbf{z}]$ , the complement  $(\text{cl}\mathcal{I}(f))^c$  of the imaginary projection (1.1) consists of finitely many convex components [15]. In the special case of a non-constant polynomial  $f = \sum_{i=1}^n a_i z_i + a_0$  with real coefficients  $a_1, \dots, a_n \in \mathbb{R}$  and  $a_0 \in \mathbb{C}$ , we have  $\mathcal{I}(f) = \{\mathbf{y} \in \mathbb{R}^n : \sum_{i=1}^n a_i y_i + \text{Im}(a_0) = 0\}$ .

A homogeneous polynomial  $f \in \mathbb{C}[\mathbf{z}]$  is called *hyperbolic* in a real direction  $\mathbf{e} \in \mathbb{R}^n$  if  $f(\mathbf{e}) \neq 0$  and for every  $\mathbf{x} \in \mathbb{R}^n$  the function  $t \mapsto f(\mathbf{x} + t\mathbf{e})$  has only real roots. Denote by  $C(\mathbf{e}) = \{\mathbf{x} \in \mathbb{R}^n : f(\mathbf{x} + t\mathbf{e}) = 0 \Rightarrow t < 0\}$  the *hyperbolicity cone* of  $f$  with respect to  $\mathbf{e}$ . Then,  $C(\mathbf{e})$  is convex,  $f$  is hyperbolic with respect to every point  $\mathbf{e}'$  in its hyperbolicity cone, and  $C(\mathbf{e}) = C(\mathbf{e}')$  (see [9]).

For a homogeneous polynomial  $f \in \mathbb{C}[\mathbf{z}]$ , the hyperbolicity cones of  $f$  coincide with the components of  $\mathcal{I}(f)^c$  [14].

### 2.3 The Siegel upper half-space

The *Siegel upper half-space* (or *Siegel upper half-plane*)  $\mathcal{H}_g$  of degree  $g$  (or genus  $g$ ) is defined as

$$\mathcal{H}_g = \{A \in \mathbb{C}^{g \times g} \text{ symmetric} : \text{Im}(A) \text{ is positive definite}\},$$

where  $\text{Im}(A) = (\text{Im}(a_{ij}))_{g \times g}$  ([23], see also, e.g., [25, Section 2]). The Siegel upper half-space constitutes the domain on which the Siegel theta functions are defined. It can be used to parameterize polarized varieties (see also, for example, [13, Vol. 1, Section 3.I] and for the use in elliptic curve cryptography [8, Section 5.1]).

### 3 K-stability and psd-stability

Let  $K$  be a *proper* cone in  $\mathbb{R}^n$ , that is, a full-dimensional, closed and pointed convex cone in  $\mathbb{R}^n$ . We consider the following generalization of stability. Let  $\mathcal{S}_n$  be the set of real symmetric  $n \times n$ -matrices, and let  $\mathcal{S}_n^+$  and  $\mathcal{S}_n^{++}$  denote its subsets of positive semidefinite and positive definite matrices.

**Definition 3.1** A polynomial  $f \in \mathbb{C}[\mathbf{z}]$  is called *K-stable*, if  $f(\mathbf{z}) \neq 0$  whenever  $\text{Im}(\mathbf{z}) \in \text{int}K$ .

If  $f \in \mathbb{C}[Z]$  on the symmetric matrix variables  $Z = (z_{ij})_{n \times n}$  is  $\mathcal{S}_n^+$ -stable, then  $f$  is called *positive semidefinite-stable* (for short, *psd-stable*).

Equivalently, a polynomial  $f \in \mathbb{C}[Z]$  on the symmetric matrix variables  $Z = (z_{ij})_{n \times n}$  is psd-stable if there does not exist a matrix  $Z$  in the Siegel upper half-space  $\mathcal{H}_n$  with  $f(Z) = 0$ . Note that psd-stability generalizes the usual stability in the sense that a polynomial  $f(z_1, \dots, z_n)$  is stable if and only if  $f(\text{diag}(z_1, \dots, z_n))$  is psd-stable.

*Example 3.2* (i) Let  $f \in \mathbb{R}[\mathbf{z}]$  be given by  $f(\mathbf{z}) = \mathbf{a}^T \mathbf{z} + b$ , where  $\mathbf{a}$  is a real  $n$ -dimensional vector and  $b \in \mathbb{R}$ . Then,  $f$  is  $K$ -stable if and only if  $\mathbf{a} \in \text{int}K^*$  or  $-\mathbf{a} \in \text{int}K^*$ , where  $K^* = \{\mathbf{y} \in \mathbb{R}^n : \langle \mathbf{x}, \mathbf{y} \rangle \geq 0 \text{ for all } \mathbf{x} \in K\}$  denotes the dual cone of  $K$  and  $\langle \cdot, \cdot \rangle$  is the Euclidean dot product on  $\mathbb{R}^n$ .

Namely, if  $\mathbf{a} \in \text{int}K^*$  or  $-\mathbf{a} \in \text{int}K^*$ , say,  $\mathbf{a} \in \text{int}K^*$ , then for any  $\mathbf{z} = \mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int}K$  we have

$$f(\mathbf{z}) = \langle \mathbf{a}, \mathbf{x} \rangle + i\langle \mathbf{a}, \mathbf{y} \rangle + b,$$

because  $\mathbf{a}$  is real. Hence,  $\text{Im} f(\mathbf{z}) \neq 0$ , and thus  $f$  is  $K$ -stable.

Conversely, let  $f$  be  $K$ -stable. Assuming  $\mathbf{a} \notin \pm \text{int} K^*$ , there exists  $\mathbf{y}' \in \text{int} K$  with  $\langle \mathbf{a}, \mathbf{y}' \rangle \leq 0$  and  $\mathbf{y}'' \in \text{int} K$  with  $\langle \mathbf{a}, \mathbf{y}'' \rangle \geq 0$ . Hence, there exists some  $\mathbf{y} \in \text{int} K$  with  $\langle \mathbf{a}, \mathbf{y} \rangle = 0$ . Choosing  $\mathbf{x} \in \mathbb{R}^n$  with  $\langle \mathbf{a}, \mathbf{x} \rangle + b = 0$  gives a contradiction to the stability of  $f$ .

For usual stability, this implies the well-known statement that  $f = \mathbf{a}^T \mathbf{z} + b$  is stable if and only if  $\mathbf{a} \in (\mathbb{R}_{>0})^n$  or  $-\mathbf{a} \in (\mathbb{R}_{>0})^n$ . For psd-stability, this implies that  $f(Z) = \langle Z, A \rangle + b$  with  $A \in \mathcal{S}_n$  is psd-stable if and only if  $A \succ 0$  or  $-A \succ 0$ ; here, the scalar product is  $\langle Z, A \rangle = \text{tr}(A^H Z) = \text{tr}(AZ)$  and  $A^H$  is the Hermitian transpose of  $A$ .

(ii) As an example for psd-stability, the polynomial  $f(Z) = \det Z$  on the set of symmetric  $n \times n$ -matrices is psd-stable. We postpone the proof to Example 3.7.

*Example 3.3* For the polynomial

$$\begin{aligned} f(z_1, z_2, z_3) &= \det \left( \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} z_1 + \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} z_2 + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} z_3 \right) \\ &= (z_1 + z_3)^2 - z_2^2 \\ &= (z_1 + z_3 - z_2)(z_1 + z_3 + z_2), \end{aligned}$$

Section 2.2 implies that the imaginary projection is

$$\mathcal{I}(f) = \{ \mathbf{y} \in \mathbb{R}^3 : y_1 - y_2 + y_3 = 0 \} \cup \{ \mathbf{y} \in \mathbb{R}^3 : y_1 + y_2 + y_3 = 0 \}.$$

Since, for example,  $(\frac{1}{2}, 1, \frac{1}{2}) \in \mathcal{I}(f) \cap \mathbb{R}_{>0}^3$ ,  $f$  is not stable. In contrast to this, setting  $Z = \begin{pmatrix} z_1 & z_2 \\ z_2 & z_3 \end{pmatrix}$ , the polynomial  $f(Z) = f(z_1, z_2, z_3)$  is psd-stable. Namely, for  $\mathbf{y} \in \mathcal{I}(f)$ , we have

$$\det \begin{pmatrix} y_1 & \pm(y_1 + y_3) \\ \pm(y_1 + y_3) & y_3 \end{pmatrix} = y_1 y_3 - (y_1 + y_3)^2 \leq 0$$

as a consequence of the arithmetic-geometric mean inequality, hence  $\mathbf{y} \notin \text{int} \mathcal{S}_2^+$ .

The following lemma allows to reduce multivariate  $K$ -stability to univariate stable polynomials.

**Lemma 3.4** *A polynomial  $f \in \mathbb{C}[\mathbf{z}] \setminus \{0\}$  is  $K$ -stable if and only if for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  with  $\mathbf{y} \in \text{int} K$  the univariate polynomial  $t \mapsto f(\mathbf{x} + t\mathbf{y})$  is stable.*

*Proof* If  $f$  is not  $K$ -stable, then there exists  $\mathbf{x} \in \mathbb{R}^n$  and  $\mathbf{y} \in \text{int} K$  with  $f(\mathbf{x} + i\mathbf{y}) = 0$ . Hence,  $i$  is a zero of the univariate polynomial  $t \mapsto f(\mathbf{x} + t\mathbf{y})$  and thus that univariate polynomial is not stable.

Conversely, if  $t \mapsto f(\mathbf{x} + t\mathbf{y})$  is not stable for  $\mathbf{y} \in \text{int} K$ , then there is some  $\alpha + i\beta \in \mathbb{C}$  with  $\beta > 0$  and  $0 = f(\mathbf{x} + (\alpha + i\beta)\mathbf{y}) = f(\mathbf{x} + \alpha\mathbf{y} + i\beta\mathbf{y})$ . Since  $\beta\mathbf{y} \in \text{int} K$ ,  $f$  is not  $K$ -stable. □

As reviewed in Sect. 2.2, for a homogeneous polynomial  $f$ , every component in the complement of the imaginary projection  $\mathcal{I}(f)$  is a hyperbolicity cone. In particular,  $f$  is stable if and only if  $f$  is hyperbolic with respect to every point in the positive orthant [9]. This generalizes as follows.

**Theorem 3.5** *Let  $f \in \mathbb{C}[\mathbf{z}]$  be homogeneous. Then, the following are equivalent:*

- (1)  $f$  is  $K$ -stable.
- (2)  $\mathcal{I}(f) \cap \text{int } K = \emptyset$ .
- (3)  $f$  is hyperbolic with respect to every point in  $\text{int } K$ .

*Proof* The equivalence (1)  $\Leftrightarrow$  (2) is clear.

“(3)  $\Rightarrow$  (1)” If  $f$  is not  $K$ -stable, then there exists  $\mathbf{x} \in \mathbb{R}$  and  $\mathbf{e} \in \text{int } K$  with  $f(\mathbf{x} + i\mathbf{e}) = 0$ . Hence,  $i$  is a root of  $t \mapsto f(\mathbf{x} + t\mathbf{e})$ , so that  $f$  is not hyperbolic with respect to  $\mathbf{e}$ .

“(1)  $\Rightarrow$  (3)” Assume  $t \mapsto f(\mathbf{x} + t\mathbf{e})$  is not hyperbolic for  $\mathbf{e} \in \text{int } K$ . In case  $f(\mathbf{e}) = 0$ , the point  $i\mathbf{e}$  is a root of the homogeneous polynomial  $f$  as well, so that  $f$  is not hyperbolic then. Hence,  $f(\mathbf{e}) \neq 0$  and there is  $\mathbf{x} \in \mathbb{R}$  and  $\alpha + i\beta \in \mathbb{C}$  with  $\beta \neq 0$  and  $f(\mathbf{x} + (\alpha + i\beta)\mathbf{e}) = 0$ . We can assume that  $\beta > 0$ , since  $-\mathbf{x} - (\alpha + i\beta)\mathbf{e}$  is a zero of  $f$ , too. Hence,  $0 = f(\mathbf{x} + \alpha\mathbf{e} + i\beta\mathbf{e})$  and  $\beta\mathbf{e} \in \text{int } K$ , so that  $f$  is not  $K$ -stable.  $\square$

The following consequence of the connection between  $K$ -stability and the imaginary projection explains that the convexity assumption in the stability notion is natural.

**Corollary 3.6** *If  $f$  is  $K$ -stable for a non-convex cone  $K$  with non-empty, connected interior, then it is  $\text{cl}(\text{conv}(\text{int}(K)))$ -stable, where  $\text{cl}(\cdot)$  and  $\text{conv}(\cdot)$  denote the closure and the convex hull.*

*Proof* If  $f$  is  $K$ -stable, then  $\mathcal{I}(f) \cap \text{int } K = \emptyset$ , that is,  $\text{int } K \subseteq \mathcal{I}(f)^c$ . Since  $\text{int } K$  is connected, it is contained in one of the connected components of  $(\text{cl } \mathcal{I}(f))^c$ . Denote this component by  $C$ . The convexity of any component in  $(\text{cl } \mathcal{I}(f))^c$  (see Sect. 2.2) implies that for  $K' := \text{cl}(\text{conv}(\text{int}(K)))$ , we have  $\text{int } K' \subseteq \text{conv int } K \subseteq C$ . Since  $C \subseteq (\text{cl } \mathcal{I}(f))^c$ ,  $f$  is  $K'$ -stable.  $\square$

*Example 3.7* We complete Example 3.2 and show that  $f(A) = \det A$  on the space of (complex) symmetric matrices is psd-stable.

Let  $B \in \mathcal{S}_n$  be positive definite and consider the univariate polynomial  $t \mapsto f(A + tB)$ . Its roots are the eigenvalues of the symmetric matrix  $-B^{-1/2}AB^{-1/2}$ , where  $B^{-1/2}$  denotes the unique square root of  $B^{-1}$ . Hence, it is real-rooted. Thus,  $f$  is hyperbolic with respect to any positive definite matrix. By Theorem 3.5,  $f$  is psd-stable.

The following fact generalizes the specialization property of stable polynomials (see, e.g., [26, Lemma 2.4]). We will use it for the special case  $K_1 = \mathbb{R}_{\geq 0}$ .

**Fact 3.8** *Let  $K = K_1 \times K_2 \subset \mathbb{R}^n \times \mathbb{R}^m$  be a cone. If  $f(\mathbf{z}_1, \mathbf{z}_2)$  is  $K$ -stable, then  $f(\mathbf{a} + i\mathbf{b}, \mathbf{z}_2)$  is  $K_2$ -stable for any  $\mathbf{a} \in \mathbb{R}^n$  and  $\mathbf{b} \in \text{int } K_1$ .*

#### 4 A conic generalization of the HKO theorem

We show that the Theorem of Hermite–Takeya–Obreschkoff as given in Proposition 2.3 can be generalized to conic stability.

For  $f, g \in \mathbb{R}[\mathbf{z}]$ , we write  $f \ll_K g$  if  $g + if$  is  $K$ -stable. First, we generalize the auxiliary result in [4, Lemma 2.8] (see also [26, Proposition 2.7]).

**Theorem 4.1** *Let  $f$  and  $g$  be real polynomials in  $\mathbf{z} = (z_1, \dots, z_n)$ . Then,  $g + if$  is  $K$ -stable if and only if  $g + wf \in \mathbb{R}[\mathbf{z}, w]$  is  $K'$ -stable, where  $K' = K \times \mathbb{R}_{\geq 0}$ .*

*Proof* “ $\Leftarrow$ ” This follows from Fact 3.8, setting  $w = i$ .

“ $\Rightarrow$ ” Let  $g + if$  be  $K$ -stable. By Lemma 3.4, the univariate polynomial

$$t \mapsto g(\mathbf{x} + t\mathbf{y}) + if(\mathbf{x} + t\mathbf{y})$$

is stable for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  with  $\mathbf{y} \in \text{int}K$ . For fixed  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  with  $\mathbf{y} \in \text{int}K$ , we write  $\tilde{f}(t) = f(\mathbf{x} + t\mathbf{y})$  and  $\tilde{g}(t) = g(\mathbf{x} + t\mathbf{y})$  as polynomials in  $\mathbb{R}[t]$ . By the univariate Hermite–Biehler Theorem 2.2,  $\tilde{f}$  interlaces  $\tilde{g}$  properly, in particular,  $\tilde{f}$  and  $\tilde{g}$  are real stable. Let  $w = \alpha + i\beta$  with  $\alpha \in \mathbb{R}$  and  $\beta > 0$ . By Lemma 3.4, we have to show that the univariate polynomial

$$t \mapsto \tilde{g} + \alpha\tilde{f} + i\beta\tilde{f} = \tilde{g} + (\alpha + i\beta)\tilde{f} \tag{4.1}$$

is stable. By the univariate Hermite–Kakeya–Obreschkoff Theorem 2.3, the linear combination  $\beta\tilde{f} + \alpha\tilde{g}$  is real stable. Then  $W(\beta\tilde{f}, \tilde{g} + \alpha\tilde{f}) = \beta W(\tilde{f}, \tilde{g}) \leq 0$  on  $\mathbb{R}$ , and thus we can deduce  $\beta\tilde{f} \ll \tilde{g} + \alpha\tilde{f}$ . Invoking again the univariate Hermite–Biehler Theorem 2.2 shows that the univariate polynomial (4.1) is stable. This completes the proof.  $\square$

**Proposition 4.2** *For every  $K$ -stable polynomial  $h = g + if$  with  $g, f \in \mathbb{R}[\mathbf{z}]$  the polynomials  $f$  and  $g$  are  $K$ -stable or identically zero.*

*Proof* By Theorem 4.1, a nonzero polynomial  $g + if$  is  $K$ -stable if and only if  $g + yf$  is  $K'$ -stable with  $K' = K \times \mathbb{R}_{\geq 0}$ . Using Hurwitz’s Theorem 2.4, sending  $y \rightarrow 0$  and  $y \rightarrow \infty$ , respectively, gives that  $g$  and  $f$  are  $K$ -stable polynomials or identically zero.  $\square$

Now, we show the following HKO generalization for  $K$ -stability.

**Theorem 4.3** (Conic HKO Theorem) *Let  $f, g \in \mathbb{R}[\mathbf{z}]$ . Then,  $\lambda f + \mu g$  is either  $K$ -stable or the zero polynomial for all  $\lambda, \mu \in \mathbb{R}$  if and only if  $f + ig$  or  $g + if$  is  $K$ -stable or  $f \equiv g \equiv 0$ .*

*Proof* “ $\Leftarrow$ ” Let  $g + if$  be  $K$ -stable and let  $\lambda, \mu \in \mathbb{R}$  (the case  $f + ig$  can be treated analogously). By Proposition 4.2, we can assume  $\mu \neq 0$ , and hence, by factoring  $\mu$ , it suffices to consider  $g + \lambda f$ .

By Theorem 4.1, the polynomial  $g + yf$  is  $K \times \mathbb{R}_{\geq 0}$ -stable. Using Fact 3.8, we set  $y = \lambda + i$ , which gives the  $K$ -stable polynomial  $(g + \lambda f) + if$ . With Proposition 4.2, the  $K$ -stability of  $g + \lambda f$  follows.

“ $\Rightarrow$ ” Assume that  $\lambda f + \mu g$  is either  $K$ -stable or identically zero for all  $\lambda, \mu \in \mathbb{R}$ . Let  $\mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int}K$ . We write  $\tilde{f}(t) = f(\mathbf{x} + t\mathbf{y})$  and  $\tilde{g}(t) = g(\mathbf{x} + t\mathbf{y})$ . Due to Lemma 3.4, the univariate polynomial  $\lambda\tilde{f} + \mu\tilde{g}$  is stable. The univariate HKO Theorem 2.3 implies that  $\tilde{f}$  and  $\tilde{g}$  interlace.

First, assume that  $\tilde{f}$  interlaces  $\tilde{g}$  properly for all  $\mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int}K$ . By the Hermite–Biehler Theorem 2.2,  $\tilde{g} + i\tilde{f}$  is stable for all  $\mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int}K$ , which implies  $K$ -stability by Lemma 3.4. The case where  $\tilde{g}$  interlaces  $\tilde{f}$  properly for all  $\mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int}K$  is treated analogously.

It remains the case where  $\tilde{f}$  interlaces  $\tilde{g}$  properly for one  $\mathbf{x}_1 + i\mathbf{y}_1 \in \mathbb{C}^n$  with  $\mathbf{y}_1 \in \text{int}K$  and  $\tilde{g}$  interlaces  $\tilde{f}$  properly for another  $\mathbf{x}_2 + i\mathbf{y}_2 \in \mathbb{C}^n$  with  $\mathbf{y}_2 \in \text{int}K$ . For  $0 \leq \tau \leq 1$ , we consider the homotopies

$$\mathbf{x}_\tau = \tau\mathbf{x}_1 + (1 - \tau)\mathbf{x}_2, \quad \mathbf{y}_\tau = \tau\mathbf{y}_1 + (1 - \tau)\mathbf{y}_2.$$

The roots of  $\tilde{f}$  and  $\tilde{g}$  vary continuously with  $\tau$ . Since  $\tilde{f}$  and  $\tilde{g}$  interlace for all  $x + iy \in \mathbb{C}^n$  with  $y \in \text{int}K$ , there must be some  $\tau \in [0, 1]$  such that the roots of  $f(\mathbf{x}_\tau + t\mathbf{y}_\tau)$  and the roots of  $g(\mathbf{x}_\tau + t\mathbf{y}_\tau)$  coincide. Hence, there is a  $c \in \mathbb{R}$  such that  $cf(\mathbf{x}_\tau + t\mathbf{y}_\tau) \equiv g(\mathbf{x}_\tau + t\mathbf{y}_\tau)$ .



Let  $h = cf - g$ . Then,  $h(\mathbf{x}_\tau + t\mathbf{y}_\tau) \equiv 0$ , which implies in particular  $h(\mathbf{x}_\tau + i\mathbf{y}_\tau) = 0$ . Due to the initial hypothesis, the polynomial  $h = cf - g$  is either  $K$ -stable or identically zero. Since the point  $\mathbf{x}_\tau + i\mathbf{y}_\tau \in \mathbb{C}^n$  is a root of the polynomial  $h$  with  $\mathbf{y}_\tau \in \text{int } K$ ,  $h$  must be identically zero. This implies  $cf \equiv g$ . Since by assumption,  $f$  and  $g$  are  $K$ -stable, and since  $K$ -stable polynomials remain  $K$ -stable under multiplication with a complex scalar,  $f + ig$  and  $g + if$  are  $K$ -stable as well or  $f \equiv g \equiv 0$ .  $\square$

For  $f, g \in \mathbb{C}[\mathbf{z}]$ , we denote by  $W_{\mathbf{v}}(f, g) := \partial_{\mathbf{v}}f \cdot g - f \cdot \partial_{\mathbf{v}}g$  the  $\mathbf{v}$ -Wronskian of  $f$  and  $g$ , where  $\partial_{\mathbf{v}}$  denotes the directional derivative with respect to  $\mathbf{v}$ . This allows to give a generalization of [4, Theorem 2.9] (see also [26, Corollary 2.10]) for polyhedral and non-polyhedral cones in terms of the directional  $\mathbf{v}$ -Wronskian.

**Theorem 4.4** *For  $f, g \in \mathbb{R}[\mathbf{z}] \setminus \{0\}$ , the following are equivalent.*

- (1)  $g + if$  is  $K$ -stable.
- (2)  $g + yf$  is  $K \times \mathbb{R}_{\geq 0}$ -stable.
- (3)  $\lambda g + \mu f$  is  $K$ -stable or the zero polynomial for all  $\lambda, \mu \in \mathbb{R}$  and  $W_{\mathbf{v}}(f, g) \leq 0$  on  $\mathbb{R}^n$  for all  $\mathbf{v} \in \text{int } K$ .

If  $K$  is a polyhedral cone  $K = \text{cone}(\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(k)})$ , the statements are also equivalent to

- (4)  $\lambda g + \mu f$  is  $K$ -stable or the zero polynomial for all  $\lambda, \mu \in \mathbb{R}$  and  $W_{\mathbf{v}^{(j)}}(f, g) \leq 0$  on  $\mathbb{R}^n$  for all  $j = 1, \dots, k$ .

*Proof* “(1) $\Leftrightarrow$ (2)” follows by Theorem 4.1.

“(1) $\Rightarrow$ (3)” The first part follows by the conic HKO Theorem 4.3. For the second part, let  $\mathbf{x} + i\mathbf{v} \in \mathbb{C}^n$  with  $\mathbf{v} \in \text{int } K$ . By Lemma 3.4, the univariate restriction

$$t \mapsto g(\mathbf{x} + t\mathbf{v}) + if(\mathbf{x} + t\mathbf{v})$$

is stable. The univariate Hermite–Biehler Theorem 2.2 implies  $f(\mathbf{x} + t\mathbf{v}) \ll g(\mathbf{x} + t\mathbf{v})$ , i.e.,

$$0 \geq W(f(\mathbf{x} + t\mathbf{v}), g(\mathbf{x} + t\mathbf{v})) = g(\mathbf{x} + t\mathbf{v})^2 \frac{d}{dt} \left( \frac{f(\mathbf{x} + t\mathbf{v})}{g(\mathbf{x} + t\mathbf{v})} \right)$$

for all  $t \in \mathbb{R}$ . Now, the claim follows from

$$W_{\mathbf{v}}(f, g)(\mathbf{x}) = W(f(\mathbf{x} + t\mathbf{v}), g(\mathbf{x} + t\mathbf{v}))|_{t=0} \leq 0.$$

“(3) $\Rightarrow$ (1)” By the conic HKO Theorem 4.3,  $f + ig$  or  $g + if$  is  $K$ -stable. And by Lemma 3.4 and the univariate Hermite–Biehler Theorem 2.2, for all  $\mathbf{x} + i\mathbf{v} \in \mathbb{C}^n$  with  $\mathbf{v} \in \text{int } K$ , the univariate real polynomials  $\tilde{f}(t) = f(\mathbf{x} + t\mathbf{v})$  and  $\tilde{g}(t) = g(\mathbf{x} + t\mathbf{v})$  interlace. Moreover, the elementary rule  $\frac{d}{dt}h(\mathbf{x} + t\mathbf{v}) = \frac{\partial}{\partial \mathbf{v}}h(\mathbf{z})|_{\mathbf{z}=\mathbf{x}+t\mathbf{v}}$  gives

$$W(\tilde{f}(t), \tilde{g}(t)) = W_{\mathbf{v}}(f(\mathbf{z}), g(\mathbf{z}))|_{\mathbf{z}=\mathbf{x}+t\mathbf{v}} \leq 0$$

by assumption. Hence, the univariate restrictions

$$t \mapsto g(\mathbf{x} + t\mathbf{v}) + if(\mathbf{x} + t\mathbf{v})$$

are stable, and thus by Lemma 3.4,  $g + if$  is  $K$ -stable.

“(3) $\Rightarrow$ (4)” Since  $K = \text{cone}(\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(k)})$ , this implication follows immediately from a continuity argument.

“(4) $\Rightarrow$ (1)” Let  $\mathbf{x} + i\mathbf{y} \in \mathbb{C}^n$  with  $\mathbf{y} \in \text{int } K$ . We can assume that  $\mathbf{y} \in \text{cone}(\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(n)})$  with linearly independent vectors  $\mathbf{v}^{(1)}, \dots, \mathbf{v}^{(n)}$ . Let  $\lambda_1, \dots, \lambda_n \geq 0$  with  $\mathbf{y} = \sum_{j=1}^n \lambda_j \mathbf{v}^{(j)}$ .



By the precondition,  $f$  and  $g$  are  $K$ -stable, and, by Theorem 4.3,  $g + if$  or  $f + ig$  is  $K$ -stable. By Lemma 3.4, the univariate restriction to  $t \mapsto \mathbf{x} + t\mathbf{y}$  is stable. Its Wronskian fulfils

$$W(f(\mathbf{x} + \mathbf{y}t), g(\mathbf{x} + \mathbf{y}t)) = g(\mathbf{x} + t\mathbf{v})^2 \frac{d}{dt} \left( \frac{f(\mathbf{x} + \mathbf{y}t)}{g(\mathbf{x} + \mathbf{y}t)} \right).$$

Expressing this via  $\frac{d}{dt} h(\mathbf{x} + \mathbf{y}t) = \sum_{j=1}^n \lambda_j \frac{\partial}{\partial v^{(j)}} h(\mathbf{z}) \Big|_{\mathbf{z}=\mathbf{x}+\mathbf{y}t}$  in terms of directional derivatives, we obtain

$$\begin{aligned} W(f(\mathbf{x} + \mathbf{y}t), g(\mathbf{x} + \mathbf{y}t)) &= g(\mathbf{x} + t\mathbf{v})^2 \sum_{j=1}^n \lambda_j \frac{\partial}{\partial v^{(j)}} \left( \frac{f(\mathbf{z})}{g(\mathbf{z})} \right) \Big|_{\mathbf{z}=\mathbf{x}+\mathbf{y}t} \\ &= \sum_{j=1}^n \lambda_j W_{\mathbf{v}^{(j)}}(f, g)(\mathbf{x} + \mathbf{y}t) \leq 0. \end{aligned}$$

Hence,  $f(\mathbf{x} + \mathbf{y}t) \ll g(\mathbf{x} + \mathbf{y}t)$ , and thus by the Hermite–Biehler Theorem 2.2,  $g(\mathbf{x} + \mathbf{y}t) + if(\mathbf{x} + \mathbf{y}t)$  is stable. By Lemma 3.4,  $g + if$  is  $K$ -stable. □

### 5 psd-Stability

In this section, we consider the cone  $K = \mathcal{S}_n^+$  of positive semidefinite matrices. In many settings, this cone provides a natural generalization of the non-negative cone (see, e.g., [2]). In Theorem 5.3, we provide a generalization of a stability criterion for determinantal polynomials to the psd-stability.

Recall that for two matrices,  $A = (a_{ij})_{n_1 \times n_2}$  and  $B = (b_{ij})_{k_1 \times k_2}$ , the *Kronecker product* (or *tensor product*)  $A \otimes B$  is the  $n_1 k_1 \times n_2 k_2$  block matrix  $C = (C_{ij})_{n_1 \times n_2}$  with blocks  $C_{ij} = a_{ij} B$ . A generalization of the Kronecker product is the Khatri–Rao product, which is defined as follows.

**Definition 5.1** Let  $A = (A_{ij})_{n_1 \times n_2}$  and  $B = (B_{ij})_{n_1 \times n_2}$  be block matrices with  $n_1 \times n_2$  blocks of size  $p_1 \times p_2$  and  $q_1 \times q_2$ , respectively. The *Khatri–Rao* product of  $A$  and  $B$  is defined as

$$A * B = (A_{ij} \otimes B_{ij})_{n_1 \times n_2},$$

which is a block matrix with  $n_1 \times n_2$  blocks of size  $p_1 q_1 \times p_2 q_2$ .

Note that in the case of  $p_1 = p_2 = 1$ , the Khatri–Rao product provides a scalar multiplication of the blocks  $B_{ij}$  by the scalars  $a_{ij}$ . And in the case  $n_1 = n_2 = 1$ ,  $A$  and  $B$  only consist of a single block and  $A * B$  gives the usual Kronecker product.

While it is classically known that the Kronecker product of positive semidefinite matrices is positive semidefinite [12], the following result on the Khatri–Rao product will be relevant in our situation [18].

**Proposition 5.2** (Liu, Theorems 5 and 6 in [18]). *Let  $A = (A_{ij})_{n_1 \times n_2}$  and  $B = (B_{ij})_{n_1 \times n_2}$  be block matrices with the same block structure  $n_1 \times n_2$ . If  $A$  and  $B$  are positive semidefinite, then  $A * B$  is positive semidefinite. If  $A$  is positive semidefinite with positive definite blocks on the diagonal and  $B$  is positive definite, then  $A * B$  is positive definite.*

Note that the positive semidefiniteness of  $A$  implies that its blocks satisfy  $A_{ij} = A_{ji}^H$ , where  $A_{ij}^H$  denotes the Hermitian transpose of  $A_{ij}$ . Now we show the following generalization of Proposition 2.6 to psd-stability.

**Theorem 5.3** *Let  $A = (A_{ij})_{n \times n}$  be a block matrix with  $n \times n$  blocks of size  $d \times d$ . If  $A$  is positive semidefinite and  $B$  is a Hermitian  $d \times d$ -matrix, then the polynomial  $f(Z) = \det(\sum_{i,j=1}^n A_{ij}z_{ij} + B)$  on the set of symmetric  $n \times n$ -matrices is psd-stable or identically zero.*

*Proof* We write  $I_d$  for the  $d \times d$  identity matrix and  $\mathbb{1}_{m_1 \times m_2}$  for the all-ones matrix of size  $m_1 \times m_2$ . First consider the case where  $A$  is positive semidefinite with positive definite blocks on the diagonal.

Let  $X, Y \in \mathcal{S}_n$  with  $Y \succ 0$ . In view of Lemma 3.4, we have to show that the univariate polynomial  $t \mapsto f(X + tY)$  has only real roots.

We can interpret  $Y$  as a block matrix with blocks of size  $1 \times 1$ . Using the Khatri–Rao product,  $Y * A$  is a block matrix whose  $(i, j)$ th block is  $y_{ij}A_{ij}$ , and we obtain the identity

$$\sum_{i,j=1}^n y_{ij}A_{ij} = (\mathbb{1}_{1 \times n} \otimes I_d) \cdot (Y * A) \cdot (\mathbb{1}_{n \times 1} \otimes I_d). \tag{5.1}$$

Note that the multiplication by the matrices from left and right in (5.1) provides a block-wise summation of all the blocks in  $Y * A$ .

By Proposition 5.2,  $Y * A$  is positive definite. Hence, for  $\mathbf{v} \in \mathbb{R}^d \setminus \{0\}$ , we have

$$\begin{aligned} & \mathbf{v}^T ((\mathbb{1}_{1 \times n} \otimes I_d) \cdot (Y * A) \cdot (\mathbb{1}_{n \times 1} \otimes I_d)) \mathbf{v} \\ &= (\mathbf{v}^T \cdots \mathbf{v}^T)(Y * A)(\mathbf{v}^T \cdots \mathbf{v}^T)^T > 0. \end{aligned}$$

This implies that the Hermitian matrix  $Q := \sum_{i,j=1}^n A_{ij}y_{ij}$  is positive definite.

The positive definite matrix  $Q$  has a square root  $Q^{1/2}$ . Set  $H = \sum_{i,j=1}^n A_{ij}x_{ij} + B$ . Then, for any real symmetric  $n \times n$ -matrix  $X$ , the univariate polynomial

$$\begin{aligned} t \mapsto f(X + tY) &= \det \left( \sum_{i,j=1}^n A_{ij}(x_{ij} + ty_{ij}) + B \right) \\ &= \det(H + tQ) \\ &= \det(Q) \det(Q^{-1/2}HQ^{-1/2} + tI_d) \end{aligned}$$

has only real roots, since they are the negatives of the eigenvalues of the Hermitian matrix  $Q^{-1/2}HQ^{-1/2}$ .

Now, for the general case, let  $A$  be a positive semidefinite matrix. Let  $A^{(k)} = (A_{ij}^{(k)})_{n \times n}$  be a sequence of positive semidefinite block matrices with positive definite blocks on the diagonal, which approximate  $A$ . Then the polynomials  $f^{(k)}(Z) = \det(\sum_{i,j=1}^n A_{ij}^{(k)}z_{ij} + B)$  are psd-stable and hence have no root in the (open) Siegel upper half-plane. Due to Hurwitz’s Theorem 2.4, the limit polynomial  $f$  is either identically zero or also non-vanishing on the Siegel upper half-plane, i.e., it is psd-stable.  $\square$

For an example of Theorem 5.3, observe that choosing  $A$  as the block matrix with  $2 \times 2$  blocks of size  $2 \times 2$ ,

$$A_{11} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A_{12} = A_{21} = \begin{pmatrix} 0 & 1/2 \\ 1/2 & 0 \end{pmatrix}, \quad A_{22} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

and  $B = 0$  results in Example 3.3. Since  $A = (A_{ij})$  has the double eigenvalues  $1/2$  and  $3/2$ , it is positive semidefinite, so that Theorem 5.3 implies the psd-stability of

$$f(Z) = \det(A_{11}z_{11} + 2A_{12}z_{12} + A_{22}z_{22}), \quad \text{where } Z = \begin{pmatrix} z_{11} & z_{12} \\ z_{12} & z_{22} \end{pmatrix}.$$

The criterion stated in Theorem 5.3 is sufficient, but not necessary. The following is a counterexample.

*Example 5.4* Let  $Z = (z_{ij})_{2 \times 2}$  be symmetric and

$$f(Z) = \det \left( \sum_{i,j=1}^2 A_{ij}z_{ij} \right) = \det \left( \begin{pmatrix} 1 & 0 \\ 0 & 5 \end{pmatrix} z_{11} + 2 \begin{pmatrix} 0 & 2 \\ 2 & 0 \end{pmatrix} z_{12} + \begin{pmatrix} 5 & 0 \\ 0 & 1 \end{pmatrix} z_{22} \right).$$

We claim that  $f$  is psd-stable. Namely, for a real matrix  $Y = (y_{ij})_{2 \times 2} \succ 0$ , we have

$$f(Y) = (y_{11} + 5y_{22})(5y_{11} + y_{22}) - 16y_{12}^2 > 5(y_{11}^2 + y_{22}^2) + (26 - 16)y_{11}y_{22} > 0,$$

since  $y_{11}, y_{22} > 0$ . Hence,  $\sum_{i,j=1}^2 A_{ij}y_{ij} \succ 0$ , and thus, by Example 3.2 (ii),  $f$  is psd-stable. However, the matrix

$$A = (A_{ij}) = \begin{pmatrix} 1 & 0 & 0 & 2 \\ 0 & 5 & 2 & 0 \\ 0 & 2 & 5 & 0 \\ 2 & 0 & 0 & 1 \end{pmatrix}$$

is not positive semidefinite, since already the  $2 \times 2$ -minor with indices  $(1, 4)$  is negative.

We note that already the most simple case of a  $2 \times 2$ -matrix  $Z$  and diagonal coefficient matrices  $A_{ij}$  provides nonlinear conditions as the following statement shows.

**Proposition 5.5** *Let  $A_{ij} = \text{diag}(a_1^{(ij)}, \dots, a_d^{(ij)})$  be diagonal  $d \times d$ -matrices,  $1 \leq i, j \leq n$ . Then the block matrix  $A = (A_{ij})$  is positive semidefinite if and only if for every  $k \in \{1, \dots, d\}$  the matrix  $(a_k^{(ij)})_{1 \leq i, j \leq n}$  is positive semidefinite.*

*For  $n = 2$  and a Hermitian block matrix  $A = (A_{ij})$ , the criterion becomes*

$$a_k^{(11)}, a_k^{(22)} \geq 0, \text{ and } a_k^{(11)} a_k^{(22)} - |a_k^{(12)}|^2 \geq 0, \quad k = 1, \dots, d.$$

*Proof* Since the blocks of  $A$  are diagonal matrices, every row of  $A$  contains at most  $n$  nonzero entries. Namely, in the  $k$ th row of the  $i$ -th block row, these are the elements  $a_k^{(ij)}$ ,  $j = 1, \dots, n$ . By reordering the rows and columns of  $A$  using a permutation matrix  $P$ , the resulting matrix  $P^T A P$  has block diagonal structure with blocks  $A_k := (a_k^{(ij)})_{i,j}$  of size  $n \times n$ . Thus,  $A$  is positive semidefinite if and only if each block  $A_k$  is positive semidefinite.

For  $n = 2$ , these blocks are of size 2. Hence, the minors of  $A$  consist of factors of the form  $a_k^{(11)} a_k^{(22)} - a_k^{(12)} a_k^{(21)}$  together with diagonal elements. □

### 6 Conclusion and outlook

We have introduced the concept of conic stability for multivariate polynomials in  $\mathbb{C}[\mathbf{z}]$  and showed generalizations of some core results for stable polynomials to the conic stability. These positive results also show that the conic generalization of the stability notion appears to be very natural and fruitful. In particular, this raises the general question to which extent the theory of stable polynomials, for instance stability preserving operators, can be

generalized to the conic stability. With regard to the theorems on conic stability in this article, a question is if they can be further extended to even more general types of stability regions.

#### Acknowledgements

The authors thank Bernd Sturmfels for an inspiring conversation which became a trigger for the current paper. Moreover, thanks to an anonymous referee for helpful suggestions.

Received: 7 December 2017 Accepted: 4 May 2018 Published online: 15 May 2018

#### References

- Anari, N., Oveis Gharan, S.: A generalization of permanent inequalities and applications in counting and optimization. In: Proceedings of the Symposium on Theory of Computing, Montreal. ACM (2017)
- Blekherman, G., Parrilo, P.A., Thomas, R.R.: Semidefinite Optimization and Convex Algebraic Geometry. SIAM, Philadelphia (2013)
- Borcea, J., Brändén, P.: Applications of stable polynomials to mixed determinants: Johnson's conjectures, unimodality, and symmetrized Fischer products. *Duke Math. J.* **143**(2), 205–223 (2008)
- Borcea, J., Brändén, P.: The Lee-Yang and Pólya-Schur programs. I. Linear operators preserving stability. *Invent. Math.* **177**(3), 541–569 (2009)
- Borcea, J., Brändén, P.: The Lee-Yang and Pólya-Schur programs. II. Theory of stable polynomials and applications. *Commun. Pure Appl. Math.* **62**(12), 1595–1631 (2009)
- Borcea, J., Brändén, P.: Multivariate Pólya-Schur classification problems in the Weyl algebra. *Proc. Lond. Math. Soc.* **101**(1), 73–104 (2010)
- Brändén, P.: Obstructions to determinantal representability. *Adv. Math.* **226**(2), 1202–1212 (2011)
- Frey, G., Lange, T.: Varieties over special fields. In: Cohen, H., et al. (eds.) Handbook of Elliptic and Hyperelliptic Curve Cryptography, pp. 87–113. Chapman & Hall, Boca Raton (2006)
- Gårding, L.: An inequality for hyperbolic polynomials. *J. Math. Mech.* **8**, 957–965 (1959)
- Gurvits, L.: Van der Waerden/Schrijver-Valiant like conjectures and stable (aka hyperbolic) homogeneous polynomials: one theorem for all. *Electron. J. Comb.* **15**(1), R66 (2008)
- Helton, J.W., Vinnikov, V.: Linear matrix inequality representation of sets. *Commun. Pure Appl. Math.* **60**(5), 654–674 (2007)
- Horn, R.A., Johnson, C.R.: Matrix Analysis, 2nd edn. Cambridge University Press, Cambridge (2013)
- Itô, K. (ed.): Encyclopedic Dictionary of Mathematics, vol. I, 2nd edn. MIT Press, Cambridge (1987)
- Jörgens, T., Theobald, T.: Hyperbolicity cones and imaginary projections. *Proc. Am. Math. Soc.* (2018, To appear)
- Jörgens, T., Theobald, T., de Wolff, T.: Imaginary projections of polynomials (2016). [arXiv:1602.02008](https://arxiv.org/abs/1602.02008) (Preprint)
- Kummer, M., Plaumann, D., Vinzant, C.: Hyperbolic polynomials, interlacers, and sums of squares. *Math. Program.* **153**(1, Ser. B), 223–245 (2015)
- Lewis, A.S., Parrilo, P.A., Ramana, M.V.: The Lax conjecture is true. *Proc. Am. Math. Soc.* **133**, 2495–2499 (2005)
- Liu, S.: Matrix results on the Khatri-Rao and Tracy-Singh products. *Linear Algebra Appl.* **289**(1), 267–277 (1999)
- Marcus, A.W., Spielman, D.A., Srivastava, N.: Interlacing families I: bipartite Ramanujan graphs of all degrees. *Ann. Math.* **182**(1), 307–325 (2015)
- Marcus, A.W., Spielman, D.A., Srivastava, N.: Interlacing families II: mixed characteristic polynomials and the Kadison-Singer problem. *Ann. Math.* **182**(1), 327–350 (2015)
- Pemantle, R.: Hyperbolicity and stable polynomials in combinatorics and probability. In: Jerison, D., Mazur, B., Mrowka, T., et al. (eds.) Current Development in Mathematics 2011, pp. 57–124. Int. Press, Somerville, MA (2012)
- Rahman, Q.I., Schmeisser, G.: Analytic Theory of Polynomials, Volume 26 of London Mathematical Society Monographs. Clarendon Press, Oxford (2002)
- Siegel, C.L.: Einführung in die Theorie der Modulformen  $n$ -ten Grades. *Math. Ann.* **116**, 617–657 (1939)
- Straszak, D., Vishnoi, N.K.: Real stable polynomials and matroids: optimization and counting. In Proceedings of the Symposium on Theory of Computing, Montreal. ACM (2017)
- van der Geer, G.: Siegel modular forms and their applications. In: The 1-2-3 of Modular Forms, Universitext, pp. 181–245. Springer, Berlin (2008)
- Wagner, D.G.: Multivariate stable polynomials: theory and applications. *Bull. Am. Math. Soc.* **48**(1), 53–84 (2011)