



# Multiple-attribute decision-making based on picture fuzzy Archimedean power Maclaurin symmetric mean operators

Yuchu Qin<sup>1</sup> · Xiaolan Cui<sup>2</sup> · Meifa Huang<sup>1</sup> · Yanru Zhong<sup>3</sup> · Zhemin Tang<sup>1</sup> · Peizhi Shi<sup>4</sup>

Received: 6 April 2020 / Accepted: 14 June 2020 / Published online: 16 July 2020  
© Springer Nature Switzerland AG 2020

## Abstract

In this paper, a novel multiple-attribute decision-making method based on a set of Archimedean power Maclaurin symmetric mean operators of picture fuzzy numbers is proposed. The Maclaurin symmetric mean operator, power average operator, and operational rules based on Archimedean T-norm and T-conorm are introduced into picture fuzzy environment to construct the aggregation operators. The formal definitions of the aggregation operators are presented. Their general and specific expressions are established. The properties and special cases of the aggregation operators are, respectively, explored and discussed. Using the presented aggregation operators, a method for solving the multiple-attribute decision-making problems based on picture fuzzy numbers is designed. The method is illustrated through example and experiments and validated by comparisons. The results of the comparisons show that the proposed method is feasible and effective that can provide the generality and flexibility in aggregation of values of attributes and consideration of interactions among attributes and the capability to lower the negative effect of biased attribute values on the result of aggregation.

**Keywords** Multiple-attribute decision-making · Picture fuzzy set · Aggregation operator · Maclaurin symmetric mean operator · Power average operator · Archimedean T-norm and T-conorm

## 1 Introduction

Multiple-attribute decision-making (MADM) is the process of finding the best option through comprehensively assessing the values of multiple attributes of all options. There are two essential tasks in this process. One task is to describe the values of attributes and the other task is to fuse the described values to determine the best option. One of

the most important tools used in the first task is fuzzy set. Among the existing different types of fuzzy sets (Bustince et al. 2016), the fuzzy set (FS) presented by Zadeh (1965) is a classic type of fuzzy sets which leverages a degree of positive membership  $\mu$  ( $0 \leq \mu \leq 1$ ) to describe the degree of satisfaction. It is sufficient for fuzzy information description in some practical applications (Chen et al. 2009, 2012; Chen and Niou 2011; Chen and Chen 2014; Chen and Adam 2017; Castillo et al. 2019). However, FS is insufficient to express the fuzzy information consisting of the degrees of satisfaction, dissatisfaction, and hesitancy.

In response to this limitation, Atanassov (1986) extended FS and presented the intuitionistic fuzzy set (IFS), which provides a degree of positive membership  $\mu$  and a degree of negative membership  $\nu$  ( $0 \leq \mu \leq 1$ ;  $0 \leq \nu \leq 1$ ;  $0 \leq \mu + \nu \leq 1$ ). The two degrees can, respectively, quantify the degrees of satisfaction and dissatisfaction, and therefore, the degree of hesitancy is indirectly quantified by  $1 - \mu - \nu$ . Due to such capability, IFSs have been widely applied to describe attribute values in MADM. Various research topics regarding IFSs in MADM, such as calculus for IFSs (Lei and Xu 2016, 2017; Ai and Xu 2018),

✉ Meifa Huang  
meifahuang@yeah.net

<sup>1</sup> School of Mechanical and Electrical Engineering, Guilin University of Electronic Technology, Guilin 541004, People's Republic of China

<sup>2</sup> School of Management, Huazhong University of Science and Technology, Wuhan 430074, People's Republic of China

<sup>3</sup> Guangxi Key Laboratory of Intelligent Processing of Computer Images and Graphic, Guilin University of Electronic Technology, Guilin 541004, People's Republic of China

<sup>4</sup> School of Computing and Engineering, University of Huddersfield, Huddersfield HD1 3DH, UK

intuitionistic preference relations (Liao and Xu 2014; Liao et al. 2015; Zhang and Pedrycz 2017a, Zhang and Pedrycz 2018; Zhang et al. 2020), operations for IFSs (Jamkhaneh and Garg 2018; Dutta 2019; Dutta and Saikia 2019), information measures for IFSs (Chen et al. 2016a; Garg and Kumar 2018, 2019; Song et al. 2019; Tan et al. 2020), aggregation operators (AOs) of intuitionistic fuzzy numbers (IFNs) for MADM (Xu and Yager 2011; Chen and Chang 2016; Liu and Chen 2017; Liu and Tang 2018; Zhang et al. 2019; Seikh and Mandal 2019; Liu et al. 2020), and MADM methods based on IFSs (Wang and Zhang 2013; Chen et al. 2016b; Garg 2017a; Zhang and Pedrycz 2017b; Kumar and Garg 2018; Rani et al. 2019; Zeng et al. 2019), have received widespread attention.

Even though IFS has showed great capability in MADM, it cannot be used to describe more complex fuzzy information. IFS does not provide an approach to express the fuzzy information including the degree of neutrality. Aiming at this shortcoming, Hinde et al. (2007) introduced the theory of picture fuzzy set (PFS). A PFS extends an IFS with a degree of neutral membership  $\eta$ . Obviously, PFS is the generalisation of FS and IFS, since a PFS will reduce to a FS when its  $\eta = \nu = 0$  and will become an IFS if its  $\eta = 0$ . It is obvious that PFSs have the greatest expressive capability compared to FSs and IFSs. Because of this, PFSs and their application in MADM have also received a lot of attention. A variety of related research topics, such as correlation coefficients of PFSs (Singh 2015), distance measure of PFSs (Son 2016), picture fuzzy clustering (Thong 2016), application of PFSs in decision-making (Wang et al. 2018a, b; Ju et al. 2019), AOs of picture fuzzy numbers (PFNs) for MADM (Garg 2017b; Wei 2017, 2018; Wei et al. 2018a; Zhang et al. 2018; Jana et al. 2019; Xu et al. 2019), MADM methods based on PFSs (Wang et al. 2018a; Liang et al. 2018), and extensions of PFSs (Wei et al. 2018b; Mahmood et al. 2019; Khalil et al. 2019) are gaining importance.

MADM problems are generally solved using traditional methods or AOs. AOs are capable of solving MADM problems in a more effective way since they can produce summary values and rankings of options, and traditional methods only output rankings (Liu and Liu 2018; Liu and Wang 2018; Qin et al. 2019a, 2020a). So far, a number of MADM methods based on AOs of PFNs have been presented, such as the methods presented by Wei (2017), Garg (2017b), Wei (2018), Jana et al. (2019), Wei et al. (2018a), Zhang et al. (2018), and Xu et al. (2019). The AOs of PFNs on which these methods are based are listed in Table 1. To the best of the knowledge, there is not yet a method that has flexibility in the aggregation of values of attributes and generality in the handling of interactions of attributes and, meanwhile, can lower the negative effect of biased values of attributes.

In practical MADM problems, the preferences of decision makers usually change dynamically and various interactions always exist among different considered attributes. To generate reasonable results for these problems, the used AOs should be general and flexible enough to capture the preferences and interactions when aggregating the values of attributes (Liu and Wang 2019). Among the existing AOs of PFNs, the Archimedean WA, Archimedean OWA, and Archimedean HA operators in (Garg 2017b) can provide flexibility in the aggregation of values of attributes. But they are only applicable for the situation where all of the considered attributes are independent of each other. The weighted Heronian mean operator in (Wei et al. 2018a) and the Dombi weighted Heronian mean and Dombi weighted dual Heronian mean operators in (Zhang et al. 2018) can work normally under the condition that there are no interactions among attributes or there are interactions between two attributes. But they could produce unreasonable results when there are interactions among multiple attributes. The weighted Muirhead mean and weighted dual Muirhead mean operators in (Xu et al. 2019) can make up for this deficiency. But they are not general

**Table 1** The AOs of PFNs on Which the Seven MADM Methods Are Based

Method	AOs of PFNs	Operation
Wei (2017)	Weighted averaging (WA); Weighted geometric (WG); Ordered WA (OWA); Ordered WG (OWG); Hybrid averaging (HA); Hybrid geometric (HG)	Algebraic
Garg (2017b)	Archimedean WA; Archimedean OWA; Archimedean HA	Archimedean
Wei (2018)	Hamacher WA, OWA, HA, WG, OWG, HG, correlated averaging (CA) and correlated geometric (CG); Induced Hamacher OWA, OWG, CA and CG; Hamacher weighted prioritised average, prioritised geometric, power average and power geometric	Hamacher
Jana et al. (2019)	Dombi WA; Dombi OWA; Dombi HA; Dombi WG; Dombi OWG; Dombi HG	Dombi
Wei et al. (2018a)	Weighted Heronian mean	Algebraic
Zhang et al. (2018)	Dombi weighted Heronian mean; Dombi weighted dual Heronian mean	Dombi
Xu et al. (2019)	Weighted Muirhead mean, Weighted dual Muirhead mean	Algebraic

and flexible enough for aggregating attribute values. Besides, the attribute values are mostly evaluated by experts. The absolute objectivity of this way is usually difficult to be ensured. This means that a few experts will provide some biased attribute values (Liu and Liu 2017). To obtain reasonable aggregation results under this circumstance, it is required to lower the negative impact of biased values of attributes. However, none of the existing AOs of PFNs have such capability. Based on the analysis above, the motivations of this paper are outlined as follows:

- 1 To construct a flexible AO of PFNs, Archimedean T-norm and T-conorm (ATT) (Klement et al. 2000; Xia et al. 2012) are introduced into PFSs. The ATT operations are important tools for generalising logical conjunction and disjunction to fuzzy logic. They can be used to develop versatile rules for the operations between fuzzy numbers. The AOs using such operational rules are flexible (Liu and Wang 2019; Zhong et al. 2019a; Qin et al. 2019b, 2020b).
- 2 To make the AO general in capturing the interactions of attributes, Maclaurin symmetric mean (MSM) operator (Maclaurin 1729) is chosen as its core component. The MSM operator, a generalisation of the arithmetic average (AA) operator, Bonferroni mean (BM) operator, and geometric average (GA) operator, is a general AO for describing the interactions of the aggregated arguments. It is applicable for the situations where there are no interactions among all arguments, where there are interactions between two arguments, and where there are interactions among multiple arguments.
- 3 To make the AO capable to reduce the negative effect of unduly high or unduly low attribute values on the aggregation result, power average (PA) operator (Yager 2001) is combined with the MSM operator of PFNs under the ATT operations. The PA operator is an AO which can assign weights to the aggregated arguments via computing the degrees of support between the arguments. This makes it capable to reduce the negative effect of unreasonable argument values (Liu et al. 2018; Teng et al. 2018; Zhong et al. 2019b; Qin et al. 2020c).

In a word, the objective of the paper is to present a set of Archimedean power MSM operators of PFNs for solving the MADM problems based on PFNs. This objective is achieved via combining the MSM operator and the PA operator under the ATT operations in the context of PFSs. The major contribution of the paper is the development of an MADM method based on picture fuzzy Archimedean power MSM operators. This method can provide the generality in aggregation of values of attributes, the flexibility in handling of interactions among attributes, and the

capability to lower the negative impact of biased attribute values.

The rest of the paper is organised as follows. A brief introduction of some basic concepts is provided in Sect. 2. Section 3 describes the details of the presented operators. The specific process of the proposed method is described in Sect. 4. Section 5 demonstrates and evaluates the proposed method. Section 6 ends the paper with a conclusion.

## 2 Preliminaries

### 2.1 PFS theory

PFS can be seen as the generalisation of FS (Zadeh 1965) and IFS (Atanassov 1986). Its formal definition is given by Hinde et al. (2007):

**Definition 1** APFS  $S$  in a finite domain of discourse  $X$  is  $S = \{ \langle x, \mu_S(x), \eta_S(x), \nu_S(x) \rangle \mid x \in X \}$ , where  $\mu_S: X \rightarrow [0, 1]$  is the degree of positive membership of  $x \in X$  to  $S$ ,  $\eta_S: X \rightarrow [0, 1]$  is the degree of neutral membership of  $x \in X$  to  $S$ , and  $\nu_S: X \rightarrow [0, 1]$  is the degree of negative membership of  $x \in X$  to  $S$ , such that  $0 \leq \mu_S(x) + \eta_S(x) + \nu_S(x) \leq 1$ . The degree of refusal membership of  $x \in X$  to  $S$  is  $\pi_S(x) = 1 - \mu_S(x) - \eta_S(x) - \nu_S(x)$ .

A triple  $\langle \mu_S(x), \eta_S(x), \nu_S(x) \rangle$  is called a PFN and is usually denoted as  $\langle \mu, \eta, \nu \rangle$ . Two PFNs can be compared using their score values and accuracy values. Jana et al. (2019) introduced a function to calculate the score value of a PFN and Wei (2017) introduced a function to calculate the accuracy value of a PFN:

**Definition 2** Suppose  $\alpha = \langle \mu, \eta, \nu \rangle$  is a PFN, its score value can be calculated via  $S(\alpha) = 0.5 \times (1 + \mu - \nu)$ .

**Definition 3** Suppose  $\alpha = \langle \mu, \eta, \nu \rangle$  is a PFN, its accuracy value can be calculated via  $A(\alpha) = \mu + \eta + \nu$ .

Based on the score and accuracy values, Wei (2017) introduced the rules for comparing two PFNs:

**Definition 4** Suppose  $\alpha_1 = \langle \mu_1, \eta_1, \nu_1 \rangle$  and  $\alpha_2 = \langle \mu_2, \eta_2, \nu_2 \rangle$  are two PFNs,  $S(\alpha_1)$  and  $S(\alpha_2)$  are their score values, and  $A(\alpha_1)$  and  $A(\alpha_2)$  are their accuracy values. Then: (1)  $\alpha_1 > \alpha_2$  if  $S(\alpha_1) > S(\alpha_2)$ ; (2)  $\alpha_1 > \alpha_2$  if  $S(\alpha_1) = S(\alpha_2)$  and  $A(\alpha_1) > A(\alpha_2)$ ; (3)  $\alpha_1 = \alpha_2$  if  $S(\alpha_1) = S(\alpha_2)$  and  $A(\alpha_1) = A(\alpha_2)$ .

The distance of two PFNs can be calculated using a distance measure of PFNs. Cuong (2014) introduced a normalised Hamming distance measure of PFNs:

**Definition 5** Suppose  $\alpha_1 = \langle \mu_1, \eta_1, \nu_1 \rangle$  and  $\alpha_2 = \langle \mu_2, \eta_2, \nu_2 \rangle$  are two PFNs. Their normalised Hamming distance is  $d(\alpha_1, \alpha_2) = 0.5(|\mu_1 - \mu_2| + |\eta_1 - \eta_2| + |\nu_1 - \nu_2|)$ .

## 2.2 Operational rules

Based on ATT, Garg (2017b) introduced a set of operational rules of PFNs:

**Definition 6** Suppose  $\alpha = \langle \mu, \eta, v \rangle$ ,  $\alpha_1 = \langle \mu_1, \eta_1, v_1 \rangle$ , and  $\alpha_2 = \langle \mu_2, \eta_2, v_2 \rangle$  are three PFNs, and  $c$  and  $d$  are two positive real numbers. The operations of PFNs based on  $T(x, y) = \varphi^{-1}(\varphi(x) + \varphi(y))$  and  $T^C(x, y) = \psi^{-1}(\psi(x) + \psi(y))$  are defined as follows:

$$\begin{aligned} \alpha_1 \oplus \alpha_2 &= \langle T^C(\mu_1, \mu_2), T(\eta_1, \eta_2), T(v_1, v_2) \rangle \\ &= \langle \psi^{-1}(\psi(\mu_1) + \psi(\mu_2)), \varphi^{-1}(\varphi(\eta_1) + \varphi(\eta_2)), \varphi^{-1}(\varphi(v_1) + \varphi(v_2)) \rangle, \end{aligned} \tag{1}$$

$$\begin{aligned} \alpha_1 \otimes \alpha_2 &= \langle T(\mu_1, \mu_2), T^C(\eta_1, \eta_2), T^C(v_1, v_2) \rangle \\ &= \langle \varphi^{-1}(\varphi(\mu_1) + \varphi(\mu_2)), \psi^{-1}(\psi(\eta_1) + \psi(\eta_2)), \psi^{-1}(\psi(v_1) + \psi(v_2)) \rangle, \end{aligned} \tag{2}$$

$$c\alpha = \langle \psi^{-1}(c\psi(\mu)), \varphi^{-1}(c\varphi(\eta)), \varphi^{-1}(c\varphi(v)) \rangle, \tag{3}$$

$$\alpha^d = \langle \varphi^{-1}(d\varphi(\mu)), \psi^{-1}(d\psi(\eta)), \psi^{-1}(d\psi(v)) \rangle. \tag{4}$$

## 2.3 MSM operator

The MSM operator was introduced by Maclaurin (1729). This operator can be formally defined as:

**Definition 7** Suppose  $a_1, a_2, \dots, a_n$  are  $n$  crisp numbers and  $k$  is an integer such that  $1 \leq k \leq n$ . If  $(i_1, i_2, \dots, i_k)$  traverse all the  $k$ -tuple combinations of  $(1, 2, \dots, n)$ , then

$$\text{MSM}^{(k)}(a_1, a_2, \dots, a_n) = \left( \frac{k!(n-k)!}{n!} \sum_{1 \leq i_1 < \dots < i_k \leq n} \prod_{h=1}^k a_{i_h} \right)^{1/k} \tag{5}$$

is called the MSM operator.

## 2.4 PA operator

The PA operator was introduced by Yager (2001). This operator can be formally defined as:

**Definition 8** Suppose  $a_1, a_2, \dots, a_n$  are  $n$  crisp numbers,  $S(a_i, a_j) = 1 - d(a_i, a_j)$  ( $i, j = 1, 2, \dots, n$  and  $j \neq i$ ;  $d(a_i, a_j)$  is the distance between  $a_i$  and  $a_j$ ) is the degree of support for  $a_i$  from  $a_j$  which satisfies  $0 \leq S(a_i, a_j) \leq 1$ ,  $S(a_i, a_j) = S(a_j, a_i)$ , and  $S(a_i, a_j) \geq S(a_p, a_q)$  if  $|a_i, a_j| \leq |a_p, a_q|$ , and

$$T(a_i) = \sum_{j=1, j \neq i}^n S(a_i, a_j),$$

then

$$\text{PA}(a_1, a_2, \dots, a_n) = \sum_{i=1}^n ((1 + T(a_i))a_i) / \sum_{i=1}^n (1 + T(a_i)) \tag{6}$$

is called the PA operator.

## 3 Archimedean power MSM operators

### 3.1 PFAPMSM operator

A picture fuzzy Archimedean power MSM (PFAPMSM) operator is a power MSM operator of PFNs, in which the operations are carried out via the operational rules of PFNs based on ATT. This operator can be formally defined as:

**Definition 9** Suppose  $\alpha_1, \alpha_2, \dots, \alpha_n$  ( $\alpha_i = \langle \mu_i, \eta_i, v_i \rangle$ ,  $i = 1, 2, \dots, n$ ) are  $n$  PFNs,  $k$  is an integer which meets  $1 \leq k \leq n$ ,  $\alpha_p \oplus \alpha_q$  and  $\alpha_p \otimes \alpha_q$  ( $p, q = 1, 2, \dots, n$  and  $p \neq q$ ) and  $c\alpha_r$  and  $\alpha_s^d$  ( $r, s = 1, 2, \dots, n$  and  $r \neq s$ ;  $c, d > 0$ ) are the operations of PFNs based on ATT, and  $S(\alpha_p, \alpha_q) = 1 - d(\alpha_p, \alpha_q)$  ( $d(\alpha_p, \alpha_q)$  is the distance between  $\alpha_p$  and  $\alpha_q$ ) be the degree of support for  $\alpha_p$  from  $\alpha_q$  which satisfies  $0 \leq S(\alpha_p, \alpha_q) \leq 1$ ,  $S(\alpha_p, \alpha_q) = S(\alpha_q, \alpha_p)$ , and  $S(\alpha_p, \alpha_q) \geq S(\alpha_r, \alpha_s)$  if  $|\alpha_p - \alpha_q| \leq |\alpha_r - \alpha_s|$ , and

$$T(\alpha_p) = \sum_{q=1, q \neq p}^n S(\alpha_p, \alpha_q).$$

If  $(i_1, i_2, \dots, i_k)$  traverse all the  $k$ -tuple combinations of  $(1, 2, \dots, n)$ , then

$$\begin{aligned} \text{PFAPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) &= \left( \frac{1}{C_n^k} \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k \left( \frac{n(1 + T(\alpha_{i_h}))}{\sum_{j=1}^n (1 + T(\alpha_j))} \alpha_{i_h} \right) \right)^{1/k} \end{aligned} \tag{7}$$

is called the PFAPMSM operator.

The general expression of the PFAPMSM is constructed in the following theorem:

**Theorem 1** Suppose  $\alpha_1, \alpha_2, \dots, \alpha_n$  ( $\alpha_i = \langle \mu_i, \eta_i, v_i \rangle$ ,  $i = 1, 2, \dots, n$ ) are  $n$  PFNs. Then

$$\text{PFAPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) = \langle \mu, \eta, v \rangle, \tag{8}$$

where

$$\begin{aligned} \mu &= \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi \left( \psi^{-1} \left( (n\omega_{i_h}) \psi(\mu_{i_h}) \right) \right) \right) \right) \right) \right) \right) \\ \eta &= \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi \left( \varphi^{-1} \left( (n\omega_{i_h}) \varphi(\eta_{i_h}) \right) \right) \right) \right) \right) \right) \right) \\ v &= \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi \left( \varphi^{-1} \left( (n\omega_{i_h}) \varphi(v_{i_h}) \right) \right) \right) \right) \right) \right) \right) \\ \omega_{i_h} &= (1 + T(\alpha_{i_h})) / \sum_{j=1}^n (1 + T(\alpha_j)) \end{aligned}$$

and PFAPMSM<sup>(k)</sup>(α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub>) is still a PFN.

The proof of this theorem is provided in App. A. The properties of the PFAPMSM operator are stated in the following theorems:

**Theorem 2 (Idempotency)** Suppose α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub> (α<sub>i</sub> = ⟨μ<sub>i</sub>, η<sub>i</sub>, v<sub>i</sub>⟩, i = 1, 2, ..., n) are n PFNs. If α<sub>i</sub> = α = ⟨μ<sub>α</sub>, η<sub>α</sub>, v<sub>α</sub>⟩ for all i = 1, 2, ..., n, then PFAPMSM<sup>(k)</sup>(α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub>) = α = ⟨μ<sub>α</sub>, η<sub>α</sub>, v<sub>α</sub>⟩.

**Theorem 3 (Commutativity)** Suppose α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub> (α<sub>i</sub> = ⟨μ<sub>i</sub>, η<sub>i</sub>, v<sub>i</sub>⟩, i = 1, 2, ..., n) are n PFNs. If (β<sub>1</sub>, β<sub>2</sub>, ..., β<sub>n</sub>) is any permutation of (α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub>), then PFAPMSM<sup>(k)</sup>(β<sub>1</sub>, β<sub>2</sub>, ..., β<sub>n</sub>) = PFAPMSM<sup>(k)</sup>(α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub>).

**Theorem 4 (Boundedness).** Suppose α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub> (α<sub>i</sub> = ⟨μ<sub>i</sub>, η<sub>i</sub>, v<sub>i</sub>⟩, i = 1, 2, ..., n) are n PFNs, α<sup>+</sup> = ⟨max(μ<sub>i</sub>), min(η<sub>i</sub>), min(v<sub>i</sub>)⟩, and α<sup>-</sup> = ⟨min(μ<sub>i</sub>), max(η<sub>i</sub>), max(v<sub>i</sub>)⟩. Then α<sup>-</sup> ≤ PFAPMSM<sup>(k)</sup>(α<sub>1</sub>, α<sub>2</sub>, ..., α<sub>n</sub>) ≤ α<sup>+</sup>.

The proofs of these theorems are, respectively, provided in App.es B, C, and D. Some special cases of the PFAPMSM operator are as follows:

- 1 If k = 1, the PFAPMSM operator will become

$$\left\langle \psi^{-1} \left( \frac{1}{n} \sum_{i=1}^n ((n\omega_i) \psi(\mu_i)) \right), \varphi^{-1} \left( \frac{1}{n} \sum_{i=1}^n ((n\omega_i) \varphi(\eta_i)) \right), \varphi^{-1} \left( \frac{1}{n} \sum_{i=1}^n ((n\omega_i) \varphi(v_i)) \right) \right\rangle = \text{PFAPAA}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{9}$$

which is a picture fuzzy Archimedean power AA (PFAPAA) operator.

- 2 If k = 2, the PFAPMSM operator will become

$$\begin{aligned} &\left\langle \varphi^{-1} \left( \frac{1}{2} \varphi \left( \psi^{-1} \left( \frac{1}{n(n-1)} \sum_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \psi \left( \varphi^{-1} \left( \varphi \left( \psi^{-1} \left( (n\omega_{i_1}) \psi(\mu_{i_1}) \right) \right) + \varphi \left( \psi^{-1} \left( (n\omega_{i_2}) \psi(\mu_{i_2}) \right) \right) \right) \right) \right) \right) \right) \right) \\ &\psi^{-1} \left( \frac{1}{2} \psi \left( \varphi^{-1} \left( \frac{1}{n(n-1)} \sum_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \varphi \left( \psi^{-1} \left( \psi \left( \varphi^{-1} \left( (n\omega_{i_1}) \varphi(\eta_{i_1}) \right) \right) + \psi \left( \varphi^{-1} \left( (n\omega_{i_2}) \varphi(\eta_{i_2}) \right) \right) \right) \right) \right) \right) \right) \right) \\ &\psi^{-1} \left( \frac{1}{2} \psi \left( \varphi^{-1} \left( \frac{1}{n(n-1)} \sum_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \varphi \left( \psi^{-1} \left( \psi \left( \varphi^{-1} \left( (n\omega_{i_1}) \varphi(v_{i_1}) \right) \right) + \psi \left( \varphi^{-1} \left( (n\omega_{i_2}) \varphi(v_{i_2}) \right) \right) \right) \right) \right) \right) \right) \right) \right) \\ &= \text{PFAPBM}^{(1,1)}(\alpha_1, \alpha_2, \dots, \alpha_n), \end{aligned} \tag{10}$$

which is a picture fuzzy Archimedean power BM (PFAPBM) operator.

3 If  $k = n$ , the PFAPMSM operator will become

$$\left\langle \varphi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \varphi(\psi^{-1}((n\omega_i)\psi(\mu_i))) \right), \right. \\ \left. \psi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \psi(\varphi^{-1}((n\omega_i)\varphi(\eta_i))) \right), \right. \\ \left. \psi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \psi(\varphi^{-1}((n\omega_i)\varphi(v_i))) \right) \right\rangle, \tag{11}$$

$$= \text{PFAPGA}(\alpha_1, \alpha_2, \dots, \alpha_n)$$

(a) If  $k = 1$ , the PFAAPMSM operator will become

$$\left\langle 1 - \prod_{i=1}^n (1 - \mu_i)^{\omega_i}, \prod_{i=1}^n \eta_i^{\omega_i}, \prod_{i=1}^n v_i^{\omega_i} \right\rangle \\ = \text{PFPA}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{13}$$

which is a picture fuzzy power AA (PFPA) operator.

(b) If  $k = 2$ , the PFAAPMSM operator will become

$$\left\langle \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n (1 - (1 - (1 - \mu_{i_1})^{n\omega_{i_1}})(1 - (1 - \mu_{i_2})^{n\omega_{i_2}}))^{\frac{1}{n(n-1)}} \right)^{1/2}, 1 - \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n (1 - (1 - \eta_{i_1}^{n\omega_{i_1}})(1 - \eta_{i_2}^{n\omega_{i_2}}))^{\frac{1}{n(n-1)}} \right)^{1/2}, \right. \\ \left. 1 - \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n (1 - (1 - v_{i_1}^{n\omega_{i_1}})(1 - v_{i_2}^{n\omega_{i_2}}))^{\frac{1}{n(n-1)}} \right)^{1/2} \right\rangle = \text{PFPA}^{(1,1)}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{14}$$

which is a picture fuzzy Archimedean power GA (PFAPGA) operator.

The specific expressions of the PFAPMSM operator are constructed as follows:

1 If Algebraic T-norm and T-conorm are applied to Eq. (8), a picture fuzzy Archimedean Algebraic power MSM (PFAAPMSM) operator can be obtained as:

$$\text{PFAAPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\ = \left\langle \left( 1 - \left( \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( 1 - \prod_{h=1}^k (1 - (1 - \mu_{i_h})^{n\omega_{i_h}}) \right) \right) \right)^{1/C_n^k} \right\rangle^{1/k}, \\ 1 - \left( 1 - \left( \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( 1 - \prod_{h=1}^k (1 - \eta_{i_h}^{n\omega_{i_h}}) \right) \right) \right)^{1/C_n^k} \right\rangle^{1/k}, \\ 1 - \left( 1 - \left( \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( 1 - \prod_{h=1}^k (1 - v_{i_h}^{n\omega_{i_h}}) \right) \right) \right)^{1/C_n^k} \right\rangle^{1/k}. \tag{12}$$

Some special cases of this operator are as follows:

which is a picture fuzzy power BM (PFPA) operator.

(c) If  $k = n$ , the PFAAPMSM operator will become

$$\left\langle \prod_{i=1}^n (1 - (1 - \mu_i)^{n\omega_i})^{1/n}, 1 - \prod_{i=1}^n (1 - \eta_i^{n\omega_i})^{1/n}, \right. \\ \left. 1 - \prod_{i=1}^n (1 - v_i^{n\omega_i})^{1/n} \right\rangle = \text{PFPGA}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{15}$$

which is a picture fuzzy power GA (PFPGA) operator.

2 If Einstein T-norm and T-conorm are applied to Eq. (8), a picture fuzzy Archimedean Einstein power MSM (PFAEPMSM) operator can be obtained as:

$$\begin{aligned} & \text{PFAEPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\ &= \left\langle \left( 2(\mu'' - 1)^{1/k} \right) / \left( (\mu'' + 3)^{1/k} + (\mu'' - 1)^{1/k} \right), \right. \\ & \left. \left( (\eta'' + 3)^{1/k} - (\eta'' - 1)^{1/k} \right) / \left( (\eta'' + 3)^{1/k} + (\eta'' - 1)^{1/k} \right), \right. \\ & \left. \left( (\nu'' + 3)^{1/k} - (\nu'' - 1)^{1/k} \right) / \left( (\nu'' + 3)^{1/k} + (\nu'' - 1)^{1/k} \right) \right\rangle, \end{aligned} \tag{16}$$

where

$$\begin{aligned} \mu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\mu' + 3) / (\mu' - 1))^{1/C_n^k}, \quad \mu' = \prod_{h=1}^k \left( \frac{(1 + \mu_{i_h})^{n\omega_{i_h}} + 3(1 - \mu_{i_h})^{n\omega_{i_h}}}{(1 + \mu_{i_h})^{n\omega_{i_h}} - (1 - \mu_{i_h})^{n\omega_{i_h}}} \right), \\ \eta'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\eta' + 3) / (\eta' - 1))^{1/C_n^k}, \quad \eta' = \prod_{h=1}^k \left( \frac{(2 - \eta_{i_h})^{n\omega_{i_h}} + 3\eta_{i_h}^{n\omega_{i_h}}}{(2 - \eta_{i_h})^{n\omega_{i_h}} - \eta_{i_h}^{n\omega_{i_h}}} \right), \\ \nu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\nu' + 3) / (\nu' - 1))^{1/C_n^k}, \quad \nu' = \prod_{h=1}^k \left( \frac{(2 - \nu_{i_h})^{n\omega_{i_h}} + 3\nu_{i_h}^{n\omega_{i_h}}}{(2 - \nu_{i_h})^{n\omega_{i_h}} - \nu_{i_h}^{n\omega_{i_h}}} \right). \end{aligned}$$

(PFAFPMSM) operator can be obtained as:

$$\begin{aligned} & \text{PFAFPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\ &= \left\langle \log_\varepsilon \left( 1 + (\varepsilon^{\mu''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right), \right. \\ & \quad \left. 1 - \log_\varepsilon \left( 1 + (\varepsilon^{1-\eta''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right), \right. \\ & \quad \left. 1 - \log_\varepsilon \left( 1 + (\varepsilon^{1-\nu''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right) \right\rangle, \end{aligned} \tag{18}$$

3 If Hamacher T-norm and T-conorm are applied to Eq. (8), a picture fuzzy Archimedean Hamacher power MSM (PFAHPMSM) operator can be obtained as:

$$\begin{aligned} & \text{PFAHPMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\ &= \left\langle \left( \lambda(\mu'' - 1)^{1/k} \right) / \left( (\mu'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\mu'' - 1)^{1/k} \right), \right. \\ & \quad \left( (\eta'' + \lambda^2 - 1)^{1/k} - (\eta'' - 1)^{1/k} \right) / \left( (\eta'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\eta'' - 1)^{1/k} \right), \\ & \quad \left( (\nu'' + \lambda^2 - 1)^{1/k} - (\nu'' - 1)^{1/k} \right) / \left( (\nu'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\nu'' - 1)^{1/k} \right) \right\rangle, \end{aligned} \tag{17}$$

where  $\lambda > 0$  and

$$\begin{aligned} \mu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\mu' + \lambda^2 - 1) / (\mu' - 1))^{1/C_n^k}, \\ \mu' &= \prod_{h=1}^k \frac{(\lambda + (1 - \lambda)(1 - \mu_{i_h}))^{n\omega_{i_h}} + (\lambda^2 - 1)(1 - \mu_{i_h})^{n\omega_{i_h}}}{(\lambda + (1 - \lambda)(1 - \mu_{i_h}))^{n\omega_{i_h}} - (1 - \mu_{i_h})^{n\omega_{i_h}}}, \\ \eta'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\eta' + \lambda^2 - 1) / (\eta' - 1))^{1/C_n^k}, \\ \eta' &= \prod_{h=1}^k \left( \frac{((\lambda + (1 - \lambda)\eta_{i_h})^{n\omega_{i_h}} + (\lambda^2 - 1)\eta_{i_h}^{n\omega_{i_h}})}{((\lambda + (1 - \lambda)\eta_{i_h})^{n\omega_{i_h}} - \eta_{i_h}^{n\omega_{i_h}})} \right), \\ \nu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\nu' + \lambda^2 - 1) / (\nu' - 1))^{1/C_n^k}, \\ \nu' &= \prod_{h=1}^k \left( \frac{((\lambda + (1 - \lambda)\nu_{i_h})^{n\omega_{i_h}} + (\lambda^2 - 1)\nu_{i_h}^{n\omega_{i_h}})}{((\lambda + (1 - \lambda)\nu_{i_h})^{n\omega_{i_h}} - \nu_{i_h}^{n\omega_{i_h}})} \right). \end{aligned}$$

4 If Frank T-norm and T-conorm are applied to Eq. (8), a picture fuzzy Archimedean Frank power MSM

where  $\varepsilon > 1$  and

$$\begin{aligned} \mu''' &= 1 - \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\varepsilon - 1) / (\varepsilon^{1-\mu''} - 1))^{1/C_n^k} \right), \\ \mu'' &= \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k ((\varepsilon - 1) / (\varepsilon^{\mu'} - 1)) \right), \\ \mu' &= 1 - \log_\varepsilon \left( 1 + ((\varepsilon^{1-\mu_h} - 1)^{n\omega_{i_h}} / (\varepsilon - 1)^{n\omega_{i_h}-1}) \right), \\ \eta''' &= \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\varepsilon - 1) / (\varepsilon^{\eta''} - 1))^{1/C_n^k} \right), \\ \eta'' &= 1 - \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k ((\varepsilon - 1) / (\varepsilon^{1-\eta'} - 1)) \right), \\ \eta' &= \log_\varepsilon \left( 1 + (\varepsilon^{\eta_h} - 1)^{n\omega_{i_h}} / (\varepsilon - 1)^{n\omega_{i_h}-1} \right), \\ \nu''' &= \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\varepsilon - 1) / (\varepsilon^{\nu''} - 1))^{1/C_n^k} \right), \\ \nu'' &= 1 - \log_\varepsilon \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k ((\varepsilon - 1) / (\varepsilon^{1-\nu'} - 1)) \right), \\ \nu' &= \log_\varepsilon \left( 1 + (\varepsilon^{\nu_h} - 1)^{n\omega_{i_h}} / (\varepsilon - 1)^{n\omega_{i_h}-1} \right). \end{aligned}$$

### 3.2 PFAPWMSM operator

To capture the relative importance of the aggregated PFNs, a picture fuzzy Archimedean power weighted MSM (PFAPWMSM) operator is presented. This operator can be formally defined as:

**Definition 10** Suppose  $w_1, w_2, \dots, w_n$  ( $0 \leq w_1, w_2, \dots, w_n \leq 1$  and  $w_1 + w_2 + \dots + w_n = 1$ ) are, respectively, the

weights of  $\alpha_1, \alpha_2, \dots, \alpha_n$  that, respectively, denote the relative importance of  $\alpha_1, \alpha_2, \dots, \alpha_n$ . Then on the basis of Def. 9,

$$PFAPWMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) = \left( \frac{1}{C_n^k} \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k \left( \frac{nw_{i_h}(1 + T(\alpha_{i_h}))}{\sum_{j=1}^n (w_j(1 + T(\alpha_j)))} \alpha_{i_h} \right) \right)^{1/k} \tag{19}$$

is called the PFAPWMSM operator.

The general expression of the PFAPWMSM is constructed in the following theorem:

**Theorem 5** Suppose  $\alpha_1, \alpha_2, \dots, \alpha_n$  ( $\alpha_i = \langle \mu_i, \eta_i, \nu_i \rangle, i = 1, 2, \dots, n$ ) are  $n$  PFNs. Then

$$PFAPWMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) = \langle \mu, \eta, \nu \rangle, \tag{20}$$

where

$$\begin{aligned} \mu &= \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi \left( \psi^{-1} \left( \left( \frac{nw_{i_h} \omega_{i_h}}{\sum_{t=1}^n (w_t \omega_t)} \right) \psi(\mu_{i_h}) \right) \right) \right) \right) \right) \right) \right) \right), \\ \eta &= \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi \left( \varphi^{-1} \left( \left( \frac{nw_{i_h} \omega_{i_h}}{\sum_{t=1}^n (w_t \omega_t)} \right) \varphi(\eta_{i_h}) \right) \right) \right) \right) \right) \right) \right) \right), \\ \nu &= \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi \left( \varphi^{-1} \left( \left( \frac{nw_{i_h} \omega_{i_h}}{\sum_{t=1}^n (w_t \omega_t)} \right) \varphi(\nu_{i_h}) \right) \right) \right) \right) \right) \right) \right) \right), \\ \omega_{i_h} &= (1 + T(\alpha_{i_h})) / \sum_{j=1}^n (1 + T(\alpha_j)), \\ \omega_t &= (1 + T(\alpha_t)) / \sum_{j=1}^n (1 + T(\alpha_j)). \end{aligned}$$

and  $PFAPWMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n)$  is still a PFN.

This theorem can be proved like proof Theorem 1 in App. A. The properties of the PFAPWMSM operator are stated in the following theorems:

**Theorem 6** (Commutativity) Suppose  $\alpha_1, \alpha_2, \dots, \alpha_n$  ( $\alpha_i = \langle \mu_i, \eta_i, \nu_i \rangle, i = 1, 2, \dots, n$ ) are  $n$  PFNs. If  $(\beta_1, \beta_2, \dots, \beta_n)$  is any permutation of  $(\alpha_1, \alpha_2, \dots, \alpha_n)$ , then  $PFAPWMSM^{(k)}(\beta_1, \beta_2, \dots, \beta_n) = PFAPWMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n)$ .

**Theorem 7** (Boundedness) Suppose  $\alpha_1, \alpha_2, \dots, \alpha_n$  ( $\alpha_i = \langle \mu_i, \eta_i, \nu_i \rangle, i = 1, 2, \dots, n$ ) are  $n$  PFNs,

$\alpha^+ = \langle \max(\mu_i), \min(\eta_i), \min(\nu_i) \rangle$ , and  $\alpha^- = \langle \min(\mu_i), \max(\eta_i), \max(\nu_i) \rangle$ . Then  $\alpha^- \leq PFAPWMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \leq \alpha^+$ .

These theorems can be respectively proved like proof of Theorem 3 in App. C and proof of Theorem 4 in App. D. Some special cases of the PFAPWMSM operator are as follows:

1 If  $k = 1$ , the PFAPWMSM operator will become

$$\begin{aligned} &\left\langle \psi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \left( \left( \frac{nw_i \omega_i}{\sum_{t=1}^n (w_t \omega_t)} \right) \psi(\mu_i) \right) \right), \right. \\ &\varphi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \left( \left( \frac{nw_i \omega_i}{\sum_{t=1}^n (w_t \omega_t)} \right) \varphi(\eta_i) \right) \right), \\ &\left. \varphi^{-1} \left( \frac{1}{n} \sum_{i=1}^n \left( \left( \frac{nw_i \omega_i}{\sum_{t=1}^n (w_t \omega_t)} \right) \varphi(\nu_i) \right) \right) \right\rangle \\ &= PFAPWAA(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{21} \end{aligned}$$

which is a picture fuzzy Archimedean power weighted

AA (PFAPWAA) operator.

2 If  $k = 2$ , the PFAPWMSM operator will become





(a) If  $k = 1$ , the PFAAPWMSM operator will become

$$\left\langle 1 - \prod_{i=1}^n (1 - \mu_i)^{\frac{(w_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}}, \prod_{i=1}^n \eta_i^{\frac{(w_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}}, \prod_{i=1}^n v_i^{\frac{(w_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}} \right\rangle = \text{PFPWAA}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{25}$$

which is a picture fuzzy power weighted AA (PFPWAA) operator.

(b) If  $k = 2$ , the PFAAPWMSM operator will become

$$\left\langle \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \left( 1 - \left( 1 - (1 - \mu_{i_1})^{\frac{(nw_{i_1} \omega_{i_1})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \left( 1 - (1 - \mu_{i_2})^{\frac{(nw_{i_2} \omega_{i_2})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \right)^{\frac{1}{n(n-1)}} \right)^{1/2}, \right. \\ \left. 1 - \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \left( 1 - \left( 1 - \eta_{i_1}^{\frac{(nw_{i_1} \omega_{i_1})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \left( 1 - \eta_{i_2}^{\frac{(nw_{i_2} \omega_{i_2})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \right)^{\frac{1}{n(n-1)}} \right)^{1/2}, \right. \\ \left. 1 - \left( 1 - \prod_{\substack{i_1, i_2=1 \\ i_2 \neq i_1}}^n \left( 1 - \left( 1 - v_{i_1}^{\frac{(nw_{i_1} \omega_{i_1})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \left( 1 - v_{i_2}^{\frac{(nw_{i_2} \omega_{i_2})}{\sum_{i=1}^n (w_i \omega_i)}} \right) \right)^{\frac{1}{n(n-1)}} \right)^{1/2} \right\rangle \\ = \text{PFPWBM}^{(1,1)}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{26}$$

which is a picture fuzzy power weighted BM (PFPWBM) operator.

(c) If  $k = n$ , the PFAAPWMSM operator will become

$$\left\langle \prod_{i=1}^n \left( 1 - (1 - \mu_i)^{\frac{(nw_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}} \right)^{1/n}, \right. \\ \left. 1 - \prod_{i=1}^n \left( 1 - \eta_i^{\frac{(nw_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}} \right)^{1/n}, \right. \\ \left. 1 - \prod_{i=1}^n \left( 1 - v_i^{\frac{(nw_i \omega_i)}{\sum_{i=1}^n (w_i \omega_i)}} \right)^{1/n} \right\rangle \\ = \text{PFPWGA}(\alpha_1, \alpha_2, \dots, \alpha_n), \tag{27}$$

which is a picture fuzzy power weighted GA (PFPWGA) operator.

5 If Einstein T-norm and T-conorm are applied to Eq. (20), a picture fuzzy Archimedean Einstein power weighted MSM (PFAEPWMSM) operator can be obtained as:

$$\begin{aligned} & \text{PFAEPWMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\ &= \left\langle \left( 2(\mu'' - 1)^{1/k} \right) / \left( (\mu'' + 3)^{1/k} + (\mu'' - 1)^{1/k} \right), \right. \\ & \left( (\eta'' + 3)^{1/k} - (\eta'' - 1)^{1/k} \right) / \left( (\eta'' + 3)^{1/k} + (\eta'' - 1)^{1/k} \right), \\ & \left. \left( (v'' + 3)^{1/k} - (v'' - 1)^{1/k} \right) / \left( (v'' + 3)^{1/k} + (v'' - 1)^{1/k} \right) \right\rangle, \end{aligned} \tag{28}$$

where

$$\begin{aligned} \mu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\mu' + 3) / (\mu' - 1))^{1/C_n^k}, \\ \mu' &= \prod_{h=1}^k \frac{(1 + \mu_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} + 3(1 - \mu_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}{(1 + \mu_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} - (1 - \mu_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}, \\ \eta'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((\eta' + 3) / (\eta' - 1))^{1/C_n^k}, \\ \eta' &= \prod_{h=1}^k \frac{(2 - \eta_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} + 3\eta_{i_h}^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}{(2 - \eta_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} - \eta_{i_h}^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}, \\ v'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} ((v' + 3) / (v' - 1))^{1/C_n^k}, \\ v' &= \prod_{h=1}^k \frac{(2 - v_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} + 3v_{i_h}^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}{(2 - v_{i_h})^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}} - v_{i_h}^{\frac{(nw_{i_h} \omega_{i_h})}{\sum_{i=1}^n (w_i \omega_i)}}}. \end{aligned}$$

6 If Hamacher T-norm and T-conorm are applied to Eq. (20), a picture fuzzy Archimedean Hamacher power weighted MSM (PFAHPWMSM) operator can be obtained as:

$$\begin{aligned}
 & \text{PFAHPWMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\
 &= \left\langle \left( \frac{\lambda(\mu'' - 1)^{1/k}}{(\mu'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\mu'' - 1)^{1/k}} \right), \right. \\
 & \quad \left( \frac{(\eta'' + \lambda^2 - 1)^{1/k} - (\eta'' - 1)^{1/k}}{(\eta'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\eta'' - 1)^{1/k}} \right), \\
 & \quad \left( \frac{(\nu'' + \lambda^2 - 1)^{1/k} - (\nu'' - 1)^{1/k}}{(\nu'' + \lambda^2 - 1)^{1/k} + (\lambda - 1)(\nu'' - 1)^{1/k}} \right) \right\rangle,
 \end{aligned} \tag{29}$$

where

$$\begin{aligned}
 \mu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( \frac{\mu' + \lambda^2 - 1}{\mu' - 1} \right)^{1/C_n^k}, \\
 \mu' &= \prod_{h=1}^k \frac{(\lambda + (1 - \lambda)(1 - \mu_{i_h}))^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} + (\lambda^2 - 1)(1 - \mu_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}{(\lambda + (1 - \lambda)(1 - \mu_{i_h}))^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} - (1 - \mu_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}, \\
 \eta'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( \frac{\eta' + \lambda^2 - 1}{\eta' - 1} \right)^{1/C_n^k}, \\
 \eta' &= \prod_{h=1}^k \frac{(\lambda + (1 - \lambda)\eta_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} + (\lambda^2 - 1)\eta_{i_h}^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}{(\lambda + (1 - \lambda)\eta_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} - \eta_{i_h}^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}, \\
 \nu'' &= \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( \frac{\nu' + \lambda^2 - 1}{\nu' - 1} \right)^{1/C_n^k}, \\
 \nu' &= \prod_{h=1}^k \frac{(\lambda + (1 - \lambda)\nu_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} + (\lambda^2 - 1)\nu_{i_h}^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}{(\lambda + (1 - \lambda)\nu_{i_h})^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)} - \nu_{i_h}^{(nw_{i_h} \omega_{i_h}) / \sum_{t=1}^n (w_t \omega_t)}}.
 \end{aligned}$$

7 If Frank T-norm and T-conorm are applied to Eq. (20), a picture fuzzy Archimedean Frank power weighted MSM (PFAFPWMSM) operator can be obtained as:

$$\begin{aligned}
 & \text{PFAFPWMSM}^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \\
 &= \left\langle \log_\varepsilon \left( 1 + (\varepsilon^{\mu''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right), \right. \\
 & \quad \left. 1 - \log_\varepsilon \left( 1 + (\varepsilon^{1-\eta''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right), \right. \\
 & \quad \left. 1 - \log_\varepsilon \left( 1 + (\varepsilon^{1-\nu''} - 1)^{1/k} / (\varepsilon - 1)^{1/k-1} \right) \right\rangle,
 \end{aligned} \tag{30}$$

where

$$\begin{aligned} \mu''' &= 1 - \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( (\varepsilon - 1) / (\varepsilon^{1 - \mu''} - 1) \right)^{1/C_n^k} \right), \quad \mu'' = \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k \left( (\varepsilon - 1) / (\varepsilon^{\mu^h} - 1) \right) \right), \\ \mu' &= 1 - \log_{\varepsilon} \left( 1 + \left( (\varepsilon^{1 - \mu_h} - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i) \right) / (\varepsilon - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i - 1) \right), \\ \eta''' &= \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( (\varepsilon - 1) / (\varepsilon^{\eta''} - 1) \right)^{1/C_n^k} \right), \quad \eta'' = 1 - \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k \left( (\varepsilon - 1) / (\varepsilon^{1 - \eta^h} - 1) \right) \right), \\ \eta' &= \log_{\varepsilon} \left( 1 + (\varepsilon^{\eta_h} - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i) \right) / (\varepsilon - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i - 1), \\ \nu''' &= \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{1 \leq i_1 < \dots < i_k \leq n} \left( (\varepsilon - 1) / (\varepsilon^{\nu''} - 1) \right)^{1/C_n^k} \right), \quad \nu'' = 1 - \log_{\varepsilon} \left( 1 + (\varepsilon - 1) / \prod_{h=1}^k \left( (\varepsilon - 1) / (\varepsilon^{1 - \nu^h} - 1) \right) \right), \\ \nu' &= \log_{\varepsilon} \left( 1 + (\varepsilon^{\nu_h} - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i) \right) / (\varepsilon - 1)^{(nw_{i_h} \omega_{i_h})} / \sum_{i=1}^n (w_i \omega_i - 1). \end{aligned}$$

#### 4 MADM method

In an MADM problem based on PFNs, decision makers need to select the most appropriate option from a certain number of options. The selection criterion is usually based on a certain number of attributes, whose relative important is measured by weights. The values of the attributes of each option are given by PFNs. The basic components of an MADM problem based on PFNs include a set of options  $O = \{O_1, O_2, \dots, O_m\}$ , a set of attributes  $A = \{A_1, A_2, \dots, A_n\}$ , a vector of weights of attributes  $w = [w_1, w_2, \dots, w_n]$  ( $0 \leq w_1, w_2, \dots, w_n \leq 1, w_1 + w_2 + \dots + w_n = 1$ ), and a picture fuzzy decision matrix  $M = [\alpha_{i,j}]_{m \times n}$  ( $i = 1, 2, \dots, m; j = 1, 2, \dots, n; \alpha_{i,j} = \langle \mu_{i,j}, \eta_{i,j}, \nu_{i,j} \rangle$  is a PFN that is the evaluation value of  $A_j$  of  $O_i$ ). The objective of the MADM problem is to determine the best option from the options in  $O$  on the basis of  $M$  and  $w$ . This objective can be achieved through the following steps:

**Step 1** Normalise the picture fuzzy decision matrix  $M$ . In general, an MADM problem may contain benefit and cost attributes, which have opposite influences on the aggregation result. To unify the influences, a complement rule is usually applied to normalise the PFNs expressing the values of cost attributes. Using this rule, the picture fuzzy decision matrix  $M = [\alpha_{i,j}]_{m \times n}$  is normalised according to the following equation:

$$\begin{aligned} M_N &= [\alpha_{i,j}]_{m \times n} \\ &= \begin{cases} \langle \mu_{i,j}, \eta_{i,j}, \nu_{i,j} \rangle_{m \times n}, & \text{if } A_j \text{ is a benefit attribute} \\ \langle \nu_{i,j}, \eta_{i,j}, \mu_{i,j} \rangle_{m \times n}, & \text{if } A_j \text{ is a cost attribute} \end{cases} \end{aligned} \quad (31)$$

Step 2 Compute the weights of  $\alpha_{i,j}$ . Based on Def.s 9 and 10 and Theorem 5, the weights of  $\alpha_{i,j}$  are calculated using

$$\begin{aligned} \varpi_{i,j} &= (w_j \omega_j) / \sum_{t=1}^n (w_t \omega_t) \\ &= \left( w_j \left( 1 + \sum_{p=1, p \neq j}^n (1 - d(\alpha_{i,j}, \alpha_{i,p})) \right) \right) \\ & / \sum_{t=1}^n \left( w_t \left( 1 + \sum_{q=1, q \neq t}^n (1 - d(\alpha_{i,t}, \alpha_{i,q})) \right) \right), \end{aligned} \tag{32}$$

where  $d(\alpha_{i,j}, \alpha_{i,p})$  and  $d(\alpha_{i,t}, \alpha_{i,q})$  are, respectively, the normalised Hamming distances of  $\alpha_{i,j}$  and  $\alpha_{i,p}$  and  $\alpha_{i,t}$  and  $\alpha_{i,q}$ . They can be calculated according to Def. 5.

Step 3 Compute the summary values of  $\alpha_{i,j}$ . The summary values of  $\alpha_{i,j}$  are calculated via

$$\begin{aligned} \alpha_i &= \langle \mu_i, \eta_i, \nu_i \rangle \\ &= PFAPWMSM^{(k)}(\alpha_{i,1}, \alpha_{i,2}, \dots, \alpha_{i,n}) \end{aligned} \tag{33}$$

where  $PFAPWMSM$  is a specific PFAPWMSM operator, such as the specific operators in Eqs. (24), (28), (29), and (30).

Step 4 Compute the score and accuracy values of  $\alpha_i$ . According to Defs. 2 and 3, the score and accuracy values of  $\alpha_i$  are, respectively, calculated

Step 5 Rank  $O_i$ . According to Def. 4 and the score and accuracy values of  $\alpha_i$ , all  $O_i$  are ranked

Step 6 Determine the best option. The best option is generally the option ranked first

method. The five viable enterprises are  $E_1, E_2, E_3, E_4$ , and  $E_5$ . There are four attributes for decision-making, which are the technical advancement ( $A_1$ ), the potential market ( $A_2$ ), the industrialisation framework, human resources, and financial investments ( $A_3$ ), and the employment formation and progress of science and technology ( $A_4$ ). The weights of these attributes are given by  $w = [0.2, 0.1, 0.3, 0.4]$ . To provide enough freedom in the evaluation of attribute values, PFNs were used. The evaluation results form a picture fuzzy decision matrix  $M = [\alpha_{i,j}]_{5 \times 4}$ , whose elements are listed in Table 2.

According to the conditions above and the proposed MADM method, selection of the best emerging technology enterprise can be carried out as follows:

Step 1 Normalise the picture fuzzy decision matrix  $M$ . Since all of the four attributes are benefit attributes, the normalised picture fuzzy decision matrix  $M_N = M = [\alpha_{i,j}]_{5 \times 4}$

Step 2 Compute the weights of  $\alpha_{i,j}$ . On the basis of Eq. (32), the weights of  $\alpha_{i,j}$  are calculated as:

$$[\varpi_{i,j}]_{5 \times 4} = \begin{bmatrix} 0.2296 & 0.0921 & 0.3342 & 0.3441 \\ 0.2031 & 0.1000 & 0.2874 & 0.4095 \\ 0.1979 & 0.1094 & 0.2549 & 0.4378 \\ 0.1905 & 0.1006 & 0.2968 & 0.4121 \\ 0.1894 & 0.0994 & 0.2917 & 0.4195 \end{bmatrix}.$$

Step 3 Compute the summary values of  $\alpha_{i,j}$ . The summary values of  $\alpha_{i,j}$  can be calculated via Eq. (33). Here the specific operator in Eq. (29) ( $\lambda = 3$  and  $k = 3$ ) is leveraged in Eq. (33). The calculated summary values are as follows:

$$\begin{aligned} \alpha_1 &= \langle 0.3965, 0.4461, 0.1450 \rangle, \alpha_2 = \langle 0.3205, 0.4746, 0.2026 \rangle, \alpha_3 = \langle 0.3366, 0.3767, 0.1287 \rangle, \\ \alpha_4 &= \langle 0.2318, 0.5861, 0.1435 \rangle, \alpha_5 = \langle 0.2895, 0.5283, 0.1641 \rangle \end{aligned}$$

## 5 Example, experiments, and comparisons

### 5.1 Example

An MADM example about selection of the best emerging technology enterprise from five viable enterprises (Jana et al. 2019) is introduced to illustrate the proposed MADM

**Table 2** The elements of the picture fuzzy decision matrix  $M$

	$A_1$	$A_2$	$A_3$	$A_4$
$E_1$	$\langle 0.56, 0.34, 0.10 \rangle$	$\langle 0.90, 0.07, 0.03 \rangle$	$\langle 0.40, 0.33, 0.19 \rangle$	$\langle 0.09, 0.79, 0.03 \rangle$
$E_2$	$\langle 0.70, 0.10, 0.09 \rangle$	$\langle 0.10, 0.66, 0.20 \rangle$	$\langle 0.06, 0.81, 0.12 \rangle$	$\langle 0.72, 0.14, 0.09 \rangle$
$E_3$	$\langle 0.88, 0.09, 0.03 \rangle$	$\langle 0.08, 0.10, 0.06 \rangle$	$\langle 0.05, 0.83, 0.09 \rangle$	$\langle 0.65, 0.25, 0.07 \rangle$
$E_4$	$\langle 0.80, 0.07, 0.04 \rangle$	$\langle 0.70, 0.15, 0.11 \rangle$	$\langle 0.03, 0.88, 0.05 \rangle$	$\langle 0.07, 0.82, 0.05 \rangle$
$E_5$	$\langle 0.85, 0.06, 0.03 \rangle$	$\langle 0.64, 0.07, 0.22 \rangle$	$\langle 0.06, 0.88, 0.05 \rangle$	$\langle 0.13, 0.77, 0.09 \rangle$




Fig. 1 The result of Experiment 1

- Step 4 Compute the score and accuracy values of  $\alpha_i$ .  
 According to Def.s 2 and 3, the score and accuracy values of  $\alpha_i$  are calculated as:  
 $S(\alpha_1) = 0.6258, S(\alpha_2) = 0.5589, S(\alpha_3) = 0.6039,$   
 $S(\alpha_4) = 0.5441, S(\alpha_5) = 0.5627$   
 $A(\alpha_1) = 0.9876, A(\alpha_2) = 0.9978, A(\alpha_3) = 0.8421,$   
 $A(\alpha_4) = 0.9614, A(\alpha_5) = 0.9819$
- Step 5 Rank  $E_i$ . On the basis of Def. 4, all  $E_i$  are ranked as:  
 $E_1 \succ E_3 \succ E_5 \succ E_2 \succ E_4$
- Step 6 Determine the best enterprise. The best enterprise is  $E_1$  as it is ranked first

### 5.2 Experiments

To study the impact of different ATT operations and different argument values on the aggregation result, the following experiments were conducted:

- Experiment 1. This experiment aims to study the impact of different ATT operations on the aggregation result. In this, the specific AOs using different ATT operations in Eqs. (24), (28), (29), and (30) with  $k = 3$  and  $\lambda = \varepsilon = 3$  were, respectively, used in the numerical example. The experiment results, as depicted in Fig. 1, are the score values of  $\alpha_i$  and the rankings of  $E_i$ . As can be seen from Fig. 1, the generated rankings of  $E_i$  under different ATT operations just have difference at the fourth and fifth places, which indicates that different ATT operations have no obvious impact on the aggregation result for the numerical example. Please note that this does not mean that an arbitrary ATT operation can be used in all MADM problems based on PFNs. Whether an ATT operation is suitable for a specific problem should be judged according to the characteristics of the problem.
- Experiment 2. This experiment aims to study the impact of different values of  $k$  on the aggregation result. In this, the presented specific AOs in Eqs. (24), (28), (29), and (30) with  $k = 1, 2, 3, 4$  were, respectively, used in the numerical example. The experiment results, as listed in Table 3, are the score values of  $\alpha_i$  and the rankings of  $E_i$ . From Table 3, the

Table 3 The results of Experiment 2

Specific AO used in Eq. (33)	Value of $k$	Value of $\lambda$ ( $\varepsilon$ )	The calculated score values of the five enterprises					The generated ranking	The best enterprise
			$S(\alpha_1)$	$S(\alpha_2)$	$S(\alpha_3)$	$S(\alpha_4)$	$S(\alpha_5)$		
PFAAPWMSM	1	$\square$	0.6900	0.7210	0.7660	0.6605	0.6742	$E_3 \succ E_2 \succ E_1 \succ E_5 \succ E_4$	$E_3$
	2	$\square$	0.6542	0.6126	0.6735	0.5933	0.6040	$E_3 \succ E_1 \succ E_2 \succ E_5 \succ E_4$	$E_3$
	3	$\square$	0.6370	0.5303	0.5708	0.5502	0.5643	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$
	4	$\square$	0.6170	0.4878	0.5343	0.5151	0.5210	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$
PFAEPWMSM	1	$\square$	0.6752	0.7065	0.7490	0.6352	0.6499	$E_3 \succ E_2 \succ E_1 \succ E_5 \succ E_4$	$E_3$
	2	$\square$	0.6443	0.6171	0.6736	0.5873	0.6009	$E_3 \succ E_1 \succ E_2 \succ E_5 \succ E_4$	$E_3$
	3	$\square$	0.6304	0.5476	0.5914	0.5473	0.5635	$E_1 \succ E_3 \succ E_5 \succ E_2 \succ E_4$	$E_1$
	4	$\square$	0.6111	0.4996	0.5441	0.5128	0.5251	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$
PFAHPWMSM	1	$\lambda = 3$	0.6673	0.6972	0.7386	0.6205	0.6366	$E_3 \succ E_2 \succ E_1 \succ E_5 \succ E_4$	$E_3$
	2	$\lambda = 3$	0.6385	0.6211	0.6747	0.5821	0.5982	$E_3 \succ E_1 \succ E_2 \succ E_5 \succ E_4$	$E_3$
	3	$\lambda = 3$	0.6258	0.5589	0.6039	0.5441	0.5627	$E_1 \succ E_3 \succ E_5 \succ E_2 \succ E_4$	$E_1$
	4	$\lambda = 3$	0.6068	0.5078	0.5509	0.5116	0.5279	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$
PFAPWMSM	1	$\varepsilon = 3$	0.6790	0.7110	0.7536	0.6433	0.6568	$E_3 \succ E_2 \succ E_1 \succ E_5 \succ E_4$	$E_3$
	2	$\varepsilon = 3$	0.6469	0.6172	0.6746	0.5907	0.6027	$E_3 \succ E_1 \succ E_2 \succ E_5 \succ E_4$	$E_3$
	3	$\varepsilon = 3$	0.6324	0.5443	0.5874	0.5490	0.5641	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$
	4	$\varepsilon = 3$	0.6124	0.4971	0.5420	0.5134	0.5238	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1$




Fig. 2 The result of Experiment 3 when PFAHPWMSM is used




Fig. 3 The result of Experiment 3 when PFAFPWMSM is used

score value of each enterprise become smaller and smaller and the generated rankings and best enterprises are different from  $k = 1$  to  $k = 4$  for each AO. This indicates that the argument  $k$  in each AO reflects the risk attitude and the risk attitude changes from optimism to pessimism as  $k$  changes from 1 to 4. It is worth nothing that MSM will reduce to AA and all attributes are independent of each other when  $k = 1$ , MSM will reduce to BM and there are interactions between any two attributes when  $k = 2$ , and MSM will reduce to GA and there are interactions among the four attributes when  $k = 4$ . Thus the order of the optimism of AA, BM, MSM, and GA is  $AA \succ BM \succ MSM \succ GA$  from the results of the experiment. In practical MADM problems, the value of  $k$  should be assigned on the basis of the interactions of attributes.

3 Experiment 3. This experiment aims to study the impact of different values of  $\lambda$  ( $\varepsilon$ ) on the aggregation result. In this, the PFAHPWMSM operator in Eq. (29) with  $k = 3$  and  $0.01 \leq \lambda \leq 20.00$  and the PFAFPWMSM operator in Eq. (30) with  $k = 3$  and  $1.01 \leq \varepsilon \leq 20.00$  were, respectively, leveraged in the numerical example. The experiment results, as depicted in Figs. 2 and 3, are the score values of  $\alpha_i$  and the rankings of  $E_i$ . From Figs. 2 and 3, the scores calculated by the two operators decrease or increase and the generated rankings have minor changes as the values of  $\lambda$  and  $\varepsilon$  gradually increase. This shows that there is no fixed rule for the influence of  $\lambda$  and  $\varepsilon$  on the aggregation result for the numerical example, although the two arguments can be used as risk attitude factors in other practical MADM problems. In general, an appropriate  $\lambda$  ( $\varepsilon$ ) value (e.g.  $\lambda = 1, 2, 3; \varepsilon = 2, 3, 4$ ) is recommended when the PFAHPWMSM (PFAFPWMSM) operator is used.

### 5.3 Comparisons

Representative MADM methods based on AOs of PFNs are the methods presented by Wei (2017), Garg (2017b), Wei (2018), Jana et al. (2019), Wei et al. (2018a), Zhang et al. (2018), and Xu et al. (2019). A qualitative comparison and a quantitative comparison between them and the proposed MADM method are, respectively, carried out to validate the proposed method:

- 1 Qualitative comparison. The qualitative comparison was made through comparing the features of the AOs. For the eight methods, the flexibility in the aggregation of attribute values, the generality in the consideration of attribute interactions, and the capability to reduce the negative impact of biased attribute values are used as the comparison characteristics. The comparison results are listed in Table 4:
  - (a) Flexibility in the aggregation of attribute values. The methods of Wei (2017), Wei et al. (2018a), and Xu et al. (2019) realise the aggregation through the operational rules based on Algebraic T-norm and T-conorm, their flexibility is relatively limited. The methods of Wei (2018), Jana et al. (2019), and Zhang et al. (2018) leverage the Hamacher, Dombi, and Dombi T-norms and T-conorms to carry out their aggregations, respectively. Their flexibility can be seen as moderate because each of the two T-norms and T-conorms can provide a flexible argument for aggregation. The flexibility of the method of Garg (2017b) and the proposed method are

**Table 4** The results of the qualitative comparison

MADM method	Flexibility in aggregation	Generality in consideration of interactions			Capability to reduce influence
		Independent	Two	Multiple	
Wei (2017)	Limited	Yes	No	No	No
Garg (2017b)	Satisfying	Yes	No	No	No
Wei (2018)	Moderate	Yes	No	No	Yes
Jana et al. (2019)	Moderate	Yes	No	No	No
Wei et al. (2018a)	Limited	Yes	Yes	No	No
Zhang et al. (2018)	Moderate	Yes	Yes	No	No
Xu et al. (2019)	Limited	Yes	Yes	Yes	No
The proposed method	Satisfying	Yes	Yes	Yes	Yes

satisfying since their aggregations can be realised via the operational rules based on any family of ATTs.

- (b) Generality in the consideration of attribute interactions. All of the comparison methods can handle the case where all attributes are independent of each other. The methods of Wei et al. (2018a) and Zhang et al. (2018) are applicable for the independence situation and the situation in which there are interactions between any two attributes because of the use of Heronian mean operator in them. The method of Xu et al. (2019) and the proposed method have the most desirable generality in the consideration of attribute interactions because of the use of two all-in-one AOs for capturing relationships, i.e. the Muirhead mean and MSM operators.
  - (c) Capability to reduce the negative impact of biased attribute values. Among all of the eight methods, the method of Wei (2018) and the proposed method have this capability since both of them combine the PA operator.
- 2 Quantitative comparison. The quantitative comparison was made leveraging the numerical examples in Ref. (Jana et al. 2019) (Example 1), Ref. (Wei 2017) (Example 2), and Ref. (Garg 2017b) (Example 3) as the benchmarks. The WA, Hamacher WA, Hamacher WA, Dombi WA, weighted Heronian mean, Dombi weighted Heronian mean, weighted Muirhead mean, and PFAHPWMSM operators were, respectively, used in the methods in Wei (2017), Garg (2017b), Wei (2018), Jana et al. (2019), Wei et al. (2018a), Zhang et al. (2018), Xu et al. (2019), and the present paper. In addition, the same function for calculating the score values was used in all of these methods for to unify the

comparison. The comparison results are listed in Table 5.

From Table 5, the ranking of the proposed method has no significant difference with the rankings of the methods of Jana et al. (2019), Wei et al. (2018a), Zhang et al. (2018), and Xu et al. (2019) for Example 1, with the rankings of the methods of Wei et al. (2018a) and Xu et al. (2019) for Example 2, and with the rankings of all remaining methods for Example 3. This demonstrates that for practical MADM problems based on PFNs, the proposed method is feasible and effective. From Table 4, the method of Xu et al. (2019) and the proposed method are the most similar in characteristics. From Table 5, the rankings of the two methods are exactly identical or have no obvious difference with each other for all examples, which also validates the proposed method. As can also be seen from Table 5, the ranking and best option of the proposed method are fully different with the ranking and best option of the methods of Wei (2017), Garg (2017b), Wei (2018) for Example 1 and of the methods of Wei (2017), Garg (2017b), Wei (2018), Jana et al. (2019), and Zhang et al. (2018) for Example 2. This is because the specific AOs on which these methods are based are different from the specific AO in the proposed method, so do their mathematical properties.

It can be concluded from the results in Table 4 that the proposed MADM method can provide satisfying flexibility in the aggregation of attribute values and the consideration of attribute interactions and concurrently has the capability to reduce the negative impact of biased attribute values on aggregation result. The former characteristic is obvious. But the latter one is not intuitive. To show the difference between a method that has the capability to reduce negative impact and a method that does not have such capability, an additional quantitative comparison was carried out. In this comparison, the weighted Muirhead mean operator in the



**Table 5** The results of the quantitative comparison

Benchmark	MADM method	Value of arguments	The calculated score values of all options					The generated ranking	The best option
			$S(\alpha_1)$	$S(\alpha_2)$	$S(\alpha_3)$	$S(\alpha_4)$	$S(\alpha_5)$		
Example 1	Wei (2017)	$\square$	0.6884	0.7175	0.7587	0.6649	0.6801	$O_3 \succ O_2 \succ O_1 \succ O_5 \succ O_4$	$O_3$
	Garg (2017b), Wei (2018)	$\gamma = 3$	0.6630	0.6929	0.7286	0.6245	0.6419	$O_3 \succ O_2 \succ O_1 \succ O_5 \succ O_4$	$O_3$
	Jana et al. (2019)	$R = 3$	0.8849	0.7894	0.8828	0.8294	0.8618	$O_1 \succ O_3 \succ O_5 \succ O_4 \succ O_2$	$O_1$
	Wei et al. (2018a)	$p = 1, q = 2$	0.9147	0.8880	0.8988	0.8933	0.8964	$O_1 \succ O_3 \succ O_5 \succ O_4 \succ O_2$	$O_1$
	Zhang et al. (2018)	$\lambda = 3, p = 1, q = 2$	0.7965	0.6834	0.7929	0.7295	0.7627	$O_1 \succ O_3 \succ O_5 \succ O_4 \succ O_2$	$O_1$
	Xu et al. (2019)	$P = (1, 2, 3, 0)$	0.6485	0.5948	0.6398	0.5934	0.6052	$O_1 \succ O_3 \succ O_5 \succ O_2 \succ O_4$	$O_1$
	The proposed method	$\lambda = 3, k = 3$	0.6258	0.5589	0.6039	0.5441	0.5627	$O_1 \succ O_3 \succ O_5 \succ O_2 \succ O_4$	$O_1$
Example 2	Wei (2017)	$\square$	0.6896	0.7263	0.7804	0.6837	0.7057	$O_3 \succ O_2 \succ O_5 \succ O_1 \succ O_4$	$O_3$
	Garg (2017b), Wei (2018)	$\gamma = 3$	0.6648	0.6993	0.7478	0.6374	0.6603	$O_3 \succ O_2 \succ O_1 \succ O_5 \succ O_4$	$O_3$
	Jana et al. (2019)	$R = 3$	0.8819	0.8022	0.9117	0.8584	0.9038	$O_3 \succ O_5 \succ O_1 \succ O_4 \succ O_2$	$O_3$
	Wei et al. (2018a)	$p = 1, q = 2$	0.9140	0.8907	0.9040	0.9004	0.9044	$O_1 \succ O_5 \succ O_3 \succ O_4 \succ O_2$	$O_1$
	Zhang et al. (2018)	$\lambda = 3, p = 1, q = 2$	0.7929	0.6990	0.8360	0.7639	0.8222	$O_3 \succ O_5 \succ O_1 \succ O_4 \succ O_2$	$O_3$
	Xu et al. (2019)	$P = (1, 2, 3, 0)$	0.6473	0.5920	0.6433	0.6026	0.6168	$O_1 \succ O_3 \succ O_5 \succ O_4 \succ O_2$	$O_1$
	The proposed method	$\lambda = 3, k = 3$	0.6293	0.5529	0.6058	0.5525	0.5739	$O_1 \succ O_3 \succ O_5 \succ O_2 \succ O_4$	$O_1$
Example 3	Wei (2017)	$\square$	0.4558	0.2868	0.4260	0.6038	$\square$	$O_4 \succ O_1 \succ O_3 \succ O_2$	$O_4$
	Garg (2017b), Wei (2018)	$\gamma = 3$	0.4438	0.2822	0.4183	0.5904	$\square$	$O_4 \succ O_1 \succ O_3 \succ O_2$	$O_4$
	Jana et al. (2019)	$R = 3$	0.5420	0.3301	0.4928	0.6913	$\square$	$O_4 \succ O_1 \succ O_3 \succ O_2$	$O_4$
	Wei et al. (2018a)	$p = 1, q = 2$	0.8098	0.7441	0.8090	0.8800	$\square$	$O_4 \succ O_1 \succ O_3 \succ O_2$	$O_4$
	Zhang et al. (2018)	$\lambda = 3, p = 1, q = 2$	0.3940	0.2220	0.3653	0.5626	$\square$	$O_4 \succ O_1 \succ O_3 \succ O_2$	$O_4$
	Xu et al. (2019)	$P = (1, 2, 3, 0)$	0.3737	0.2796	0.4007	0.5378	$\square$	$O_4 \succ O_3 \succ O_1 \succ O_2$	$O_4$
	The proposed method	$\lambda = 3, k = 3$	0.3701	0.2714	0.3907	0.5191	$\square$	$O_4 \succ O_3 \succ O_1 \succ O_2$	$O_4$


method of Xu et al. (2019) with  $P = (1, 1, 1, 0)$  and the PFAHPWMSM operator in Eq. (29) with  $\lambda = 1$  and  $k = 3$  were selected to solve the MADM problem in Example 1. Please note that the only difference of the two operators under such conditions is whether combines the PA operator (i.e. whether has the capability). Suppose the value of attribute  $A_2$  of enterprise  $E_1$  is a biased value. The degree of positive membership of this value was constantly reduced based on Table 6. It can be guessed that this adjustment will have influence on the ranking of  $E_1$ , which could be dropped from the first enterprise to the last one.

To confirm this conjecture, Table 6 and Fig. 4 list the variation of the places of  $E_1$  in the rankings generated by the method of Xu et al. (2019) and the proposed method.

From Table 6 and Fig. 4, the results of both methods fit the conjecture. This demonstrates the effectiveness of the proposed method indirectly. In addition, the place of  $E_1$  in the rankings of the method of Xu et al. (2019) descends faster than the place of  $E_1$  in the rankings of the proposed method, which shows the capability to reduce the negative impact of biased attribute values intuitively.

**Table 6** The results of the additional quantitative comparison

Value of $A_2$ of $E_1$	Notation	Ranking of the method of Xu et al. (2019)	Ranking of the proposed method
$\langle 0.90, 0.07, 0.03 \rangle$	A	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.80, 0.07, 0.03 \rangle$	B	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.70, 0.07, 0.03 \rangle$	C	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.60, 0.07, 0.03 \rangle$	D	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.50, 0.07, 0.03 \rangle$	E	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.40, 0.07, 0.03 \rangle$	F	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.30, 0.07, 0.03 \rangle$	G	$E_3 \succ E_5 \succ E_1 \succ E_4 \succ E_2$	$E_1 \succ E_3 \succ E_5 \succ E_4 \succ E_2$
$\langle 0.20, 0.07, 0.03 \rangle$	H	$E_3 \succ E_5 \succ E_1 \succ E_4 \succ E_2$	$E_3 \succ E_5 \succ E_1 \succ E_4 \succ E_2$
$\langle 0.10, 0.07, 0.03 \rangle$	I	$E_3 \succ E_5 \succ E_4 \succ E_1 \succ E_2$	$E_3 \succ E_5 \succ E_4 \succ E_1 \succ E_2$
$\langle 0.01, 0.07, 0.03 \rangle$	J	$E_3 \succ E_5 \succ E_4 \succ E_1 \succ E_2$	$E_3 \succ E_5 \succ E_4 \succ E_1 \succ E_2$



**Fig. 4** The change of score value and ranking of  $E_1$  in the additional quantitative comparison

As can be summarised from the comparisons above, the main advantages of the proposed method over the existing methods are providing the flexibility in the aggregation of picture fuzzy information and the consideration of the interactions among attributes and the capability to reduce the negative impact of some biased attribute values.

### 6 Conclusion

In the present paper, a PFAPMSM operator and a PFAPWMSM operator have been proposed to address the MADM problems based on PFNs. The formal definitions and the general expressions of these AOs have been presented. Their properties and special cases have been explored. The specific expressions of the two AOs have been established applying the operational rules of PFNs based on four families of ATTs. Using the established specific PFAPWMSM operators, a method to solve the

PFNs based MADM problems has been proposed. The paper has also introduced a numerical example to illustrate the proposed method and carried out a set of comparisons to demonstrate its features, feasibility, and effectiveness. The comparison results show that the method is feasible and effective that provides the flexibility in aggregation of values of attributes, the generality in consideration of interactions among attributes, and the capability to reduce the negative impact of extreme values of attributes. Future work will focus especially on extending the proposed method from the aspects of dealing with more complex interactions among attributes and applying the method in solving practical MADM problems based on PFNs.

**Funding** This study was funded by the National Natural Science Foundation of PR China (No. 51765012 and No. 61562016) and the Key Laboratory Project of Guangxi (No. GIIP1805).

**Data availability** The Java implementation code of all quantitative comparison methods and related data used to support the findings of this study have been deposited in the GitHub repository (<https://github.com/YuchuChingQin/MADMAOsOfPFNs>).

**Ethical approval** This article does not contain any studies with human participants or animals performed by any of the authors.

### Compliance with ethical standards

**Conflict of interest** Yuchu Qin declares that he has no conflict of interest. Xiaolan Cui declares that she has no conflict of interest. Meifa Huang declares that he has no conflict of interest. Yanru Zhong declares that she has no conflict of interest. Zhemin Tang declares that he has no conflict of interest. Peizhi Shi declares that he has no conflict of interest.

## Appendixes

### A. Proof of Theorem 1

**Proof**

- 1 The following equations are successively derived from the operational rules in Eqs. (1), (2), (3), and (4)

$$\begin{aligned}
 (n\omega_{i_h})\alpha_{i_h} &= \langle \psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})), \varphi^{-1}((n\omega_{i_h})\varphi(\eta_{i_h})), \varphi^{-1}((n\omega_{i_h})\varphi(v_{i_h})) \rangle, \\
 \bigotimes_{h=1}^k ((n\omega_{i_h})\alpha_{i_h}) &= \left\langle \varphi^{-1} \left( \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \right), \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(\eta_{i_h}))) \right), \right. \\
 &\quad \left. \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(v_{i_h}))) \right) \right\rangle, \\
 \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k ((n\omega_{i_h})\alpha_{i_h}) &= \left\langle \psi^{-1} \left( \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \right) \right) \right), \right. \\
 &\quad \left. \varphi^{-1} \left( \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(\eta_{i_h}))) \right) \right) \right), \right. \\
 &\quad \left. \varphi^{-1} \left( \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(v_{i_h}))) \right) \right) \right) \right\rangle, \\
 \frac{1}{C_n^k} \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k ((n\omega_{i_h})\alpha_{i_h}) &= \left\langle \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \right) \right) \right), \right. \\
 &\quad \left. \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(\eta_{i_h}))) \right) \right) \right), \right. \\
 &\quad \left. \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(v_{i_h}))) \right) \right) \right) \right\rangle, \\
 \left( \frac{1}{C_n^k} \bigoplus_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k ((n\omega_{i_h})\alpha_{i_h}) \right)^{1/k} &= \left\langle \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \right) \right) \right) \right) \right), \right. \\
 &\quad \left. \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(\eta_{i_h}))) \right) \right) \right) \right) \right), \right. \\
 &\quad \left. \psi^{-1} \left( \frac{1}{k} \psi \left( \varphi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \varphi \left( \psi^{-1} \left( \sum_{h=1}^k \psi(\varphi^{-1}((n\omega_{i_h})\varphi(v_{i_h}))) \right) \right) \right) \right) \right) \right\rangle.
 \end{aligned}$$

2 The proof of “ $PFAPMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n)$  is a PFN” is equivalent to the proof of “ $0 \leq \mu \leq 1, 0 \leq \eta \leq 1, 0 \leq v \leq 1$ , and  $0 \leq \mu + \eta + v \leq 1$ ”. The following is the proof of “ $0 \leq \mu \leq 1$ ”:

- (i) According to the conditions  $0 \leq \mu_{ih} \leq 1$ ,  $\varphi(t)$  and  $\varphi^{-1}(t)$  are monotonically decreasing, and  $\psi(t)$  and  $\psi^{-1}(t)$  are monotonically increasing, the following inequalities are successively derived

$$\begin{aligned} (n\omega_{i_h})\psi(0) &\leq (n\omega_{i_h})\psi(\mu_{i_h}) \leq (n\omega_{i_h})\psi(1), \\ \psi^{-1}((n\omega_{i_h})\psi(0)) &\leq \psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})) \leq \psi^{-1}((n\omega_{i_h})\psi(1)), \\ \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(0))) &\geq \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \geq \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(1))), \\ \varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(0)))\right) &\leq \varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right) \leq \varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(1)))\right), \\ \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(0)))\right)\right) &\leq \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right) \\ &\leq \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(1)))\right)\right). \end{aligned}$$

- That is  $0 \leq \mu \leq 1$ . “ $0 \leq \eta \leq 1$ ” and “ $0 \leq v \leq 1$ ” can be proved in a similar way.
- (ii) The following is the proof of “ $0 \leq \mu + \eta + v \leq 1$ ”:

Since  $0 \leq \mu \leq 1, 0 \leq \eta \leq 1$ , and  $0 \leq v \leq 1$ , it is obtained that  $0 \leq \mu + \eta + v \leq 3$ . According to the definition of a PFN in Def. 1, it is further obtained that  $\mu_{ih} + \eta_{ih} + v_{ih} \leq 1$  and  $\mu_{ih} \leq 1 - (\eta_{ih} + v_{ih})$ .

According to the conditions  $\varphi(t)$  and  $\varphi^{-1}$

Since

$$\begin{aligned} \sum_{1 \leq i_1 < \dots < i_k \leq n} \sum_{h=1}^k (n\omega_{i_h}) &= n \sum_{1 \leq i_1 < \dots < i_k \leq n} \\ (\omega_{i_1} + \omega_{i_2} + \dots + \omega_{i_k}) &= n \frac{1}{n} = 1. \end{aligned}$$

The following inequalities are successively obtained

$(t)$  are monotonically decreasing,  $\psi(t)$  and  $\psi^{-1}(t)$  are monotonically increasing,  $\psi(1 - t) = \varphi(t)$ ,  $\psi^{-1}(t) = 1 - \varphi^{-1}(t)$ ,  $\varphi(1 - t) = \psi(t)$ , and  $\varphi^{-1}(t) = 1 - \psi^{-1}(t)$ , the following inequalities are successively derived

$$\begin{aligned} \psi(\varphi^{-1}(k\varphi(0))) &\leq \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right) \leq \psi(\varphi^{-1}(k\varphi(1))), \\ \varphi^{-1}(k\varphi(0)) &\leq \psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right) \leq \varphi^{-1}(k\varphi(1)), \\ \varphi(0) &\geq \frac{1}{k} \varphi\left(\psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right)\right) \geq \varphi(1), \\ 0 &\leq \varphi^{-1}\left(\frac{1}{k} \varphi\left(\psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right)\right)\right) \leq 1. \end{aligned}$$



That is,  $\mu + \eta + \nu \leq 1$ .

Since  $0 \leq \mu + \eta + \nu \leq 3$  and  $\mu + \eta + \nu \leq 1$  have been proved,  $0 \leq \mu + \eta + \nu \leq 1$  can be obtained.

### B. Proof of Theorem 2

**Proof** Since  $\alpha_i = \alpha = \langle \mu_\alpha, \eta_\alpha, \nu_\alpha \rangle$  for all  $i = 1, 2, \dots, n$ ,  $d(\alpha_i, \alpha_j) = 0$  for all  $j = 1, 2, \dots, n$  and  $j \neq i$ . Based on the expression of  $T(\alpha_p)$  in Def. 9, it is obtained that

$$n\omega_{i_h} = n \left( \sum_{1 \leq i_1 < \dots < i_k \leq n} \sum_{h=1}^k (1 + T(\alpha_{i_h})) \right) / \sum_{j=1}^n (1 + T(\alpha_j)) = n(1 + (n - 1)) / (n(1 + (n - 1))) = 1.$$

Based on this and Theorem 1, it can be obtained that

$$\mu = \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) \right) \right) \right) \right).$$

Since  $\mu_i = \mu_\alpha$ , then

$$\sum_{h=1}^k \varphi(\mu_{i_h}) = \sum_{h=1}^k \varphi(\mu_\alpha) = k\varphi(\mu_\alpha) \text{ and } \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) = \varphi^{-1}(k\varphi(\mu_\alpha)).$$

The following equations are successively obtained

$$\begin{aligned} \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) \right) &= \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi(\varphi^{-1}(k\varphi(\mu_\alpha))) = \psi(\varphi^{-1}(k\varphi(\mu_\alpha))), \\ \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) \right) \right) &= \psi^{-1}(\psi(\varphi^{-1}(k\varphi(\mu_\alpha)))) = \varphi^{-1}(k\varphi(\mu_\alpha)), \\ \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) \right) \right) \right) &= \frac{1}{k} \varphi(\varphi^{-1}(k\varphi(\mu_\alpha))) = \varphi(\mu_\alpha), \\ \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\mu_{i_h}) \right) \right) \right) \right) \right) &= \varphi^{-1}(\varphi(\mu_\alpha)) = \mu_\alpha, \\ \varphi^{-1} \left( \frac{1}{k} \varphi \left( \psi^{-1} \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi \left( \varphi^{-1} \left( \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \right) \right) \right) \right) \right) &= \mu_\alpha. \end{aligned}$$

This is  $\mu = \mu_\alpha$ . Similarly, it can be proved that  $\eta = \eta_\alpha$  and  $\nu = \nu_\alpha$ . Thus,  $PFAPMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) = \langle \mu_\alpha, \eta_\alpha, \nu_\alpha \rangle$ .

### C. Proof of Theorem 3

**Proof** It can be derived from the definition of the PFAPMSM operator that

$$\begin{aligned} &\left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k \left( \frac{n(1 + T(\beta_{i_h}))}{\sum_{j=1}^n (1 + T(\beta_j))} \beta_{i_h} \right) \right)^{1/k} \\ &= \left( \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \bigotimes_{h=1}^k \left( \frac{n(1 + T(\alpha_{i_h}))}{\sum_{j=1}^n (1 + T(\alpha_j))} \alpha_{i_h} \right) \right)^{1/k}. \end{aligned}$$

According to Def. 9, it is obtained that  $PFAPMSM^{(k)}(\beta_1, \beta_2, \dots, \beta_n) = PFAPMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n)$ .

### D. Proof of Theorem 4

**Proof** Based on Theorem 2, it can be obtained that  $PFAPMSM^{(k)}(\alpha^-, \alpha^-, \dots, \alpha^-) = \alpha\alpha^-$ ,  $PFAPMSM^{(k)}(\alpha^+, \alpha^+, \dots, \alpha^+) = \alpha^+$ , and  $n\zeta_{i_h} = 1$  for both  $PFAPMSM^{(k)}(\alpha^-, \alpha^-, \dots, \alpha^-)$  and  $PFAPMSM^{(k)}(\alpha^+, \alpha^+, \dots, \alpha^+)$ . According to the conditions  $\mu^- \leq \mu_{i_h} \leq \mu^+$ ,  $\varphi(t)$  and  $\varphi^{-1}(t)$  are monotonically decreasing, and  $\psi(t)$  and  $\psi^{-1}(t)$  are monotonically increasing, the following inequalities are successively derived

$$\begin{aligned} \psi(\mu^-) &\leq (n\omega_{i_h})\psi(\mu_{i_h}) \leq \psi(\mu^+), \\ \mu^- &\leq \psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})) \leq \mu^+, \\ k\varphi(\mu^-) &\geq \sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h}))) \geq k\varphi(\mu^+), \\ \varphi^{-1}(k\varphi(\mu^-)) &\leq \varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right) \leq \varphi^{-1}(k\varphi(\mu^+)), \\ \psi(\varphi^{-1}(k\varphi(\mu^-))) &\leq \frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right) \leq \psi(\varphi^{-1}(k\varphi(\mu^+))), \\ \varphi^{-1}(k\varphi(\mu^-)) &\leq \psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right) \leq \varphi^{-1}(k\varphi(\mu^+)), \\ \varphi(\mu^-) &\geq \frac{1}{k} \varphi\left(\psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right)\right) \geq \varphi(\mu^+), \\ \mu^- &\leq \varphi^{-1}\left(\frac{1}{k} \varphi\left(\psi^{-1}\left(\frac{1}{C_n^k} \sum_{1 \leq i_1 < \dots < i_k \leq n} \psi\left(\varphi^{-1}\left(\sum_{h=1}^k \varphi(\psi^{-1}((n\omega_{i_h})\psi(\mu_{i_h})))\right)\right)\right)\right)\right) \leq \mu^+. \end{aligned}$$

That is  $\mu^- \leq \mu \leq \mu^+$ . Similarly, it can be proved that  $\eta^- \geq \eta \geq \eta^+$  and  $v^- \geq v \geq v^+$ .

According to Def. 2, it is obtained that

$$PFAPMSM^{(k)}(\alpha^-, \alpha^-, \dots, \alpha^-) \leq PFAPMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \leq PFAPMSM^{(k)}(\alpha^+, \alpha^+, \dots, \alpha^+)$$

Therefore,  $\alpha^- \leq PFAPMSM^{(k)}(\alpha_1, \alpha_2, \dots, \alpha_n) \leq \alpha^+$ .

## References

- Ai Z, Xu Z (2018) Multiple Definite Integrals of Intuitionistic Fuzzy Calculus and Isomorphic Mappings. *IEEE Trans Fuzzy Syst* 26(2):670–680
- Atanassov KT (1986) Intuitionistic fuzzy sets. *Fuzzy Sets Syst* 20(1):87–96
- Bustince H, Barrenechea E, Pagola M, Fernandez J, Xu Z, Bedregal B, Montero J, Hagrass H, Herrera F, De Baets B (2016) A historical account of types of fuzzy sets and their relationships. *IEEE Trans Fuzzy Syst* 24(1):179–194
- Castillo O, Cervantes L, Melin P, Pedrycz W (2019) A new approach to control of multivariable systems through a hierarchical aggregation of fuzzy controllers. *Granular Comput* 4(1):1–13
- Chen SM, Adam SI (2017) Adaptive fuzzy interpolation based on general representative values of polygonal fuzzy sets and the shift and modification techniques. *Inf Sci* 414:147–157
- Chen SM, Chang CH (2016) Fuzzy multiattribute decision making based on transformation techniques of intuitionistic fuzzy values and intuitionistic fuzzy geometric averaging operators. *Inf Sci* 352:133–149
- Chen SM, Chen SW (2014) Fuzzy forecasting based on two-factors second-order fuzzy-trend logical relationship groups and the probabilities of trends of fuzzy logical relationships. *IEEE Trans Cybern* 45(3):391–403
- Chen SM, Niou SJ (2011) Fuzzy multiple attributes group decision-making based on fuzzy preference relations. *Expert Syst Appl* 38(4):3865–3872
- Chen SM, Ko YK, Chang YC, Pan JS (2009) Weighted fuzzy interpolative reasoning based on weighted increment transformation and weighted ratio transformation techniques. *IEEE Trans Fuzzy Syst* 17(6):1412–1427
- Chen SM, Chu HP, Sheu TW (2012) TAIEX forecasting using fuzzy time series and automatically generated weights of multiple factors. *IEEE Trans Syst Man Cybern Part A Syst Hum* 42(6):1485–1495
- Chen SM, Cheng SH, Lan TC (2016a) Multicriteria decision making based on the TOPSIS method and similarity measures between intuitionistic fuzzy values. *Inf Sci* 367:279–295
- Chen SM, Cheng SH, Chiou CH (2016b) Fuzzy multiattribute group decision making based on intuitionistic fuzzy sets and evidential reasoning methodology. *Inf Fusion* 27:215–227
- Cuong BC (2014) Picture fuzzy sets. *J Comput Sci Cybern* 30(4):409–420
- Dutta P (2019) Multi-criteria decision making under uncertainty via the operations of generalized intuitionistic fuzzy numbers. *Granular Comput*. <https://doi.org/10.1007/s41066-019-00189-z>
- Dutta P, Saikia B (2019) Arithmetic operations on normal semi elliptic intuitionistic fuzzy numbers and their application in decision-making. *Granular Comput*. <https://doi.org/10.1007/s41066-019-00175-5>
- Garg H (2017a) Novel intuitionistic fuzzy decision making method based on an improved operation laws and its application. *Eng Appl Artif Intell* 60:164–174

- Garg H (2017b) Some picture fuzzy aggregation operators and their applications to multicriteria decision-making. *Arab J Sci Eng* 42(12):5275–5290
- Garg H, Kumar K (2018) An advanced study on the similarity measures of intuitionistic fuzzy sets based on the set pair analysis theory and their application in decision making. *Soft Comput* 22(15):4959–4970
- Garg H, Kumar K (2019) Improved possibility degree method for ranking intuitionistic fuzzy numbers and their application in multiattribute decision-making. *Granul Comput* 4(2):237–247
- Hinde CJ, Patching RS, McCoy SA (2007) Inconsistent Intuitionistic Fuzzy Sets and Mass Assignment. In: Atanassov KT, Baczyński M, Drewniak J, Kacprzyk J, Krawczak M, Szmidt E, Wygralak M, Zadrożny S (Eds.) *Developments in Fuzzy Sets, Intuitionistic Fuzzy Sets, Generalized Nets and Related Topics*, vol. I: Foundations, Warsaw: SRI PAS/IBS PAN, pp 133 – 153
- Jamkhaneh EB, Garg H (2018) Some new operations over the generalized intuitionistic fuzzy sets and their application to decision-making process. *Granul Comput* 3(2):111–122
- Jana C, Senapati T, Pal M, Yager RR (2019) Picture fuzzy Dombi aggregation operators: Application to MADM process. *Applied Soft Computing* 74:99–109
- Ju Y, Ju D, Gonzalez EDS, Giannakis M, Wang A (2019) Study of site selection of electric vehicle charging station based on extended GRP method under picture fuzzy environment. *Comput Ind Eng* 135:1271–1285
- Khalil AM, Li SG, Garg H, Li H, Ma S (2019) New operations on interval-valued picture fuzzy set, interval-valued picture fuzzy soft set and their applications. *IEEE Access* 7:51236–51253
- Klement EP, Mesiar R, Pap E (2000) *Triangular Norms*. Springer, Dordrecht
- Kumar K, Garg H (2018) Connection number of set pair analysis based TOPSIS method on intuitionistic fuzzy sets and their application to decision making. *Appl Intell* 48(8):2112–2119
- Lei Q, Xu Z (2016) Chain and substitution rules of intuitionistic fuzzy calculus. *IEEE Trans Fuzzy Syst* 24(3):519–529
- Lei Q, Xu Z (2017) A unification of intuitionistic fuzzy calculus theories based on subtraction derivatives and division derivatives. *IEEE Trans Fuzzy Syst* 25(5):1023–1040
- Liang W, Zhao G, Luo S (2018) An integrated EDAS-ELECTRE method with picture fuzzy information for cleaner production evaluation in gold mines. *IEEE Access* 6:65747–65759
- Liao H, Xu Z (2014) Priorities of intuitionistic fuzzy preference relation based on multiplicative consistency. *IEEE Trans Fuzzy Syst* 22(6):1669–1681
- Liao H, Xu Z, Zeng XJ, Merigó JM (2015) Framework of group decision making with intuitionistic fuzzy preference information. *IEEE Trans Fuzzy Syst* 23(4):1211–1227
- Liu P, Chen SM (2017) Group decision making based on Heronian aggregation operators of intuitionistic fuzzy numbers. *IEEE Trans Cybern* 47(9):2514–2530
- Liu P, Chen SM, Wang P (2018) Multiple-attribute group decision-making based on q-rung orthopair fuzzy power Maclaurin symmetric mean operators. *IEEE Trans Syst Man Cybern Syst*. <https://doi.org/10.1109/tsmc.2018.2852948>
- Liu P, Chen SM, Wang Y (2020) Multiattribute group decision making based on intuitionistic fuzzy partitioned Maclaurin symmetric mean operators. *Inf Sci* 512:830–854
- Liu P, Tang G (2018) Some intuitionistic fuzzy prioritized interactive einstein choquet operators and their application in decision making. *IEEE Access* 6:72357–72371
- Liu P, Liu X (2017) Multiattribute group decision making methods based on linguistic intuitionistic fuzzy power Bonferroni mean operators. *Complexity* 2017:3571459
- Liu P, Liu J (2018) Some q-rung orthopair fuzzy Bonferroni mean operators and their application to multi-attribute group decision making. *Int J Intell Syst* 33(2):315–347
- Liu P, Wang P (2018) Some q-rung orthopair fuzzy aggregation operators and their applications to multiple-attribute decision making. *Int J Intell Syst* 33(2):259–280
- Liu P, Wang P (2019) Multiple-attribute decision making based on Archimedean Bonferroni operators of q-rung orthopair fuzzy numbers. *IEEE Trans Fuzzy Syst* 27(5):834–848
- Maclaurin C (1729) A second letter to Martin Folkes, Esq.; Concerning the roots of equations, with the demonstration of other rules in algebra. *Philos Trans R Soc Lond* 36(1):59–96
- Mahmood T, Ullah K, Khan Q, Jan N (2019) An approach toward decision-making and medical diagnosis problems using the concept of spherical fuzzy sets. *Neural Comput Appl* 31(11):7041–7053
- Qin Y, Qi Q, Scott PJ, Jiang X (2019a) Multi-criteria group decision making based on Archimedean power partitioned Muirhead mean operators of q-rung orthopair fuzzy numbers. *PLoS One* 14(9):e0221759
- Qin Y, Cui X, Huang M, Zhong Y, Tang Z, Shi P (2019b) Archimedean muirhead aggregation operators of q-rung orthopair fuzzy numbers for multicriteria group decision making. *Complexity* 2019:3103741
- Qin Y, Cui X, Huang M, Zhong Y, Tang Z, Shi P (2020a) Linguistic interval-valued intuitionistic fuzzy archimedean power muirhead mean operators for multiattribute group decision-making. *Complexity* 2020:2373762
- Qin Y, Qi Q, Shi P, Scott PJ, Jiang X (2020b) Linguistic interval-valued intuitionistic fuzzy Archimedean prioritised aggregation operators for multi-criteria decision making. *J Intell Fuzzy Syst* 38(4):4643–4666
- Qin Y, Qi Q, Scott PJ, Jiang X (2020c) Multiple criteria decision making based on weighted Archimedean power partitioned Bonferroni aggregation operators of generalised orthopair membership grades. *Soft Comput* 24(16):12329–12355
- Rani P, Jain D, Hooda DS (2019) Extension of intuitionistic fuzzy TODIM technique for multi-criteria decision making method based on shapley weighted divergence measure. *Granul Comput* 4(3):407–420
- Seikh MR, Mandal U (2019) Intuitionistic fuzzy Dombi aggregation operators and their application to multiple attribute decision-making. *Granul Comput*. <https://doi.org/10.1007/s41066-019-00209-y>
- Singh P (2015) Correlation coefficients for picture fuzzy sets. *J Intell Fuzzy Syst* 28(2):591–604
- Son LH (2016) Generalized picture distance measure and applications to picture fuzzy clustering. *Appl Soft Comput* 46:284–295
- Song Y, Fu Q, Wang YF, Wang X (2019) Divergence-based cross entropy and uncertainty measures of Atanassov's intuitionistic fuzzy sets with their application in decision making. *Appl Soft Comput* 84:105703
- Tan A, Shi S, Wu WZ, Li J, Pedrycz W (2020) Granularity and entropy of intuitionistic fuzzy information and their applications. *IEEE Trans Cybern*. <https://doi.org/10.1109/tycy.2020.2973379>
- Teng F, Liu Z, Liu P (2018) Some power Maclaurin symmetric mean aggregation operators based on Pythagorean fuzzy linguistic numbers and their application to group decision making. *Int J Intell Syst* 33(9):1949–1985
- Thong PH (2016) A novel automatic picture fuzzy clustering method based on particle swarm optimization and picture composite cardinality. *Knowl-Based Syst* 109:48–60
- Wang JQ, Zhang HY (2013) Multicriteria decision-making approach based on Atanassov's intuitionistic fuzzy sets with incomplete certain information on weights. *IEEE Trans Fuzzy Syst* 21(3):510–515



- Wang L, Zhang H, Wang J, Li L (2018a) Picture fuzzy normalized projection-based VIKOR method for the risk evaluation of construction project. *Appl Soft Comput* 64:216–226
- Wang L, Peng J, Wang J (2018b) A multi-criteria decision-making framework for risk ranking of energy performance contracting project under picture fuzzy environment. *J Clean Prod* 191:105–118
- Wei G (2017) Picture fuzzy aggregation operators and their application to multiple attribute decision making. *J Intell Fuzzy Syst* 33(2):713–724
- Wei G (2018) Picture fuzzy Hamacher aggregation operators and their application to multiple attribute decision making. *Fundamenta Informaticae* 157(3):271–320
- Wei G, Lu M, Gao H (2018a) Picture fuzzy Heronian mean aggregation operators in multiple attribute decision making. *Int J Knowl-Based Intell Eng Syst* 22(3):167–175
- Wei G, Alsaadi FE, Hayat T, Alsaedi A (2018b) Picture 2-tuple linguistic aggregation operators in multiple attribute decision making. *Soft Comput* 22(3):989–1002
- Xia M, Xu Z, Zhu B (2012) Some issues on intuitionistic fuzzy aggregation operators based on Archimedean t-conorm and t-norm. *Knowl-Based Syst* 31:78–88
- Xu Y, Shang X, Wang J, Zhang R, Li W, Xing Y (2019) A method to multi-attribute decision making with picture fuzzy information based on Muirhead mean. *J Intell Fuzzy Syst* 36(4):3833–3849
- Xu Z, Yager RR (2011) Intuitionistic fuzzy Bonferroni means. *IEEE Trans Syst Man Cybern-Part B Cybern* 41(2):568–578
- Yager RR (2001) The power average operator. *IEEE Trans Syst Man Cybern-Part A Syst Hum* 31(6):724–731
- Zadeh LA (1965) Fuzzy sets. *Inf Control* 8(3):338–353
- Zeng S, Chen SM, Kuo LW (2019) Multiattribute decision making based on novel score function of intuitionistic fuzzy values and modified VIKOR method. *Inf Sci* 488:76–92
- Zhang FW, Huang WW, Sun J, Liu ZD, Zhu YH, Li KT, Xu SH, Li Q (2019) Generalized fuzzy additive operators on intuitionistic fuzzy sets and interval-valued intuitionistic fuzzy sets and their application. *IEEE Access* 7:45734–45743
- Zhang H, Zhang R, Huang H, Wang J (2018) Some picture fuzzy Dombi Heronian mean operators with their application to multi-attribute decision-making. *Symmetry* 10(11):593
- Zhang Z, Pedrycz W (2017a) Models of mathematical programming for intuitionistic multiplicative preference relations. *IEEE Trans Fuzzy Syst* 25(4):945–957
- Zhang Z, Pedrycz W (2017b) Intuitionistic multiplicative group analytic hierarchy process and its use in multicriteria group decision-making. *IEEE Trans Cybern* 48(7):1950–1962
- Zhang Z, Pedrycz W (2018) Goal programming approaches to managing consistency and consensus for intuitionistic multiplicative preference relations in group decision making. *IEEE Trans Fuzzy Syst* 26(6):3261–3275
- Zhang Z, Chen SM, Wang C (2020) Group decision making with incomplete intuitionistic multiplicative preference relations. *Inf Sci* 516:560–571
- Zhong Y, Guo X, Gao H, Qin Y, Huang M, Luo X (2019a) A new multi-criteria decision-making method based on Pythagorean hesitant fuzzy Archimedean Muirhead mean operators. *J Intell Fuzzy Syst* 37(4):5551–5571
- Zhong Y, Gao H, Guo X, Qin Y, Huang M, Luo X (2019b) Dombi power partitioned Heronian mean operators of q-rung orthopair fuzzy numbers for multiple attribute group decision making. *PLoS One* 14(10):e0222007

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.