



Extreme Rainfall Events over Accra, Ghana, in Recent Years

Michael Padi^{1,2} · Bennet Atsu K. Foli^{3,4}  · Ebenezer S. Nyadjro⁵ · Kwadwo Owusu^{2,6} · George Wiafe^{3,4}

Received: 30 June 2021 / Revised: 7 October 2021 / Accepted: 27 October 2021 / Published online: 9 November 2021
© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2021

Abstract

This study examines the recent changes in extreme rainfall events over Accra, Ghana. For this study, an extreme rainfall event is defined as a day with rainfall equal to or exceeding the 1980–2019 95th percentile. Knowing extreme rainfall events help to identify the years with the likelihood of rainfall-related disasters in Accra. In addition, it helps to identify the years with the likelihood of drought or severe dryness which are critical for the livelihoods and economic activities of the people. The study used rainfall data from rain gauge for Accra and satellite-derived winds at the 850 hPa level over southern Ghana from 1980 to 2019. It compares these climatic parameters for both pre-2000 and post-2000 to find out the changes that have occurred throughout the study period. Results show that the frequency and magnitude of extreme rainfall have generally increased during the post-2000 period than during the pre-2000 period, causing increases in mortalities and damages to properties. Seasonally, extreme rainfall events were most intense in July during the pre-2000 period but have changed to June during the post-2000 period. Notably, more intense rainfall events have also occurred during post-2000 winter than pre-2000 winter, consistent with increased warming in the study area. Monthly mean meridional winds at the 850 hPa level were stronger (weaker) in the northerly (southerly) direction during the pre-2000 period but have changed to be stronger (weaker) in the southerly (northerly) direction during the post-2000 period.

Keywords Extreme rainfall · 850 hPa Meridional winds · Climate Change · Ghana · Accra

1 Introduction

Precipitation is one of the atmospheric parameters that can be used to assess climate change [36]. It can cause worldwide natural hazards when it exceeds tolerable levels [19]. It is expected that warming climates will significantly induce changes in the distribution of extreme weather events across the globe, thus affecting precipitation patterns. Such usually

devastating events include severe drought, extreme rainfall, and flooding [27]. The early part of this twenty-first century saw several research publications on the impacts of climate change and vulnerabilities on humans and natural resources by the Intergovernmental Panel on Climate Change (IPCC) [16]. Extreme rainfall has caused interruptions to power supply, destructions of farm produce and infrastructure, displacement of communities as well as the outbreak of diseases across many parts of the world. Flooding has led to food insecurity in Asia [14] and needs innovative developments to cope. According to estimates by Insurance Companies, worldwide losses of property recorded through severe storms, flooding, droughts, as well as climate-related fires reached \$60 billion in 1996 and \$89 billion in 1998 [8].

Food security in West Africa heavily depends on rainfall. However, in extreme rainfall, so much havoc is caused, leading to food scarcity. Several communities in Nigeria for example have recently experienced devastating flooding which was attributed to the combined effects of climate change and the neglect of implementing building laws which lead to poor community planning [20]. The magnitude and frequency of extreme rainfall due to climate change have

✉ Bennet Atsu K. Foli
bentsufo@yahoo.co.uk; bafoli@ug.edu.gh

¹ Ghana Meteorological Agency (GMet), Accra, Ghana

² Center for Climate Change and Sustainability Studies (C3SS), University of Ghana, Legon, Ghana

³ Department of Marine and Fisheries Sciences, University of Ghana, Legon, Ghana

⁴ Global Monitoring for Environment and Security (GMES) and Africa Programme, University of Ghana, Legon, Ghana

⁵ Northern Gulf Institute, Mississippi State University, Stennis Space Center, MS, USA

⁶ Department of Geography and Resource Development, University of Ghana, Legon, Ghana

increased, warranting intense studies to understand the factors that drive such occurrences [13, 26]. The chronology of rainfall, such as consecutive dry or wet days, is of high importance for most activities especially those related to water resources. Some societies have tried to reduce the consequences of extreme rainfall events through the improvement in technology and social organization [22]. The consequences include major loss of population, out-migration, and even the collapse of societies. Ogega et al. [29] assessed the performance of a model run to investigate heavy precipitation events over East Africa concerning climate change and concluded that there is a possibility of frequent occurrence of extreme precipitation by the end of this twenty-first century. It has therefore become necessary for similar studies to be conducted in Ghana to examine the potential impacts that changes in the climate will have, since the perennial flooding in Accra, the largest city and capital of Ghana, usually trap people at unwanted locations whenever they occur [44].

Increased urbanization in Accra [11] has exacerbated the effects of climate change and this is evident through the perennial flooding of the city. Changes in rainfall patterns over Accra have been attributed to climate change by several authors (e.g. [1, 4, 10, 46]). The strength of winds that accompany rainstorms have increased in recent times, becoming more destructive over Accra [35] and its environs. Consequently, in this study, we examine the extreme rainfall events that have been occurring over Accra in recent years. Our results indicate that the frequency and intensity of extreme rainfall events as well as annual rainfall amounts have increased during the post-2000 compared to the pre-2000 climatic era. Meanwhile, mean monthly temperatures for the post-2000 era were also estimated to be above the long term mean of 26 °C whilst the pre-2000 era was mostly below the mean.

During the pre-2000 era, the heaviest rainfall was reported to have occurred on 3rd July 1995. According to the United Nations Department of Humanitarian Affairs (UNDHA) [48], a heavy rainstorm entered into Ghana from the east and affected almost the whole country overnight and exited the country in the morning. It rendered the city of Accra to severe flooding. Official reports indicated that more than 1,500 people were rendered homeless through that severe rainfall event. Other amenities like telecommunication and electrical systems were severely damaged for almost 4 days. The report indicated that there was a lack of preparedness which hindered the provision of relief to the affected people. Meanwhile the post-2000 era recorded its highest rainfall over Accra on 3rd June 2015. The World Bank Group [49], described the 3rd June 2015 rain disaster as the worst flood in recent history of Accra. Over 154 Ghanaians were reported to have been killed by the downpour and its associated fire that occurred in the city [15, 31]. This study therefore provides insight into the occurrence of such

extreme rainfall events and some associated causative factors, so as to aid in advance prediction of such events for planning and preparedness purposes.


2 Effect of Meridional Winds on Rainfall over Ghana

Due to the geographical location of Ghana (Fig. 1), the country is affected by two distinct air masses: the dry continental air mass which brings about the harmattan or the dry season and the maritime air mass which is moist and brings about rain and active storms [47]. Wind directions in this study conform to meteorological conventions. The continental air mass is usually northerly in direction, but mostly northeasterly whilst the maritime air mass is southerly, usually southwesterly in direction [17]. However, these air masses are controlled by high-pressure systems that develop on both sides of the hemisphere, either on the continent of Africa or around the continent [41]. Ghana is usually under the influence of a low-pressure usually termed as the equatorial trough because it is not too far from the equator. As a Meteorological norm, winds blow from the high-pressure zone to the low-pressure zone and that is why two different air masses alternatively affect the country.

Ghana is divided into two weather and climate zones which comprise the northern sector and the southern sector [34]. The southern sector has two rainy seasons; the major and minor rainy seasons, whilst the northern sector has only one rainy season [32, 34]. These rainy seasons are then followed by the dry harmattan season, which starts in November and eventually affect the whole country till the beginning of the rains in March [7]. The major rainy season in the south starts from March to July with a break in August which is usually termed as the “little dry season” [2]. Mostly, farmers take advantage of that period to harvest their crops and dry cereals and prepare their lands for the minor rainy season. The minor rainy season then starts in September and ends in November. However, the northern rainy season is prolonged and starts from May and ends in October to give way to the dry harmattan season. There had, however, been times when the harmattan resurfaces during the latter part of March or early April [25].

In weather forecasting, the use of meridional (v) and zonal (u)-winds is very important and play significant roles in ensemble forecasts and ocean models [17, 38]. At the beginning of the major rainy season, winds change from the northerly direction to become southerly and create perturbations in the atmosphere for massive cloud developments that result in heavy rains [34]. These rains are very erratic from the beginning of the rainy season, they are mostly accompanied by very strong winds but as the season grows, the windy conditions reduce while the rain amounts increase. There is

Fig. 1 Map of Ghana and Accra, showing the study area



a clear indication that both northerly and southerly meridional winds strongly influence rainfall over West Africa in general and has a predictive ability that can be harnessed for forecasting purposes [39].

Meteorologists have used satellite and model-derived meridional winds at the 850 hPa for forecasting and it has worked well over the years [43]. Winds at this level influence rainfall over West Africa as they control moisture distribution over the sub-region and also have potential for forecasting. Pineda & Willems [37] used the US National Centers for Environmental Prediction/ National Center for Atmospheric Research (NCEP/NCAR) reanalysis wind data at the 850 hPa level to determine drivers of extreme rainfall with complex terrains extreme value (EV) models. Meridional wind components have been the main drivers of tropical extreme rainfall events whilst zonal winds have been responsible for the extra-tropics [37]. Boos & Kuang [6], observed that zonal winds break down towards the east of India and become meridional due to the effect of the Tibetan Plateau and produce heavy rains in the summer season. The use of meridional winds, even though widely used in other places around the world, has not been well investigated for use in Ghana with regards to analyzing extreme rainfall events. This study therefore seeks to examine the role that meridional winds play in generating extreme rainfall events over Accra.

According to de Boer et al. [12], meridional wind flow causes overturning in the Atlantic Ocean and are the main features of the global overturning circulation which give rise to massive cloud developments. The ocean would therefore not support deep convection if there were no

winds or vertical diffusivity that are high enough to initiate overturning in the atmosphere. There had been suggestions that the fluctuations in the Southern Hemisphere winds can change the pathways of the Atlantic meridional overturning circulation as well as the properties of the water and the associated heat and transportation of freshwater [42]. Wind patterns have changed recently over southern Ghana due to climate change [4]. There is therefore the need to study the recent wind patterns that initiate perturbations for atmospheric overturning to take place along the coast of Ghana.

3 Data and Methods

Rainfall and meridional wind data for 40 years, 1980–2019, have been chosen for this study and centering on the year 2000 to find out how climate change has affected extreme rainfall events over Accra. The year 2000 has been used in this study as a baseline to divide two perceived climatic regimes based on observations, similarly as applied by some previous studies for examining a possible shift in rainfall trends in the West African region [32–34]. This is based on the fact that from the year 2000, the frequency of extreme rainfalls was observed to have increased. Consequently, for our study, two epochs are defined, 1980–1999 and 2000–2019, to enable us to examine the changes in the rainfall and wind patterns on either side of the year 2000.

3.1 Study Area

The study area for this work is Accra, the capital city of Ghana. It is located along the Guinea Coast of West Africa, approximately between 0.3°W to 0.1°W and 5.5°N to 5.7°N, at an approximate altitude of 68 m above mean sea-level (Fig. 1). However, a wider area covering 3.0°W to 3.0°E and 0.0° to 8.0°N, which forms part of southern Ghana has also been considered. In our analyses, there are also instances where a much larger area, comprising of the North and South Atlantic Oceans as well as portions of the western parts of the Indian Ocean are considered for their teleconnection impacts on western Africa. The coast of Ghana can be divided into three geographical zones, the Western, Central, and the Eastern coasts. The Western coastline is about 95 km long and ends at the border with Cote d'Ivoire while the Central Coast is about 321 km. The Eastern coast, where Accra belongs, is about 149 km [5] and ends at the border with the Republic of Togo. The study area is urbanized and also noted for commercial fishing activities at the local, coastal communities. Accra is a low-lying coastal city and is mostly affected by flooding whenever it rains heavily [3, 31].

The Gulf of Guinea, which is located to the south of the country, has been the main source of moisture supply for massive cloud developments and rain formation over Ghana. For this reason, it has become interesting to investigate wind components at the atmospheric boundary layer (i.e., 850 hPa level). This is to examine how these winds influence the moisture content of the airmass that affects the coast of Ghana and generate extreme rainfall occurrences.

3.2 Data

In this study, we used quality-controlled homogeneous daily rainfall data collected with rain gauge from the Ghana Meteorological Agency (GMet) sampling station at the Kotoka International Airport Meteorological Office (KIAMO), Accra, Ghana. The data covered the period 1980–2019.

Daily 2.5° × 2.5° gridded 850 hPa meridional wind, as well as surface temperature reanalysis data were obtained from the National Oceanic and Atmospheric Administration (NOAA) National Center for Environmental Prediction [21]. Top of atmosphere (TOA) brightness temperature data on a 0.07° × 0.07° grid, used as a proxy to track storms in the study area, were obtained from NOAA National Centers for Environmental Information [23].

3.3 Methods

In this study, we defined extreme rainfall events as done in Seleshi & Camberlin [40] and Krishnamurthy et al. [24], days with rainfall equal to or exceeding the 1980–2019 95th percentile (i.e., 44.40 mm). Effectively, only days with rainfall

in the top 5% of all days with rainfall (i.e., 142 days) are considered. We examine two components of the extreme events; (a) frequency, defined as the number of days with extreme rainfall events and (b) intensity, defined as the daily mean of extreme rainfall events. A test of significance between the rainfall and meridional winds has been investigated using Student's *t*-test. The *t*-test has been used to compare the two climatic means, those of the pre-2000s and the post-2000s to study the differences that exist between their rainfall distributions. The mean (\bar{X}) is given by the formula, $\bar{X} = \frac{\sum x_i}{n}$, where n is the number of samples and x_i is the n^{th} term of the distribution. The sample standard deviation (σ_s) is estimated with Eq. 1 (Eq. 1) with a degree of freedom, $df = n - 1$. The *t*-statistic is thus computed using Eq. 2.

$$\sigma_s = \sqrt{\frac{\sum (X_i - \bar{X})^2}{n - 1}} \quad (1)$$

$$t = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{(n_1 - 1)\sigma_1^2 + (n_2 - 1)\sigma_2^2}{n_1 + n_2 - 2}} \times \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \quad (2)$$

The *p*-value is used to examine the statistical significance of the differences between the parameters during the two climatic periods examined.

4 Results and Discussions

The amount of rainfall over Accra has generally recorded an increasing trend since 1980 (Fig. 2). Rainfalls have been generally higher during post-2000 than they were during pre-2000 (Table 1). The *t*-test conducted at 95% confidence level between the two climatic periods indicates that all the respective months had *p*-values higher than 0.05, suggesting




Fig. 2 Annual rainfall totals (mm) for Accra

Table 1 Mean, standard deviation, and the *p*-values for *t*-test of the monthly rainfall totals (mm) in Accra for the pre-2000 and post-2000

Monthly rainfall totals over Accra					
Months	Mean (mm)		Standard deviation (mm)		<i>p</i> -value <i>t</i> -test
	1980–1999	2000–2019	1980–1999	2000–2019	
Jan	5.3	15.4	7.3	26.3	0.1126
Feb	13.0	31.0	18.1	43.1	0.09816
Mar	54.9	59.2	48.4	48.5	0.7799
Apr	84.7	85.4	65.9	56.8	0.9731
May	146.8	153.0	70.4	80.0	0.7952
Jun	158.3	196.2	94.2	91.1	0.2033
Jul	67.6	54.0	74.1	37.3	0.4694
Aug	23.9	23.6	26.1	26.9	0.9744
Sep	49.4	52.1	62.2	32.7	0.8655
Oct	65.9	84.8	47.3	54.5	0.2488
Nov	27.7	36.4	22.6	29.7	0.3088
Dec	17.6	33.6	24.5	37.3	0.1185

that the difference in monthly total rainfall between pre-2000 and post-2000 for Accra is not statistically different. Nevertheless, the overall results show that the month of the most intense rainfall events has shifted from July during the pre-2000 era to June during the post-2000 era (Fig. 3a). The mean rainfall had been higher for June during post-2000 than pre-2000. Although January had the most intense rainfall event (112.5 mm) during post-2000, this was a one-time event and could be considered an anomaly (Fig. 3). The shift in rainfall patterns in general can lead to crop failures. Additionally, it affects water resource sharing in the region as well as operation of infrastructure such as dams, which contribute to hydro-electric generation and irrigation. Thus, information on the variability (shift) in extreme rainfall is essential for use in planning towards the risks as well as the opportunities associated with such climate hazards.

The frequency of extreme rainfall events in the post-2000 was higher than in the pre-2000 (Fig. 3b). Notably, there have been more extreme rainfall events during post-2000 winter than pre-2000 winter. For example, there were respectively 3, 1, and 6 extreme rainfall events during December, January, and February of post-2000 compared to none during December and January and 1 during February of pre-2000 (Fig. 3b). Consequently, mortalities associated with extreme rainfall have also increased during the post-2000 over Accra than the pre-2000 with the increased mortalities (Table 2) also aided by land surface changes, human behaviors, and congestions [15, 48].

There had also been differences in the meridional wind pattern at the 850 hPa level between the two climatic periods for all the months, with *p*-values lower than 0.05, except for June, July, and August which had the *p*-values of more




Fig. 3 a Monthly mean rainfall intensity, and standard deviation (mm) and b monthly frequency of extreme rain events over Accra during pre-2000 (black line) and post-2000 (red line)

than 0.05 between the pre-2000 and the post-2000 (Table 3). Northerly winds (negative meridional winds) were stronger during the pre-2000 than the post-2000 whilst southerly winds (positive meridional winds) were stronger during the post-2000 than the pre-2000 (Table 3).

4.1 Temperature Changes

Monthly mean surface temperature distribution over southern Ghana has been estimated to have a mean value of 26 °C for the study period, 1980 to 2019 (Fig. 4a). However, a trendline indicates that the pre-2000 surface temperature was below the average while the post-2000 surface temperature was above the average. This suggests a possible effect of climate change and warming over southern Ghana. Comparing temperatures of both periods, the pre-2000 had most of its months with mean temperatures below 26 °C while the post-2000 period only had fewer months below this value. The trend in observed temperatures over the periods positively agrees with the trend in meridional winds at the 850 hPa level (Fig. 4a and 4b). This suggests the influence of changing temperatures on meridional winds, which in turn affects rainfall distribution pattern and intensity in the

Table 2 Effects of reported extreme rainfall occurrence in Accra pre- and post-2000 [30]

Date of extreme rain	Effect/impact	Period
July 5, 1995	Storm caused flooding in low areas in the Accra metropolis; commuters and vehicles and electric substation were affected and resulted in power cuts	Pre-2000
June 13, 1997	Hours of an intermittent downpour for two continuous days in Accra caused floods	Pre-2000
June 28, 2001	Over 300,000 people affected in an early morning downpour that submerged portions of the city	Post-2000
May 5, 2010	Central Accra and other areas deeply submerged in water after 2 h of stormy rains	Post-2000
June 22, 2010	Death toll of 35 recorded after one of the nation's worst flood disasters	Post-2000
February 24, 2011	A downpour wreaked extensive havoc on properties in most parts of Accra and some of its surrounding communities	Post-2000
November 1, 2011	43,087 people affected, with 14 deaths recorded after a downpour occurred in Accra	Post-2000
May 31, 2013	Heavy rains caused flooding in many parts of Accra	Post-2000
June 3, 2015	Over 160 lives lost with property after a heavy downpour in Accra caused flooding, triggering a fuel station explosion	Post-2000

Table 3 Mean, standard deviation, and the p -values for t -test of the mean monthly meridional winds (ms^{-1}) at the 850 hPa level for southern Ghana for the pre-2000 and post-2000

Months	Monthly meridional winds over southern Ghana				
	Mean (m/s)		Standard deviation (m/s)		p -value
	1980–1999	2000–2019	1980–1999	2000–2019	t -test
Jan	−1.66025	−0.56085	0.727699	0.667922	0.00001447
Feb	−1.17115	0.02745	0.665026	0.755117	0.000004966
Mar	0.14685	1.08375	0.594539	0.57255	0.0000105
Apr	1.02225	1.65730	0.554261	0.260709	0.00008076
May	1.23060	1.69675	0.45064	0.396349	0.001315
Jun	0.60775	0.90605	0.678951	0.725613	0.1874
Jul	−0.09870	−0.22305	0.726778	0.682128	0.5802
Aug	−0.63430	−0.58645	0.59137	0.51057	0.7857
Sep	−0.97085	−0.61615	0.356568	0.464909	0.01035
Oct	−1.25130	−0.57015	0.56598	0.500674	0.0002621
Nov	−1.53590	−0.78105	0.491086	0.57316	0.00007056
Dec	−1.61855	−0.89095	0.382913	0.513702	0.00001253

region. Studies (e.g. [9, 18] have shown positive relationships between temperature and meridional winds.


4.2 Meridional Wind Changes

Monthly mean meridional winds at the 850 hPa level during the pre-2000 period were dominated by strong northerly winds while the post-2000 period was dominated by strong southerly winds (Fig. 4b). This suggests that the northerly winds became gradually weakened during the late 1990s and southerly winds started to become stronger during the early 2000s. The strongest mean monthly meridional wind recorded in the pre-2000 was a northerly wind with a speed of 3.6 m/s in January 1983 (Fig. 4b). That year experienced the poorest annual rainfall in the history of Accra (Fig. 2). Conversely, the strongest mean monthly meridional wind in the post-2000 was southerly

with a speed of 2.7 m/s, recorded in May 2008 and 2019 (Fig. 4b). Notably, the year 2008 recorded the highest annual rainfall in Accra during the study period (Fig. 2). There had been no month in the pre-2000 with a mean southerly wind of more than 2.0 m/s whilst there had also not been a month in the post-2000 with a northerly mean wind speed of up to 2.0 m/s (Fig. 4b).

Changes in the pattern of meridional winds along the Gulf of Guinea coast affects the position of the intertropical convergence zone (ITCZ), thus affecting the distribution of rain. According to [28], the ITCZ lies between the transition between the northerly wind and the southerly wind and migrates from around latitude 9°N in January and 20°N in August. The location of this ITCZ is a position on the surface of the earth where maximum heating and convective activities take place, therefore, rainfall is usually abundant and torrent.

Fig. 4 Monthly mean **a** surface temperature ($^{\circ}\text{C}$) and **b** meridional winds (ms^{-1}) over southern Ghana. Dashed red lines show linear trend. Dashed blue line in **a** marks the mean surface temperature (i.e., 26°C)




4.3 Pre-2000 Extreme Rainfalls over Accra

The highest rainfall during the pre-2000 over Accra is observed to occur on 3rd July 1995 with an amount of 243.9 mm (figure not shown). Satellite information on meridional winds at the 850 hPa level on that 3rd July 1995 indicated strong southerly winds of about 9 m/s over the South Atlantic Ocean (Fig. 5a). This wind strength had the potential to transport enough moisture from the sea towards the coast of Ghana. They however weakened as they approached the coast of Ghana to deposit their moisture content as rain. An opposing northerly wind, therefore, prevailed over the Sahel with about the same strength, 9 m/s and weakened to about 2 m/s as they reached the coastal sector of Ghana (Fig. 5a). This created a wind convergence zone. The maritime winds were restricted from moving further inland into the West African sub-region by the opposing continental winds. The two air masses mixed and were lifted

for massive condensation to occur and form massive clouds. The storm that was created by this scenario stagnated over the Gulf of Guinea Coast and produced heavy rainfall for a long period. Satellite imageries indicated that the storm stayed over the whole country for more than 12 h (Fig. 5).

This storm was sighted on satellite over Nigeria at 1130 UTC on 3rd July 1995 (Fig. 5b) propagating westward, and 6 h later, it was located over the eastern half of Ghana. As at 2330 UTC, the storm occupied almost the whole of Ghana (Fig. 5c) and start to exit the country into La Cote d'Ivoire around 0530 UTC the following day, on 4th July 1995 (Fig. 5d). This particular storm is quite spectacular; it developed an “eye-like” structure, a phenomenon usually associated with tropical cyclones, which is very rare over the West African sub-region. The “eye” can be visible at 1130 UTC when it was over Nigeria (Fig. 5b) and also when it was exiting Ghana into Cote d'Ivoire at 0530 UTC on 4th July 1995 (Fig. 5d). It took the storm about 6 h to propagate

Fig. 5 **a** 850 hPa meridional winds (ms^{-1}) averaged for 2–3 July 1995; **b** top of atmosphere (TOA) brightness temperature (K) at 1130 UTC on 3 July 1995, **c** at 2330 UTC on 3 July 1995, and **d** at 0530 UTC on 4 July 1995




from the western borders of Nigeria to Ghana and stayed over Ghana for about 12 h. The shape of the storm has a spiral nature which led to the development of the “eye-like” structure and the stagnation.

In 1983, the lowest extreme monthly rainfall in the pre-2000 over Accra was recorded with an amount of 46.3 mm. Meanwhile, it has also been the year with the poorest record

of rainfall in Accra during the study period (Fig. 2). That highest amount of rain for that year was recorded on the 19th of June (figure not shown). From NOAA satellite information, the entire country was occupied by northerly meridional winds at the 850 hPa level as well as over the South Atlantic Ocean, the moisture source for West Africa (Fig. 6a). Strong northerly winds of up to 10 m/s were

Fig. 6 **a** 850 hPa meridional winds (ms^{-1}) averaged for 18–19 June 1983; **b** TOA brightness temperature (K) at 0530 UTC on 20 June 1983



observed over the South Atlantic Ocean with just a few portions experiencing weak “pockets” of southerly winds. The Sahel was therefore dominated by strong northerlies up to 6 m/s which brought drier winds to Ghana and made it impossible to sustain massive cloud developments. Considering annual total rainfall amounts, the year 1983 was associated with the lowest rainfall, dry spells, drought, and very hot weather conditions with famine in Ghana [45]. A weak storm was spotted on the Meteosat-2 IR satellite imagery on the 20th June 1983 at around 0530 UTC which produced this highest rainfall for the year. The storm, though propagated from the east of the country to the west, the center was slightly north of the coastline (Fig. 6b).

4.4 Post-2000 Extreme Rainfall over Accra

The highest rainfall event in the post-2000 occurred on June 3, 2015, over Accra, with a recorded amount of 212.8 mm. Satellite information from NOAA indicated that strong southerly meridional winds of about 15 m/s were observed over the South Atlantic Ocean and as they approached the

coast of Ghana, they weakened to about 3 m/s (Fig. 7a). This is an indication of sufficient moisture transport from the Atlantic to the coast of West Africa. As at 0900 UTC on the 3rd of June 2015, a storm developed just along the coastline, stretching from the west of Nigeria to Togo and moved westward. Three hours later, the storm entered Ghana over the southeast and started raining heavily (Fig. 7).

The lowest extreme rainfall for the post-2000 was recorded on 24th June 2012 with an amount of 47.9 mm. At that time, satellite information from NOAA indicated that though southerly winds prevailed over the south Atlantic Ocean, the speeds were not strong, about 6 m/s, with a large “pool” of strong northerly winds prevailing over the Gulf of Guinea and extending far into North Africa (Fig. 8a). For this reason, the convergence zone between the southerlies and the northerlies occurs offshore, far away over the sea. As a result, massive cloudiness did not occur close to the land, the northerly winds, therefore, prevented sufficient moisture from reaching the coast of West Africa. Nonetheless, on the 24th of June 2012, the highest rainfall amount of the year was recorded from a weak storm cell that tracked across the

Fig. 7 **a** 850 hPa meridional winds (ms^{-1}) averaged for 2–3 June 2015; **b** TOA brightness temperature (K) at 2100 UTC on 3 June 2015





Fig. 8 **a** 850 hPa meridional winds (ms^{-1}) averaged for 23–24 June 2012; **b** TOA brightness temperature (K) at 0900 UTC on 24 June 2012



inland areas of the country with a small storm cell surviving along the coast (Fig. 8b).

From the results, both southerly winds and northerly winds at the 850 hPa level have their peculiar ways of influencing extreme rainfalls over Accra. When strong southerly winds over the South Atlantic Ocean become weak as they approach the coast of West Africa, they produce excessive rains over Accra. Conversely, when strong northerly winds affect the coast of West Africa without becoming weak, they reduce rainfall amounts over Accra. It has also been noted that if the southerly winds are too strong over southern Ghana, most of the moisture tends to be transported further inland, leaving the coast with little amounts of rain. The highest monthly total rainfall amounts for both periods were recorded in June (Table 1) with southerly wind speeds of 0.6 m/s and 0.9 m/s during 1980–1999 and 2000–2019 respectively (Table 3). This implies that for the post-2000 era when meridional winds are southerly along the coast of Ghana with speeds around 0.9 m/s they are capable of producing intensive rainfall.

Because Accra is located along the coast, when southerly winds are very strong, they tend to drive moisture further away into the inland, depriving the coast from heavy rains. When southerly winds are weak as they approach the coast of West Africa, they tend to dampen moisture along the coast and help to form massive clouds that produce heavy rains. Similarly, when northerly winds are very strong, they tend to bring in dry air from the land or drive away the available moisture further into the Gulf of Guinea leaving the coast dry. However, if the northerly winds become weak as they approach the coast, they tend to serve as a blockade to prevent moisture from the ocean from leaving the coast further inland.

5 Conclusion

The frequency, intensity and variability of rainfall are important to the socio-economic livelihood of many across the globe. Extreme rainfall can be devastating, causing loss of lives and damages to properties. In this study, we investigated the recent changes in extreme rainfall over Accra using rain gauge and satellite-derived data. The study covered 1980–2019 and was divided into two epochs: 1980–1999 and 2000–2019. Our results suggest that possible climate change has affected the monthly rainfall distribution for June, such that post-2000 June rainfall totals are higher than the pre-2000. During the pre-2000 era, the most intense rainfalls occurred during July. This shift is important for planning purposes to help mitigate against the associated negative impacts. Even though mean monthly meridional winds at the 850 hPa level were southerly for both the post-2000 and the pre-2000 for June, the values were higher during the

post-2000 than those of the pre-2000 (Table 3). The findings show that both southerly winds and northerly winds at the 850 hPa level have forecast potential and can be used in addition to other parameters to predict extreme rainfall events over the study area. It is therefore recommended that further studies be done on how meridional winds at the surface influence and affect the weather conditions along the coastal sector of Ghana.

Acknowledgements The authors acknowledge and express their sincere gratitude to the Global Monitoring for Environment and Security (GMES) and Africa programme for providing funding support for this study under the “Marine and Coastal Areas Management in Western Africa” theme through the University of Ghana Regional Marine Centre. The various sources of freely available data used are also duly acknowledged, i.e., the Ghana Meteorological Agency and the National Oceanic and Atmospheric Administration (NOAA) National Centers for Environmental Prediction (NCEP).

Funding Financial support was provided by the Global Monitoring for Environment and Security (GMES) and Africa programme under the “Marine and Coastal Areas Management in Western Africa” theme implemented by the University of Ghana (Grant number: HRST/ST/G&A/CALL1/2017).

Data availability Rainfall data were collected from the Kotoka International Meteorological Office (KIAMO) in Accra by the Ghana Meteorological Agency, <https://www.meteo.gov.gh/gmet/>. Meridional winds at 850 hPa pressure level and surface temperature data were obtained from the National Oceanic and Atmospheric Administration (NOAA) NCEP Reanalysis Derived data provided by the NOAA/OAR/ESRL PSL, Boulder, Colorado, USA, from their Web site at <https://psl.noaa.gov/data/timeseries/>

Code Availability None.

Declarations

Conflict of Interest The authors declare no competing interests.

References

1. Abbam T, Johnson FA, Dash J, Padmadas SS (2018) Spatiotemporal variations in rainfall and temperature in Ghana over the twentieth century, 1900–2014. *Earth and Space Science* 5(4):120–132. <https://doi.org/10.1002/2017EA000327>
2. Adejwon JO, Odekunle TO (2006) Variability and the severity of the “Little Dry Season” in southwestern Nigeria. *J Clim* 19(3):483–493. <https://doi.org/10.1175/JCLI3642.1>
3. Appeaning-Addo K (2013) Assessing coastal vulnerability index to climate change: the case of Accra-Ghana. *J Coastal Res* 165:1892–1897. <https://doi.org/10.2112/si65-320.1>
4. Asante FA, Amuakwa-Mensah F (2015) Climate change and variability in Ghana: stocktaking. *Climate* 3(1):78–99. <https://doi.org/10.3390/cli3010078>
5. Boateng I, Wiafe G, Jayson-Quashigah P-N (2017) Mapping vulnerability and risk of Ghana’s coastline to sea level rise. *Mar Geodesy* 40(1):23–39. <https://doi.org/10.1080/01490419.2016.1261745>

6. Boos WR, Kuang Z (2010) Dominant control of the South Asian monsoon by orographic insulation versus plateau heating. *Nature* 463(7278):218–222. <https://doi.org/10.1038/nature08707>
7. Breuning-Madsen H, Awadzi TW (2005) Harmattan dust deposition and particle size in Ghana. *CATENA* 63(1):23–38. <https://doi.org/10.1016/j.catena.2005.04.001>
8. Brunner RD (2001) Science and the climate change regime. *Policy Sci* 34(1):1–33. <https://doi.org/10.1023/A:1010393101905>
9. Cho Y-M, Shepherd GG, Shepherd MG, Hocking WK, Mitchell NJ, Won Y-I (2011) A study of temperature and meridional wind relationships at high northern latitudes. *J Atmos Solar Terr Phys* 73:936–943. <https://doi.org/10.1016/j.jastp.2010.08.011>
10. Codjoe SNA, Owusu G, Burkett V (2014) Perception, experience, and indigenous knowledge of climate change and variability: the case of Accra, a sub-Saharan African city. *Reg Environ Change* 14(1):369–383. <https://doi.org/10.1007/s10113-013-0500-0>
11. Danquah, I. O. (2013). Climate change and its impacts on flooding in Accra-Greater Accra Metropolitan Assembly [Kwame Nkrumah University of Science and Technology]. Retrieved June 26, 2021, from <http://ir.knuist.edu.gh:8080/handle/123456789/5415>
12. de Boer AM, Toggweiler JR, Sigman DM (2008) Atlantic dominance of the meridional overturning circulation. *J Phys Oceanogr* 38(2):435–450. <https://doi.org/10.1175/2007JPO3731.1>
13. Della-Marta, P. M., Haylock, M. R., Luterbacher, J., & Wanner, H. (2007). Doubled length of western European summer heat waves since 1880. *J Geophys Res Atmos* 112(15). <https://doi.org/10.1029/2007JD008510>
14. Douglas I (2009) Climate change, flooding and food security in south Asia. *Food Security* 1(2):127–136. <https://doi.org/10.1007/s12571-009-0015-1>
15. Emmanuel, K. (2018). Ghana remembers June 3 nightmare. *Pulse Ghana*. Retrieved June 26, 2021, from <https://www.pulse.com.gh/ece-frontpage/june-3-disaster-ghana-remembers-june-3-nightmare/15523bs>
16. Few R (2003) Flooding, vulnerability and coping strategies: local responses to a global threat. *Prog Dev Stud* 3(1):43–58. <https://doi.org/10.1191/1464993403ps049ra>
17. Foli BAK, Appeaning Addo K, Ansong JK, Wiafe G (2021) Evaluation of ECMWF and NCEP reanalysis wind fields for long-term historical analysis and ocean wave modelling in West Africa. *Remote Sensing in Earth Syst Sci*. <https://doi.org/10.1007/s41976-021-00052-3>
18. Gurubaran S, Sridharan R, Raghavarao R (1995) Effects of neutral temperature on meridional winds estimated from ionospheric data. *J Atmos Terr Phys* 57(10):1095–1101. [https://doi.org/10.1016/0021-9169\(94\)00125-8](https://doi.org/10.1016/0021-9169(94)00125-8)
19. Hallegatte S, Green C, Nicholls RJ, Corfee-Morlot J (2013) Future flood losses in major coastal cities. *Nat Clim Chang* 3(9):802–806. <https://doi.org/10.1038/nclimate1979>
20. Iroaganachi N, Ufere JK (2013) Flooding in Nigeria and sustainable land development: case of Delta State. *J Environ Earth Sci* 3(5):38–42
21. Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Leetmaa, A., ... Joseph, D. (1996). The NCEP/NCAR 40-year reanalysis project. *Bulletin of the American Meteorological Society*, 77(3), 437–472
22. Kates, R. W., Colten, C. E., Laska, S., & Leatherman, S. P. (2006). Reconstruction of New Orleans after Hurricane Katrina: a research perspective. 10.1073/pnas.0605726103
23. Knapp KR (2008) Scientific data stewardship of international satellite cloud climatology project B1 global geostationary observations. *J Appl Remote Sens* 2(1):023548. <https://doi.org/10.1117/1.3043461>
24. Krishnamurthy CKB, Lall U, Kwon HH (2009) Changing frequency and intensity of rainfall extremes over India from 1951 to 2003. *J Clim* 22(18):4737–4746. <https://doi.org/10.1175/2009JCLI2896.1>
25. Lyngsie G, Awadzi T, Breuning-Madsen H (2011) Origin of Harmattan dust settled in Northern Ghana-long transported or local dust? *Geoderma* 167–168:351–359. <https://doi.org/10.1016/j.geoderma.2011.07.026>
26. Mahjabin, T., & Abdul-Aziz, O. I. (2020). Trends in the magnitude and frequency of extreme rainfall regimes in Florida. *Water (Switzerland)*, 12(9). <https://doi.org/10.3390/W12092582>
27. Martel JL, Mailhot A, Brissette F, Caya D (2018) Role of natural climate variability in the detection of anthropogenic climate change signal for mean and extreme precipitation at local and regional scales. *J Clim* 31(11):4241–4263. <https://doi.org/10.1175/JCLI-D-17-0282.1>
28. Nicholson SE, Grist JP (2003) The seasonal evolution of the atmospheric circulation over West Africa and equatorial Africa. *J Clim* 16(7):1013–1030. [https://doi.org/10.1175/1520-0442\(2003\)016%3c1013:TSEOTA%3e2.0.CO;2](https://doi.org/10.1175/1520-0442(2003)016%3c1013:TSEOTA%3e2.0.CO;2)
29. Ogega OM, Koske J, Kung'u, J. B., Scoccimarro, E., Endris, H. S., & Mistry, M. N. (2020) Heavy precipitation events over East Africa in a changing climate: results from CORDEX RCMS. *Clim Dyn* 55(3–4):993–1009. <https://doi.org/10.1007/s00382-020-05309-z>
30. Osei-Tutu, E. (2020). Accra floods; 63 years of inertness! ModernGhana. Retrieved September 29, 2021, from <https://www.modernghana.com/news/1035276/accra-floods-63-years-of-inertness.html>
31. Owusu, K., & Obour, P. B. (2021). Urban Flooding, Adaptation Strategies, and Resilience: Case Study of Accra, Ghana. In W. Leal Filho, N. Ogue, D. Ayal, L. Adeleke, & I. da Silva (Eds.), *African handbook of climate change adaptation* (pp. 2387–2403). Springer, Cham. https://doi.org/10.1007/978-3-030-45106-6_249
32. Owusu K, Waylen P (2009) Trends in spatio-temporal variability in annual rainfall in Ghana (1951–2000). *Weather* 64(5):115–120. <https://doi.org/10.1002/wea.255>
33. Owusu K, Waylen P, Qiu Y (2008) Changing rainfall inputs in the Volta basin: implications for water sharing in Ghana. *GeoJournal* 71(4):201–210. <https://doi.org/10.1007/s10708-008-9156-6>
34. Owusu K, Waylen PR (2013) The changing rainy season climatology of mid-Ghana. *Theoret Appl Climatol* 112(3–4):419–430. <https://doi.org/10.1007/s00704-012-0736-5>
35. Padi M (2017) Citation: Padi M (2017) Weather transition periods in Ghana. *J Climatol Weather Forecasting* 5(2):211. <https://doi.org/10.4172/2332-2594.1000211>
36. Pereira SC, Marta-Almeida M, Carvalho AC, Rocha A (2020) Extreme precipitation events under climate change in the Iberian Peninsula. *Int J Climatol* 40(2):1255–1278. <https://doi.org/10.1002/joc.6269>
37. Pineda LE, Willems P (2018) Rainfall extremes, weather and climate drivers in complex terrain: A data-driven approach based on signal enhancement methods and EV modeling. *J Hydrol* 563(February):283–302. <https://doi.org/10.1016/j.jhydrol.2018.05.062>
38. Pinson P (2012) Adaptive calibration of (u, v)-wind ensemble forecasts. *Q J R Meteorol Soc* 138(666):1273–1284. <https://doi.org/10.1002/qj.1873>
39. Raj J, Hamza, ·, Bangalath, K., & Stenichikov, G. (2019) West African Monsoon: current state and future projections in a high-resolution AGCM. *Clim Dyn* 52:6441–6461. <https://doi.org/10.1007/s00382-018-4522-7>
40. Seleshi Y, Camberlin P (2006) Recent changes in dry spell and extreme rainfall events in Ethiopia. *Theoret Appl Climatol* 83(1–4):181–191. <https://doi.org/10.1007/s00704-005-0134-3>

41. Singleton AT, Reason CJC (2007) Variability in the characteristics of cut-off low pressure systems over subtropical southern Africa. *Int J Climatol* 27(3):295–310. <https://doi.org/10.1002/joc.1399>
42. Speich, S., Blanke, B., & Cai, W. (2007). Atlantic meridional overturning circulation and the Southern Hemisphere supergyre. *Geophysical Research Letters*, 34(23). <https://doi.org/10.1029/2007GL031583>
43. Speich, S., Blanke, B., & Cai, W. (2007). Atlantic meridional overturning circulation and the Southern Hemisphere supergyre. *Geophysical Research Letters*, 34(23)<https://doi.org/10.1155/2016/1940456>
44. Tabiri, M. O. (2015). Perennial problem of floods in Accra: what is the way forward? *International Journal of Advanced Research in Education & Technology (IJARET)*, 2(3), 198–201. www.ijaret.com
45. Tan, C. M., & Rockmore, M. (2019). Famine in Ghana and its impact. In V. R. Preedy & V. B. Patel (Eds.), *Handbook of famine, starvation, and nutrient deprivation: from biology to policy* (pp. 31–45). Springer, Cham. https://doi.org/10.1007/978-3-319-55387-0_95
46. Tettey, M., Oduro, F. T., Adedia, D., & Abaye, D. A. (2017). Markov chain analysis of the rainfall patterns of five geographical locations in the south eastern coast of Ghana. *Earth Perspectives*, 4(1). <https://doi.org/10.1186/s40322-017-0042-6>
47. Toledano C, Cachorro VE, De Frutos AM, Torres B, Berjón A, Sorribas M, Stone RS (2009) Airmass classification and analysis of aerosol types at El Arenosillo (Spain). *J Appl Meteorol Climatol* 48(5):962–981. <https://doi.org/10.1175/2008JAMC2006.1>
48. UN DHA. (1995). Ghana-Floods Situation Report No. 1. Reliefweb. Retrieved June 26, 2021, from <https://reliefweb.int/report/ghana/ghana-floods-situation-report-no-1>
49. World Bank. (2017). Enhancing urban resilience in the Greater Accra Metropolitan Area. In *Global Practice on Social, Urban, Rural and Resilience* (Issue May). <https://doi.org/10.1596/27516>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.